Map of *The Pentagon, Machine-Checked: Abel's five-term relation for the dilogarithm in Lean 4*

Special functions · the dilogarithm

1 results | ✅ **1** formalized in Lean | ◐ **0** with a part in Lean | **0** checked by a program | written proof only | (a result can carry several tags)

Dependencies in the written proof: an arrow A → B means that this proof of B uses A. Every result rests on its written proof; the tags record what was checked besides, and each statement's "Evidence & scope" block says how far. The colour of a box is its strongest tag, in the order above; the tags in the box give the rest. Boxes with a thick border are the main results; click a box to read the statement.

Q2: can the generator be reached by the same three-move recipe?

Theorem 1
Abel's five-term relation for the Rogers L-function on $(0,1)^2$
✅ Lean

> \begin{itemize} \item A sorry-free Lean~4 proof of Abel's five-term relation for the Rogers L-function, $L(x)+L(y)=L(xy)+L\big(\tfrac{x(1-y)}{1-xy}\big)+L\big(\tfrac{y(1-x)}{1-xy}\big)$ for $x,y\in(0,1)$ (Theorem~1). \item The route is the classical one: the five-term difference has zero derivative in x (nine logarithms collapse into a basis of five), so it is constant, and its value is read off as $x\to0^+$, where $\log t\cdot\log(1-t)\to0$ by a squeeze. \item Three reusable blocks on the dilogarithm of the companion paper: the derivative of L on $(0,1)$, $L(t)\to0$ as $t\to0^+$, and continuity of L on $(0,1)$. \item The mathematics is classical (Spence, Abel, Rogers); the paper claims the formalization, after a documented search of proof-assistant libraries. \end{itemize}

🚀 **Start here: Theorem 1**: Abel's five-term relation for the Rogers L-function on $(0,1)^2$.

Main results:

- **Theorem 1**: Abel's five-term relation for the Rogers L-function on $(0,1)^2$

> ## 🧭 Fields: where the paper lives and what it connects
>
> **Tools taken from other branches**
>
> - **Real analysis** → vanishing derivative on (0,1) gives a constant (mean-value theorem); the boundary limit $\log t \cdot \log(1-t) \to 0$ by the squeeze $2(-t \log t)$ — Theorem 1
> - **Formal verification** → Lean 4 + Mathlib: the nine-into-five logarithm cancellation, closed by field_simp and ring — Theorem 1
>
> **What the results serve**
>
> - **Functional equations of the dilogarithm** ← by Wojtkowiak and de Jeu, every rational functional equation of Li_2 follows from this one (cited, not formalized) — Theorem 1
> - **Bloch group, cluster algebras, quantum dilogarithm** ← the defining relation of the Bloch group and the pentagon of cluster mutations (context; not formalized here)