

Manual for Package: mathematics

Revision 19M

Karl Kästner

June 12, 2022

Contents

1	calendar	1
1.1	days_per_month	1
1.2	isnight	1
2	mathematics	1
2.1	cast_byte_to_integer	1
3	complex-analysis	1
3.1	complex_exp_product_im_im	1
3.2	complex_exp_product_im_re	2
3.3	complex_exp_product_re_im	2
3.4	complex_exp_product_re_re	2
3.5	croots	3
3.6	root_complex	3
3.7	test_imroots	3
4	derivation	3
4.1	derive_acfar1	3
4.2	derive_ar1_spectral_density	4
4.3	derive_ar2param	4
4.4	derive_arc_length	4
4.5	derive_fourier_power	4
4.6	derive_fourier_power_exp	4
4.7	derive_laplacian_curvilinear	4
4.8	derive_laplacian_fourier_piecewise_linear	4
4.9	derive_logtripdf	4
4.10	derive_smooth1d_parametric	4
4.11	derive_spectral_density_bandpass_initial_condition	4
5	derivation/master	5

5.1	derive_bc_one_sided	5
5.2	derive_convergence	5
5.3	derive_error_fdm	5
5.4	derive_fdm_poly	5
5.5	derive_fdm_power	5
5.6	derive_fdm_taylor	5
5.7	derive_fdm_vargrid	5
5.8	derive_fem_2d_mass	5
5.9	derive_fem_error_2d	5
5.10	derive_fem_error_3d	6
5.11	derive_fem_sym_2d	6
5.12	derive_grid_constants	6
5.13	derive_interpolation	6
5.14	derive_laplacian	6
5.15	derive_limit	6
5.16	derive_nc_1d	6
5.17	derive_nc_1d_	6
5.18	derive_nc_2d	6
5.19	derive_nonuniform_symmetric	6
5.20	derive_richardson	7
5.21	derive_sum	7
5.22	nn	7
5.23	test_derive	7
5.24	test_derive_fdm_poly	7
5.25	test_filter	7
5.26	test_vargrid	7
6	derivation	7
6.1	simplify_atan	7
7	mathematics	8
7.1	entropy	8
8	finance	8
8.1	derive_skewrnd_walsh_paramter	8
8.2	gbm_cdf	8
8.3	gbm_fit	8
8.4	gbm_fit_old	8
8.5	gbm_inv	8
8.6	gbm_mean	8
8.7	gbm_median	8
8.8	gbm_pdf	9
8.9	gbm_simulate	9
8.10	gbm_skewness	9

8.11	gbm_std	9
8.12	gbm_transform_time_step	9
8.13	put_price_black_scholes	9
8.14	skewgbm_simulate	9
8.15	skewrnd_walsh	9
9	finance/test	9
9.1	test_gbm	9
9.2	test_gbm_pdf	10
9.3	test_skewrnd_walsh	10
10	fourier/@STFT	10
10.1	STFT	10
10.2	itransform	10
10.3	stft_	10
10.4	stftmat	10
10.5	transform	10
11	fourier	11
11.1	amplitude_from_peak	11
11.2	dftmtx_man	11
11.3	example_fourier_window	11
11.4	fft_man	11
11.5	fftsmooth	12
11.6	fix_fourier	12
11.7	fourier_axis	12
11.8	fourier_axis_2d	13
11.9	fourier_cesaro_correction	13
11.10	fourier_coefficient_piecewise_linear	13
11.11	fourier_coefficient_piecewise_linear_1	13
11.12	fourier_coefficient_ramp3	14
11.13	fourier_coefficient_ramp_pulse	14
11.14	fourier_coefficient_ramp_step	14
11.15	fourier_coefficient_square_pulse	14
11.16	fourier_cubic_interaction_coefficients	14
11.17	fourier_derivative	14
11.18	fourier_derivative_matrix_1d	15
11.19	fourier_derivative_matrix_2d	15
11.20	fourier_expand	15
11.21	fourier_fit	15
11.22	fourier_interpolate	15
11.23	fourier_matrix	15
11.24	fourier_matrix2	15
11.25	fourier_matrix3	16

11.26	fourier_matrix_exp	16
11.27	fourier_multiplicative_interaction_coefficients	16
11.28	fourier_power	16
11.29	fourier_power_exp	16
11.30	fourier_predict	17
11.31	fourier_quadratic_interaction_coefficients	17
11.32	fourier_random_phase_walk	17
11.33	fourier_range	17
11.34	fourier_regress	17
11.35	fourier_resampled_fit	17
11.36	fourier_resampled_predict	17
11.37	fourier_signed_square	18
11.38	fourier_transform	18
11.39	fourier_transform_fractional	18
11.40	hyperbolic_fourier_box	18
11.41	idftmtx_man	18
11.42	laplace_2d_pwlinear	19
11.43	mean_fourier_power	19
11.44	moments_fourier_power	19
11.45	nanfft	19
11.46	peaks	19
11.47	roots_fourier	20
11.48	spectral_density	20
11.49	std_fourier_power	20
11.50	test_complex_exp_product	20
11.51	test_fourier_filter	20
11.52	test_idftmtx	20
11.53	var_fourier_power	20
12 mathematics		21
12.1	gaussfit_quantile	21
13 geometry/@Geometry		21
13.1	Geometry	21
13.2	arclength	21
13.3	arclength_old	21
13.4	arclength_old2	21
13.5	base_point	21
13.6	base_point_limited	22
13.7	centroid	22
13.8	cosa_min_max	22
13.9	cross2	22
13.10	curvature	22
13.11	ddot	22

13.12	distance	22
13.13	distance2	22
13.14	dot	23
13.15	edge_length	23
13.16	enclosed_angle	23
13.17	enclosing_triangle	23
13.18	hexagon	23
13.19	inPolygon	23
13.20	inTetra	23
13.21	inTetra2	23
13.22	inTriangle	24
13.23	intersect	24
13.24	lineintersect	24
13.25	lineintersect1	24
13.26	minimum_distance_lines	24
13.27	mittenpunkt	24
13.28	nagelpoint	24
13.29	onLine	24
13.30	orthocentre	25
13.31	plumb_line	25
13.32	poly_area	25
13.33	poly_edges	25
13.34	poly_set	25
13.35	poly_width	25
13.36	polyxpoly	25
13.37	project_to_curve	26
13.38	quad_isconvex	26
13.39	random_disk	26
13.40	random_simplex	26
13.41	sphere_volume	26
13.42	tetra_volume	26
13.43	tobarycentric	26
13.44	tobarycentric1	26
13.45	tobarycentric2	27
13.46	tobarycentric3	27
13.47	tri_angle	27
13.48	tri_area	27
13.49	tri_centroid	27
13.50	tri_distance_opposit_midpoint	27
13.51	tri_edge_length	27
13.52	tri_edge_midpoint	27
13.53	tri_excircle	28
13.54	tri_height	28
13.55	tri_incircle	28

13.56	tri_isacute	28
13.57	tri_isobtuse	28
13.58	tri_semiperimeter	28
13.59	tri_side_length	28
14	geometry	29
14.1	Polygon	29
14.2	bounding_box	29
14.3	curvature_1d	29
14.4	cvt	29
14.5	deg_to_frac	29
14.6	ellipse	30
14.7	ellipseX	30
14.8	ellipseY	30
14.9	first_intersect	30
14.10	golden_ratio	30
14.11	hypot3	30
14.12	meanangle	30
14.13	meanangle2	30
14.14	meanangle3	31
14.15	meanangle4	31
14.16	medianangle	31
14.17	medianangle2	31
14.18	pilim	31
14.19	streamline_radius_of_curvature	31
15	histogram/@Histogram	32
15.1	2x	32
15.2	Histogram	32
15.3	bimodes	32
15.4	cdf	32
15.5	cdfS	32
15.6	chi2test	32
15.7	cmoment	32
15.8	cmomentS	32
15.9	entropy	32
15.10	entropyS	33
15.11	export_csv	33
15.12	iquantile	33
15.13	kstest	33
15.14	kurtosis	33
15.15	kurtosisS	33
15.16	mean	33
15.17	meanS	33

15.18	median	33
15.19	medianS	33
15.20	mode	34
15.21	modeS	34
15.22	moment	34
15.23	momentS	34
15.24	pdf	34
15.25	quantile	34
15.26	quantileS	34
15.27	resample	34
15.28	setup	34
15.29	skewness	34
15.30	skewnessS	35
15.31	stairs	35
15.32	stairsS	35
15.33	std	35
15.34	stdS	35
15.35	var	35
15.36	varS	35
16	histogram	35
16.1	hist_man	35
16.2	histadapt	35
16.3	histconst	36
16.4	pdf_poly	36
16.5	plotcdf	36
16.6	test_histogram	36
17	mathematics	36
17.1	imrotmat	36
18	linear-algebra	36
18.1	averaging_matrix_2	36
18.2	colnorm	36
18.3	condest_	36
18.4	connectivity_matrix	37
19	linear-algebra/coordinate-transformation	37
19.1	barycentric2cartesian	37
19.2	barycentric2cartesian3	37
19.3	cartesian2barycentric	37
19.4	cartesian_to_unit_triangle_basis	37
19.5	ellipsoid2geoid	37
19.6	example_approximate_utm_conversion	37

19.7	latlon2utm	37
19.8	latlon2utm_simple	38
19.9	lowrance_mercator_to_wgs84	38
19.10	nmea2utm	38
19.11	sn2xy	38
19.12	unit_triangle_to_cartesian	38
19.13	utm2latlon	38
19.14	xy2nt	38
19.15	xy2sn	39
19.16	xy2sn_java	39
19.17	xy2sn_old	39
20	linear-algebra	39
20.1	det2x2	39
20.2	det3x3	39
20.3	det4x4	39
20.4	diag2x2	39
20.5	down	40
20.6	eig2x2	40
21	linear-algebra/eigenvalue	40
21.1	eig_bisection	40
21.2	eig_inverse	40
21.3	eig_inverse_iteration	40
21.4	eig_power_iteration	40
22	linear-algebra/eigenvalue/jacobi-davidson	40
22.1	afun_jdm	40
22.2	davidson	40
22.3	jacobi_davidson	40
22.4	jacobi_davidson_qr	41
22.5	jacobi_davidson_qz	41
22.6	jacobi_davidson_simple	41
22.7	jdqr	41
22.8	jdqr_sleijpen	44
22.9	jdqr_vorst	48
22.10	jdqz	51
22.11	mfunc_jdm	55
22.12	mgs	55
22.13	minres_	55
22.14	mv_jacobi_davidson	55
23	linear-algebra	56
23.1	first	56

23.2	gershgorin_circle	56
23.3	hausdorff	56
23.4	ieig2x2	56
23.5	inv2x2	56
23.6	inv3x3	56
23.7	inv4x4	56
23.8	kernel2matrix	57
24	linear-algebra/lanczos	57
24.1	arnoldi	57
24.2	arnoldi_new	57
24.3	eigs_lanczos_man	57
24.4	lanczos	57
24.5	lanczos_	57
24.6	lanczos_biorthogonal	57
24.7	lanczos_biorthogonal_improved	57
24.8	lanczos_ghep	57
24.9	mv_lanczos	58
24.10	reorthogonalise	58
24.11	test_lanczos	58
25	linear-algebra	58
25.1	left	58
26	linear-algebra/linear-systems	58
26.1	gmres_man	58
26.2	minres_recycle	58
27	linear-algebra	58
27.1	lpmean	58
27.2	lpnorm	58
27.3	matvec3	59
27.4	max2d	59
27.5	mid	59
27.6	mpoweri	59
27.7	mtimes2x2	59
27.8	mtimes3x3	59
27.9	nannorm	59
27.10	nanshift	59
27.11	nl	60
27.12	normalise	60
27.13	normalize1	60
27.14	normrows	60
27.15	orth2	60

27.16	orth_man	60
27.17	orthogonalise	60
27.18	padd2	61
27.19	paddext	61
27.20	paddval1	61
27.21	paddval2	61
28	linear-algebra/polynomial	61
28.1	chebychev	61
28.2	piecewise_polynomial	61
28.3	roots1	61
28.4	roots2	62
28.5	roots2poly	62
28.6	roots3	62
28.7	roots4	62
28.8	roots_piecewise_linear	62
28.9	test_roots4	62
28.10	vanderi_1d	62
29	linear-algebra	62
29.1	randrot	62
29.2	right	63
29.3	rot2	63
29.4	rot2dir	63
29.5	rot3	63
29.6	rotR	63
29.7	rownorm	63
29.8	simmilarity_matrix	63
29.9	spnorm	63
29.10	spzeros	63
29.11	test_roots3	64
29.12	transform_minmax	64
29.13	transpose3	64
29.14	transposeall	64
29.15	up	64
29.16	vander_nd	64
29.17	vanderd_2d	64
30	logic	64
30.1	bitor_man	64
31	master/plot	65
31.1	attach_boundary_value	65
31.2	cartesian_polar	65

31.3	img_vargrid	65
31.4	plot_basis_functions	65
31.5	plot_convergence	65
31.6	plot_dof	65
31.7	plot_eigenbar	65
31.8	plot_error_estimation	65
31.9	plot_error_estimation_2	65
31.10	plot_error_fem	66
31.11	plot_fdm_kernel	66
31.12	plot_fdm_vs_fem	66
31.13	plot_fem_accuracy	66
31.14	plot_function_and_grid	66
31.15	plot_hat	66
31.16	plot_hydrogen_wf	66
31.17	plot_mesh	66
31.18	plot_mesh_2	66
31.19	plot_refine	66
31.20	plot_refine_3d	67
31.21	plot_runtime	67
31.22	plot_spectrum	67
31.23	plot_wavefunction	67
32	master/ported	67
32.1	assemble_2d_phi_phi	67
32.2	assemble_3d_dphi_dphi	67
32.3	assemble_3d_phi_phi	67
32.4	dV_2d_	67
32.5	derivative_2d	67
32.6	derivative_3d	68
32.7	element_neighbour_2d	68
32.8	prefetch_2d_	68
32.9	promote_2d_3_10	68
32.10	promote_2d_3_15	68
32.11	promote_2d_3_21	68
32.12	promote_2d_3_6	68
32.13	promote_3d_4_10	68
32.14	promote_3d_4_20	68
32.15	promote_3d_4_35	68
32.16	vander_2d	69
32.17	vander_3d	69
33	number-theory	69
33.1	ceiln	69
33.2	digitsb	69

33.3	floorn	69
33.4	iseven	69
33.5	multichoosek	69
33.6	nchoosek_man	70
33.7	pythagorean_triple	70
33.8	roundn	70
34	numerical-methods	70
34.1	advect_analytic	70
35	numerical-methods/differentiation	70
35.1	derivative1	70
35.2	derivative2	70
36	numerical-methods	70
36.1	diffuse_analytic	70
37	numerical-methods/finite-difference	71
37.1	cdiff	71
37.2	cdiffb	71
37.3	central_difference	71
37.4	cmean	71
37.5	cmean2	71
37.6	derivative_matrix_1_1d	71
37.7	derivative_matrix_2_1d	71
37.8	derivative_matrix_2d	72
37.9	derivative_matrix_curvilinear	72
37.10	derivative_matrix_curvilinear_2	72
37.11	difference_kernel	72
37.12	diffusion_matrix_2d_anisotropic	72
37.13	diffusion_matrix_2d_anisotropic2	72
37.14	distmat	72
37.15	downwind_difference	72
37.16	gradpde2d	73
37.17	laplacian	73
37.18	laplacian_fdm	73
37.19	lrmean	73
38	numerical-methods/finite-difference/master	73
38.1	fdm_adaptive_grid	73
38.2	fdm_adaptive_refinement_old	73
38.3	fdm_assemble_d1_2d	73
38.4	fdm_assemble_d2_2d	73
38.5	fdm_confinement	74

38.6	fdm_d_vargrid	74
38.7	fdm_h_unstructured	74
38.8	fdm_hydrogen_vargrid	74
38.9	fdm_mark_unstructured_2d	74
38.10	fdm_plot	74
38.11	fdm_plot_series	74
38.12	fdm_refine_2d	74
38.13	fdm_refine_3d	74
38.14	fdm_refine_unstructured_2d	74
38.15	fdm_schroedinger_2d	75
38.16	fdm_schroedinger_3d	75
38.17	relocate	75
39 numerical-methods/finite-difference		75
39.1	mid	75
39.2	pwmid	75
39.3	ratio	75
39.4	steplength	75
39.5	swapoddeven	75
39.6	test_derivative_matrix_2d	76
39.7	test_derivative_matrix_curvilinear	76
39.8	test_difference_kernel	76
39.9	upwind_difference	76
40 numerical-methods/finite-element		76
40.1	Mesh_2d.java	76
40.2	Tree_2d.java	76
40.3	assemble_1d_dphi_dphi	76
40.4	assemble_1d_phi_phi	76
40.5	assemble_2d_dphi_dphi.java	76
40.6	assemble_2d_phi_phi.java	77
40.7	assemble_3d_dphi_dphi.java	77
40.8	assemble_3d_phi_phi.java	77
40.9	boundary_1d	77
40.10	boundary_2d	77
40.11	boundary_3d	77
40.12	check_area_2d	77
40.13	circmesh	77
40.14	cropradius	77
40.15	display_2d	77
40.16	display_3d	78
40.17	distort	78
40.18	err_2d	78
40.19	estimate_err_2d_3	78

40.20	example_1d	78
40.21	example_2d	78
40.22	explode	78
40.23	fem_2d	78
40.24	fem_2d_heuristic_mesh	78
40.25	fem_get_2d_radial	78
40.26	fem_interpolation	79
40.27	fem_plot_1d	79
40.28	fem_plot_1d_series	79
40.29	fem_plot_2d	79
40.30	fem_plot_2d_series	79
40.31	fem_plot_3d	79
40.32	fem_plot_3d_series	79
40.33	fem_plot_confine_series	79
40.34	fem_radial	79
40.35	flip_2d	80
40.36	get_mesh_arrays	80
40.37	hashkey	80
41 numerical-methods/finite-element/int		80
41.1	int_1d_gauss	80
41.2	int_1d_gauss_1	80
41.3	int_1d_gauss_2	80
41.4	int_1d_gauss_3	80
41.5	int_1d_gauss_4	80
41.6	int_1d_gauss_5	80
41.7	int_1d_gauss_6	81
41.8	int_1d_gauss_lobatto	81
41.9	int_1d_gauss_n	81
41.10	int_1d_nc_2	81
41.11	int_1d_nc_3	81
41.12	int_1d_nc_4	81
41.13	int_1d_nc_5	81
41.14	int_1d_nc_6	81
41.15	int_1d_nc_7	81
41.16	int_1d_nc_7_hardy	81
41.17	int_2d_gauss_1	82
41.18	int_2d_gauss_12	82
41.19	int_2d_gauss_13	82
41.20	int_2d_gauss_16	82
41.21	int_2d_gauss_19	82
41.22	int_2d_gauss_25	82
41.23	int_2d_gauss_3	82
41.24	int_2d_gauss_33	82

41.25	int_2d_gauss_4	82
41.26	int_2d_gauss_6	82
41.27	int_2d_gauss_7	83
41.28	int_2d_gauss_9	83
41.29	int_2d_nc_10	83
41.30	int_2d_nc_15	83
41.31	int_2d_nc_21	83
41.32	int_2d_nc_3	83
41.33	int_2d_nc_6	83
41.34	int_3d_gauss_1	83
41.35	int_3d_gauss_11	83
41.36	int_3d_gauss_14	83
41.37	int_3d_gauss_15	84
41.38	int_3d_gauss_24	84
41.39	int_3d_gauss_4	84
41.40	int_3d_gauss_45	84
41.41	int_3d_gauss_5	84
41.42	int_3d_nc_11	84
41.43	int_3d_nc_4	84
41.44	int_3d_nc_6	84
41.45	int_3d_nc_8	84
42 numerical-methods/finite-element		85
42.1	interpolation_matrix	85
42.2	mark	85
42.3	mark_1d	85
42.4	mesh_1d_uniform	85
42.5	mesh_3d_uniform	85
42.6	mesh_interpolate	85
42.7	neighbour_1d	85
42.8	old	85
42.9	pdeeig_1d	85
42.10	pdeeig_2d	86
42.11	pdeeig_3d	86
42.12	polynomial_derivative_1d	86
42.13	potential_const	86
42.14	potential_coulomb	86
42.15	potential_harmonic_oscillator	86
42.16	project_circle	86
42.17	project_rectangle	86
42.18	promote_1d_2_3	86
42.19	promote_1d_2_4	86
42.20	promote_1d_2_5	87
42.21	promote_1d_2_6	87

42.22	quadrilateate	87
42.23	recalculate_regularity_2d	87
42.24	refine_1d	87
42.25	refine_2d_21	87
42.26	refine_2d_structural	87
42.27	regularity_1d	87
42.28	regularity_2d	87
42.29	regularity_3d	88
42.30	relocate_2d	88
42.31	test_circmesh	88
42.32	test_hermite	88
42.33	tri_assign_points	88
42.34	triangulation_uniform	88
42.35	vander_1d	88
42.36	vanderd_1d	88
42.37	vanderi_1d	88
43	numerical-methods/finite-volume/@Advection	89
43.1	Advection	89
43.2	dot_advection	89
44	numerical-methods/finite-volume/@Burgers	89
44.1	burgers_split	89
44.2	dot_burgers_fdm	89
44.3	dot_burgers_fft	89
45	numerical-methods/finite-volume/@Finite_Volume	89
45.1	Finite_Volume	89
45.2	apply_bc	90
45.3	solve	90
45.4	step_split_strang	90
45.5	step_unsplit	90
46	numerical-methods/finite-volume/@Flux_Limiter	90
46.1	Flux_Limiter	90
46.2	beam_warming	90
46.3	fromm	91
46.4	lax_wendroff	91
46.5	minmod	91
46.6	monotized_central	91
46.7	muscl	91
46.8	superbee	91
46.9	upwind	91
46.10	vanLeer	92

47 numerical-methods/finite-volume/@KDV	92
47.1 dot_kdv_fdm	92
47.2 dot_kdv_fft	92
47.3 kdv_split	92
48 numerical-methods/finite-volume/@Reconstruct_Average_Evolve	92
48.1 Reconstruct_Average_Evolve	92
48.2 advect_highres	93
48.3 advect_lowress	93
49 numerical-methods/finite-volume	93
49.1 Godunov	93
49.2 Lax_Friedrich	93
49.3 Measure	93
49.4 Roe	93
49.5 fv_swe	94
49.6 staggered_euler	94
49.7 staggered_grid	94
50 numerical-methods	94
50.1 grid2quad	94
51 numerical-methods/integration	94
51.1 cumintL	94
51.2 cumintR	94
51.3 int_trapezoidal	94
52 numerical-methods/interpolation/@Kriging	95
52.1 Kriging	95
52.2 estimate_semivariance	95
52.3 interpolate_	95
53 numerical-methods/interpolation/@RegularizedInterpolator1	95
53.1 RegularizedInterpolator1	95
53.2 init	95
54 numerical-methods/interpolation/@RegularizedInterpolator2	96
54.1 RegularizedInterpolator2	96
54.2 init	96
55 numerical-methods/interpolation/@RegularizedInterpolator3	96
55.1 RegularizedInterpolator3	96
55.2 init	96
56 numerical-methods/interpolation	96

56.1	IDW	96
56.2	IPoly	96
56.3	IRBM	97
56.4	ISparse	97
56.5	Inn	97
56.6	Interpolator	97
56.7	fixnan	97
56.8	idw1	97
56.9	idw2	97
56.10	inner2outer	98
56.11	inner2outer2	98
56.12	interp1_circular	98
56.13	interp1_limited	98
56.14	interp1_man	98
56.15	interp1_piecewise_linear	98
56.16	interp1_save	98
56.17	interp1_slope	99
56.18	interp1_smooth	99
56.19	interp1_unique	99
56.20	interp2_man	99
56.21	interp_angle	99
56.22	interp_fourier	99
56.23	interp_fourier_batch	99
56.24	interp_sn	100
56.25	interp_sn2	100
56.26	interp_sn3	100
56.27	interp_sn_	100
56.28	limit_by_distance_1d	100
56.29	resample1	100
56.30	resample_d_min	100
56.31	resample_vector	101
56.32	test_interp1_limited	101
57	numerical-methods	101
57.1	inverse_complex	101
57.2	maccormack_step	101
57.3	minmod	101
58	numerical-methods/multigrid	101
58.1	mg_interpolate	101
58.2	mg_restrict	101
59	numerical-methods/ode/@BVPS_Characteristic	101
59.1	BVPS_Characteristic	101

59.2	assemble1_A	102
59.3	assemble1_A_Q	102
59.4	assemble2_A	102
59.5	assemble_AA	102
59.6	assemble_AAA	102
59.7	assemble_Ic	102
59.8	bvp1c	102
59.9	check_arguments	102
59.10	couple_junctions	103
59.11	derivative	103
59.12	init	103
59.13	inner2outer_bvp2c	103
59.14	reconstruct	103
59.15	resample	103
59.16	solve	103
59.17	test_assemble1_A	104
59.18	test_assemble2_A	104
60 numerical-methods/ode/@Time_Stepper		104
60.1	Time_Stepper	104
60.2	solve	104
61 numerical-methods/ode		104
61.1	bvp2fdm	104
61.2	bvp2wavetrain	105
61.3	bvp2wavetwopass	105
61.4	ivp_euler_forward	105
61.5	ivp_euler_forward2	105
61.6	ivprk2	105
61.7	ode2_matrix	105
61.8	ode2characteristic	105
61.9	step_trapezoidal	106
61.10	test_bvp2	106
62 numerical-methods/optimisation		106
62.1	aitken_iteration	106
62.2	anderson_iteration	106
62.3	armijo_stopping_criterion	106
62.4	astar	106
62.5	binsearch	106
62.6	bisection	106
62.7	box1	107
62.8	box2	107
62.9	cauchy	107

62.10	cauchy2	107
62.11	directional_derivative	107
62.12	dud	107
62.13	extreme3	107
62.14	extreme_quadratic	108
62.15	ftest	108
62.16	fzero_bisect	108
62.17	fzero_newton	108
62.18	grad	108
62.19	hessian	108
62.20	hessian_from_gradient	109
62.21	hessian_projected	109
62.22	line_search	109
62.23	line_search2	109
62.24	line_search_polynomial	109
62.25	line_search_polynomial2	110
62.26	line_search_quadratic	110
62.27	line_search_quadratic2	110
62.28	line_search_wolfe	110
62.29	ls_bgfs	110
62.30	ls_broyden	111
62.31	ls_generalized_secant	111
62.32	nlcg	111
62.33	nlls	111
62.34	picard	111
62.35	poly_extrema	112
62.36	quadratic_function	112
62.37	quadratic_programming	112
62.38	quadratic_step	112
62.39	rosenbrock	112
62.40	sqrt_heron	112
62.41	test_directional_derivative	112
62.42	test_dud	112
62.43	test_fzero_newton	112
62.44	test_line_search_quadratic2	113
62.45	test_ls_generalized_secant	113
62.46	test_nlcg_6_order	113
62.47	test_nlls	113
63	numerical-methods/pde	113
63.1	laplacian2d_fundamental_solution	113
64	numerical-methods/piecewise-polynomials	113
64.1	Hermite1	113

64.2	hp2_fit	113
64.3	hp2_predict	114
64.4	hp_predict	114
64.5	hp_regress	114
64.6	lp_count	114
64.7	lp_predict	114
64.8	lp_regress	114
64.9	lp_regress_	114
65	numerical-methods	114
65.1	test_adams_bashforth	114
66	patterns	115
66.1	band_pattern	115
66.2	hexagonal_pattern	115
67	regression/@PolyOLS	115
67.1	PolyOLS	115
67.2	coefftest	115
67.3	detrend	115
67.4	fit	115
67.5	fit_	115
67.6	predict	116
67.7	predict_	116
67.8	slope	116
68	regression/@PowerLS	116
68.1	PowerLS	116
68.2	fit	116
68.3	predict	116
68.4	predict_	116
69	regression/@Theil	117
69.1	Theil	117
69.2	detrend	117
69.3	fit	117
69.4	predict	117
69.5	slope	117
70	regression	117
70.1	Theil_Multivariate	117
70.2	areg	118
70.3	ginireg	118
70.4	hesssimplereg	118
70.5	l1lin	118

70.6	lsq_sparam	118
70.7	polyfitd	118
70.8	regression_method_of_moments	119
70.9	robustlinreg	119
70.10	theil2	119
70.11	theil_generalised	119
70.12	total_least_squares	119
70.13	weighted_median_regression	119
71	set-theory	120
71.1	issubset	120
72	mathematics	120
72.1	shuffle_index	120
73	signal-processing	120
73.1	acfar1	120
73.2	acfar1_2	120
73.3	acfar2	120
73.4	acfar2_2	121
73.5	ar1_cutoff_frequency	121
73.6	ar1_effective_sample_size	121
73.7	ar1_mse_mu_single_sample	121
73.8	ar1_mse_pop	121
73.9	ar1_mse_range	121
73.10	ar1_spectrum	121
73.11	ar1_to_tikhonov	121
73.12	ar1_var_factor	122
73.13	ar1_var_factor_	122
73.14	ar1_var_range2	122
73.15	ar1delay	122
73.16	ar1delay_old	122
73.17	ar2_acf2c	122
73.18	ar2conv	123
73.19	ar2dof	123
73.20	ar2param	123
73.21	asymwin	123
74	signal-processing/autocorr	123
74.1	autocorr2	123
74.2	autocorr_bandpass	123
74.3	autocorr_brownian_phase	123
74.4	autocorr_brownian_phase_2d	123
74.5	autocorr_brownian_phase_across	124

74.6	autocorr_decay_rate	124
74.7	autocorr_effective_sample_size	124
74.8	autocorr_fft	124
74.9	autocorr_genton	124
74.10	autocorr_highpass	124
74.11	autocorr_lowpass	124
74.12	autocorr_periodic_additive_noise	124
74.13	autocorr_periodic_windowed	124
74.14	autocorr_radial	125
75	signal-processing	125
75.1	average_wave_shape	125
75.2	bandpass	125
75.3	bandpass1d	125
75.4	bandpass1d_fft	125
75.5	bandpass1d_implicit	125
75.6	bandpass2	125
75.7	bandpass2d	125
75.8	bandpass2d_2	126
75.9	bandpass2d_fft	126
75.10	bandpass2d_ideal	126
75.11	bandpass2d_implicit	126
75.12	bandpass2d_iso	126
75.13	bandpass_arg	126
75.14	bandpass_f0_to_rho	126
75.15	bandpass_max	126
75.16	bandpass_max2	126
75.17	bartlett	127
75.18	bin1d	127
75.19	bin2d	127
75.20	binormrnd	127
75.21	coherence	127
75.22	conv1_man	127
75.23	conv2_man	128
75.24	conv2z	128
75.25	conv30	128
75.26	conv_	128
75.27	conv_centered	128
75.28	convz	128
75.29	cosexpdelay	128
75.30	csmooth	128
75.31	daniell_window	129
75.32	danielle_window	129
75.33	db2neper	129

75.34	db2power	129
75.35	derive_bandpass_normalization_and_zeros	129
75.36	derive_danielle_weight	129
75.37	derive_limit_0_acfar	129
75.38	detect_peak	129
75.39	determine_phase_shift	130
75.40	determine_phase_shift1	130
75.41	digital_low_pass_filter	130
75.42	doublesum_ij	130
75.43	effective_sample_size_to_ar1	130
75.44	filt_hodges_lehman	130
75.45	filter1	130
75.46	filter2	130
75.47	filter_	131
75.48	filter_r_to_f0	131
75.49	filter_rho_to_f0	131
75.50	filter_twosided	131
75.51	filteriir	131
75.52	filterp	132
75.53	filterp1	132
75.54	filterstd	132
75.55	firls_man	132
75.56	fit_spectral_density	132
75.57	fit_spectral_density_2d	132
75.58	fit_spectral_density_radial	132
75.59	flattopwin	133
75.60	frequency_response_boxcar	133
75.61	freqz_boxcar	133
75.62	gaussfilt1	133
75.63	hanchangewin	133
75.64	hanchangewin2	133
75.65	hanwin	133
75.66	hanwin_	133
75.67	high_pass_1d_simple	134
75.68	highpass	134
75.69	highpass1d_fft_cos	134
75.70	highpass1d_implicit	134
75.71	highpass2d_fft	134
75.72	highpass2d_ideal	134
75.73	highpass2d_implicit	134
75.74	highpass_arg	134
75.75	highpass_fc_to_rho	134
75.76	jackknife_block	135
75.77	kaiserwin	135

75.78	kalman	135
75.79	lanczoswin	135
75.80	last	135
75.81	lowpass	135
75.82	lowpass1d_fft	135
75.83	lowpass1d_implicit	135
75.84	lowpass2	135
75.85	lowpass2d_2	136
75.86	lowpass2d_anisotropic	136
75.87	lowpass2d_fft	136
75.88	lowpass2d_ideal	136
75.89	lowpass2d_implicit	136
75.90	lowpass_arg	136
75.91	lowpass_fc_to_rho	136
75.92	lowpass_iir	136
75.93	lowpass_iir_symmetric	136
75.94	lowpassfilter2	137
75.95	maxfilt1	137
75.96	meanfilt1	137
75.97	meanfilt2	137
75.98	medfilt1_man	137
75.99	medfilt1_man2	137
75.100	medfilt1_padded	137
75.101	medfilt1_reduced	137
75.102	mid_term_single_sample	138
75.103	minfilt1	138
75.104	mu2ar1	138
75.105	mysmooth	138
75.106	nanautocorr	138
75.107	nanmedfilt1	138
75.108	neper2db	138
75.109	oscillator_noisy	138
75.110	peaks_man	139
75.111	periodogram	139
75.112	periodogram_2d	139
75.113	periodogram_annular	139
75.114	periodogram_bartlett	139
75.115	periodogram_bootstrap	139
75.116	periodogram_confidence_interval	139
75.117	periodogram_median	139
75.118	periodogram_p_value	139
75.119	periodogram_qq	140
75.120	periodogram_quantiles	140
75.121	periodogram_radial	140

75.122	periodogram_std	140
75.123	periodogram_test_periodicity	140
75.124	periodogram_test_periodicity_2d	141
75.125	periodogram_test_stationarity	141
75.126	periodogram_welsh	141
75.127	polyfilt1	141
75.128	qmedfilt1	141
75.129	radial_window	142
75.130	randar1	142
75.131	randar1_dual	142
75.132	randar2	142
75.133	randarp	142
75.134	range_window	142
75.135	rectwin	142
75.136	recursive_sum	142
75.137	select_range	143
75.138	smooth1d_parametric	143
75.139	smooth2	143
75.140	smooth_man	143
75.141	smooth_parametric	143
75.142	smooth_parametric2	143
75.143	smooth_with_splines	143
75.144	smoothfft	143
76 signal-processing/spectral-density		144
76.1	spectral_density_ar2	144
76.2	spectral_density_area	144
76.3	spectral_density_bandpass2d_ideal	144
76.4	spectral_density_bandpass_2d	144
76.5	spectral_density_bandpass_2d_scale	144
76.6	spectral_density_bandpass_2d_scale_old	144
76.7	spectral_density_bandpass_continuous	144
76.8	spectral_density_bandpass_continuous_max	145
76.9	spectral_density_bandpass_continuous_max2par	145
76.10	spectral_density_bandpass_continuous_scale	145
76.11	spectral_density_bandpass_discrete	145
76.12	spectral_density_brownian_phase	145
76.13	spectral_density_brownian_phase_2d	145
76.14	spectral_density_brownian_phase_across	145
76.15	spectral_density_brownian_phase_mode	145
76.16	spectral_density_brownian_phase_mode2par	146
76.17	spectral_density_brownian_phase_scale	146
76.18	spectral_density_estimate_2d	146
76.19	spectral_density_flat	146

76.20	spectral_density_highpass	146
76.21	spectral_density_highpass2d_ideal	146
76.22	spectral_density_highpass_2d	146
76.23	spectral_density_highpass_cos	146
76.24	spectral_density_lorentzian	146
76.25	spectral_density_lorentzian_max	147
76.26	spectral_density_lorentzian_max2par	147
76.27	spectral_density_lorentzian_scale	147
76.28	spectral_density_lowpass	147
76.29	spectral_density_lowpass2d_ideal	147
76.30	spectral_density_lowpass_2d	147
76.31	spectral_density_lowpass_one_sided	147
76.32	spectral_density_periodic_additive_noise	147
76.33	spectral_density_wperiodic	147
77	signal-processing	148
77.1	spectrogram	148
77.2	std_window	148
77.3	sum_i_lag	148
77.4	sum_ii	148
77.5	sum_ii_	148
77.6	sum_ij	148
77.7	sum_ij_	148
77.8	sum_ij_partial_	149
77.9	sum_multivar	149
77.10	test_acfar1	149
77.11	test_acfar1_2	149
77.12	test_acfar1_3	149
77.13	test_acfar1_4	149
77.14	test_acfar2	149
77.15	test_ar1_var_factor	149
77.16	test_ar1_var_factor_2	149
77.17	test_ar1_var_mu_single_sample	149
77.18	test_ar1_var_pop	150
77.19	test_ar1_var_pop_1	150
77.20	test_ar1delay	150
77.21	test_bivariate_covariance_term	150
77.22	test_convexity	150
77.23	test_lanczoswin	150
77.24	test_madcorr	150
77.25	test_randar1	150
77.26	test_randar1_multivariate	150
77.27	test_randar2	150
77.28	test_sum_ij	151

77.29	test_sum_multivar	151
77.30	test_trifilt1	151
77.31	test_wautocorr	151
77.32	test_wavelet_transform	151
77.33	test_wordfilt	151
77.34	test_xar1_mid_term	151
77.35	tikhonov_to_ar1	151
77.36	trapwin	151
77.37	trifilt1	152
77.38	trifilt2	152
77.39	triwin	152
77.40	triwin2	152
77.41	varar1	152
77.42	welch_spectrogram	152
77.43	wfilt	152
77.44	winbandpass	153
77.45	window2d	153
77.46	window_make_odd	153
77.47	winfilt0	153
77.48	winlength	153
77.49	wmeanfilt	153
77.50	wmedfilt	153
77.51	wordfilt	153
77.52	wordfilt_edgeworth	154
77.53	wrapphase	154
77.54	xar1	154
77.55	xcorr_man	154
78	sorting	154
78.1	sort2	154
78.2	sort2d	154
79	special-functions	154
79.1	bessel_sphere	154
79.2	digamma_man	154
79.3	exp10	155
79.4	hankel_sphere	155
79.5	hermite	155
79.6	legendre_man	155
79.7	neumann_sphere	155
80	statistics	155
80.1	atan_s2	155
80.2	beta_kurt	156

80.3	beta_mean	156
80.4	beta_mode_to_parameter	156
80.5	beta_skew	156
80.6	beta_std	156
80.7	chi2_kurt	156
80.8	chi2_mean	156
80.9	chi2_skew	156
80.10	chi2_std	156
80.11	coefficient_of_determination	156
80.12	conditional_expectation_normal	157
80.13	correlation_confidence_pearson	157
81	statistics/distributions	157
81.1	PDF	157
81.2	binorm_separation_coefficient	157
81.3	binormcdf	157
81.4	binormfit	157
81.5	binormpdf	157
81.6	edgeworth_cdf	157
81.7	edgeworth_pdf	158
81.8	gam_moment2param	158
81.9	logn_mean	158
81.10	logn_mode	158
81.11	logn_moment2param	158
81.12	logn_param2moment	158
81.13	logn_std	158
81.14	lognpdf_	158
81.15	pdfsample	159
81.16	t2cdf	159
81.17	t2inv	159
82	statistics	159
82.1	example_standard_error_of_sample_quantiles	159
82.2	f_var_finite	159
82.3	fisher_mean	159
82.4	fisher_std	159
82.5	gam_mean	159
82.6	gam_std	160
82.7	gamma_mode_to_parameter	160
82.8	gamma_stirling	160
82.9	gaussfit3	160
82.10	gaussfit_quantile	160
82.11	geoserr	160
82.12	geostd	160

82.13	hodges_lehmann_correlation	160
82.14	hodges_lehmann_dispersion	161
83	statistics/information-theory	161
83.1	akaike_information_criterion	161
83.2	bayesian_information_criterion	161
84	statistics	161
84.1	kurtncdf	161
84.2	kurtnpdf	161
84.3	kurtosis_bias_corrected	162
84.4	limit	162
84.5	logfactorial	162
84.6	loglogpdf	162
84.7	lognfit_quantile	162
84.8	logskewcdf	162
84.9	logskewpdf	162
85	statistics/logu	162
85.1	lambertw_numeric	162
85.2	logtrialtcd	163
85.3	logtrialtinv	163
85.4	logtrialtmean	163
85.5	logtrialtpdf	163
85.6	logtrialtrnd	163
85.7	logtricdf	163
85.8	logtriinv	164
85.9	logtrimean	164
85.10	logtripdf	164
85.11	logtrirnd	164
85.12	logucdf	164
85.13	loguclm	164
85.14	loguinv	164
85.15	logumean	164
85.16	logupdf	165
85.17	logurnd	165
85.18	loguvar	165
85.19	medlogu	165
85.20	test_logurnd	165
85.21	tricdf	165
85.22	triinv	165
85.23	trimean	165
85.24	tripdf	166
85.25	trirnd	166

86	statistics	166
86.1	max_exprnd	166
86.2	maxnnormals	166
86.3	mean_generalized_gampdf	166
86.4	midrange	166
86.5	minavg	166
86.6	mode_man	167
87	statistics/moment-statistics	167
87.1	autocorr_man3	167
87.2	autocorr_man4	167
87.3	autocorr_man5	167
87.4	blockserr	167
87.5	comoment	167
87.6	corr_man	168
87.7	cov_man	168
87.8	dof	168
87.9	edgeworth_quantile	168
87.10	effective_sample_size	168
87.11	f_correlation	169
87.12	f_finite	169
87.13	lmean	169
87.14	lmoment	169
87.15	maskmean	169
87.16	masknanmean	169
87.17	mean1	169
87.18	mean_man	169
87.19	mse	170
87.20	nanautocorr_man1	170
87.21	nanautocorr_man2	170
87.22	nanautocorr_man4	170
87.23	nancorr	170
87.24	nancumsum	170
87.25	nanlmean	170
87.26	nanr2	171
87.27	nanrms	171
87.28	nanrmse	171
87.29	nanserr	171
87.30	nanwmean	171
87.31	nanwstd	171
87.32	nanwvar	171
87.33	nanxcorr	172
87.34	pearson	172
87.35	pearson_to_kendall	172

87.36	pool_samples	172
87.37	qmean	172
87.38	range_mean	172
87.39	rmse_	172
87.40	serr	172
87.41	serr1	173
87.42	test_qskew	173
87.43	test_qstd_qskew_optimal_p	173
87.44	wautocorr	173
87.45	wcorr	173
87.46	wcov	173
87.47	wdof	173
87.48	wkurt	174
87.49	wmean	174
87.50	wrms	174
87.51	wserr	174
87.52	wskew	174
87.53	wstd	174
87.54	wvar	174
88 statistics		175
88.1	nangeomean	175
88.2	nangeostd	175
89 statistics/nonparametric-statistics		175
89.1	kernel1d	175
89.2	kernel2d	175
90 statistics		175
90.1	normmoment	175
90.2	normpdf2	176
91 statistics/order-statistics		176
91.1	hodges_lehmann_location	176
91.2	kendall	176
91.3	kendall_to_pearson	176
91.4	mad2sd	176
91.5	madcorr	176
91.6	median2_holder	177
91.7	median_ci	177
91.8	median_man	177
91.9	mediani	177
91.10	nanmadcorr	177
91.11	nanwmedian	177

91.12	nanwquantile	177
91.13	oja_median	178
91.14	qkurtosis	178
91.15	qmoments	178
91.16	qskew	178
91.17	qskewq	178
91.18	qstdq	179
91.19	quantile1_optimisation	179
91.20	quantile2_breckling	179
91.21	quantile2_chaudhuri	179
91.22	quantile2_projected	179
91.23	quantile2_projected2	179
91.24	quantile_envelope	179
91.25	quantile_regression_simple	179
91.26	ranking	180
91.27	spatial_median	180
91.28	spatial_quantile	180
91.29	spatial_quantile2	180
91.30	spatial_quantile3	180
91.31	spatial_rank	180
91.32	spatial_sign	180
91.33	spatial_signed_rank	180
91.34	spearman	181
91.35	spearman_rank	181
91.36	spearman_to_pearson	181
91.37	wmedian	181
91.38	wquantile	181
92	statistics	181
92.1	qstd	181
92.2	quantile_extrap	181
92.3	quantile_sin	181
93	statistics/random-number-generation	182
93.1	laplacernd	182
93.2	randc	182
93.3	skewness2param	182
93.4	skewpdf_central_moments	182
93.5	skewrnd	182
93.6	skewrnd2	182
94	statistics	182
94.1	range	182
94.2	resample_with_replacement	182

95	statistics/resampling-statistics/@Jackknife	183
95.1	Jackknife	183
95.2	estimated_STATIC	183
95.3	matrix1_STATIC	183
95.4	matrix2	183
96	statistics/resampling-statistics	184
96.1	block_jackknife	184
96.2	jackknife_moments	184
96.3	moving_block_jackknife	184
96.4	randblockserr	184
96.5	resample	184
97	statistics	185
97.1	scale_quantile_sd	185
97.2	sd_sample_quantiles	185
97.3	skew_generalized_normal_fit	185
97.4	skew_generalized_normpdf	185
97.5	skewcdf	185
97.6	skewparam_to_central_moments	185
97.7	skewpdf	185
97.8	spatialrnd	186
97.9	test_mean_generalized_gampdf	186
97.10	test_skew_generalized_normpdf	186
97.11	trimmed_mean	186
97.12	ttest2_man	186
97.13	ttest_man	186
97.14	ttest_paired	186
97.15	wgeomean	187
97.16	wgeovar	187
97.17	wharmean	187
97.18	wharstd	187
97.19	wharvar	187
97.20	wnormpdf	187
98	stochastic	187
98.1	brownian_noise_1d_acf	187
98.2	brownian_noise_1d_fft	187
98.3	brownian_noise_2d_fft	188
99	mathematics	188
99.1	ternary_diagram	188
100	test/master	188

100.1	dat_test_lanczos_3d_k_20_n_40	188
100.2	poisson2d.blk	188
100.3	qr_implicit_givens_2	188
100.4	spectral_derivative_2d	188
100.5	test_2d_eigensolver_hydrogen	188
100.6	test_2d_refine	188
100.7	test_3d_eigensolver_hydrogen	189
100.8	test_FEM	189
100.9	test_Mesh_3d	189
100.10	test_arnoldi	189
100.11	test_arpackc	189
100.12	test_assemble	189
100.13	test_assembly_performance	189
100.14	test_bc_one_sided	189
100.15	test_compare_solvers	189
100.16	test_complete	189
100.17	test_convergence	190
100.18	test_convergence_b	190
100.19	test_df_2d	190
100.20	test_eig_algs	190
100.21	test_eig_inverse	190
100.22	test_eigs_lanczos	190
100.23	test_eigs_lanczos_1	190
100.24	test_eigs_lanczos_2	190
100.25	test_eigs_lanczos_performance	190
100.26	test_fdm	190
100.27	test_fdm_d_vargrid	191
100.28	test_fdm_spectral	191
100.29	test_fem	191
100.30	test_fem_1d	191
100.31	test_fem_1d_higher_order	191
100.32	test_fem_2d_adaptive	191
100.33	test_fem_2d_higher_order	191
100.34	test_fem_3d_higher_order	191
100.35	test_fem_3d_refine	191
100.36	test_fem_b	191
100.37	test_fem_derivative	192
100.38	test_fem_quadrature	192
100.39	test_final	192
100.40	test_fix_substitution	192
100.41	test_forward	192
100.42	test_get_sparse_arrays	192
100.43	test_harmonic_oscillator	192
100.44	test_high_order_fdm_periodic_bc	192

100.45	test_hydrogen_wf	192
100.46	test_ichol	192
100.47	test_interpolation	193
100.48	test_inverse_problem	193
100.49	test_it_vs_exact	193
100.50	test_jama	193
100.51	test_jd	193
100.52	test_jdqz	193
100.53	test_lanczos_2	193
100.54	test_lanczos_biorthogonal	193
100.55	test_laplacian	193
100.56	test_laplacian_non_uniform	193
100.57	test_laplacian_simple	194
100.58	test_mesh_2d_uniform	194
100.59	test_mesh_2d_uniform_2	194
100.60	test_mesh_circle	194
100.61	test_mesh_generation	194
100.62	test_mesh_interpolate	194
100.63	test_mg	194
100.64	test_minres_recycle	194
100.65	test_multigrid	194
100.66	test_nc	194
100.67	test_nonuniform_symmetric	195
100.68	test_pde	195
100.69	test_permutation	195
100.70	test_poisson_fem	195
100.71	test_polar	195
100.72	test_potential	195
100.73	test_powers	195
100.74	test_precondition	195
100.75	test_project_rectangle	195
100.76	test_qr	195
100.77	test_quantum_well	196
100.78	test_radial_adaptive	196
100.79	test_radial_confinement	196
100.80	test_radial_fixes	196
100.81	test_refine_2d	196
100.82	test_refine_2d_b	196
100.83	test_refine_3d	196
100.84	test_refine_structural	196
100.85	test_regularisation	196
100.86	test_round_off	196
100.87	test_schrödinger_potentials	197
100.88	test_uniform_mesh	197

100.89	test_vargrid	197
101	test	197
101.1	test_bandpass2d	197
101.2	test_gaussfit3	197
101.3	test_geoserr	197
101.4	test_hexagonal_pattern	197
101.5	test_lognfit_quantile	197
101.6	test_lowpass1d_implicit	197
101.7	test_lowpass2d_anisotropic	198
101.8	test_lowpass2d_fft	198
101.9	test_max_normal	198
101.10	test_moments_fourier_power	198
101.11	test_mtimes3x3	198
102	wavelet	198
102.1	contiuous_wavelet_transform	198
102.2	cwt_man	198
102.3	example_wavelets	198
102.4	phaseswap	199
102.5	test_cwt_man	199
102.6	test_phaseswap	199
102.7	test_wavelet	199
102.8	test_wavelet2	199
102.9	test_wavelet_analysis	199
102.10	test_wavelet_reconstruct	199
102.11	test_wtc	199
102.12	wavelet	199
102.13	wavelet_reconstruct	200
102.14	wavelet_transform	200

1 calendar

1.1 days_per_month

1.2 isnight

2 mathematics

mathematical functions of various kind

2.1 cast_byte_to_integer

cast byte to integer

3 complex-analysis

operations on complex numbers

3.1 complex_exp_product_im_im

product of the imaginary part of two complex exponentials

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.2 complex_exp_product_im_re

product of the imaginary part of one and the real part of a second complex exponential

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.3 complex_exp_product_re_im

the product has two frequency components

product of the imaginary part of one and the real part of a second
complex exponential

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.4 complex_exp_product_re_re

product of the real part of two complex exponentials

$$\begin{aligned} \text{re}(c1 \exp(i o1 x)) * \text{re}(c2 \exp(i o2 x)) = \\ 1/2 * (\text{real}(c1 * c2 * \exp(i * (n1 + n2) * o * x)) \dots \\ + \text{real}(\text{conj}(c1) * c2 * \exp(i * (n2 - n1) * o * x))) \end{aligned}$$

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.5 croots

nth-roots of a complex number

input:
c : complex number
n : order of root
 n must be rational, to obtain n solutions
 otherwise no finite set of solutions exists

r : roots of the complex number

3.6 root_complex

root of a complex number

3.7 test_imroots

4 derivation

derivation of several functions by means of symbolic computation

4.1 derive_acfar1

4.2 derive_ar1_spectral_density

4.3 derive_ar2param

4.4 derive_arc_length

4.5 derive_fourier_power

4.6 derive_fourier_power_exp

4.7 derive_laplacian_curvilinear

4.8 `derive_laplacian_fourier_piecewise_linear`

4.9 `derive_logtripdf`

4.10 `derive_smooth1d_parametric`

4.11 `derive_spectral_density_bandpass_initial_condition`

5 `derivation/master`

5.1 `derive_bc_one_sided`

5.2 `derive_convergence`

5.3 `derive_error_fdm`

5.4 `derive_fdm_poly`

5.5 `derive_fdm_power`

5.6 `derive_fdm_taylor`

5.7 `derive_fdm_vargrid`

5.8 `derive_fem_2d_mass`

5.9 `derive_fem_error_2d`

5.10 `derive_fem_error_3d`

5.11 `derive_fem_sym_2d`

5.12 `derive_grid_constants`

5.13 `derive_interpolation`

5.14 `derive_laplacian`

5.15 `derive_limit`

5.16 `derive_nc_1d`

5.17 `derive_nc_1d_`

5.18 `derive_nc_2d`

5.19 `derive_nonuniform_symmetric`

%

5.20 `derive_richardson`

5.21 `derive_sum`

5.22 `nn`

5.23 `test_derive`

5.24 `test_derive_fdm_poly`

5.25 `test_filter`

5.26 test_vargrid

6 derivation

derivation of several functions by means of symbolic computation

6.1 simplify_atan

symbolic simplification of the arcus tangent

7 mathematics

mathematical functions of various kind

7.1 entropy

8 finance

8.1 derive_skewrnd_walsh_paramter

8.2 gbm_cdf

8.3 gbm_fit

8.4 gbm_fit_old

8.5 `gbm_inv`

8.6 `gbm_mean`

8.7 `gbm_median`

8.8 `gbm_pdf`

8.9 `gbm_simulate`

8.10 `gbm_skewness`

8.11 `gbm_std`

8.12 `gbm_transform_time_step`

8.13 `put_price_black_scholes`

8.14 `skewgbm_simulate`

8.15 skewrnd_walsh

9 finance/test

9.1 test_gbm

9.2 test_gbm_pdf

9.3 test_skewrnd_walsh

10 fourier/@STFT

10.1 STFT

class for short time fourier transform

Note: the interval T_i should be set to at least $2 \cdot \max(T)$, as otherwise coefficients

tend to oscillate in the presence of noise

Note: for convenience, the independent variable is labeled as time (t),

but the independent variable is arbitrary, so it works likewise in space

10.2 itransform

inverse of the short time fourier transform

10.3 stft_

static wrapper for STFT

10.4 stftmat

transformation matrix for the short time fourier transform

10.5 transform

short time fourier transform

11 fourier

support and analysis functions both for the discrete (fast) fourier
transform (dft/fft)
and continuous fourier analysis (fourier series)

11.1 amplitude_from_peak

amplitude and standard deviation of the amplitude of a frequency
component

represented by a peak in the fourier domain

input :

h : peak height

w : peak width at half height

output:

a : amplitude in real space

s : standard deviation of the frequency (!)

11.2 dftmtx_man

fourier matrix in matlab style with a limited number of rows,
columns of higher frequencies are omitted

input :

n : number of samples

nr : number of columns

output :

F : fourier matrix

11.3 example_fourier_window

11.4 fft_man

fast fourier transform for complex input data

input:
F : data in real space

output :

F : fourier transformation of F

11.5 fftsmooth

smooth the fourier transform and determine upper and lower bound confidence intervals

input :
f :
sfunc : a smoothing function (for example fir convolution with rectangular window)
 returns filtered (mean) value and normalized fir window
nf : window length
nsigma : number of standard deviations for confidence intervals

output :
ff : filtered fourier transform
l : lower bound
u : upper bound

11.6 fix_fourier

fill gaps (missing data) by means of fourier extrapolation

fix periodic data series with fourier interpolation
longest gap should not exceed 1/2 of the shortest time span of interest (1/cutoff frequency)
note: this limit equals the position of first side lobe of the ft of a rectangular window with gap length

11.7 `fourier_axis`

return axis of frequencies and periods for the discrete fourier transform
as computed by fft (matlab-style)

input:

X : sample locations (equal interval)

L : length of samples

n : number of samples

output :

f : frequencies

T : periods

mask : mask for 1/2 of the fourier transform
(as both halves are complex conjugates)

N : frequency id

11.8 `fourier_axis_2d`

11.9 `fourier_cesaro_correction`

11.10 `fourier_coefficient_piecewise_linear`

fourier series coefficients of a piecewise linear function
(not coefficient of discrete fourier transform)
function can be discontinuous between intervals
scales domain length to 2π

input :

l,r : end points of piecewise linear function

lval, rval : values at end points

L : length of domain

n : number of samples/highest frequency

output :

a, b : coefficients for frequency components

11.11 `fourier_coefficient_piecewise_linear_1`

fourier series coefficients of a piecewise linear function
(not coefficient of discrete fourier transform)
function can be discontinuous between intervals
scales domain length to 2π

input :
X : end points of piecewise linear function
Y : values at end points

output :
ab : coefficients for frequency components

11.12 `fourier_coefficient_ramp3`

fourier series coefficient of a ramp

11.13 `fourier_coefficient_ramp_pulse`

fourier series coefficient of a ramp pules

11.14 `fourier_coefficient_ramp_step`

fourier coefficient of a ramp-step

11.15 `fourier_coefficient_square_pulse`

fourier series coefficients of a square pulse

11.16 `fourier_cubic_interaction_coefficients`

11.17 `fourier_derivative`

derivative via fourier transform
exponential convergence for periodic functions
results in spurious oscillations for aperiodic functions

input:

`x` : data, sampled in equal intervals

`k` : order of the derivative

`dx` : kth-derivative of `x`

note : 1) the derivative converges with spectral accuracy, i.e. is
exact up to rounding condition for `L` sufficiently large
and `x` being periodic
2) the derivative converges with order `p`, when `x` has only
`p`-continuous derivatives, including discontinuous
derivatives
over the boundary
3) discontinuous derivatives result in gibbs phenomenon

11.18 `fourier_derivative_matrix_1d`

11.19 `fourier_derivative_matrix_2d`

11.20 `fourier_expand`

expand values of fourier series

11.21 `fourier_fit`

fit a fourier series to a set of sample points that are not spaced
in
equal intervals

11.22 `fourier_interpolate`

interpolate samples `y` sampled at moments (location) `t` to locations
`ti`

11.23 `fourier_matrix`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.24 `fourier_matrix2`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.25 `fourier_matrix3`

transformation matrix for the continuous fourier transform
this is a matrix with $(2n+1)$ real columns

11.26 `fourier_matrix_exp`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.27 `fourier_multiplicative_interaction_coefficients`

11.28 `fourier_power`

powers of a continuous fourier series in sin/cos form

powers of $a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \delta))$ ^p
phase of first component assumed 0

frequencies higher than 2ω ignored in input
frequencies higher than 3ω not computed

11.29 `fourier_power_exp`

powers of the continuous fourier series

$$a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \delta)) ^p$$

phase of first component assumed 0

higher orders than 2 ignored input

higher order than 3 not computed in output

$$y = a_0 + \sum (a_j \sin(j\omega t) + b_j \cos(j\omega t))$$

$$= \text{Real}(\sum_{i=0}^{\infty} c_i \exp(i\omega t)), \quad c_i = a_i + b_i$$

NOT the alternative $\sum_{i=-\infty}^{\infty} \tilde{c}_i$, tile $c_j = 1/2 a_j + 1/2i b_j$

11.30 `fourier_predict`

expand a continuous fourier series at times t

11.31 `fourier_quadratic_interaction_coefficients`

11.32 `fourier_random_phase_walk`

evaluate fourier series where the phase undergoes a brownian motion

11.33 `fourier_range`

approximate range of a continuous Fourier series with 2 components

$$\text{range}(y) = \max(y) - \min(y)$$

11.34 `fourier_regress`

fit a continuous fourier series to a set of sample points not
sampled
at equal intervals

11.35 `fourier_resampled_fit`

fits coefficients of a continuous fourier transform,
but stores them as resampled values

11.36 `fourier_resampled_predict`

interpolates a continuous fourier series that has been stored as
values
at their support points

11.37 `fourier_signed_square`

coefficients of the fourier series of $|\cos a + \cos t|$ ($\cos a + \cos t$)
in general
 $\cos a$ is midrange
 $\cos t$ is tidal variation
c.f Dronkers

11.38 `fourier_transform`

continuous fourier transformation of y
(not discrete fourier transformation dft/fft)

input:
 b : data sampled at equal intervals
 T : length of data in time or space, i.e. position of last
 sample if
 position of first sample is 0
 T_max : maximum period to include

output :
 A : fourier matrix
 p : fourier transformation of b
 tt : TODO

11.39 `fourier_transform_fractional`

11.40 hyperbolic_fourier_box

11.41 idftmtx_man

inverse matrix for the discrete fourier transform in matlab style
with a limited number of columns, thus ignoring higher frequencies
keep $2nc+1$ columns (mean and conj-complex pairs of nc frequencies)

11.42 laplace_2d_pwlinear

solution to the Laplacian in two dimensions for a finite
rectangular domain
with piecewise constant boundary conditions
linear system with 4 unknowns per frequency component
these are coefficients of s, c, sh, ch
 $(pu*(s + c) + qu*(s' + c'))*(shu + chu) = ru$ % upper bc
 $(pd*(s + c) + qd*(s' + c'))*(shd + chd) = rd$ % lower bc
 $((sl + cl)*(pl*(shl + chl) + ql*(shl' + chl'))) = rl$ % left
bc
 $((sr + cr)*(pr*(shr + chr) + qr*(shr' + chr'))) = rr$ % right
bc

least squares with piecewise integration
[x0,p,q,r] piecewise linear polynomials at the boundaries

11.43 mean_fourier_power

11.44 moments_fourier_power

11.45 nanfft

discrete fourier transform of a data series with gaps

11.46 peaks

peaks of the power spectrum of a discrete fourier transform

rule for peaks: there is no higher value left or right of the "peak"

until the signal drops to $p \cdot y_{\text{peak}}$, $p = 0.5$

works best, when spectrum has been smoothened

input :

f : frequency

y : absolute value of fourier transform (power spectrum)

L : length in space or time of series

output :

a0 : amplitude

s0 : standard deviation (error?) of amplitude

w0 : width of peak

lambda = wave length (period?)

pdx : index of peak

f : frequency (if not given as input)

11.47 roots_fourier

zeros of continuous fourier series series

$$f = a_0 + \sum_{j=1}^n a_j \cos(j x) + b_j \sin(j x)$$

11.48 spectral_density

spectral density

11.49 std_fourier_power

11.50 test_complex_exp_product

11.51 test_fourier_filter

11.52 test_idftmtx

11.53 var_fourier_power

12 mathematics

mathematical functions of various kind

12.1 gaussfit_quantile

13 geometry/@Geometry

13.1 Geometry

13.2 arclength

arc length of a two dimensional curve

8th order accurate

does not require the segments length to vary smoothly

note: the curve can be considered parametric, e.g. $x = x(t)$, $y = y(t)$
and

and $t = t(s)$, but the error term contains derivatives of t ,
thus a non smooth t (strongly varying distance between points)
requires the scaling as done below

13.3 arclength_old

arc length of a two dimensional function

13.4 arclength_old2

arc length of a two dimensional function

13.5 base_point

base point (fusspunkt), i.e. point on a line with shortest distance to another point

13.6 base_point_limited

base point (Fusspunkt) of a point on a line

13.7 centroid

centroid pf a polygone

13.8 cosa_min_max

13.9 cross2

cross product in two dimensions

13.10 curvature

curvature of a function in two dimensions

13.11 ddot

sum of squares of cos of inner angles of triangle

13.12 distance

euclidian distance between two points

13.13 distance2

euclidean distance between two points
this function requires a and be of equal dimensions, or the least
the first pair or second pair to be a scalar

13.14 dot

dot product

13.15 edge_length

edge length

13.16 enclosed_angle

angle enclosed between two lines

13.17 enclosing_triangle

smallest enclosing equilateral triangle with bottom side parallel to
X axis

13.18 hexagon

coordinates of a hexagon, scaled and rotated

13.19 inPolygon

flag points contained in a polygon
much faster than matlab internal function

13.20 inTetra

flag points contained in tetrahedron

13.21 inTetra2

flag points contained in tetrahedron

13.22 inTriangle

flag points contained in triangle
function [flag, c] = inTriangle(P1,P2,P3,P0)

13.23 intersect

intersect between two lines

13.24 lineintersect

intersect of two lines

13.25 lineintersect1

intersect of two lines

13.26 minimum_distance_lines

minimum distance of two lines in three dimensions

13.27 mittenpunkt

mittenpunkt of a triangle

13.28 nagelpoint

nagelpoint of a triangle

13.29 onLine

13.30 orthocentre

orthocentre of triangle

13.31 plumb_line

13.32 poly_area

area of a polygon
function A = poly_area(x,y)

13.33 poly_edges

edges of a polygon

13.34 poly_set

associate point at arbitrary location with a polygon it is contained
in
and assign the value of the polygon to it

13.35 poly_width

width of polygon width holes by surface normals
holes / islands separated with NaN
order of points of outer boundary must be cw
order of points of holes must be ccw
note that this function does not give the true width for expanding
sections
use voronoi polygons for this

13.36 polyxpoly

intersections of two polygons

13.37 project_to_curve

closest point on a curve with respect to a point at distance to the
curve

13.38 quad_isconvex

13.39 random_disk

draw random points on the unit disk

13.40 random_simplex

random point inside of a triangle

13.41 sphere_volume

volume of a sphere
function v = sphere_volume(r)

13.42 tetra_volume

volume of a tetrahedron

13.43 tobarycentric

cartesian to barycentric coordinates

13.44 tobarycentric1

cartesian to barycentric coordinates

13.45 tobarycentric2

cartesian to barycentric coordinates

13.46 tobarycentric3

cartesian to barycentric coordinates

13.47 tri_angle

cos of angles of a triangle

13.48 tri_area

angle of a triangle

13.49 tri_centroid

centroid of a triangle

13.50 tri_distance_opposit_midpoint

distance between corner of a triangle and its opposing mid-point

13.51 tri_edge_length

edge length of a triangle

13.52 tri_edge_midpoint

mid point of a triangle

13.53 tri_excircle

excircle of a triangle

13.54 tri_height

height of a triangle

13.55 tri_incircle

incircle of a triangle

13.56 tri_isacute

flag acute triangles

13.57 tri_isobtuse

flag obtuse triangles

13.58 tri_semiperimeter

semiperimeter of a triangle

13.59 tri_side_length

edge length of triangle

14 geometry

14.1 Polygon

Simple 2D polygon class

Polygon properties:

- x - x coordinates of polygon
- y - y coordinates of polygon
- nnodes - number of nodes in the polygon

Polygon methods:

- in - checks whether given points lie inside, on the edge, or outside of the polygon
- area - returns the area of the polygon
- centerline - computes the centerline of the river
- iscw - check whether polygon is clockwise
- reverse - reverse the order of the polygon

14.2 bounding_box

bounding box of X

14.3 curvature_1d

curvature of a sampled parametric curve in two dimensions

14.4 cvt

centroidal voronoi tessellation

14.5 deg_to_frac

degree, minutes and seconds to fractions

14.6 ellipse

return points on an ellipse
n : number of points
ci : confidence interval, i.e. for 1 sigma

14.7 ellipseX

x-coordinates of y-coordinates of an ellipse

14.8 ellipseY

14.9 first_intersect

get first intersection between lines in A and B

14.10 golden_ratio

golden ratio

14.11 hypot3

hypotenuse in 3D

14.12 meanangle

weighted mean of angles

14.13 meanangle2

mean angle

14.14 meanangle3

mean angle

14.15 meanangle4

mean angle

14.16 medianangle

median angle
angle, that has the smallest squared distance to all others

14.17 medianangle2

median angle

input
alpha : x*m, [rad] angle

output
ma : 1*m, [rad] median angle
sa : 1*m, [rad] standard error of median angle for uncorrelated
error

14.18 pilim

limit to $\pm \pi$

14.19 streamline_radius_of_curvature

streamline radius of curvature
simplifies when rotate to streamwise coordinates to $R = 1/dv/ds * u$

15 histogram/@Histogram

15.1 2x

15.2 Histogram

15.3 bimodes

15.4 cdf

15.5 cdfS

15.6 chi2test

15.7 cmoment

15.8 cmomentS

15.9 entropy

15.10 entropyS

15.11 export_csv

15.12 iquantile

15.13 kstest

15.14 kurtosis

15.15 kurtosisS

15.16 mean

15.17 meanS

15.18 median

15.19 medianS

15.20 mode

15.21 modeS

15.22 moment

15.23 momentS

15.24 pdf

15.25 quantile

15.26 quantileS

15.27 resample

15.28 setup

15.29 **skewness**

15.30 **skewnessS**

15.31 **stairs**

15.32 **stairsS**

15.33 **std**

15.34 **stdS**

15.35 **var**

15.36 **varS**

16 **histogram**

16.1 **hist_man**

16.2 **histadapt**

16.3 **histconst**

16.4 **pdf_poly**

16.5 **plotcdf**

16.6 **test_histogram**

17 **mathematics**

mathematical functions of various kind

17.1 **imrotmat**

18 **linear-algebra**

18.1 **averaging_matrix_2**

18.2 **colnorm**

norms of columns

18.3 condest_

estimation of the condition number

18.4 connectivity_matrix

19 linear-algebra/coordinate-transformation

19.1 barycentric2cartesian

barycentric to cartesian coordinates

19.2 barycentric2cartesian3

convert barycentric to cartesian coordinates

19.3 cartesian2barycentric

cartesian to barycentric coordinates

19.4 cartesian_to_unit_triangle_basis

transform coordinates into unit triangle

19.5 ellipsoid2geoid

19.6 example_approximate_utm_conversion

19.7 latlon2utm

transform latitude and longitude to WGS84 UTM

19.8 latlon2utm_simple

19.9 lowrance_mercator_to_wgs84

convert lowrance coordinates to wgs84

based on spreadsheet by D Whitney King and Patty B at Lowrance

19.10 nmea2utm

convert nmea messages to utm coordinates

19.11 sn2xy

convert sn to xy coordinates

19.12 unit_triangle_to_cartesian

transform coordinates in unit triangle to cartesian coordinates

19.13 utm2latlon

convert wgs84 utm to latitude and longitude

19.14 xy2nt

project all points onto the cross section and assign them nz-coordinates

transform coordinate into N-T reference
rotate coordinate, so that cross section goes along x-axis
then x and y are n and t respectively scaled by width
N and T coordinates

19.15 xy2sn

convert cartesian to streamwise coordiantes

19.16 xy2sn.java

use java port for speed up

19.17 xy2sn_old

transform points from cartesian into streamwise coordinates

NOTE : prefer the java version, this has some problems with round off

20 linear-algebra

20.1 det2x2

2x2 matrix inverse of 2x2 matrices stacked along dim 3

20.2 det3x3

determinant of stacked 3x3 matrices

20.3 det4x4

determinant of stacked 4x4 matrices

20.4 diag2x2

diagonal of stacked 2x2 matrices

20.5 down

20.6 eig2x2

eigenvalues of stacked 2x2 matrices

21 linear-algebra/eigenvalue

21.1 eig_bisection

21.2 eig_inverse

21.3 eig_inverse_iteration

21.4 eig_power_iteration

22 linear-algebra/eigenvalue/jacobi-davidson

22.1 afun_jdm

22.2 davidson

22.3 jacobi_davidson

22.4 jacobi_davidson_qr

22.5 jacobi_davidson_qz

22.6 jacobi_davidson_simple

22.7 jdqr

```
% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal, A*V=W*R+Qschur*E, R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%   V*V=eye(j), Qschur'*V=0,
%   W=A*V, M=V'*W
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
```

```

%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%   V*V=eye(j), Qschur'*V=0, W'*W=eye(j), Qschur'*W=0
%   (A*V-tau*V)=W*R+Qschur*E, E=Qschur'*(A*V-tau*V), M=W'*V
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur, AV=A*V-
%   tau*V
%   V*V=eye(j), W'*W=eye(j), Qschur'*V=0, Qschur'*W=0,
%   (I-Qschur*Qschur')*AV=W*R, M=W'*V; R=W'*AV;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%

```

```

%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
  W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
  W=V*M; V(:,j+1)=[];R=R'*R; M=H(1:j,:)'
%===== ARNOLDI (for initializing spaces)
=====
%===== END ARNOLDI
=====

% not accurate enough M=Rw'\(M/Rv);
%===== COMPUTE SORTED JORDAN FORM
=====

% compute vectors and matrices for skew projection
% solve preconditioned system
% 0 step of bicgstab eq. 1 step of bicgstab

```

```

% Then x is a multiple of b
% HIST=[0,1];
    explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
    implicit preconditioning
% collect the updates for x in l-space
% but, do the orth to Q implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Q explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow, pause
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down
% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
r=r/rho; rho=1;
% HIST=rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
%----- compute schur form -----
A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----

```



```
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults
%% 'v'
```

22.8 jdqr_sleijpen

```
% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal,  $A*V=W*R+Qschur*E$ , R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%    $V*V=eye(j)$ ,  $Qschur'*V=0$ ,
%    $W=A*V$ ,  $M=V'*W$ 
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%    $V*V=eye(j)$ ,  $Qschur'*V=0$ ,  $W'*W=eye(j)$ ,  $Qschur'*W=0$ 
%    $(A*V-tau*V)=W*R+Qschur*E$ ,  $E=Qschur'*(A*V-tau*V)$ ,  $M=W'*V$ 
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
```

```

%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur, AV=A*V-
%   tau*V
%   V*V=eye(j), W'*W=eye(j), Qschur'*V=0, Qschur'*W=0,
%   (I-Qschur*Qschur')*AV=W*R, M=W'*V; R=W'*AV;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%

```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
W=V*H; V(:,j+1)=[];R=R'*R; M=H(1:j,:)' ;
%===== ARNOLDI (for initializing spaces)
=====
%===== END ARNOLDI
=====
% not accurate enough M=Rw' \ (M/Rv);
%===== COMPUTE SORTED JORDAN FORM
=====
% compute vectors and matrices for skew projection
% solve preconditioned system
% 0 step of bicgstab eq. 1 step of bicgstab
% Then x is a multiple of b
% HIST=[0,1];
% explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% implicit preconditioning
% collect the updates for x in l-space
% but, do the orth to Q implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Q explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow, pause
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down

```

```

% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
r=r/rho; rho=1;
% HIST=rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
%----- compute schur form -----
A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults
%% 'v'

```

22.9 jdqr_vorst

```

% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal, A*V=W*R+Qschur*E, R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%   V*V=eye(j), Qschur'*V=0,
%   W=A*V, M=V'*W
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%

```

```

%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%   V*V=eye(j), Qschur'*V=0, W'*W=eye(j), Qschur'*W=0
%   (A*V-tau*V)=W*R+Qschur*E, E=Qschur'*(A*V-tau*V), M=W'*V
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur, AV=A*V-
%   tau*V
%   V*V=eye(j), W'*W=eye(j), Qschur'*V=0, Qschur'*W=0,
%   (I-Qschur*Qschur')*AV=W*R, M=W'*V; R=W'*AV;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%

```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
  W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
  W=V*H; V(:,j+1)=[];R=R'*R; M=H(1:j,:)' ;
%===== ARNOLDI (for initializing spaces)
%=====
%===== END ARNOLDI
%=====

% not accurate enough M=Rw'\(M/Rv);
%===== COMPUTE SORTED JORDAN FORM
%=====

% accepted separation between eigenvalues:
% no preconditioning
% solve left preconditioned system
% compute vectors and matrices for skew projection
% precondition and project r
% solve preconditioned system

```

```

% no preconditioning
% solve two-sided expl. preconditioned system
% compute vectors and matrices for skew projection
% precondition and project r
% solve preconditioned system
% "unprecondition" solution
%%%% u(:,j+1)=Atilde*u(:,j)
%%%% r(:,j+1)=Atilde*r(:,j)
%----- compute schur form -----
A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults

```

22.10 jdqz

```

% Read/set parameters
% Return if eigenvalueproblem is trivial
% Initialize target, test space and interaction matrices
% V=RepGS(Qschur,V); [AV,BV]=MV(V); %%% more stability??
% W=RepGS(Zschur,eval(testspace)); %%% dangerous if sigma~lambda
% Solve the preconditioned correction equation
% Expand the subspaces and the interaction matrices
% Check for stagnation
% Solve projected eigenproblem
% Compute approximate eigenpair and residual
%=== an alternative, but less stable way of computing z =====
% display history
% save history
% check convergence
% EXPAND Schur form
% Expand preconditioned Schur matrix MinvZ=M\Zschur
% check for conjugate pair
% To detect whether another eigenpair is accurate enough
% restart if dim(V)> jmax
% Initialize target, test space and interaction matrices
% additional stabilisation. May not be needed
% V=RepGS(Zschur,V); [AV,BV]=MV(V);
% end add. stab.
% Solve the preconditioned correction equation
% expand the subspaces and the interaction matrices

```

```

% Check for stagnation
% compute approximate eigenpair
% Compute approximate eigenpair and residual
% display history
% save history
% check convergence
% expand Schur form
% ZastQ=Z'*Q0
% the final Qschur
% check for conjugate pair
% t perp Zschur, t in span(Q0,imag(q))
% To detect whether another eigenpair is accurate enough
% restart if dim(V)> jmax
%===== END JDQZ
=====
%=====

%===== PREPROCESSING
=====
%=====

%===== ARNOLDI (for initial spaces)
=====
%% then precondition=I and target = 0: apply Arnoldi with A
%===== END ARNOLDI
=====
%=====

%===== POSTPROCESSING
=====
%=====

%===== SORT QZ DECOMPOSITION INTERACTION MATRICES
=====
%===== COMPUTE SORTED JORDAN FORM
=====
%===== END JORDAN FORM
=====
%===== OUTPUT
=====
%=====

%===== UPDATE PRECONDITIONED SCHUR VECTORS
=====
%=====

%=====

```



```

%===== SOLVE CORRECTION EQUATION
=====
%=====

% solve preconditioned system
%=====

%===== LINEAR SOLVERS
=====
%=====

% [At,Bt]=MV(x); At=theta(2)*At-theta(1)*Bt;
% xtol=norm(r-At+Z*(Z'*At))/norm(r);
%===== Iterative methods
=====
% 0 step of bicgstab eq. 1 step of bicgstab
% Then x is a multiple of b
% HIST=[0,1];
% explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/srm]];
% sufficient accuracy. No need to update r,u
% implicit preconditioning
% collect the updates for x in l-space
% but, do the orth to Z implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/srm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Z explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down
% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow
%===== END SOLVE CORRECTION EQUATION
=====
%=====

```

```

%===== BASIC OPERATIONS
=====

%=====

y(1:5,1), pause
%===== COMPUTE r AND z
=====
% E*u=Q*sigma, sigma(1,1)>sigma(2,2)
%===== END computation r and z
=====

%=====

%===== Orthogonalisation
=====

%=====

%===== END Orthogonalisation
=====

%=====

%===== Sorts Schur form
=====

%=====

kappa=max(norm(A,inf)/max(norm(B,inf),1.e-12),1);
kappa=2^(round(log2(kappa)));
%----- compute the qz factorization -----
%----- scale the eigenvalues -----
%----- sort the eigenvalues -----
%----- swap the qz form -----
% repeat SwapQZ if angle is too small
%=====

%=====

% i>j, move ith eigenvalue to position j
% compute q s.t. C*q=(t(i,1)*S-s(i,1)*T)*q=0
C*P=Q*R
check whether last but one diag. elt r nonzero
C*q
% end computation q
%===== END sort QZ decomposition interaction matrices
=====

%=====

%===== INITIALIZATION
=====

%=====

```

```

%=====

% defaults          %%%% search for 'xx' in fieldnames
%% 'ma'
%% 'sch'
%% 'to'
%% 'di'
% jmin=nselect+p0 %%%% 'jmi'
% jmax=jmin+p1 %%%% 'jma'
%% 'te'
%% 'pai'
%% 'av'
%% 'tr'
%% 'fix'
%% 'ns'
%% 'ch'
%% 'lso'
%% 'ls_m'
%% 'ls_t'
%% 'ls_e'
%% 'ty'
%% 'l_'
%% 'u_'
%% 'p_'
%% 'sca'
%% 'v0'
initiation
'standard'
'harmonic'
'searchspace'
%=====

% or Operator_Form=3 or Operator_Form=5???
%=====

%===== DISPLAY FUNCTIONS
=====
%=====

%=====

%=====

%=====

```

22.11 mfunc_jdm

22.12 mgs

22.13 minres_

22.14 mv_jacobi_davidson

23 linear-algebra

23.1 first

23.2 gershgorin_circle

range of eigenvalues determined by the gershgorin circle theorem

23.3 haussdorff

haussdorf dimension

box counting: count cectangles passed through by line (covered by
polygon)

Koch snow flake 3:4 -> 1.2619

Kantor set 2:3, (4:9) -> 0.6309

quadrat 4:2, 9:3, 16:4 -> 2

23.4 ieig2x2

reconstruct matrix from eigenvalue decomposition

23.5 inv2x2

2x2 inverse of stacked matrices

23.6 inv3x3

23.7 inv4x4

inverse of stacked 4x4 matrices

23.8 kernel2matrix

24 linear-algebra/lanczos

24.1 arnoldi

24.2 arnoldi_new

24.3 eigs_lanczos_man

24.4 lanczos

24.5 lanczos_

24.6 lanczos_biorthogonal

24.7 lanczos_biorthogonal_improved

24.8 lanczos_ghep

24.9 mv_lanczos

24.10 reorthogonalise

24.11 test_lanczos

25 linear-algebra

25.1 left

left element of vector, leftmost column is extrapolated

26 linear-algebra/linear-systems

26.1 gmres_man

break on convergence

26.2 minres_recycle

27 linear-algebra

27.1 lpmean

mean of pth-power of a

27.2 lpnorm

norm of lth-power of a

27.3 matvec3

matrix-vector product of stacked matrices and vectors

27.4 max2d

maximum value and i-j index for matrix

27.5 mid

mid point between neighbouring vector elements

27.6 mpoweri

approximation of A^p , where p is not integer by quadratic interpolation

27.7 mtimes2x2

27.8 mtimes3x3

product of stacked 3x3 matrices

27.9 nannorm

norm of a vector, skips nan-values

27.10 nanshift

shift vector, but set out of range values to NaN

27.11 nl

number rows (lines) of a matrix

analogue to unix nl command

27.12 normalise

normalise a vector or the columns of a matrix

note that the columns are independently normalised, and hence not necessarily

orthogonal to each other use the gram schmidt algorithm for this (qr or orth)

27.13 normalize1

normalize columns in x to [-1,1]

27.14 normrows

27.15 orth2

make matrix A orthogonal to B

27.16 orth_man

orthogonalize the columns of A

27.17 orthogonalise

make x orthogonal to Y

27.18 padd2

padd values around a 2d (image) matrix, constant extrapolation

27.19 paddext

padd values to vactor
not suitable for noisy data
order = 0 : constant extrapolation (hold)
order = 1 : linear extrapolation

27.20 paddval1

padd values at end of x

27.21 paddval2

padd values to x

28 linear-algebra/polynomial

28.1 chebychev

chebycheff polynomials

28.2 piecewise_polynomial

evaluate piecewise polynomial

28.3 roots1

roots of linear functions

28.4 roots2

roots of quadratic function
 $c_1 x^2 + c_2 x + c_3 = 0$

28.5 roots2poly

28.6 roots3

28.7 roots4

28.8 roots_piecewise_linear

28.9 test_roots4

28.10 vanderi_1d

vandermonde matrix of an integral

29 linear-algebra

29.1 randrot

random rotation matrix

29.2 right

get right column by shifting columns to left
extrapolate rightmost column

29.3 rot2

rotation matrix from angle

29.4 rot2dir

rotation matrix from direction vector

29.5 rot3

29.6 rotR

29.7 rownorm

29.8 simmilarity_matrix

29.9 spnorm

frobenius norm

29.10 spzeros

allocate a sparze matrix of zeros

29.11 test_roots3

29.12 transform_minmax

29.13 transpose3

transpose stacked 3x3 matrices

29.14 transposeall

29.15 up

29.16 `vander_nd`

29.17 `vanderd_2d`

30 `logic`

bitwise operations on integers

30.1 `bitor_man`

bitwise OR of the numbers of the columns of A

input:
 A (positive integer)

31 `master/plot`

31.1 `attach_boundary_value`

31.2 `cartesian_polar`

31.3 `img_vargrid`

31.4 `plot_basis_functions`

31.5 `plot_convergence`

31.6 `plot_dof`

31.7 `plot_eigenbar`

31.8 `plot_error_estimation`

31.9 `plot_error_estimation_2`

31.10 `plot_error_fem`

31.11 `plot_fdm_kernel`

31.12 `plot_fdm_vs_fem`

31.13 `plot_fem_accuracy`

31.14 `plot_function_and_grid`

31.15 `plot_hat`

31.16 `plot_hydrogen_wf`

31.17 `plot_mesh`

31.18 `plot_mesh_2`

31.19 `plot_refine`

31.20 `plot_refine_3d`

31.21 `plot_runtime`

31.22 `plot_spectrum`

31.23 `plot_wavefunction`

32 `master/ported`

32.1 `assemble_2d_phi_phi`

32.2 assemble_3d_dphi_dphi

32.3 assemble_3d_phi_phi

32.4 dV_2d_

32.5 derivative_2d

32.6 derivative_3d

32.7 element_neighbour_2d

32.8 prefetch_2d_

32.9 promote_2d_3_10

32.10 promote_2d_3_15

32.11 promote_2d_3_21

32.12 `promote_2d_3_6`

32.13 `promote_3d_4_10`

32.14 `promote_3d_4_20`

32.15 `promote_3d_4_35`

32.16 `vander_2d`

32.17 `vander_3d`

33 `number-theory`

33.1 `ceiln`

`floor to leading n-digits`

33.2 `digitsb`

`number of digits with respect to specified base`

33.3 `floorn`

`floor to n-digits`

33.4 iseven

true for even numbers in X

33.5 multichoosek

all combinations of length k from set values with repetitions
c.f. nchoosek, combinations without repetition

input :
 x : scalar integer or vector of arbitrary numbers
 k : length of subsets
output :
 if x scalar : number of combinations
 if x vector : the exact combinations

33.6 nchoosek_man

vectorised binomial coefficient
 $b = N! / K! (N-K)!$

33.7 pythagorean_triple

pythagorean triple

33.8 roundn

round to n digits

34 numerical-methods

34.1 advect_analytic

35 numerical-methods/differentiation

35.1 derivative1

first derivative on variable mesh
second order accurate

35.2 derivative2

second derivative on a variable mesh

36 numerical-methods

36.1 diffuse_analytic

37 numerical-methods/finite-difference

37.1 cdiff

differences of columns of X
degree = 1 : central first order differences
degreee = 2 : central second order differences

37.2 cdiffb

differences of columns of X
degree = 1 : central first order differences
degreee = 2 : central second order differences
TODO use difference matrix function for simplicity

37.3 central_difference

37.4 cmean

single gaussian smoothing step with kernel $1/4*[1,2,1]$

37.5 cmean2

37.6 derivative_matrix_1_1d

finite difference matrix of first derivative in one dimensions
n : number of grid points
h = L/(n+1) constant step with

function [D1, d1] = derivative_matrix_1d(n,L,order)

37.7 derivative_matrix_2_1d

finite derivative matrix of second derivative in one dimension

37.8 derivative_matrix_2d

finite difference derivative matrix in two dimensions

37.9 derivative_matrix_curvilinear

derivative matrix on a curvilinear grid

37.10 derivative_matrix_curvilinear_2

derivative matrix on a two dimensional curvilinear grid
the grid has not necessarily to be orthogonal

37.11 difference_kernel

difference kernels for equispaced grids
c.f. Computing the Spectrum of the Confined Hydrogen Atom, Kastner,
2012

37.12 diffusion_matrix_2d_anisotropic

37.13 diffusion_matrix_2d_anisotropic2

37.14 distmat

distance matrix for a 2 dimensional rectangular matrix

37.15 downwind_difference

37.16 gradpde2d

objective function gradiend on two dimensional regular grid
numeric gradient for non-linear least squares optimisation
of a PDE on a rectangular grid
 $x_* = \min(f(x))$
 $f = (v(x) - v(x_*))^2 = f(x) + A \, dx + O(dx^2)$
 $a_{ij} = df_i/dx_j$

37.17 laplacian

37.18 laplacian_fdm

finite difference matrix of the laplacian
BC

37.19 lrmean

mean of the left and right element

38 numerical-methods/finite-difference/master

38.1 fdm_adaptive_grid

38.2 fdm_adaptive_refinement_old

38.3 fdm_assemble_d1_2d

38.4 fdm_assemble_d2_2d

38.5 fdm_confinement

38.6 fdm_d_vargrid

38.7 fdm_h_unstructured

38.8 fdm_hydrogen_vargrid

38.9 `fdm_mark_unstructured_2d`

38.10 `fdm_plot`

38.11 `fdm_plot_series`

38.12 `fdm_refine_2d`

38.13 `fdm_refine_3d`

38.14 `fdm_refine_unstructured_2d`

38.15 `fdm_schroedinger_2d`

38.16 `fdm_schroedinger_3d`

38.17 `relocate`

39 numerical-methods/finite-difference

39.1 mid

mid point between neighbouring vector elements

39.2 pwmid

segment end point to segment mid point transformation for regular 1
d grids

39.3 ratio

ratio of two subsequent values

39.4 steplength

step length of a vector if it were equispaced

39.5 swapoddeven

swap odd and even elements in a vector

39.6 test_derivative_matrix_2d

39.7 test_derivative_matrix_curvilinear

39.8 test_difference_kernel

39.9 upwind_difference

40 numerical-methods/finite-element

40.1 Mesh_2d.java

40.2 Tree_2d.java

40.3 assemble_1d_dphi_dphi

40.4 assemble_1d_phi_phi

40.5 assemble_2d_dphi_dphi.java

40.6 assemble_2d_phi_phi.java

40.7 assemble_3d_dphi_dphi.java

40.8 assemble_3d_phi_phi.java

40.9 `boundary_1d`

40.10 `boundary_2d`

40.11 `boundary_3d`

40.12 `check_area_2d`

40.13 `circmesh`

40.14 `cropradius`

40.15 `display_2d`

40.16 `display_3d`

40.17 `distort`

40.18 `err_2d`

40.19 `estimate_err_2d_3`

40.20 `example_1d`

40.21 `example_2d`

40.22 `explode`

40.23 `fem_2d`

40.24 `fem_2d_heuristic_mesh`

40.25 `fem_get_2d_radial`

40.26 `fem_interpolation`

40.27 `fem_plot_1d`

40.28 `fem_plot_1d_series`

40.29 fem_plot_2d

40.30 fem_plot_2d_series

40.31 fem_plot_3d

40.32 fem_plot_3d_series

40.33 fem_plot_confine_series

40.34 fem_radial

adaptive grid
constant grid

40.35 flip_2d

40.36 get_mesh_arrays

40.37 hashkey

41 numerical-methods/finite-element/int

41.1 int_1d_gauss

41.2 int_1d_gauss_1

41.3 int_1d_gauss_2

41.4 int_1d_gauss_3

41.5 int_1d_gauss_4

41.6 int_1d_gauss_5

41.7 int_1d_gauss_6

41.8 int_1d_gauss_lobatto

41.9 int_1d_gauss_n

41.10 int_1d_nc_2

41.11 int_1d_nc_3

41.12 int_1d_nc_4

41.13 int_1d_nc_5

41.14 int_1d_nc_6

41.15 int_1d_nc_7

41.16 int_1d_nc_7_hardy

41.17 int_2d_gauss_1

41.18 int_2d_gauss_12

41.19 int_2d_gauss_13

41.20 int_2d_gauss_16

41.21 int_2d_gauss_19

41.22 int_2d_gauss_25

41.23 int_2d_gauss_3

41.24 int_2d_gauss_33

41.25 int_2d_gauss_4

41.26 int_2d_gauss_6

41.27 int_2d_gauss_7

41.28 int_2d_gauss_9

41.29 int_2d_nc_10

41.30 int_2d_nc_15

41.31 int_2d_nc_21

41.32 int_2d_nc_3

41.33 int_2d_nc_6

41.34 int_3d_gauss_1

41.35 int_3d_gauss_11

41.36 int_3d_gauss_14

41.37 int_3d_gauss_15

41.38 int_3d_gauss_24

41.39 int_3d_gauss_4

41.40 int_3d_gauss_45

41.41 int_3d_gauss_5

41.42 int_3d_nc_11

41.43 int_3d_nc_4

41.44 int_3d_nc_6

41.45 int_3d_nc_8

42 numerical-methods/finite-element

42.1 interpolation_matrix

42.2 mark

42.3 mark_1d

42.4 mesh_1d_uniform

42.5 mesh_3d_uniform

42.6 mesh_interpolate

42.7 neighbour_1d

42.8 old

42.9 pdeeig_1d

42.10 pdeeig_2d

42.11 pdeeig_3d

42.12 polynomial_derivative_1d

42.13 potential_const

42.14 potential_coulomb

42.15 potential_harmonic_oscillator

42.16 project_circle

42.17 project_rectangle

42.18 promote_1d_2_3

42.19 promote_1d_2_4

42.20 promote_1d_2_5

42.21 promote_1d_2_6

42.22 quadrilaterate

42.23 recalculate_regularity_2d

42.24 refine_1d

42.25 refine_2d_21

42.26 refine_2d_structural

42.27 regularity_1d

42.28 regularity_2d

42.29 regularity_3d

```
{      T = [1 2 3 4];  
}
```

42.30 relocate_2d

42.31 test_circmesh

42.32 test_hermite

42.33 tri_assign_points

42.34 triangulation_uniform

42.35 vander_1d

van der Monde matrix

42.36 vanderd_1d

42.37 vanderi_1d

43 numerical-methods/finite-volume/@Advection

43.1 Advection

FVM treatment of the Advection equation

43.2 dot_advection

advection equation

44 numerical-methods/finite-volume/@Burgers

44.1 burgers_split

viscous Burgers' equation,
mixed analytic and numerical derivative in frequency space
by splitting scheme
 $u_t = -(0.5*u^2)_x + c*u_{xx}$

44.2 dot_burgers_fdm

viscous burgers' equation
 $u_t = -d/dx (1/2*u^2) + c d^2/dx^2 u_{xx}$

44.3 dot_burgers_fft

viscous Burgers' equation in frequency space
 $u_t + (0.5*u^2)_x = c*u_{xx}$

45 numerical-methods/finite-volume/@Finite_Volume

45.1 Finite_Volume

finite volume method for partial differential equations 1+1
dimensions
(time and space)

45.2 apply_bc

apply boundary conditions

45.3 solve

solve the the PDE by successively stepping in time
this is a trivial implmentation with constant step length
severity of diffusive error depends on dt/dx-ratio
stability depends on wave height

```
printf('Progress %2.1f%% %2.1fs\n',100*(t-Ti
(1))/(Ti(2)-Ti(1)),t_real);
```

45.4 step_split_strang

step in time, treat inhomogeneous part by Strang splitting
this scheme is not suitable for stationary solutions, for example
steady shallow water flow

45.5 step_unsplit

step in time, without splitting the inhomogeneous term

46 numerical-methods/finite-volume/@Flux_Limiter

46.1 Flux_Limiter

class of flux limiters

46.2 beam_warming

beam warming scheme

low resolution

note: works only if sign of eigenvalues point into the same
direction according to RL

46.3 fromm

fromme limiter

low res

46.4 lax_wendroff

lax wendroff scheme

second order accurate, but no tvd

this is effectively not a limiter

eq. 6.39 in randall, leveque

46.5 minmod

min-mod schock limiter

46.6 monotized_central

monotonized central flux limiter

46.7 muscl

muscl flux limiter

46.8 superbee

superbee limiter

46.9 upwind

godunov scheme
godunov, first order accurate

46.10 vanLeer

van Leer limiter

47 numerical-methods/finite-volume/@KDV

47.1 dot_kdv_fdm

korteweg de vries equation
 $u_t + (0.5*u^2)_x = c*u_{xxx}$

47.2 dot_kdv_fft

korteweg de vries equation
compute derivatives in frequency space
 $u_t + (0.5*u^2)_x = c*u_{xxx}$

47.3 kdv_split

korteweg de vries equation in frequency space,
derivative treated by splitting scheme

48 numerical-methods/finite-volume/@Reconstruct_Average_Evolve

48.1 Reconstruct_Average_Evolve

Reconstruct Average Evolve Finite Volume Method for treatment of
1+1D pdes

McCronack Scheme

err = $O(dt^2)$ + $O(dx^2)$, except as discontinuities

error:

```
h_xxx(3:end-2) = 1/dx^3*( -0.5*h(1:end-4) + h(2:end-3) - h(4:
    end-1) + 0.5*h(5:end) );
th = -1/6*dx^2*qh_.*(1 - (qh_*dt/dx).^2).*h_xxx;
```

48.2 advect_highres

single time step for the reconstruct evolve algorithm

48.3 advect_lowress

single time step

low resolution

49 numerical-methods/finite-volume

49.1 Godunov

Godunov, upwind method for systems of pdes

49.2 Lax-Friedrich

Lax-Friedrich-Method
for hyperbolic conservation laws
 $\text{err} = O(\text{dt}) + O(\text{dx})$
 $|a \text{ dt/dx}| < 1$

49.3 Measure

49.4 Roe

non linear roe solver for the SWE (randall, leveque 15.3.1)

The roe solver guarantess:
- A is diagonalisable with real eigenvalues (15.12)
- can be determined by a closed formula
- is an efficient replacement for true Rieman solver

49.5 fv_swe

wrapper for solving SWE

49.6 staggered_euler

forward euler method with staggered grid

49.7 staggered_grid

staggered grid approximation to the SWE

50 numerical-methods

50.1 grid2quad

extract rectangular elements of a structured grid
in form of an unstructured quad-mesh format

51 numerical-methods/integration

51.1 cumintL

cumulative integral from left to right

51.2 cumintR

cumulative integral from right to left

51.3 int_trapezoidal

integrate y along x with the trapezoidal rule

52 numerical-methods/interpolation/@Kriging

52.1 Kriging

class for Kriging interpolation

52.2 estimate_semivariance

estimate the parameter of the semivariance model for Kriging
interpolation
 % set up the regression matrix and solve for
 parameters

52.3 interpolate_

interpolate with Kriging method

this function may interpolate several quantities per coordinate,
using the same variogram, if the semivariance of the quantities
differs,
the user may prefer to estimate the semivariance and interpolate
each quantity
individually

Xs : source point coordinates
 Vs : value at source points
 Xt : target point coordinates
 Vt : value at target points
 E2t : squared interpolation error at target points

53 numerical-methods/interpolation/@RegularizedInterpolator

53.1 RegularizedInterpolator1

class for regularized interpolation (Thikonov) on a 1D mesh

53.2 init

initialize the interpolator with a set of sampling points

54 numerical-methods/interpolation/@RegularizedInterpolator

54.1 RegularizedInterpolator2

class for regularized interpolation on an unstructures mesh (interpolation)

54.2 init

initialize the interpolator with a set of point samples

55 numerical-methods/interpolation/@RegularizedInterpolator

55.1 RegularizedInterpolator3

class for regularized interpolation (Tikhonov) on a triangulation (unstructured mesh)

55.2 init

initialize the interpolator with a set of sampling points

56 numerical-methods/interpolation

56.1 IDW

spatial averaging by inverse distance weighting

56.2 IPoly

polynomial interpolation class

56.3 IRBM

interpolate by the radial basis function method
fprintf(1,'Progress IRBM: %d%%\n',round(100*
idx/size(Xi,1)));

56.4 ISparse

sparse interpolation class

56.5 Inn

nearest neighbour interpolation

56.6 Interpolator

interpolator super-class
fprintf(1,'Progress: %f%% %fs\n',100*
idx/size(Xt,1),t);

56.7 fixnan

fill nan-values in vector with gaps

56.8 idw1

spatial average by inverse distance weighting

56.9 idw2

spatial average by inverse distance weighting

56.10 inner2outer

linear interpolation of segment mid point to grid points at segment
ends
assumes equal grid spacing

56.11 inner2outer2

interpolate from element (segment) centres to edge points

56.12 interp1_circular

56.13 interp1_limited

interpolate values, but not beyond a certain distance
this function is idempotent, i.e. it will not extrapolate over into
gaps
exceedint the limit and thus not spuriously extend the series when
called a second time on the same data

56.14 `interp1_man`

`interpolate`

56.15 `interp1_piecewise_linear`

56.16 `interp1_save`

make interpolation save to round off errors
the matlab internal interpolation suffers from rounding errors,
which
are unacceptable when values of X and Y are large (for example UTM
coordinates)
this normalization prevents this

56.17 `interp1_slope`

quadratic interpolation returning value and derivative(s)

56.18 `interp1_smooth`

56.19 `interp1_unique`

matlab fails to interpolate, when x values are not unique
this function makes the values unique before use

56.20 `interp2_man`

nearest neighbour interpolation in two dimensions

56.21 `interp_angle`

interpolate an angle

56.22 `interp_fourier`

interpolation by the fourier method

56.23 `interp_fourier_batch`

batch interpolation by the fourier interpolation

56.24 `interp_sn`

interpolate along streamwise coordinates
This gives similar result to setting aspect ratio for sN to
infinity,
but not quite, as the input point set is not dense (scale for sN to
infinity does not work)
`sdX = sdX(sdx_);`

56.25 `interp_sn2`

interpolation in streamwise coordinates

56.26 `interp_sn3`

56.27 `interp_sn_`

56.28 limit_by_distance_1d

smooth subsequent values along a curve such that
 $v(x_0+dx) < v(x_0) + (\text{ratio}-1)*dx$
if v is the edge length in a resampled polygon, then $v_i/v_{(i+1)} <$
ratio
 $\text{ratio}^1 = \exp(a*1)$

56.29 resample1

interpolation along a parametric curve with variable step width

56.30 resample_d_min

resample a function

56.31 resample_vector

resample a track so that velocity vectors do not run into each other

56.32 test_interp1_limited

57 numerical-methods

57.1 inverse_complex

57.2 maccormack_step

57.3 minmod

58 numerical-methods/multigrid

58.1 mg_interpolate

58.2 mg_restrict

59 numerical-methods/ode/@BVPS_Characteristic

59.1 BVPS_Characteristic

solve coupled first- and second-order 1D boundary-value problems

59.2 assemble1_A

assemble the discretisation matrix for a first order ode
(mean component, zero frequency)

59.3 assemble1_A_Q

assemble the discretisation matrix for a first order ode
(mean component, zero frequency)

59.4 assemble2_A

assemble the discretisation matrix for a second-order ode
(non-zero frequency component)

59.5 assemble_AA

assemble the discretisation matrix for each channel
iteratively calls assembly for each frequency components

59.6 assemble_AAA

assemble the discretisation matrix for the entire network
iteratively calls assembly for each channel

59.7 assemble_Ic

59.8 bvp1c

59.9 check_arguments

59.10 couple_junctions

59.11 derivative

59.12 init

59.13 inner2outer_bvp2c

59.14 reconstruct

59.15 resample

59.16 solve

solve system of non-linear second order odes (in more than one variable)
as boundary value problems

odefun provides ode coefficients c:

$$c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4)$$
$$c_1 y'' + c_2 y' + c_3 y + c_4 = c_4$$

subject to the boundary conditions

bcfun provides v and p and optionally q, so that:

$$b_1 y + b_2 y' = f$$

$$q(x,1)*(p(x,1) y_l(x) + p(x,2) y_l'(x) + q(x,2)*(p(x,1) y_r(x) + p(x,2) y_r'(x) = v(x)$$

where q weighs the waves travelling from left to right and right to left (default [1 1])

59.17 test_assemble1_A

59.18 test_assemble2_A

60 numerical-methods/ode/@Time_Stepper

60.1 Time_Stepper

60.2 solve

61 numerical-methods/ode

61.1 bvp2fdm

solve system of non-linear second order odes (in more than one variable)

as boundary value problems by the finite difference method

odefun provides ode coefficients c:

$$c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4)$$
$$c_1 y'' + c_2 y' + c_3 y + c_4 = 0$$

subject to the boundary conditions

bcfun provides v and p and optionally q, so that:

$$b_1 y + b_2 y' = f$$

$$q(x,1) * (p(x,1) y_l(x) + p(x,2) y_l'(x) \\ + q(x,2) * (p(x,1) y_r(x) + p(x,2) y_r'(x) = v(x)$$

where q weighs the waves travelling from left to right and right to left (default [1 1])

61.2 bvp2wavetrain

solve second order boundary value problem by repeated integration

61.3 bvp2wavetwopass

two pass solution for the linearised wave equation

solve first for the wave number k, and then for y

61.4 ivp_euler_forward

solve initial value problem by the euler forward method

61.5 ivp_euler_forward2

61.6 ivprk2

solve initial value problem by the two step runge kutta method

61.7 ode2_matrix

transformation matrix of second order ode
to left and right going wave

```
c = odefun(x)
c1 y'' + c2' y + c3 y == 0
y = y_p + y_m, left and right going wave
d/dx [y_p, y_m] = A*[y_m, y_p]
```

61.8 ode2characteristic

second order odes
transmitted and reflected wave

61.9 step_trapezoidal

single trapezoidal step

61.10 test_bvp2

62 numerical-methods/optimisation

62.1 aitken_iteration

62.2 anderson_iteration

62.3 armijo_stopping_criterion

armijo stopping criterion for optimizations

62.4 astar

astar path finding algorithm

62.5 binsearch

binary search on a line

62.6 bisection

bisection

62.7 box1

test objective function for optimisation routines

62.8 box2

62.9 cauchy

62.10 cauchy2

solve non-linear system by cuachy's method
slower than quadratic optimisation, but does not require a hessian
fun : objective function, returns
 f : scalar, objective function value
 g : nx1, gradient
x : nx1, initial position
opt : options

62.11 directional_derivative

directional (projected) derivative
d : derivative, highest first
p : series expansion around x0

62.12 dud

optimization by the dud algorithm

62.13 extreme3

extract maxima by quadratic approximation from sampled function val
(t)
intended to be called after [mval, mid] = max(val) for refinement
of
location and maximum

input
t : sampling time (uniformly spaced)
v : values at sampling times
output:
tdx : index where extremum should be computed
t0 : location of the extremum
val0 : value of extremum

$v'(dt0) = 0$ and $v''(dt0)$ determines type of extremum

62.14 extreme_quadratic

62.15 ftest

62.16 fzero_bisect

62.17 `fzero_newton`

62.18 `grad`

numerical gradient

62.19 `hessian`

numerical hessian

62.20 `hessian_from_gradient`

numerical hessian from gradient

62.21 `hessian_projected`

numerical hessian projected to one dimension

62.22 `line_search`

bisection routine

62.23 `line_search2`

bisection method

`fun` : objective funct
`x0` : start value
`f0` : objective function value at `x0`
`g` : gradient at `x0`
`p` : search direction from `x0` (`p = g` for steepest descend)
`h` : initial step length (default 1)
`lb` : lower bound for `x`
`up` : upper bound for `x`

62.24 line_search_polynomial

```
polynomial line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

62.25 line_search_polynomial2

```
cubic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

62.26 line_search_quadratic

```
quadratic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

62.27 line_search_quadratic2

```
quadratic line search
```

62.28 line_search_wolfe

line search by wolfe method
c.f.: OPTIMIZATION THEORY AND METHODS - Nonlinear Programming, Sun,
Yuan

62.29 ls_bgfs

least squares by the bgfs method

62.30 ls_broyden

least squares by the broyden method
for rectangular / non symmetric systems
Numerical Optimization nokedal
Practical Methods of Optimization fletcher
c.f. gerber 1981
c.f. fletcher 1978 (more advanced, not used here)
c.f. Kelley 1999 ch. 4

BGFS:
Broyden 1965
Fletcher 1970
Goldfarb 1970
Shanno 1970

62.31 ls_generalized_secant

least squares by the secant method
Barnes, 1965
Wolfe, 1959
Fletcher 1980, 6.3
seber 2003
gerber

62.32 nlcg

non-linear conjugate gradient
input:
x : nx1 start vectort

opt : struct options
fdx : gradient constraint

62.33 nlls

non-linear least squares

62.34 picard

picard iteration

62.35 poly_extrema

extrema of a polynomial

62.36 quadratic_function

evaluate quadratic function in higher dimensions

62.37 quadratic_programming

optimize by quadratic programming

62.38 quadratic_step

single step of the quadratic programming

62.39 rosenbrock

rosenbrock test function

62.40 `sqrt_heron`

Heron's method for the square root

62.41 `test_directional_derivative`

62.42 `test_dud`

62.43 `test_fzero_newton`

62.44 `test_line_search_quadratic2`

62.45 `test_ls_generalized_secant`

62.46 `test_nlcg_6_order`

62.47 `test_nlls`

```
f = w'*(p*abs(x-1).^4) + w'*(1-p)*abs(x-1).^2;
```

63 numerical-methods/pde

63.1 `laplacian2d_fundamental_solution`

64 numerical-methods/piecewise-polynomials

64.1 Hermite1

hermite polynomial interpolation in 1d

64.2 hp2_fit

fit a hermite polynomial
coefficients are derivative free
x0 : left point of first segment
x1 : right point of last segment
n : number of segments
x : sample x-value
val : sample y-value
c : coefficients (values at points, no derivatives)

64.3 hp2_predict

prediction with pw hermite polynomial
c are values at support points

64.4 hp_predict

predict with piecewise hermite polynomial

64.5 hp_regress

fit piecewise hermite polynomial
coefficients are values and derivatives

64.6 lp_count

lagrangian basis for interpolation
count number of valid samples

64.7 lp_predict

lagrangian basis piecewise interpolation, predictor

64.8 lp_regress

64.9 lp_regress_

65 numerical-methods

65.1 test_adams_bashforth

66 patterns

66.1 band_pattern

66.2 hexagonal_pattern

67 regression/@PolyOLS

67.1 PolyOLS

class for polynomial least squares

67.2 coefftest

67.3 detrend

detrending by polynomial regression

67.4 fit

fit a polynomial function
like polyfit, but returns parameter error estimates
TODO automatically activate scaleflag

67.5 fit_

fit a polynomial function

67.6 predict

predict polynomial function values

67.7 predict_

67.8 slope

slope by linear regression

68 regression/@PowerLS

68.1 PowerLS

class for power law regression

68.2 fit

fit a power law
like polyfit, but returns parameter error estimates

68.3 predict

predict with power law
S2 = diag((A*obj.C)*A');
L = Y - S;
U = Y + S;

68.4 predict_

69 regression/@Theil

69.1 Theil

Kendal-Theil-Sen robust regression

69.2 detrend

linear detrending of a set of samples by the Theil-Senn Slope

69.3 fit

fit slope and intercept to a set of sample with the Theil-Sen
method

c : confidence interval $c = 2*ns*normcdf(1)$ for ns-sigma
intervals
param : itercept and slope
P : confidence interval

69.4 predict

predict values and confidence intervals with the Theil-Sen method

69.5 slope

fit the slope with the Theil-Sen method

70 regression

linear and non-linear regression

70.1 Theil_Multivariate

extension of the Theil-Senn regression to higher dimensions by means of the Gauss-Seidel iteration

70.2 areg

regression using the pth-fraction of samples with smallest residual

70.3 ginireg

gini regression

70.4 hesssimplereg

hessian, gradient and objective function value of the simple
regression
 $\text{rhs} = p(1) + p(2) x + \text{eps}$

70.5 l1lin

solve $\|Ax - b\|_{L1}$ by means of linear programming

70.6 lsq_sparam

parameter covariance of the least squares regression

fun : model function for prediction
b : sample values
 $f(p) = b$
p : parameter at point of evaluation (preferably optimum)

70.7 polyfitd

fit a polynomial of order n to a set of sampled values and sampled values of the derivative

x0 must contain at least for conditioning as otherwise the intercept cannot be determined

70.8 regression_method_of_moments

fit linear function $\|a \ b \ x = y\|_{L2}$ by the method of moments
 $y + \text{eps} = \text{alpha} + \text{beta} * x$

70.9 robustlinreg

fit a linear function by splitting the x-values at their median
 $(\text{med}(y_{\text{left}}) - \text{med}(y_{\text{right}})) / (\text{med}(x_{\text{left}}) - \text{med}(x_{\text{right}}))$
this approach performs poorly compared to the theil-senn operator

70.10 theil2

Theil senn-estimator for two dimensions (glm)

70.11 theil_generalised

generalization of the Theil-Senn operator to higher dimensions,
for arbitrary functions such as polynomials and multivariate
regression
either higher order polynomials or glm
c.f. "On theil's fitting method", Pegoraro, 1991

70.12 total_least_squares

total least squares

70.13 weighted_median_regression

weighted median regression
c.f. Scholz, 1978

71 set-theory

71.1 issubset

test if set B is subset of A in $O(n)$ -runtime

A : first set
B : second set
P : set of primes (auxiliary)

72 mathematics

mathematical functions of various kind

72.1 shuffle_index

73 signal-processing

73.1 acfar1

Autocorrelation function of the finite AR1 process

```
a_k = 1/(n-k)sum x_ix_{i+k} + (x_i + x_{i+k})mu + mu^2
      = r^k + 1/n sum_{i=j} x_i^2 + 1/n
      pause
```

73.2 acfar1_2

autocorrelation of the ar1 process

73.3 acfar2

impulse response of the ar2 process

73.4 acfar2_2

autocorrelation of the ar2 process
 $X_i + a_1 X_{i-1} + a_2 X_{i-2} = 0$

73.5 ar1_cutoff_frequency

73.6 ar1_effective_sample_size

effective sample size correction for autocorrelated series

73.7 ar1_mse_mu_single_sample

standard error of a single sample of an ar1 correlated process

73.8 ar1_mse_pop

variance of the population mean of a single realisation around zero

$E[(\mu_N - 0)^2] = E[\mu_N^2]$

73.9 ar1_mse_range

mean standard error of the mean of a range of values taken from an ar1 process

73.10 ar1_spectrum

spectrum of the ar1 process

73.11 ar1_to_tikhonov

convert ar1 correlation to tikhonovs lambda

73.12 ar1_var_factor

variance correction factor for an autocorrelated finite process
n : [1 .. inf] population size
m : [1 .. n] samples size
rho : [-1 < rho < 1 (for convergence)] correlation of samples

73.13 ar1_var_factor_

variance of an autocorrelated finite process

73.14 ar1_var_range2

variance of sub sample starting at the end of the series
from the finite length first order autocorrelated process
$$s2 = 1/m^2 \sum_i^m \sum_j^m \rho^{-|i-j|}$$

73.15 ar1delay

approximate acf by the ar1 process
acf: autocovariance or autocorrerlation function
nf : skip first samples (for mixed geometric-arithmetic series (ARMA))

73.16 ar1delay_old

autocorrelation of the residual

73.17 ar2_acf2c

determine coefficients of the ar2 process from the first two lags
of the
autocorrelation function

73.18 ar2conv

coefficients of the ar2 process determined from the two leading
correlations
of the acf [1,r1,r2,...]

73.19 ar2dof

effective samples size for the ar2 process

73.20 ar2param

ar2 parameter estimation from first two terms of acf
acf = [1 a1 a2 ...]

73.21 asymwin

creates asymmetrical filter windows
filter will always have negative weights

74 signal-processing/autocorr

74.1 autocorr2

74.2 autocorr_bandpass

74.3 autocorr_brownian_phase

74.4 autocorr_brownian_phase_2d

74.5 autocorr_brownian_phase_across

74.6 autocorr_decay_rate

estimate exponential decay of the autocorrelation

74.7 autocorr_effective_sample_size

effective sample size from acf

74.8 autocorr_fft

estimate sample autocorrelation function

74.9 autocorr_genton

autocorrelation function

74.10 autocorr_highpass

74.11 autocorr_lowpass

74.12 autocorr_periodic_additive_noise

74.13 autocorr_periodic_windowed

74.14 autocorr_radial

75 signal-processing

75.1 average_wave_shape

extract waves with varying length from a wave train and and average
their shape

75.2 bandpass

bandpass filter

75.3 bandpass1d

75.4 bandpass1d_fft

filter input vector with a spatial (two-sided) bandpass in fourier space

75.5 bandpass1d_implicit

75.6 bandpass2

bandpass filter

75.7 bandpass2d

75.8 bandpass2d_2

75.9 bandpass2d_fft

75.10 bandpass2d_ideal

75.11 bandpass2d_implicit

75.12 bandpass2d_iso

75.13 bandpass_arg

determine correlation coefficient from frequency of mode for the symmetric

75.14 bandpass_f0_to_rho

correlation coefficient for the pth-order symmetric bandpass filter with maximum at f0 (when rho_lp = rho_hp)

75.15 bandpass_max

75.16 bandpass_max2

75.17 bartlett

Effective sample size factor for bartlett window
c.f. thiebaux
c.f spectral analysis-jenkins, eq. (6.3.27)
c = acf
note: results seams always to be 1 tac too low
T : reduction factor for dof
for ar1 with $a = \rho^k = \exp(-k/L)$, $T = 2L$

75.18 bin1d

bin values of v sampled at x into bins bounded by "edges"
apply function v to it

75.19 bin2d

bin values of V sampled at X and Y into the grid structured grid ex
,ey
apply function func to all values in the bin
func = mean : default
func = sum : non-normalized frequency histogram in 2D

75.20 binormrnd

generate two correlated normally distributed vectors

75.21 coherence

75.22 conv1_man

convolutions with padding

75.23 conv2_man

convolution in 2d

75.24 conv2z

75.25 conv30

convolve with rectangular window of lenght n
circular boundaries

75.26 conv_

convolution of a with b

75.27 conv_centered

convolve x with filter window f
when length of f is even, this guarantees a symmetric result (no
off by one
displacement) by making the length of f odd at first

75.28 convz

75.29 cosexpdelay

75.30 csmooth

smooth recursively with $[1,2,1]/4$ kernel
function `x = csmooth(x,n,p,circ)`

75.31 daniell_window

Daniell window for smoothing the power spectrum
c.f. Daniell 1946
Bloomfield 2000
meko 2015

75.32 danielle_window

danielle fourier window

75.33 db2neper

convert decibel to neper

75.34 db2power

power ratio from db

75.35 derive_bandpass_normalization_and_zeros

75.36 derive_danielle_weight

75.37 derive_limit_0_acfar

75.38 detect_peak

detect peaks in a vector
requires function value to fall to p*max before new value is
allowed

75.39 determine_phase_shift

75.40 determine_phase_shift1

average phase and phase shift per time step of a train of waves

75.41 digital_low_pass_filter

design coefficients of a low pass filter with specified cut of
frequency
and sampling period
analogue low pass with pole at $s = -\omega_c = 1/\tau = 1/RC$
 $H_a = \tau / (\tau + s) = 1 / (1 + \omega_c s)$

75.42 `doublesum_ij`

double sum of r^i

75.43 `effective_sample_size_to_ar1`

convert effective sample size to ar1 correlation

75.44 `flt_hodges_lehman`

75.45 `filter1`

filter along one dimension

75.46 `filter2`

filter columns of x (matlab does only support vector input)

75.47 `filter_`

invalidate values that exceed n -times the robust standard deviation

75.48 `filter_r_to_f0`

75.49 `filter_rho_to_f0`

75.50 `filter_twosided`

75.51 filteriir

filter adcp t-n data over time

v : nz,nt : values to be filtered
H : nt,1 : depth of ensemble
last : nt,1 : last bin above bottom that can be sampled without
side lobe interference
nf : scalar : number of reweighted iterations

when samples
- distance to bed is reference (advantageous for near-bed suspended
transport)
TODO for wash load: distance to surface is more relevant
interpolate depending on z

when depth changes, neighbouring indices do not correspond to same
relative position in the water column
relative position in the column (s-coordinate) smoothes values
near the bed: absolute distance to bed is chosen
near surface: absolute distance to surface is chosen
-> cubic transformation of index

faster and avoid aliasing (smoothing along z)
resample ensemble to same number of bins in S -> filter ->
resample back
use nonlinear transform z-s coordinates
-> resampling has to be local (Hi -> H-filtered)

filtered profile coordinates to sample coordinates
zf -> zi (special transform)
corresponding indices and fractions
filtration step (update of hf and vf)
sample coordinates to updated profile coordinates
(the inverse step is actually not necessary)
write filtered value

75.52 filterp

75.53 filterp1

fir filter with some fancy extras

75.54 filterstd

75.55 firls_man

design finite impulse response filter by the least squares method

75.56 fit_spectral_density

fit spectral densities

75.57 fit_spectral_density_2d

fit spectral densities

75.58 fit_spectral_density_radial

fit spectral densities

75.59 flattopwin

the flat top window

75.60 frequency_response_boxcar

frequency response of a boxcar filter

75.61 freqz_boxcar

frequency response of a boxcar filter

75.62 `gaussfilt1`

filter data series with a gaussian window

75.63 `hanchangewin`

hanning window for change point detection

75.64 `hanchangewin2`

nanning window for chage point detection

75.65 `hanwin`

hanning filter window

75.66 `hanwin_`

hanning filter window

75.67 `high_pass_1d_simple`

75.68 `highpass`

high pass filter

75.69 `highpass1d_fft_cos`

filter the input vector with a cosine-shaped highpass in frequency space

75.70 `highpass1d_implicit`

75.71 `highpass2d_fft`

75.72 `highpass2d_ideal`

75.73 `highpass2d_implicit`

75.74 `highpass_arg`

75.75 `highpass_fc_to_rho`

75.76 `jackknife_block`

75.77 `kaiserwin`

`kaiser filter window`

75.78 `kalman`

`Kalman filter`

75.79 lanczoswin

Lanczos window

75.80 last

lake tail, but for matrices

75.81 lowpass

low pass filter

75.82 lowpass1d_fft

75.83 lowpass1d_implicit

75.84 lowpass2

design low pass filter with cutoff-frequency f1

75.85 lowpass2d_2

75.86 lowpass2d_anisotropic

75.87 lowpass2d_fft

75.88 `lowpass2d_ideal`

75.89 `lowpass2d_implicit`

75.90 `lowpass_arg`

75.91 `lowpass_fc_to_rho`

75.92 `lowpass_iir`

`iir-low pass`

75.93 `lowpass_iir_symmetric`

`two-sided iir low pass filter (for symmetry)`

75.94 `lowpassfilter2`

`low-pass filter of data`

75.95 `maxfilt1`

75.96 `meanfilt1`

`moving average filter with special treatment of the boundaries`

75.97 meanfilt2

filter with a rectangular window along both dimensions

75.98 medfilt1_man

moving median filter, supports columnwise operation

75.99 medfilt1_man2

moving median filter with special treatment of boundaries

75.100 medfilt1_padded

median filter with padding

75.101 medfilt1_reduced

median filter with padding

75.102 mid_term_single_sample

variance of single sample, mid term

75.103 minfilt1

75.104 mu2ar1

error variance of the mean of the finite length ar1 process

$(\mu)^2 = (\sum \text{epsi})^2 = \sum_i \sum_j \text{epsi}_i \text{epsi}_j = \sum_{ii}(\text{rho},n)/n^2$
this has the limit s^2 for $\text{rho} \rightarrow 1$

75.105 **mysmooth**

75.106 **nanautocorr**

autocorrelation with nan-values

75.107 **nanmedfilt1**

medfilt1, skipping nans

75.108 **neper2db**

convert neper to db

75.109 **oscillator_noisy**

75.110 **peaks_man**

peaks of a periodogram

75.111 **periodogram**

compute the normalized periodogram

75.112 **periodogram_2d**

compute the normalized periodogram in two dimensions

75.113 **periodogram_annular**

75.114 `periodogram_bartlett`

estimate the spectral density nonparametrically with Bartlett's method

75.115 `periodogram_bootstrap`

75.116 `periodogram_confidence_interval`

confidence interval for periodogram values

75.117 `periodogram_median`

75.118 `periodogram_p_value`

75.119 `periodogram_qq`

qq-plot of a spectral density estimate by smoothing against the expected beta-density

75.120 `periodogram_quantiles`

quantiles of a periodogram

75.121 `periodogram_radial`

75.122 periodogram_std

standard deviation of a periodogram

75.123 periodogram_test_periodicity

test a periodogram for hidden periodic frequency components

```
function [p, ratio, maxShat, mdx, fdx, S] = periodogram_test_periodicity  
(fx, Shat, nf, fmin, fmax, S, mode)
```

input:

```
    fx : frequengcies  
    Shat : corresponding periodogram values  
    nf  : number of bins to test for periodicity, ignored when S  
          is given  
    fmin, fmax : frequency range limits to test  
    S      : exact (a priori known theoretical spectral density,  
              must not be estimated from the periodogram)  
    mode : automatically set to "exact", when S given  
           inclusive : estimate density by smoothing including the  
                      central bin  
           exclusive : estimate density by smoothing excluding the  
                      central bin  
    note: inclusive and exclusive lead to different distribution  
          but identical p-values
```

75.124 periodogram_test_periodicity_2d

test a periodogram for hidden periodic frequency components

```
function [p, ratio, maxShat, mdx, fdx, S] = periodogram_test_periodicity  
(fx, Shat, nf, fmin, fmax, S, mode)
```

input:

```
    fx : frequengcies  
    Shat : corresponding periodogram values  
    nf  : number of bins to test for periodicity, ignored when S  
          is given  
    fmin, fmax : frequency range limits to test  
    S      : exact (a priori known theoretical spectral density,  
              must not be estimated from the periodogram)  
    mode : automatically set to "exact", when S given  
           inclusive : estimate density by smoothing including the  
                      central bin
```

exclusive : estimate density by smoothing excluding the
central bin
note: inclusive and exclusive lead to different distribution
but identical p-values

75.125 periodogram_test_stationarity

test a periodogram for stationarity
note : the method works, but is of little practical use,
as it requires about 50 periods and a small dx to detect a
frequency change by a factor of 2

75.126 periodogram_welsh

75.127 polyfilt1

polynomial filter,
can be achieved by iteratively processing the data with
a mean (zero-order) filter

75.128 qmedfilt1

medfilt1, after fitting a quadratic polynomial

75.129 radial_window

radial filter window in the 2d-frequency domain

75.130 randar1

generate random ar1 process
e1 = randar1(sigma,p,n,m)

75.131 randar1_dual

draw random variables of two correlated ar1 processes

75.132 randar2

generate ar2 process

75.133 randarp

randomly generate the instance of an ar-p process

75.134 range_window

range of values within a certain range of indices (window)

75.135 rectwin

rectangular window

75.136 recursive_sum

75.137 select_range

75.138 smooth1d_parametric

smooth position of $p_0=x_0, y_0$ between $p_1=x_1, y_1$ and $p_2=x_2, y_2$,
so that distance to p_1 and p_2 becomes equal
and the chord length remains the same

75.139 smooth2

smooth vectos of X

75.140 smooth_man

75.141 smooth_parametric

smooth a parametric function given in x-y coordinates
matvec2x2(R,[dxc;dyc])

75.142 smooth_parametric2

parametrically smooth the curve

75.143 smooth_with_splines

75.144 smoothfft

filter with fast fourier transform

76 signal-processing/spectral-density

76.1 spectral_density_ar2

76.2 spectral_density_area

integrate the spectral density

76.3 spectral_density_bandpass2d_ideal

76.4 spectral_density_bandpass_2d

76.5 spectral_density_bandpass_2d_scale

76.6 spectral_density_bandpass_2d_scale_old

76.7 spectral_density_bandpass_continuous

S : spectral density of the bandpass filter in continuous space
limit case of the discrete bandpass for $dx \rightarrow 0$

```
function [S,IS] = spectral_density_bandpass_continuous(fx,fc,order)
f      : frequency (abszissa)
fc     : central frequency, location of maximum on abszissa
order  : number of times filter is applied iteratively, not
         necessarily integer
```

76.8 spectral_density_bandpass_continuous_max

maximum of the bandpass spectral density

76.9 spectral_density_bandpass_continuous_max2par

transform mode (maxima) of the bandpass spectral density into the
parameter
of the underlying distribution

76.10 spectral_density_bandpass_continuous_scale

normalization scale of the spatial bandpass density

76.11 spectral_density_bandpass_discrete

spectral density of the discrete spatial (two-sided) bandpass filter

76.12 spectral_density_brownian_phase

spectral density of a fourier series where the phase undergoes brownian motion
with standard deviation s per unit distance

76.13 spectral_density_brownian_phase_2d

76.14 spectral_density_brownian_phase_across

76.15 spectral_density_brownian_phase_mode

mode (maximum) of the spectral density of the fourier series with brownian phase

76.16 spectral_density_brownian_phase_mode2par

transform mode to parameters of the brownian phase spectral density

76.17 spectral_density_brownian_phase_scale

normalization scale of the brownian phase spectral density

76.18 `spectral_density_estimate_2d`

76.19 `spectral_density_flat`

flat spectral density of a random vector with iid elements

76.20 `spectral_density_highpass`

76.21 `spectral_density_highpass2d_ideal`

76.22 `spectral_density_highpass_2d`

76.23 `spectral_density_highpass_cos`

consine shaped spectral density of a highpass filter

76.24 `spectral_density_lorentzian`

lorentzian spectral density

76.25 `spectral_density_lorentzian_max`

mode (maximum) of the lorentzian spectral density

76.26 `spectral_density_lorentzian_max2par`

transform maximum of the lorentzian spectral density to its
distribution parameters

76.27 `spectral_density_lorentzian_scale`

normalization scale of the lorentzian spectral density

76.28 `spectral_density_lowpass`

76.29 `spectral_density_lowpass2d_ideal`

76.30 `spectral_density_lowpass_2d`

76.31 `spectral_density_lowpass_one_sided`

76.32 `spectral_density_periodic_additive_noise`

76.33 `spectral_density_wperiodic`

77 `signal-processing`

77.1 `spectrogram`

`spectrogram`

77.2 `std_window`

moving block standard deviation

77.3 sum_i_lag

sum of ar1 matrix with lag
 $\sum_{i=1}^n \rho^{|i-k|}$

77.4 sum_ii

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^n \rho^{|i-j|}$
this is for the variance, take square root for the standard
deviation factor

77.5 sum_ii_

77.6 sum_ij

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^m \rho^{|i-j|}$

77.7 sum_ij_

77.8 sum_ij_partial_

77.9 sum_multivar

sum of matrix entries of bivariate ar1 process

77.10 test_acfar1

77.11 test_acfar1_2

77.12 test_acfar1_3

77.13 test_acfar1_4

77.14 test_acfar2

77.15 test_ar1_var_factor

77.16 test_ar1_var_factor_2

77.17 test_ar1_var_mu_single_sample

77.18 test_ar1_var_pop

77.19 test_ar1_var_pop_1

77.20 test_ar1delay

77.21 test_bivariate_covariance_term

77.22 test_convexity

77.23 test_lanczoswin

77.24 test_madcorr

77.25 test_randar1

77.26 test_randar1_multivariate

77.27 test_randar2

77.28 test_sum_ij

77.29 test_sum_multivar

77.30 test_trifilt1

77.31 `test_wautocorr`

77.32 `test_wavelet_transform`

77.33 `test_wordfilt`

77.34 `test_xar1_mid_term`

77.35 `tikhonov_to_ar1`

convert coefficient of the tikhonov regularization to correlatioon
of the ar1 process

77.36 `trapwin`

trapezoidal filter window

77.37 `trifilt1`

filter with triangular window
trifilt1 is ident to twice applying rectfilt1 (meanfilt1) with half
the domain size
note : inifinitely many convolution yield a gaussian

77.38 `trifilt2`

filter with a triangular window along both dimensions

77.39 triwin

triangular filter window

77.40 triwin2

triangular filter window

77.41 varar1

error variance of a single sample of a finite length ar1 process with respect to the mean, averaged over the population

77.42 welch_spectrogram

welch spectrogram

77.43 wfilt

filter with window

77.44 winbandpass

filter with bandpass

77.45 window2d

77.46 window_make_odd

77.47 winfilt0

filter with window

77.48 winlength

window length for desired cutoff frequency
power at f_c is halved
 $H(wf) = 1/\sqrt{2} H(f)$
if the filter window were used as a low pass filter
note: the user should prefer a windowed ideal low pass filter
TODO, relate this to DOF

77.49 wmeanfilt

mean filter with window

77.50 wmedfilt

median filter with window

77.51 wordfilt

weighted order filter

77.52 wordfilt_edgeworth

weighed order filter

77.53 wrapphase

77.54 xar1

77.55 `xcorr_man`

cross correlation of two sampled ar1 processes

78 `sorting`

78.1 `sort2`

sort two numbers

78.2 `sort2d`

sort elements of matrix in X
returns row and column index of sorted values

79 `special-functions`

79.1 `bessel_sphere`

spherical Bessel function of the first kind

79.2 `digamma_man`

79.3 `exp10`

79.4 `hankel_sphere`

spherical Hankel function for the far field (incident plane wave)
first kind

79.5 hermite

probabilistic's hermite polynomial by recurrence relation

input :
n : order
x : value

output:
f : $H_n(x)$
df : $d/dx H_n(x)$

79.6 legendre_man

legendre polynomials

79.7 neumann_sphere

spherical Neumann function
Bessel function of the second kind

80 statistics

80.1 atan_s2

stadard deviation of the arcus tangens by means of taylor expansion

80.2 beta_kurt

80.3 beta_mean

80.4 beta_mode_to_parameter

transform modes (mean and sd) to paramets of the beta function

80.5 `beta_skew`

80.6 `beta_std`

80.7 `chi2_kurt`

80.8 `chi2_mean`

80.9 `chi2_skew`

80.10 `chi2_std`

80.11 `coefficient_of_determination`

80.12 `conditional_expectation_normal`

80.13 `correlation_confidence_pearson`

confidence intervals of the correlation coefficient
c.f. Fischer 1921

81 statistics/distributions

81.1 PDF

class for quasi-distributions from a set of sampling points

81.2 binorm_separation_coefficient

separation coefficient of a bimodal normal distribution

81.3 binormcdf

bio-modal gaussian distribution

81.4 binormfit

fit sum of to normal distribution to a histogram

81.5 binormpdf

81.6 edgeworth_cdf

edgeworth expansion of an unknown cumulative distribution
with mean mu, standard deviation sigma, and third and fourth
cumulants
c.f. Rao 2010

81.7 edgeworth_pdf

probability density of and unknown distribution
with mean mu, standard deviation sigma, and third and fourth
cumulants
c.f. Rao 2010

81.8 gam_moment2param

81.9 logn_mean

81.10 logn_mode

mode (maximum) of the log-normal density

81.11 logn_moment2param

transform the mode (mu,sd) to parameters of the log normal distribution

81.12 logn_param2moment

transform parameters to mode (mu, sd) for the log normal distribution

81.13 logn_std

81.14 lognpdf_

log normal distribution called by modes rather than parameters

81.15 pdfsample

pdf from sample distribution
Note: better use kernel density estimates

81.16 t2cdf

Hotelling's T-squared cumulative distribution

81.17 t2inv

inverse of Hotelling's T-squared cumulative distribution

82 statistics

82.1 example_standard_error_of_sample_quantiles

82.2 f_var_finite

reduction of variance when sampling from a finite population
without replacement

82.3 fisher_mean

82.4 fisher_std

82.5 gam_mean

82.6 gam_std

82.7 `gamma_mode_to_parameter`

transform modes (mu,sd) to parameters of the gamma distribution

82.8 `gamma_stirling`

82.9 `gaussfit3`

82.10 `gaussfit_quantile`

82.11 `geoserr`

82.12 `geostd`

82.13 `hodges_lehmann_correlation`

`hodges_lehmann` correlatoon coefficient
c.f. Shamos 1976
c.f. Bickel and Lehmann 1976
c.f. rousseeuw 1993
c.f. Shevlyakov 2011

82.14 hodges_lehmann_dispersion

dispersion determined by the hodges lehman method
asymptotic efficiency of dispersion estimates:
standard deviation: $E(s - \hat{s})/s = 2/\sqrt{2n} \sim 0.707/\sqrt{n}$
(100%)
hodges lehmann dispersion $E(s - \hat{s})/s = (\pi/3)^2 / (\sqrt{2n}) \sim$
 $0.775/\sqrt{n}$ (91%)
mad $E(s - \hat{s})/s \sim 1.17 s/\sqrt{n}$
(60%)
c.f. Shamos 1976
c.f. Bickel and Lehmann 1976
c.f. rousseeuw 1993
nb: rousseeuw uses the 25th percentile, which is more efficient for
small sample sizes

83 statistics/information-theory

83.1 akaike_information_criterion

akaike information criterion

serr : rmse of model prediction
n : effective sample size
k : number of parameters

c.f. akaike (1974)
c.f. sugiura 1978

83.2 bayesian_information_criterion

bayesian information criterion

84 statistics

84.1 kurtncdf

84.2 kurtnpdf

84.3 kurtosis_bias_corrected

bias corrected kurtosis

84.4 limit

limit a by lower and upper bound

84.5 logfactorial

approximate log of the factorial

84.6 loglogpdf

84.7 lognfit_quantile

84.8 logskewcdf

84.9 logskewpdf

85 statistics/logu

85.1 lambertw_numeric

lambert-w function

85.2 logtrialtcdf

pdf of a logarithmic triangular distribution

85.3 logtrialtinv

inverse of the logarithmic triangular distribution

$$\begin{aligned} &= (d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \\ &\quad \log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) \\ &\quad - \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a) - 1/ \\ &\quad \log(a)}) c)^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) \\ &\quad) + d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) \\ &\quad + b \log(c) - d F \log(a) \log(c)) / (\log(a) - \log(b))) \\ x = &(d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \\ &\log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) \\ &- \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a) - 1/\log \\ &(a)}) c)^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) + \\ &d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) + b \\ &\log(c) - d F \log(a) \log(c)) / (\log(a) - \log(b))) \end{aligned}$$

85.4 logtrialtmean

mean of the logarithmic triangular distribution

85.5 logtrialtpdf

density of the logarithmic triangular distribution

85.6 logtrialtrnd

85.7 logtricdf

cumulative distribution of the logarithmic triangular distribution

85.8 logtriinv

inverse of the logarithmic triangular distribution

85.9 logtrimean

mean of the logarithmic triangular distribution

85.10 logtripdf

probability density of the logarithmic triangular distribution

85.11 logtirnd

85.12 logucdf

probability density of the logarithmic uniform distribution

85.13 logucm

central moments of the log-uniform distribution

85.14 loguinv

inverse of the log-uniform distribution

85.15 logumean

mean of the log-uniform distribution

85.16 logupdf

pdf of the log uniform distribution

85.17 logurnd

random numbers following a log-uniform distribution

85.18 loguvar

variance of the log-uniform distribution

85.19 medlogu

median of the log-uniform distribution

85.20 test_logurnd

85.21 tricdf

cumulative distribution of the log-triangular distribution

85.22 triinv

inverse of the triangular distribution

85.23 trimediam

median of the triangular distribution

85.24 tripdf

probability density of the triangular distribution

85.25 trirnd

random numbers of the triangular distribution

86 statistics

86.1 max_exprnd

86.2 maxnnormals

expected maximum of n normal variables
c.f. Wolperts
this is the median, not the mean of the maximum!
see median of gumbel

86.3 mean_generalized_gampdf

86.4 midrange

mid range of columns of X

86.5 minavg

solution of the minimum variance problem
minimise the variance of the weighted sum of n-independent
random variables with equal mean and individual variance

86.6 mode_man

87 statistics/moment-statistics

87.1 autocorr_man3

autocorrelation of the columns of X

87.2 autocorr_man4

autocorrelation for x if x is a vector, or individually for the columns of x if x is a matrix

c.f. box jenkins 2008 eq. 2.1.12

Note that it is faster to compute the acf in frequency space as done in the matlab internal function

87.3 autocorr_man5

autocorrellation of the columns of X

87.4 blockserr

estimate the standard error of potetially sequentilly correlated data

by blocking

block length should be sufficiently larger than correlation length and sufficiently smaller than data length

this uses a sliding block approach, which reduces the variation of the error estimate

87.5 comoment

non-central higher order moments of the multivariate normal
distribution

c.f. Moments and cumulants of the multivariate real and complex
Gaussian distributions

note : there seem to be some typos in the original paper,
for $x^4 c_{ii}^2$, the square seems to be missing

mu : nx1 mean vector

C : nxn covariance matrix

k : nx1 powers of variables in moments

87.6 corr_man

correlation of two vectors

87.7 cov_man

covariance matrix of two vectors

87.8 dof

mininum number of support points
for a polynomial of degree order in dim dimensions

87.9 edgeworth_quantile

inverse edgeworth expansion

c.f. cornis fisher 1937

c.f. Rao 2010

c.f. 2.50 in hall

CHERNOZHUKOV 3.3

87.10 effective_sample_size

effective sample size of the weighted mean of uncorrelated data
c.f. Kish

87.11 f_correlation

correction factor for standard error of the mean of n ar1-correlated iid samples

87.12 f_finite

reduction factor of standard error for sampling from a finite distribution without replacement

87.13 lmean

mean of $x.^l$, not of abs

87.14 lmoment

l-moment of vector x

87.15 maskmean

mean of the masked values of X

87.16 masknanmean

87.17 mean1

mean of x

87.18 mean_man

mean and standard error of X

87.19 mse

mean squared error of residual vector res
this is de-facto the std for an unbiased residual

87.20 nanautocorr_man1

autocorrelation of a vector with nan-values

87.21 nanautocorr_man2

autocorrelation of a vector with nan-values

87.22 nanautocorr_man4

compute autocorrelation for x if x is a vector, or individually
for the
columns of x if x is a matrix
box jenkins 2008 eq. 2.1.12
TODO nan is problematic!
Note that it is faster to compute the acf in frequency space
as done in the matlab internal function

87.23 nancorr

(co)-correlation matrix when samples a NaN

87.24 nancumsum

cumulative sum, setting nan values to zero

87.25 nanlmean

mean of the l-th power of the absolute value of x

87.26 nanr2

coefficient of determination when samples are invalid

87.27 nanrms

root mean square value when sample contains nan-values

87.28 nanrmse

root mean square error from vector of residuals
this is de-facto the std for an unbiased residual

87.29 nanserr

standard error of x with respect to mean when x contains nan values

87.30 nanwmean

weighted mean
 $\min_x \sum w (x - \mu)^2 \Rightarrow \mu = \sum(wx) / \sum(w)$
varargin can be dim
function [mu serr] = nanwmean(w,x)

87.31 nanwstd

weighed standard deviation

87.32 nanwvar

weighted variance of columns, corrected for degrees of freedom (bessel)

$s^2 = \sum(w*(x - \sum(wx)/\sum(w))^2) / \sum(w)$

87.33 nanxcorr

87.34 pearson

pearson correlation coefficient

87.35 pearson_to_kendall

conversion of pearson to kendall correlation coefficient
c.f. Kruskal 1958

87.36 pool_samples

pooled mean and standard deviation of several groups of different
size, mean and standard deviation

87.37 qmean

trimmed mean

87.38 range_mean

87.39 rmse_

root mean square error computed from a residual vector
this is de-facto the std for an unbiased residual

87.40 serr

standard error of the mean of a set of uncorrelated samples

87.41 `serr1`

87.42 `test_qskew`

87.43 `test_qstd_qskew_optimal_p`

87.44 `wautocorr`

autocorrelation for `x` if `x` is a vector, or individually for the columns of `x` if `x` is a matrix
samples can be weighted

c.f. box jenkins 2008 eq. 2.1.12

c.f. `autocorr_man4`

Note that it is faster to compute the acf in frequency space
as done in the matlab internal function

87.45 `wcorr`

correlation of two vectors when samples are weighted

87.46 `wcov`

covariance of two vectors when samples are weighted

87.47 `wdof`

effective degrees of freedom for weighted samples

87.48 wkurt

kurtosis with weighted samples

87.49 wmean

weighted mean

$\min_x \sum w (x - \mu)^2 \Rightarrow \mu = \sum(wx) / \sum(w)$

varargin can be dim

function [mu serr] = wmean(w,x)

87.50 wrms

weighted root mean square

87.51 wserr

weighted root mean square error

87.52 wskew

skewness of a weighted set of samples

function sk = wskew(w,x)

87.53 wstd

weighed standard deviation

87.54 wvar

weighted variance of columns, corrected for degrees of freedom (
bessel)
variance of the weighted sample mean of samples with same mean (but
not necessarily same variance)
 $s^2 = \text{sum} (w^2(x - \text{sum}(wx))^2)$

s2_mu : error of mean, s2_mu : sd of prediction

88 statistics

88.1 nangeomean

88.2 nangeostd

geometric standard deviation ignoring nan-values

89 statistics/nonparametric-statistics

89.1 kernel1d

X : ouput x axis bins
xi : samples along x
m : number of bins in X
fun : kernel function
pdf : propability density of xi

89.2 kernel2d

kernel density estimate in two dimensions

90 statistics

90.1 normmmoment

expected norm of $x.^n$, when values x in x are iid normal with mu
and sigma

90.2 normpdf2

pdf of the bivariate normal distribution

91 statistics/order-statistics

91.1 hodges_lehmann_location

hodges lehman location estimator

Asymptotic rms efficiency of location estimate:

mean: $1 s/\sqrt{n}$

hodges lehman: $\sqrt{\pi/3} * s \sim 1.0233 s/\sqrt{n}$

median: $\pi/2 s/\sqrt{n} \sim 1.25 s / \sqrt{n}$

91.2 kendall

kendall correlation coefficient

91.3 kendall_to_pearson

convert kendall rank correlation coefficient to the person product
moment
correlation coefficient

c.f. Kruska, 1985

91.4 mad2sd

transform median absolute deviation to standard deviation
for normal distributed values

91.5 madcorr

proxy correlation by median absolute deviation

91.6 median2_holder

91.7 median_ci

median and its confidence intervals under assumption of normality
`se_me = sqrt(1/2 pi) 1.25331 * sd/sqrt(n)`

91.8 median_man

median and confidence intervals
`c` is a P value for the confidence interval,
default is 0.95 (2-sigma)
median of the columns of `X`

91.9 mediani

index of median, if median is not unique, any of the values is
chosen

91.10 nanmadcorr

proxy correlation by median absolute deviation

91.11 nanwmedian

weighted median, skips nan-values

91.12 nanwquantile

weighted quantile, skips nan values

91.13 oja_median

two dimensional oja median

note: the multivariate median is not unique

oja 1983, for extension to multivariate function, see chaudhri

91.14 qkurtosis

kurtosis computed for quantiles

Note : this is a measurement of shape-tailedness and yields the same value for the

normal distribution as "kurtosis"

However, this is a separate statistic and hence requires different

methods for calculating P-values and hypothesis testing

91.15 qmoments

moments estimated from quantiles

91.16 qskew

skewness estimated from quantiles

Note : this is a measurement of shape-symmetry and yields the same value for the

skew-normal distribution as "skewness"

However, this is an own statistic and hence requires different

methods for calculating P-values and hypothesis testing

91.17 qskewq

skewness estimated by quantiles

91.18 qstdq

proxy standard deviation determined by quantiles

91.19 quantile1_optimisation

91.20 quantile2_breckling

qunatile regression

91.21 quantile2_chaudhuri

quantile regression

91.22 quantile2_projected

quantile in two dimensions

91.23 quantile2_projected2

spatial qunatile for chosen direction

91.24 quantile_envelope

91.25 quantile_regression_simple

simple quantile regression

91.26 ranking

ranking for spearman statistics

91.27 spatial_median

c.f. Oja 2008

is this the same as the oja simplex median (c.f. small 1990)?

91.28 spatial_quantile

spatial quantile

91.29 spatial_quantile2

spatial quantile

91.30 spatial_quantile3

spatial quantile

91.31 spatial_rank

unsigned rank

91.32 spatial_sign

spatial sign

91.33 spatial_signed_rank

signed rank

Note: this is only a true rank if X is normal with zero mean,
arbitrary variance

91.34 spearman

spearman's product moment coefficient

91.35 spearman_rank

91.36 spearman_to_pearson

conversion of spearman rank to person product moment correlation coefficient

91.37 wmedian

weighted median

91.38 wquantile

weighted quantile

92 statistics

92.1 qstd

92.2 quantile_extrap

92.3 quantile_sin

93 statistics/random-number-generation

93.1 laplacernd

random number of laplace distribution

93.2 randc

correlate to correlated standard normally distributed vectors

93.3 skewness2param

93.4 skewpdf_central_moments

93.5 skewrnd

random numbers of the skew normal distribution

93.6 skewrnd2

random numbers of the skew normal distribution

94 statistics

94.1 range

range and mid range of input

94.2 resample_with_replacement

95 statistics/resampling-statistics/@Jackknife

95.1 Jackknife

class for leave out 1 (delete 1) Jackknife estimates

note 1 : the 1-delete jackknife does not yield consistend estimates
for all functions,
in particular it will perform poorly on robust estimation
functions
this is overcome by the d-delete jackknife, where d has to
exceed the breakdown point
of the estimating function, for example \sqrt{n} for the
median
as this leads to unreasonably large number of repetitions,
bootstrap
is recommended for large sample cases (or blocking for
sequential data)
note 2 : as a linearisation, jackknife underestimates the error
variance in case of
dependence in the data
note 3 : studentisation and the leave out 1 jackknife are related
note 4 : the double 1 sample jackknife performs iferior to the d1
jackknife

95.2 estimated_STATIC

jackknife estimate of mean, bias and standard error
theta0 : estimate from all samples
thetad : set of estimates obtained by leaving out one data point
each
last dimension of theta is assumed to be the jackknife
dimension

95.3 matrix1_STATIC

matrix of estimation for leaving out two samples at a time

95.4 matrix2

matrix of estimations for jacknive with two samples left out

96 statistics/resampling-statistics

96.1 block_jackknife

96.2 jackknife_moments

moments determined by the jackknife

func : function of interest on the samples (e.g. mean)
A : parameter matrix
 columns : parameters
 rows : samples of the parameter sets
d : number of samples left out

96.3 moving_block_jackknife

blocked Jackknife for autocorrelated data
sliding block, statistically more efficient but computationally
expensive
note, number of blocks must be sufficiently large $h \sim \sqrt{n}$? $\ll n$

96.4 randblockterr

standard error of sequentially correlated data by blocking
block length should be sufficiently larger than correlation length
and sufficiently smaller than data length
this uses a sliding block approach, which reduces the variation of
the error estimate
TODO this does not work, randomly picking samples does not reveal
the correlation

96.5 resample

resample a vector and apply function to it

TODO, should be with replacement

n : number of samples

m : number of subsamples
cx : maximum number of combinations

97 statistics

97.1 scale_quantile_sd

scale factor for the standard deviation
of the asymptotic distribution of sample quantiles
(for normal distribution)
see cadwell, 1952

97.2 sd_sample_quantiles

97.3 skew_generalized_normal_fit

97.4 skew_generalized_normpdf

97.5 skewcdf

97.6 skewparam_to_central_moments

97.7 skewpdf

skew-normal distribution
c.f. Azzalini 1985

97.8 spatialrnd

97.9 test_mean_generalized_gampdf

97.10 test_skew_generalized_normpdf

97.11 trimmed_mean

trimmed mean

97.12 ttest2_man

two-sample t-test
here posix return value standard: h = 0 accepted, h = 1 failed
note: the matlab logic is inverse : h = 1 accepted, h = 0 failed
two sided univariate t-test

97.13 ttest_man

two-sample t-test
unequal sample size
equal variance

97.14 ttest_paired

paired t-test
unequal sample size
equal variance
more powerfull than unpaired test, as long as correlation between
x1 and x2 > 0

97.15 wgeomean

weighted geometric mean
function mu = wgeomean(w,x)

97.16 wgeovar

variance of the weighted geometric mean

97.17 wharmean

weighted harmonic mean

97.18 wharstd

97.19 wharvar

97.20 wnormpdf

wrapped normal distribution to the unit circle
c.f. stephens

98 stochastic

98.1 brownian_noise_1d_acf

98.2 brownian_noise_1d_fft

98.3 brownian_noise_2d_fft

99 mathematics

mathematical functions of various kind

99.1 ternary_diagram

100 test/master

100.1 dat_test_lanczos_3d_k_20_n_40

100.2 poisson2d_blk

100.3 qr_implicit_givens_2

100.4 spectral_derivative_2d

100.5 test_2d_eigensolver_hydrogen

100.6 test_2d_refine

100.7 test_3d_eigensolver_hydrogen

100.8 test_FEM

100.9 test_Mesh_3d

100.10 test_arnoldi

100.11 test_arpackc

100.12 test_assemble

100.13 test_assembly_performance

100.14 test_bc_one_sided

100.15 test_compare_solvers

100.16 test_complete

100.17 test_convergence

100.18 test_convergence_b

100.19 test_df_2d

100.20 test_eig_algs

100.21 test_eig_inverse

100.22 test_eigs_lanczos

100.23 test_eigs_lanczos_1

100.24 test_eigs_lanczos_2

100.25 test_eigs_lanczos_performance

100.26 test_fdm

100.27 test_fdm_d_vargrid

100.28 test_fdm_spectral

100.29 test_fem

100.30 test_fem_1d

100.31 test_fem_1d_higher_order

100.32 test_fem_2d_adaptive

100.33 test_fem_2d_higher_order

100.34 test_fem_3d_higher_order

100.35 test_fem_3d_refine

100.36 test_fem_b

100.37 test_fem_derivative

100.38 test_fem_quadrature

100.39 test_final

100.40 test_fix_substitution

100.41 test_forward

100.42 test_get_sparse_arrays

100.43 test_harmonic_oscillator

100.44 test_high_order_fdm_periodic_bc

100.45 test_hydrogen_wf

100.46 test_ichol

100.47 test_interpolation

100.48 test_inverse_problem

100.49 test_it_vs_exact

100.50 test_jama

100.51 test_jd

100.52 test_jdqz

100.53 test_lanczos_2

100.54 test_lanczos_biorthogonal

100.55 test_laplacian

100.56 test_laplacian_non_uniform

100.57 test_laplacian_simple

100.58 test_mesh_2d_uniform

100.59 test_mesh_2d_uniform_2

100.60 test_mesh_circle

100.61 test_mesh_generation

100.62 test_mesh_interpolate

100.63 test_mg

100.64 test_minres_recycle

100.65 test_multigrid

100.66 test_nc

100.67 test_nonuniform_symmetric

100.68 test_pde

100.69 test_permutation

100.70 test_poison_fem

100.71 test_polar

100.72 test_potential

100.73 test_powers

100.74 test_precondition

100.75 test_project_rectangle

100.76 test_qr

100.77 test_quantum_well

100.78 test_radial_adaptive

100.79 test_radial_confinement

100.80 test_radial_fixes

100.81 test_refine_2d

100.82 test_refine_2d_b

100.83 test_refine_3d

100.84 test_refine_structural

100.85 test_regularisation

100.86 test_round_off

100.87 test_schrödinger_potentials

100.88 test_uniform_mesh

100.89 test_vargrid

101 test

101.1 test_bandpass2d

101.2 test_gaussfit3

101.3 test_geoserr

101.4 test_hexagonal_pattern

101.5 test_lognfit_quantile

101.6 test_lowpass1d_implicit

101.7 test_lowpass2d_anisotropic

101.8 test_lowpass2d_fft

101.9 test_max_normal

101.10 test_moments_fourier_power

101.11 test_mtimes3x3

102 wavelet

102.1 continuous_wavelet_transform

continuous wavelet transform
follows "The Illustrated Wavelet Transform Handbook: Introductory
Theory and ..."

102.2 cwt_man

continuous fourier transform
as of time of implmentation, the matlab interal cwt is affected by
serious round-off errors and has issues with the scaling,
which is not the case here

102.3 example_wavelets

102.4 phasewrap

wrap the phase to $\pm \pi$

102.5 test_cwt_man

102.6 test_phasewrap

102.7 test_wavelet

102.8 test_wavelet2

102.9 test_wavelet_analysis

102.10 test_wavelet_reconstruct

102.11 test_wtc

102.12 wavelet

wavelet windows

102.13 wavelet_reconstruct

inverses wavelet transform for single frequency
(reconstruction of time series)
n : window lengths in multiples of filter period $1/f_0$

102.14 wavelet_transform

wavelet transform for single frequency
n : window lengths in multiples of filter period $1/f_0$