

Manual for Package: mathematics

Revision 2:9M

Karl Kästner

March 28, 2020

Contents

1	calendar	34
1.1	days_per_month	34
1.2	isnight	34
2	mathematics	35
2.1	cast_byte_to_integer	35
3	complex-analysis	35
3.1	complex_exp_product_im_im	35
3.2	complex_exp_product_im_re	35
3.3	complex_exp_product_re_im	35
3.4	complex_exp_product_re_re	36
3.5	croots	36
3.6	root_complex	37
3.7	test_imroots	37
4	derivation	37
4.1	derive_acfar1	37
4.2	derive_ar2param	37
4.3	derive_arc_length	37
4.4	derive_fourier_power	37
4.5	derive_fourier_power_exp	37
4.6	derive_laplacian_curvilinear	37
4.7	derive_laplacian_fourier_piecewise_linear	37
4.8	derive_logtripdf	38
4.9	derive_smooth1d_parametric	38
5	derivation/master	38
5.1	derive_bc_one_sided	38
5.2	derive_convergence	38

5.3	derive_error_fdm	38
5.4	derive_fdm_poly	38
5.5	derive_fdm_power	38
5.6	derive_fdm_taylor	38
5.7	derive_fdm_vargrid	38
5.8	derive_fem_2d_mass	39
5.9	derive_fem_error_2d	39
5.10	derive_fem_error_3d	39
5.11	derive_fem_sym_2d	39
5.12	derive_grid_constants	39
5.13	derive_interpolation	39
5.14	derive_laplacian	39
5.15	derive_limit	39
5.16	derive_nc_1d	39
5.17	derive_nc_1d_	39
5.18	derive_nc_2d	40
5.19	derive_nonuniform_symmetric	40
5.20	derive_richardson	40
5.21	derive_sum	40
5.22	nn	40
5.23	test_derive	40
5.24	test_derive_fdm_poly	40
5.25	test_filter	40
5.26	test_vargrid	40
6	derivation	41
6.1	simplify_atan	41
7	mathematics	41
7.1	exp10	41
8	finance	41
8.1	derive_skewrnd_walsh_paramter	41
8.2	gbm_cdf	41
8.3	gbm_fit	41
8.4	gbm_fit_old	41
8.5	gbm_inv	41
8.6	gbm_mean	42
8.7	gbm_median	42
8.8	gbm_pdf	42
8.9	gbm_simulate	42
8.10	gbm_skewness	42
8.11	gbm_std	42
8.12	gbm_transform_time_step	42

8.13	put_price_black_scholes	42
8.14	skewgbm_simulate	42
8.15	skewrnd_walsh	42
9	finance/test	43
9.1	test_gbm	43
9.2	test_gbm_pdf	43
9.3	test_skewrnd_walsh	43
10	fourier/@STFT	43
10.1	STFT	43
10.2	itransform	43
10.3	stft_	43
10.4	stftmat	43
10.5	transform	44
11	fourier	44
11.1	amplitude_from_peak	44
11.2	dftmtx_man	44
11.3	example_fourier_window	44
11.4	fft_derivative	45
11.5	fft_man	45
11.6	fftsmooth	45
11.7	fix_fourier	46
11.8	fourier_axis	46
11.9	fourier_cesaro_correction	46
11.10	fourier_coefficient_piecewise_linear	46
11.11	fourier_coefficient_piecewise_linear_1	47
11.12	fourier_coefficient_ramp3	47
11.13	fourier_coefficient_ramp_pulse	47
11.14	fourier_coefficient_ramp_step	47
11.15	fourier_coefficient_square_pulse	47
11.16	fourier_cubic_interaction_coefficients	47
11.17	fourier_derivative	48
11.18	fourier_expand	48
11.19	fourier_fit	48
11.20	fourier_interpolate	48
11.21	fourier_matrix	48
11.22	fourier_matrix2	48
11.23	fourier_matrix3	48
11.24	fourier_matrix_exp	49
11.25	fourier_multiplicative_interaction_coefficients	49
11.26	fourier_power	49
11.27	fourier_power_exp	49

11.28	fourier_predict	49
11.29	fourier_quadratic_interaction_coefficients	50
11.30	fourier_range	50
11.31	fourier_regress	50
11.32	fourier_resampled_fit	50
11.33	fourier_resampled_predict	50
11.34	fourier_signed_square	50
11.35	fourier_transform	51
11.36	hyperbolic_fourier_box	51
11.37	idftmtx_man	51
11.38	laplace_2d_pwlinear	51
11.39	nanfft	52
11.40	peaks	52
11.41	roots_fourier	52
11.42	spectral_density	52
11.43	test_complex_exp_product	52
11.44	test_fourier_filter	53
11.45	test_idftmtx	53
12 geometry/@Geometry		53
12.1	Geometry	53
12.2	arclength	53
12.3	arclength_old	53
12.4	arclength_old2	53
12.5	base_point	53
12.6	base_point_limited	54
12.7	centroid	54
12.8	cosa_min_max	54
12.9	cross2	54
12.10	curvature	54
12.11	ddot	54
12.12	distance	54
12.13	distance2	54
12.14	dot	55
12.15	edge_length	55
12.16	enclosed_angle	55
12.17	enclosing_triangle	55
12.18	hexagon	55
12.19	inPolygon	55
12.20	inTetra	55
12.21	inTetra2	55
12.22	inTriangle	56
12.23	intersect	56
12.24	lineintersect	56

12.25	lineintersect1	56
12.26	minimum_distance_lines	56
12.27	mittenpunkt	56
12.28	nagelpoint	56
12.29	onLine	56
12.30	orthocentre	57
12.31	plumb_line	57
12.32	poly_area	57
12.33	poly_edges	57
12.34	poly_set	57
12.35	poly_width	57
12.36	polyxpoly	57
12.37	project_to_curve	58
12.38	quad_isconvex	58
12.39	random_disk	58
12.40	random_simplex	58
12.41	sphere_volume	58
12.42	tetra_volume	58
12.43	tobarycentric	58
12.44	tobarycentric1	58
12.45	tobarycentric2	59
12.46	tobarycentric3	59
12.47	tri_angle	59
12.48	tri_area	59
12.49	tri_centroid	59
12.50	tri_distance_opposit_midpoint	59
12.51	tri_edge_length	59
12.52	tri_edge_midpoint	59
12.53	tri_excircle	60
12.54	tri_height	60
12.55	tri_incircle	60
12.56	tri_isacute	60
12.57	tri_isobtuse	60
12.58	tri_semiperimeter	60
12.59	tri_side_length	60
13 geometry		61
13.1	Polygon	61
13.2	bounding_box	61
13.3	curvature_1d	61
13.4	cvt	61
13.5	deg_to_frac	61
13.6	ellipse	61
13.7	ellipseX	62

13.8	ellipseY	62
13.9	first_intersect	62
13.10	golden_ratio	62
13.11	hypot3	62
13.12	meanangle	62
13.13	meanangle2	62
13.14	meanangle3	62
13.15	meanangle4	63
13.16	medianangle	63
13.17	medianangle2	63
13.18	pilim	63
13.19	streamline_radius_of_curvature	63
14	histogram/@Histogram	63
14.1	2x	63
14.2	Histogram	64
14.3	bimodes	64
14.4	cdf	64
14.5	cdfS	64
14.6	chi2test	64
14.7	cmoment	64
14.8	cmomentS	64
14.9	entropy	64
14.10	entropyS	64
14.11	iquantile	64
14.12	kstest	65
14.13	kurtosis	65
14.14	kurtosisS	65
14.15	mean	65
14.16	meanS	65
14.17	median	65
14.18	medianS	65
14.19	mode	65
14.20	modeS	65
14.21	moment	65
14.22	momentS	66
14.23	pdf	66
14.24	quantile	66
14.25	quantileS	66
14.26	setup	66
14.27	skewness	66
14.28	skewnessS	66
14.29	stairs	66
14.30	stairsS	66

14.31	std	66
14.32	stdS	67
14.33	var	67
14.34	varS	67
15	histogram	67
15.1	hist_man	67
15.2	histadapt	67
15.3	histconst	67
15.4	pdf_poly	67
15.5	plotcdf	67
15.6	test_histogram	67
16	linear-algebra	68
16.1	averaging_matrix_2	68
16.2	colnorm	68
16.3	condest_	68
17	linear-algebra/coordinate-transformation	68
17.1	barycentric2cartesian	68
17.2	barycentric2cartesian3	68
17.3	cartesian2barycentric	68
17.4	cartesian_to_unit_triangle_basis	68
17.5	ellipsoid2geoid	68
17.6	example_approximate_utm_conversion	69
17.7	latlon2utm	69
17.8	latlon2utm_simple	69
17.9	lowrance_mercator_to_wgs84	69
17.10	nmea2utm	69
17.11	sn2xy	69
17.12	unit_triangle_to_cartesian	69
17.13	utm2latlon	69
17.14	xy2nt	70
17.15	xy2sn	70
17.16	xy2sn_java	70
17.17	xy2sn_old	70
18	linear-algebra	70
18.1	det2x2	70
18.2	det3x3	70
18.3	det4x4	71
18.4	diag2x2	71
18.5	eig2x2	71

19	linear-algebra/eigenvalue	71
19.1	eig_bisection	71
19.2	eig_inverse	71
19.3	eig_inverse_iteration	71
19.4	eig_power_iteration	71
20	linear-algebra/eigenvalue/jacobi-davidson	71
20.1	afun_jdm	71
20.2	davidson	72
20.3	jacobi_davidson	72
20.4	jacobi_davidson_qr	72
20.5	jacobi_davidson_qz	72
20.6	jacobi_davidson_simple	72
20.7	jdqr	72
20.8	jdqr_sleijpen	76
20.9	jdqr_vorst	79
20.10	jdqz	82
20.11	mfunc_jdm	86
20.12	mgs	86
20.13	minres_	87
20.14	mv_jacobi_davidson	87
21	linear-algebra	87
21.1	first	87
21.2	gershgorin_circle	87
21.3	haussdorff	87
21.4	ieig2x2	87
21.5	inv2x2	87
21.6	inv3x3	88
21.7	inv4x4	88
22	linear-algebra/lanczos	88
22.1	arnoldi	88
22.2	arnoldi_new	88
22.3	eigs_lanczos_man	88
22.4	lanczos	88
22.5	lanczos_	88
22.6	lanczos_biorthogonal	88
22.7	lanczos_biorthogonal_improved	88
22.8	lanczos_ghep	89
22.9	mv_lanczos	89
22.10	reorthogonalise	89
22.11	test_lanczos	89

23	linear-algebra/linear-systems	89
23.1	gmres_man	89
23.2	minres_recycle	89
24	linear-algebra	89
24.1	lpmean	89
24.2	lpnorm	89
24.3	matvec3	90
24.4	max2d	90
24.5	mpoweri	90
24.6	mtimes2x2	90
24.7	mtimes3x3	90
24.8	nannorm	90
24.9	nanshift	90
24.10	nl	90
24.11	normalise	91
24.12	normalize1	91
24.13	normrows	91
24.14	orth2	91
24.15	orth_man	91
24.16	orthogonalise	91
24.17	paddext	91
24.18	paddval1	92
24.19	paddval2	92
25	linear-algebra/polynomial	92
25.1	chebychev	92
25.2	piecewise_polynomial	92
25.3	roots1	92
25.4	roots2	92
25.5	roots2poly	92
25.6	roots3	92
25.7	roots4	93
25.8	test_roots4	93
25.9	vanderi_1d	93
26	linear-algebra	93
26.1	randrot	93
26.2	right	93
26.3	rot2	93
26.4	rot2dir	93
26.5	rot3	93
26.6	rotR	94
26.7	rownorm	94

26.8	simmilarity_matrix	94
26.9	spnorm	94
26.10	spzeros	94
26.11	test_roots3	94
26.12	transform_minmax	94
26.13	transpose3	94
26.14	transposeall	94
27	logic	95
27.1	bitor_man	95
28	master/plot	95
28.1	attach_boundary_value	95
28.2	cartesian_polar	95
28.3	img_vargrid	95
28.4	plot_basis_functions	95
28.5	plot_convergence	95
28.6	plot_dof	95
28.7	plot_eigenbar	95
28.8	plot_error_estimation	96
28.9	plot_error_estimation_2	96
28.10	plot_error_fem	96
28.11	plot_fdm_kernel	96
28.12	plot_fdm_vs_fem	96
28.13	plot_fem_accuracy	96
28.14	plot_function_and_grid	96
28.15	plot_hat	96
28.16	plot_hydrogen_wf	96
28.17	plot_mesh	96
28.18	plot_mesh_2	97
28.19	plot_refine	97
28.20	plot_refine_3d	97
28.21	plot_runtime	97
28.22	plot_spectrum	97
28.23	plot_wavefunction	97
29	master/ported	97
29.1	assemble_2d_dphi_dphi	97
29.2	assemble_2d_phi_phi	97
29.3	assemble_3d_dphi_dphi	97
29.4	assemble_3d_phi_phi	98
29.5	dV_2d_	98
29.6	derivative_2d	98
29.7	derivative_3d	98

29.8	element_neighbour_2d	98
29.9	prefetch_2d_	98
29.10	promote_2d_3_10	98
29.11	promote_2d_3_15	98
29.12	promote_2d_3_21	98
29.13	promote_2d_3_6	98
29.14	promote_3d_4_10	99
29.15	promote_3d_4_20	99
29.16	promote_3d_4_35	99
29.17	vander_2d	99
29.18	vander_3d	99
30	master/sandbox	99
30.1	adapt	99
30.2	assoc_laguerre	99
30.3	assoc_legendre	99
30.4	c23	99
31	master/sandbox/cg	100
31.1	cg	100
31.2	cg_coef_to_poly	100
31.3	errmat	100
31.4	lanczos	100
31.5	laplacian_2d	100
31.6	test_cg_eigs	100
31.7	test_lanczos	100
32	master/sandbox	100
32.1	condition_number_higher_order	100
32.2	confinement_dat	100
32.3	convergence_2d_3d	101
32.4	convergence_matrix_powers	101
32.5	cut_out	101
32.6	derivative_2d	101
32.7	derivative_3d	101
32.8	dummy	101
32.9	eig_error	101
32.10	eigs_fix	101
32.11	energy_level	101
32.12	equalise	101
32.13	example_int64	102
33	master/sandbox/fem-matlab	102
33.1	boundary_circle	102

33.2	boundary_rectangle	102
33.3	geometry_circle_with_hole	102
33.4	geometry_rectangle	102
34	master/sandbox	102
34.1	fem_2d_estimate_error	102
34.2	fem_assemble_scratch	102
34.3	fem_s	103
34.4	fourier_h	103
34.5	grad_2d	103
34.6	grad_3d	103
34.7	gradient	103
34.8	harmonic_oscillator	103
34.9	hydrogen_2d_analytic	103
34.10	hydrogen_boxed	103
34.11	hydrogen_boxed_old	103
34.12	hydrogen_wave	103
34.13	hydrogen_wf	104
34.14	ichol_man	104
34.15	known_eigenvalue	104
34.16	kron_man	104
34.17	laguerre	104
34.18	laplacian_arbitrary_order_old	104
34.19	laplacian_convergence	104
34.20	laplacian_cut_out	104
34.21	laplacian_cylindrical	104
34.22	laplacian_non_uniform_old	104
34.23	laplacian_polar	105
34.24	laplacian_simple	105
34.25	lderivative_3d	105
34.26	list_dat	105
34.27	matlab-horner	105
34.28	mesh_to_grid_2d_3	105
34.29	mg_mat	105
34.30	mv	105
34.31	orth2	105
34.32	partial_derivative_2d	105
34.33	partition_function	106
34.34	partition_function_old	106
34.35	poisson	106
34.36	poisson_fem	106
34.37	potential	106
34.38	powerc	106
34.39	quick_newihbour	106

34.40	radial	106
34.41	radial_convergence	106
34.42	radial_wafefunction	106
34.43	refine_2d	107
34.44	refine_3d	107
34.45	relerr	107
34.46	restore_cw	107
34.47	runtime_bm	107
34.48	rydberg	107
34.49	s_old	107
34.50	snorm	107
34.51	spherical_harmonic	107
34.52	split_eig	107
34.53	sum1	108
34.54	sum3	108
35	master/sandbox/summation	108
35.1	acc	108
35.2	add	108
35.3	ape	108
35.4	mmul_accurately	108
35.5	sum_kahan	108
35.6	sum_pairwise	108
35.7	test_sum	108
36	master/sandbox	109
36.1	test_convergence_ill_conditioned	109
36.2	test_fem_1d	109
36.3	test_fem_2d	109
36.4	test_fem_3d	109
36.5	test_increase	109
36.6	test_lanczos_shift	109
36.7	test_ldl	109
36.8	test_power	109
36.9	trefethen_p8_fdm	109
36.10	wavefunc	110
36.11	xgrid	110
37	number-theory	110
37.1	ceiln	110
37.2	digitsb	110
37.3	floorn	110
37.4	iseven	110
37.5	multichoosek	110

37.6	nchoosek_man	111
37.7	pythagorean_triple	111
37.8	roundn	111
38	numerical-methods/differentiation	111
38.1	derivative1	111
38.2	derivative2	111
39	numerical-methods/finite-difference	111
39.1	cdiff	111
39.2	cdiffb	112
39.3	cmean	112
39.4	derivative_matrix_1_1d	112
39.5	derivative_matrix_2_1d	112
39.6	derivative_matrix_2d	112
39.7	derivative_matrix_curvilinear	112
39.8	derivative_matrix_curvilinear_2	112
39.9	difference_kernel	113
39.10	distmat	113
39.11	gradpde2d	113
39.12	laplacian	113
39.13	laplacian_fdm	113
39.14	left	113
39.15	lrmean	113
40	numerical-methods/finite-difference/master	114
40.1	fdm_adaptive_grid	114
40.2	fdm_adaptive_refinement_old	114
40.3	fdm_assemble_d1_2d	114
40.4	fdm_assemble_d2_2d	114
40.5	fdm_confinement	114
40.6	fdm_d_vargrid	114
40.7	fdm_h_unstructured	114
40.8	fdm_hydrogen_vargrid	114
40.9	fdm_mark_unstructured_2d	114
40.10	fdm_plot	115
40.11	fdm_plot_series	115
40.12	fdm_refine_2d	115
40.13	fdm_refine_3d	115
40.14	fdm_refine_unstructured_2d	115
40.15	fdm_schroedinger_2d	115
40.16	fdm_schroedinger_3d	115
40.17	relocate	115

41 numerical-methods/finite-difference	115
41.1 mid	115
41.2 pwmid	116
41.3 ratio	116
41.4 steplength	116
41.5 swapoddeven	116
41.6 test_derivative_matrix_2d	116
41.7 test_derivative_matrix_curvilinear	116
41.8 test_difference_kernel	116
 42 numerical-methods/finite-element	 116
42.1 Mesh_2d.java	116
42.2 Tree_2d.java	117
42.3 assemble_1d_dphi_dphi	117
42.4 assemble_1d_phi_phi	117
42.5 assemble_2d_dphi_dphi.java	117
42.6 assemble_2d_phi_phi.java	117
42.7 assemble_3d_dphi_dphi.java	117
42.8 assemble_3d_phi_phi.java	117
42.9 boundary_1d	117
42.10 boundary_2d	117
42.11 boundary_3d	117
42.12 check_area_2d	118
42.13 circmesh	118
42.14 cropradius	118
42.15 display_2d	118
42.16 display_3d	118
42.17 distort	118
42.18 err_2d	118
42.19 estimate_err_2d_3	118
42.20 example_1d	118
42.21 example_2d	118
42.22 explode	119
42.23 fem_2d	119
42.24 fem_2d_heuristic_mesh	119
42.25 fem_get_2d_radial	119
42.26 fem_interpolation	119
42.27 fem_plot_1d	119
42.28 fem_plot_1d_series	119
42.29 fem_plot_2d	119
42.30 fem_plot_2d_series	119
42.31 fem_plot_3d	119
42.32 fem_plot_3d_series	120
42.33 fem_plot_confine_series	120

42.34	fem_radial	120
42.35	flip_2d	120
42.36	get_mesh_arrays	120
42.37	hashkey	120
43	numerical-methods/finite-element/int	120
43.1	int_1d_gauss	120
43.2	int_1d_gauss_1	120
43.3	int_1d_gauss_2	120
43.4	int_1d_gauss_3	121
43.5	int_1d_gauss_4	121
43.6	int_1d_gauss_5	121
43.7	int_1d_gauss_6	121
43.8	int_1d_gauss_lobatto	121
43.9	int_1d_nc_2	121
43.10	int_1d_nc_3	121
43.11	int_1d_nc_4	121
43.12	int_1d_nc_5	121
43.13	int_1d_nc_6	121
43.14	int_1d_nc_7	122
43.15	int_1d_nc_7_hardy	122
43.16	int_2d_gauss_1	122
43.17	int_2d_gauss_12	122
43.18	int_2d_gauss_13	122
43.19	int_2d_gauss_16	122
43.20	int_2d_gauss_25	122
43.21	int_2d_gauss_3	122
43.22	int_2d_gauss_33	122
43.23	int_2d_gauss_6	122
43.24	int_2d_gauss_7	123
43.25	int_2d_gauss_9	123
43.26	int_2d_nc_10	123
43.27	int_2d_nc_15	123
43.28	int_2d_nc_21	123
43.29	int_2d_nc_3	123
43.30	int_2d_nc_6	123
43.31	int_3d_gauss_1	123
43.32	int_3d_gauss_11	123
43.33	int_3d_gauss_14	123
43.34	int_3d_gauss_15	124
43.35	int_3d_gauss_24	124
43.36	int_3d_gauss_4	124
43.37	int_3d_gauss_45	124
43.38	int_3d_gauss_5	124

43.39	int_3d_nc_11	124
43.40	int_3d_nc_4	124
43.41	int_3d_nc_6	124
43.42	int_3d_nc_8	124
44	numerical-methods/finite-element	125
44.1	interpolation_matrix	125
44.2	mark	125
44.3	mark_1d	125
44.4	mesh_1d_uniform	125
44.5	mesh_3d_uniform	125
44.6	mesh_interpolate	125
44.7	neighbour_1d	125
44.8	old	125
44.9	pdeeig_1d	125
44.10	pdeeig_2d	126
44.11	pdeeig_3d	126
44.12	polynomial_derivative_1d	126
44.13	potential_const	126
44.14	potential_coulomb	126
44.15	potential_harmonic_oscillator	126
44.16	project_circle	126
44.17	project_rectangle	126
44.18	promote_1d_2.3	126
44.19	promote_1d_2.4	126
44.20	promote_1d_2.5	127
44.21	promote_1d_2.6	127
44.22	quadrilaterate	127
44.23	recalculate_regularity_2d	127
44.24	refine_1d	127
44.25	refine_2d_21	127
44.26	refine_2d_structural	127
44.27	regularity_1d	127
44.28	regularity_2d	127
44.29	regularity_3d	128
44.30	relocate_2d	128
44.31	test_circmesh	128
44.32	test_hermite	128
44.33	tri_assign_points	128
44.34	triangulation_uniform	128
44.35	vander_1d	128
44.36	vanderd_1d	128
44.37	vanderi_1d	128

45	numerical-methods/finite-volume/@Advection	129
45.1	Advection	129
45.2	dot_advection	129
46	numerical-methods/finite-volume/@Burgers	129
46.1	burgers_split	129
46.2	dot_burgers_fdm	129
46.3	dot_burgers_fft	129
47	numerical-methods/finite-volume/@Finite_Volume	129
47.1	Finite_Volume	129
47.2	apply_bc	130
47.3	solve	130
47.4	step_split_strang	130
47.5	step_unsplit	130
48	numerical-methods/finite-volume/@Flux_Limiter	130
48.1	Flux_Limiter	130
48.2	beam_warming	130
48.3	fromm	131
48.4	lax_wendroff	131
48.5	minmod	131
48.6	monotized_central	131
48.7	muscl	131
48.8	superbee	131
48.9	upwind	131
48.10	vanLeer	132
49	numerical-methods/finite-volume/@KDV	132
49.1	dot_kdv_fdm	132
49.2	dot_kdv_fft	132
49.3	kdv_split	132
50	numerical-methods/finite-volume/@Reconstruct_Average_Evolve	132
50.1	Reconstruct_Average_Evolve	132
50.2	advect_highres	133
50.3	advect_lowress	133
51	numerical-methods/finite-volume	133
51.1	Godunov	133
51.2	Lax_Friedrich	133
51.3	Measure	133
51.4	Roe	133
51.5	fv_swe	134
51.6	staggered_euler	134

51.7	staggered_grid	134
52	numerical-methods	134
52.1	grid2quad	134
53	numerical-methods/integration	134
53.1	cumintL	134
53.2	cumintR	134
53.3	int_trapezoidal	134
54	numerical-methods/interpolation/@Kriging	135
54.1	Kriging	135
54.2	estimate_semivariance	135
54.3	interpolate_	135
55	numerical-methods/interpolation/@RegularizedInterpolator1	135
55.1	RegularizedInterpolator1	135
55.2	init	135
56	numerical-methods/interpolation/@RegularizedInterpolator2	136
56.1	RegularizedInterpolator2	136
56.2	init	136
57	numerical-methods/interpolation/@RegularizedInterpolator3	136
57.1	RegularizedInterpolator3	136
57.2	init	136
58	numerical-methods/interpolation	136
58.1	IDW	136
58.2	IPoly	136
58.3	IRBM	137
58.4	ISparse	137
58.5	Inn	137
58.6	Interpolator	137
58.7	fixnan	137
58.8	idw1	137
58.9	idw2	137
58.10	inner2outer	138
58.11	inner2outer2	138
58.12	interp1_limited	138
58.13	interp1_man	138
58.14	interp1_save	138
58.15	interp1_slope	138
58.16	interp1_smooth	139
58.17	interp1_unique	139

58.18	interp2_man	139
58.19	interp_angle	139
58.20	interp_fourier	139
58.21	interp_fourier_batch	139
58.22	interp_sn	139
58.23	interp_sn2	140
58.24	interp_sn3	140
58.25	interp_sn_	140
58.26	limit_by_distance_1d	140
58.27	resample1	140
58.28	resample_d_min	140
58.29	resample_vector	140
58.30	test_interp1_limited	140
59	numerical-methods	141
59.1	inverse_complex	141
60	numerical-methods/ode	141
60.1	bvp2_check_arguments	141
60.2	bvp2c	141
60.3	bvp2c2	141
60.4	bvp2fdm	142
60.5	bvp2wavetrain	142
60.6	bvp2wavetwopass	142
60.7	ivp_euler_forward	142
60.8	ivprk2	143
60.9	ode2_matrix	143
60.10	ode2characteristic	143
60.11	step_trapezoidal	143
60.12	test_bvp2	143
61	numerical-methods/optimisation	143
61.1	armijo_stopping_criterion	143
61.2	astar	143
61.3	binsearch	144
61.4	bisection	144
61.5	box1	144
61.6	box2	144
61.7	cauchy	144
61.8	cauchy2	144
61.9	directional_derivative	144
61.10	dud	145
61.11	extreme3	145
61.12	extreme_quadratic	145

61.13	ftest	145
61.14	fzero_bisect	145
61.15	fzero_newton	145
61.16	grad	146
61.17	hessian	146
61.18	hessian_from_gradient	146
61.19	hessian_projected	146
61.20	line_search	146
61.21	line_search2	146
61.22	line_search_polynomial	147
61.23	line_search_polynomial2	147
61.24	line_search_quadratic	147
61.25	line_search_quadratic2	147
61.26	line_search_wolfe	148
61.27	ls_bgfs	148
61.28	ls_broyden	148
61.29	ls_generalized_secant	148
61.30	nlcg	148
61.31	nlls	149
61.32	picard	149
61.33	poly_extrema	149
61.34	quadratic_function	149
61.35	quadratic_programming	149
61.36	quadratic_step	149
61.37	rosenbrock	149
61.38	sqrt_heron	150
61.39	test_directional_derivative	150
61.40	test_dud	150
61.41	test_fzero_newton	150
61.42	test_line_search_quadratic2	150
61.43	test_ls_generalized_secant	150
61.44	test_nlcg_6_order	150
61.45	test_nlls	150
62	numerical-methods/pde	150
62.1	laplacian2d_fundamental_solution	150
63	numerical-methods/piecewise-polynomials	151
63.1	Hermite1	151
63.2	hp2_fit	151
63.3	hp2_predict	151
63.4	hp_predict	151
63.5	hp_regress	151
63.6	lp_count	151

63.7	lp_predict	152
63.8	lp_regress	152
63.9	lp_regress_	152
64	regression/@PolyOLS	152
64.1	PolyOLS	152
64.2	coefftest	152
64.3	detrend	152
64.4	fit	152
64.5	fit_	152
64.6	predict	153
64.7	predict_	153
64.8	slope	153
65	regression/@PowerLS	153
65.1	PowerLS	153
65.2	fit	153
65.3	predict	153
65.4	predict_	153
66	regression/@Theil	154
66.1	Theil	154
66.2	detrend	154
66.3	fit	154
66.4	predict	154
66.5	slope	154
67	regression	154
67.1	Theil_Multivariate	154
67.2	areg	155
67.3	ginireg	155
67.4	hesssimplereg	155
67.5	l1lin	155
67.6	lsq_sparam	155
67.7	polyfitd	155
67.8	regression_method_of_moments	156
67.9	robustlinreg	156
67.10	theil2	156
67.11	theil_generalised	156
67.12	total_least_squares	156
67.13	weighted_median_regression	156
68	set-theory	157
68.1	issubset	157

69	signal-processing	157
69.1	acf_effective_sample_size	157
69.2	acf_genton	157
69.3	acfar1	157
69.4	acfar1_2	157
69.5	acfar2	157
69.6	acfar2_2	158
69.7	ar1_cutoff_frequency	158
69.8	ar1_effective_sample_size	158
69.9	ar1_mse_mu_single_sample	158
69.10	ar1_mse_pop	158
69.11	ar1_mse_range	158
69.12	ar1_spectrum	158
69.13	ar1_to_tikhonov	158
69.14	ar1_var_factor	159
69.15	ar1_var_factor_	159
69.16	ar1_var_range2	159
69.17	ar1delay	159
69.18	ar1delay_old	159
69.19	ar2conv	159
69.20	ar2dof	160
69.21	ar2param	160
69.22	asymwin	160
69.23	autocorr_fft	160
69.24	bandpass	160
69.25	bandpass2	160
69.26	bartlett	160
69.27	bartlett_spectrogram	161
69.28	bin1d	161
69.29	bin2d	161
69.30	binormrnd	161
69.31	conv1_man	161
69.32	conv2_man	161
69.33	conv2z	161
69.34	conv30	162
69.35	conv_	162
69.36	conv_centered	162
69.37	convz	162
69.38	cosexpdelay	162
69.39	csmooth	162
69.40	daniell_window	162
69.41	danielle_window	163
69.42	db2neper	163
69.43	db2power	163

69.44	derive_danielle_weight	163
69.45	derive_limit_0_acfar	163
69.46	detect_peak	163
69.47	digital_low_pass_filter	163
69.48	doublesum_ij	164
69.49	effective_sample_size_to_ar1	164
69.50	filt_hodges_lehman	164
69.51	filter1	164
69.52	filter2	164
69.53	filter_	164
69.54	filteriir	164
69.55	filterp	165
69.56	filterp1	165
69.57	filterstd	165
69.58	firls_man	165
69.59	flattopwin	166
69.60	frequency_response_boxcar	166
69.61	freqz_boxcar	166
69.62	gaussfilt1	166
69.63	hanchangewin	166
69.64	hanchangewin2	166
69.65	hanwin	166
69.66	hanwin_	166
69.67	highpass	167
69.68	kaiserwin	167
69.69	kalman	167
69.70	lanczoswin	167
69.71	last	167
69.72	lowpass	167
69.73	lowpass2	167
69.74	lowpass_iir	167
69.75	lowpass_iir_symmetric	168
69.76	lowpassfilter2	168
69.77	maxfilt1	168
69.78	meanfilt1	168
69.79	medfilt1_man	168
69.80	medfilt1_man2	168
69.81	medfilt1_padded	168
69.82	medfilt1_reduced	168
69.83	mid_term_single_sample	169
69.84	minfilt1	169
69.85	mu2ar1	169
69.86	mymooth	169
69.87	nanautocorr	169

69.88	nanmedfilt1	169
69.89	neper2db	169
69.90	peaks_man	169
69.91	polyfilt1	170
69.92	qmedfilt1	170
69.93	randar1	170
69.94	randar1_dual	170
69.95	randar2	170
69.96	randarp	170
69.97	range_window	170
69.98	rectwin	170
69.99	recursive_sum	171
69.100	select_range	171
69.101	smooth1d_parametric	171
69.102	smooth2	171
69.103	smooth_man	171
69.104	smooth_parametric	171
69.105	smooth_parametric2	171
69.106	smooth_with_splines	171
69.107	smoothfft	172
69.108	spectrogram	172
69.109	std_window	172
69.110	sum_i_lag	172
69.111	sum_ii	172
69.112	sum_ii_	172
69.113	sum_ij	172
69.114	sum_ij_	172
69.115	sum_ij_partial_	173
69.116	sum_multivar	173
69.117	test_acfar1	173
69.118	test_acfar1_2	173
69.119	test_acfar1_3	173
69.120	test_acfar1_4	173
69.121	test_acfar2	173
69.122	test_ar1_var_factor	173
69.123	test_ar1_var_factor_2	173
69.124	test_ar1_var_mu_single_sample	173
69.125	test_ar1_var_pop	174
69.126	test_ar1_var_pop_1	174
69.127	test_ar1delay	174
69.128	test_bivariate_covariance_term	174
69.129	test_convexity	174
69.130	test_lanczoswin	174
69.131	test_madcorr	174

69.132	test_randar1	174
69.133	test_randar1_multivariate	174
69.134	test_randar2	174
69.135	test_sum_ij	175
69.136	test_sum_multivar	175
69.137	test_trifilt1	175
69.138	test_wautocorr	175
69.139	test_wavelet_transform	175
69.140	test_wordfilt	175
69.141	test_xar1_mid_term	175
69.142	tikhonov_to_ar1	175
69.143	trapwin	175
69.144	trifilt1	176
69.145	triwin	176
69.146	triwin2	176
69.147	varar1	176
69.148	welch_spectrogram	176
69.149	wfilt	176
69.150	winbandpass	176
69.151	window_make_odd	176
69.152	winfilt0	177
69.153	winlength	177
69.154	wmeanfilt	177
69.155	wmedfilt	177
69.156	wordfilt	177
69.157	wordfilt_edgeworth	177
69.158	xar1	177
69.159	xcorr_man	178
70 sorting		178
70.1	sort2	178
70.2	sort2d	178
71 special-functions		178
71.1	bessel_sphere	178
71.2	digamma_man	178
71.3	hankel_sphere	178
71.4	hermite	179
71.5	legendre_man	179
71.6	neumann_sphere	179
72 statistics		179
72.1	atan_s2	179
72.2	beta_mode_to_parameter	179

72.3	coefficient_of_determination	179
72.4	conditional_expectation_normal	179
72.5	correlation_confidence_pearson	180
73	statistics/distributions	180
73.1	PDF	180
73.2	binorm_separation_coefficient	180
73.3	binormcdf	180
73.4	binormfit	180
73.5	binormpdf	180
73.6	edgeworth_cdf	180
73.7	edgeworth_pdf	181
73.8	logn_mode2param	181
73.9	logn_param2mode	181
73.10	lognpdf_	181
73.11	pdfsample	181
73.12	t2cdf	181
73.13	t2inv	181
74	statistics	182
74.1	example_standard_error_of_sample_quantiles	182
74.2	f_var_finite	182
74.3	gamma_mode_to_parameter	182
74.4	gaussfit3	182
74.5	gaussfit_quantile	182
74.6	hodges_lehmann_correlation	182
74.7	hodges_lehmann_dispersion	182
75	statistics/information-theory	183
75.1	akaike_information_criterion	183
75.2	bayesian_information_criterion	183
76	statistics	183
76.1	kurtncdf	183
76.2	kurtnpdf	183
76.3	kurtosis_bias_corrected	183
76.4	limit	184
76.5	logfactorial	184
76.6	loglogpdf	184
76.7	lognfit_quantile	184
76.8	logskewcdf	184
76.9	logskewpdf	184
77	statistics/logu	184

77.1	lambertw_numeric	184
77.2	logtrialtcdf	184
77.3	logtrialtinv	185
77.4	logtrialtmean	185
77.5	logtrialtpdf	185
77.6	logtrialtrnd	185
77.7	logtricdf	185
77.8	logtriinv	185
77.9	logtrimean	186
77.10	logtripdf	186
77.11	logtrirnd	186
77.12	logucdf	186
77.13	logucm	186
77.14	loguinv	186
77.15	logumean	186
77.16	logupdf	186
77.17	logurnd	187
77.18	loguvar	187
77.19	medlogu	187
77.20	test_logurnd	187
77.21	tricdf	187
77.22	triinv	187
77.23	trimean	187
77.24	tripdf	187
77.25	trirnd	188
78 statistics		188
78.1	maxnnormals	188
78.2	midrange	188
78.3	minavg	188
78.4	mode_man	188
79 statistics/moment-statistics		188
79.1	autocorr_man3	188
79.2	autocorr_man4	189
79.3	autocorr_man5	189
79.4	blockserr	189
79.5	comoment	189
79.6	corr_man	189
79.7	cov_man	190
79.8	dof	190
79.9	edgeworth_quantile	190
79.10	effective_sample_size	190
79.11	f_correlation	190

79.12	f_finite	190
79.13	lmean	190
79.14	lmoment	191
79.15	maskmean	191
79.16	masknanmean	191
79.17	mean1	191
79.18	mean_man	191
79.19	mse	191
79.20	nanautocorr_man1	191
79.21	nanautocorr_man2	191
79.22	nanautocorr_man4	192
79.23	nancorr	192
79.24	nancumsum	192
79.25	nanlmean	192
79.26	nanr2	192
79.27	nanrms	192
79.28	nanrmse	192
79.29	nanserr	193
79.30	nanwmean	193
79.31	nanwstd	193
79.32	nanwvar	193
79.33	nanxcorr	193
79.34	pearson	193
79.35	pearson_to_kendall	193
79.36	pool_samples	194
79.37	qmean	194
79.38	range_mean	194
79.39	rmse_	194
79.40	serr	194
79.41	serr1	194
79.42	test_kskew	194
79.43	test_qstd_kskew_optimal_p	194
79.44	wautocorr	195
79.45	wcorr	195
79.46	wcov	195
79.47	wdof	195
79.48	wkurt	195
79.49	wmean	195
79.50	wrms	196
79.51	wserr	196
79.52	wskew	196
79.53	wstd	196
79.54	wvar	196

80 statistics	196
80.1 nangeomean	196
80.2 nangeostd	196
81 statistics/nonparametric-statistics	197
81.1 kernel1d	197
81.2 kernel2d	197
82 statistics	197
82.1 normmoment	197
82.2 normpdf2	197
83 statistics/order-statistics	197
83.1 hodges_lehmann_location	197
83.2 kendall	198
83.3 kendall_to_pearson	198
83.4 mad2sd	198
83.5 madcorr	198
83.6 median2_holder	198
83.7 median_ci	198
83.8 median_man	198
83.9 mediani	199
83.10 nanmadcorr	199
83.11 nanwmedian	199
83.12 nanwquantile	199
83.13 oja_median	199
83.14 qkurtosis	199
83.15 qmoments	200
83.16 qskew	200
83.17 qskewq	200
83.18 qstdq	200
83.19 quantile1_optimisation	200
83.20 quantile2_breckling	200
83.21 quantile2_chaudhuri	200
83.22 quantile2_projected	201
83.23 quantile2_projected2	201
83.24 quantile_envelope	201
83.25 quantile_regression_simple	201
83.26 ranking	201
83.27 spatial_median	201
83.28 spatial_quantile	201
83.29 spatial_quantile2	201
83.30 spatial_quantile3	202
83.31 spatial_rank	202

83.32	spatial_sign	202
83.33	spatial_signed_rank	202
83.34	spearman	202
83.35	spearman_rank	202
83.36	spearman_to_pearson	202
83.37	wmedian	202
83.38	wquantile	203
84	statistics	203
84.1	qstd	203
84.2	quantile_extrap	203
85	statistics/random-number-generation	203
85.1	laplacernd	203
85.2	randc	203
85.3	skewness2param	203
85.4	skewpdf_central_moments	203
85.5	skewrnd	203
85.6	skewrnd2	204
86	statistics	204
86.1	range	204
86.2	resample_with_replacement	204
87	statistics/resampling-statistics/@Jackknife	204
87.1	Jackknife	204
87.2	estimated_STATIC	205
87.3	matrix1_STATIC	205
87.4	matrix2	205
88	statistics/resampling-statistics	205
88.1	block_jackknife	205
88.2	jackknife_moments	205
88.3	moving_block_jackknife	205
88.4	randblockserr	206
88.5	resample	206
89	statistics	206
89.1	scale_quantile_sd	206
89.2	sd_sample_quantiles	206
89.3	skewpdf	207
89.4	trimmed_mean	207
89.5	ttest2_man	207
89.6	ttest_man	207
89.7	ttest_paired	207

89.8	wgeomean	207
89.9	wgeovar	208
89.10	wharmean	208
89.11	wharstd	208
89.12	wharvar	208
90	mathematics	208
90.1	ternary_diagram	208
91	test/master	208
91.1	dat_test_lanczos_3d_k_20_n_40	208
91.2	poisson2d.blk	208
91.3	qr_implicit_givens_2	208
91.4	spectral_derivative_2d	209
91.5	test_2d_eigensolver_hydrogen	209
91.6	test_2d_refine	209
91.7	test_3d_eigensolver_hydrogen	209
91.8	test_FEM	209
91.9	test_Mesh_3d	209
91.10	test_arnoldi	209
91.11	test_arpackc	209
91.12	test_assemble	209
91.13	test_assembly_performance	209
91.14	test_bc_one_sided	210
91.15	test_compare_solvers	210
91.16	test_complete	210
91.17	test_convergence	210
91.18	test_convergence_b	210
91.19	test_df_2d	210
91.20	test_eig_algs	210
91.21	test_eig_inverse	210
91.22	test_eigs_lanczos	210
91.23	test_eigs_lanczos_1	210
91.24	test_eigs_lanczos_2	211
91.25	test_eigs_lanczos_performance	211
91.26	test_fdm	211
91.27	test_fdm_d_vargrid	211
91.28	test_fdm_spectral	211
91.29	test_fem	211
91.30	test_fem_1d	211
91.31	test_fem_1d_higher_order	211
91.32	test_fem_2d_adaptive	211
91.33	test_fem_2d_higher_order	211
91.34	test_fem_3d_higher_order	212

91.35	test_fem_3d_refine	212
91.36	test_fem_b	212
91.37	test_fem_derivative	212
91.38	test_fem_quadrature	212
91.39	test_final	212
91.40	test_fix_substitution	212
91.41	test_forward	212
91.42	test_get_sparse_arrays	212
91.43	test_harmonic_oscillator	212
91.44	test_high_order_fdm_periodic_bc	213
91.45	test_hydrogen_wf	213
91.46	test_ichol	213
91.47	test_interpolation	213
91.48	test_inverse_problem	213
91.49	test_it_vs_exact	213
91.50	test_jama	213
91.51	test_jd	213
91.52	test_jdqz	213
91.53	test_lanczos_2	213
91.54	test_lanczos_biorthogonal	214
91.55	test_laplacian	214
91.56	test_laplacian_non_uniform	214
91.57	test_laplacian_simple	214
91.58	test_mesh_2d_uniform	214
91.59	test_mesh_2d_uniform_2	214
91.60	test_mesh_circle	214
91.61	test_mesh_generation	214
91.62	test_mesh_interpolate	214
91.63	test_mg	214
91.64	test_minres_recycle	215
91.65	test_multigrid	215
91.66	test_nc	215
91.67	test_nonuniform_symmetric	215
91.68	test_pde	215
91.69	test_permutation	215
91.70	test_poison_fem	215
91.71	test_polar	215
91.72	test_potential	215
91.73	test_powers	215
91.74	test_precondition	216
91.75	test_project_rectangle	216
91.76	test_qr	216
91.77	test_quantum_well	216
91.78	test_radial_adaptive	216

91.79	test_radial_confinement	216
91.80	test_radial_fixes	216
91.81	test_refine_2d	216
91.82	test_refine_2d_b	216
91.83	test_refine_3d	216
91.84	test_refine_structural	217
91.85	test_regularisation	217
91.86	test_round_off	217
91.87	test_schrödinger_potentials	217
91.88	test_uniform_mesh	217
91.89	test_vargrid	217
92	test	217
92.1	test_gaussfit3	217
92.2	test_mtimes3x3	217
93	wavelet	218
93.1	continuous_wavelet_transform	218
93.2	cwt_man	218
93.3	example_wavelets	218
93.4	phasewrap	218
93.5	test_cwt_man	218
93.6	test_phasewrap	218
93.7	test_wavelet	218
93.8	test_wavelet2	218
93.9	test_wavelet_analysis	219
93.10	test_wavelet_reconstruct	219
93.11	test_wtc	219
93.12	wavelet	219
93.13	wavelet_reconstruct	219
93.14	wavelet_transform	219
94	mathematics	219
94.1	wrapphase	219

1 calendar

1.1 days_per_month

1.2 isnight

2 mathematics

mathematical functions of various kind

2.1 cast_byte_to_integer

cast byte to integer

3 complex-analysis

operations on complex numbers

3.1 complex_exp_product_im_im

product of the imaginary part of two complex exponentials

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.2 complex_exp_product_im_re

product of the imaginary part of one and the real part of a second complex exponential

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.3 complex_exp_product_re_im

the product has two frequency components

product of the imaginary part of one and the real part of a second complex exponential

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.4 complex_exp_product_re_re

product of the real part of two complex exponentials

$$\begin{aligned} \text{re}(c1 \exp(i o1 x)) * \text{re}(c2 \exp(i o2 x)) = \\ \frac{1}{2} * (\text{real}(c1 * c2 * \exp(i * (n1 + n2) * o * x)) \dots \\ + \text{real}(\text{conj}(c1) * c2 * \exp(i * (n2 - n1) * o * x))) \end{aligned}$$

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

3.5 croots

nth-roots of a complex number

input:
c : complex number
n : order of root
 n must be rational, to obtain n solutions
 otherwise no finite set of solutions exists

r : roots of the complex number

3.6 root_complex

root of a complex number

3.7 test_imroots

4 derivation

derivation of several functions by means of symbolic computation

4.1 derive_acfar1

4.2 derive_ar2param

4.3 derive_arc_length

4.4 derive_fourier_power

4.5 derive_fourier_power_exp

4.6 derive_laplacian_curvilinear

4.7 derive_laplacian_fourier_piecewise_linear

4.8 `derive_logtripdf`

4.9 `derive_smooth1d_parametric`

5 `derivation/master`

5.1 `derive_bc_one_sided`

5.2 `derive_convergence`

5.3 `derive_error_fdm`

5.4 `derive_fdm_poly`

5.5 `derive_fdm_power`

5.6 `derive_fdm_taylor`

5.7 `derive_fdm_vargrid`

5.8 `derive_fem_2d_mass`

5.9 `derive_fem_error_2d`

5.10 `derive_fem_error_3d`

5.11 `derive_fem_sym_2d`

5.12 `derive_grid_constants`

5.13 `derive_interpolation`

5.14 `derive_laplacian`

5.15 `derive_limit`

5.16 `derive_nc_1d`

5.17 `derive_nc_1d_`

5.18 `derive_nc_2d`

5.19 `derive_nonuniform_symmetric`

%

5.20 `derive_richardson`

5.21 `derive_sum`

5.22 `nn`

5.23 `test_derive`

5.24 `test_derive_fdm_poly`

5.25 `test_filter`

5.26 `test_vargrid`

6 derivation

derivation of several functions by means of symbolic computation

6.1 simplify_atan

symbolic simplification of the arcus tangent

7 mathematics

mathematical functions of various kind

7.1 exp10

8 finance

8.1 derive_skewrnd_walsh_paramter

8.2 gbm_cdf

8.3 gbm_fit

8.4 gbm_fit_old

8.5 gbm_inv

8.6 `gbm_mean`

8.7 `gbm_median`

8.8 `gbm_pdf`

8.9 `gbm_simulate`

8.10 `gbm_skewness`

8.11 `gbm_std`

8.12 `gbm_transform_time_step`

8.13 `put_price_black_scholes`

8.14 `skewgbm_simulate`

8.15 `skewrnd_walsh`

9 finance/test

9.1 test_gbm

9.2 test_gbm_pdf

9.3 test_skewrnd_walsh

10 fourier/@STFT

10.1 STFT

class for short time fourier transform

Note: the interval T_i should be set to at least $2 \cdot \max(T)$, as otherwise coefficients

tend to oscillate in the presence of noise

Note: for convenience, the independent variable is labeled as time (t) ,

but the independent variable is arbitrary, so it works likewise in space

10.2 itransform

inverse of the short time fourier transform

10.3 stft_

static wrapper for STFT

10.4 stftmat

transformation matrix for the short time fourier transform

10.5 transform

short time fourier transform

11 fourier

support and analysis functions both for the discrete (fast) fourier transform (dft/fft) and continuous fourier analysis (fourier series)

11.1 amplitude_from_peak

amplitude and standard deviation of the amplitude of a frequency component

represented by a peak in the fourier domain

input :

h : peak height

w : peak width at half height

output:

a : amplitude in real space

s : standard deviation of the frequency (!)

11.2 dftmtx_man

fourier matrix in matlab style with a limited number of rows, columns of higher frequencies are omitted

input :

n : number of samples

nr : number of columns

output :

F : fourier matrix

11.3 example_fourier_window

11.4 fft_derivative

derivative by fourier transform
exponential convergence for periodic functions
results in spurious oscillations for aperiodic functions

input:
x : data, sampled in equal intervals
k : order of the derivative

dx : kth-derivative of x

11.5 fft_man

fast fourier transform for complex input data

input:
F : data in real space

output :

F : fourier transformation of F

11.6 fftsmooth

smooth the fourier transform and determine upper and lower bound
confidence intervals

input :
f :
sfunc : a smoothing function (for example fir convolution with
 rectangular window)
 returns filtered (mean) value and normalized fir window
nf : window length
nsigma : number of standard deviations for confidence intervals

output :
ff : filtered fourier transform
l : lower bound
u : upper bound

11.7 fix_fourier

fill gaps (missing data) by means of fourier extrapolation

fix periodic data series with fourier interpolation

longest gap should not exceed $1/2$ of the shortest time span of interest ($1/\text{cutoff frequency}$)

note: this limit equals the position of first side lobe of the ft of a rectangular window with gap length

11.8 fourier_axis

return axis of frequencies and periods for the discrete fourier transform
as computed by fft (matlab-style)

input:

X : sample locations (equal interval)

L : length of samples

n : number of samples

output :

f : frequencies

T : periods

mask : mask for $1/2$ of the fourier transform
(as both halves are complex conjugates)

N : frequency id

11.9 fourier_cesaro_correction

11.10 fourier_coefficient_piecewise_linear

fourier series coefficients of a piecewise linear function

(not coefficient of discrete fourier transform)

function can be discontinuous between intervals

scales domain length to 2π

input :

l,r : end points of piecewise linear function

lval, rval : values at end points

L : length of domain

n : number of samples/highest frequency
output :
a, b : coefficients for frequency components

11.11 `fourier_coefficient_piecewise_linear_1`

fourier series coefficients of a piecewise linear function
(not coefficient of discrete fourier transform)
function can be discontinuous between intervals
scales domain length to 2π

input :
X : end points of piecewise linear function
Y : values at end points

output :
ab : coefficients for frequency components

11.12 `fourier_coefficient_ramp3`

fourier series coefficient of a ramp

11.13 `fourier_coefficient_ramp_pulse`

fourier series coefficient of a ramp pulses

11.14 `fourier_coefficient_ramp_step`

fourier coefficient of a ramp-step

11.15 `fourier_coefficient_square_pulse`

fourier series coefficients of a square pulse

11.16 `fourier_cubic_interaction_coefficients`

11.17 `fourier_derivative`

coefficients of the derivative of a fourier series
not of discrete fourier transform (fft)

11.18 `fourier_expand`

expand values of fourier series

11.19 `fourier_fit`

fit a fourier series to a set of sample points that are not spaced
in
equal intervals

11.20 `fourier_interpolate`

interpolate samples y sampled at moments (location) t to locations
 t_i

11.21 `fourier_matrix`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.22 `fourier_matrix2`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.23 `fourier_matrix3`

transformation matrix for the continuous fourier transform
this is a matrix with $(2n+1)$ real columns

11.24 `fourier_matrix_exp`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

11.25 `fourier_multiplicative_interaction_coefficients`

11.26 `fourier_power`

powers of a continuous fourier series in sin/cos form

powers of $a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \phi))^p$
phase of first component assumed 0

frequencies higher than 2-omega ignored in input
frequencies higher than 3-omega not computed

11.27 `fourier_power_exp`

powers of the continuous fourier series
 $a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \phi))^p$
phase of first component assumed 0

higher orders than 2 ignored input
higher order than 3 not computed in output

$y = a_0 + \sum (a_j \sin(j\omega t) + b_j \cos(j\omega t))$
 $= \text{Real}(\sum_{i=0}^{\infty} c_i \exp(i\omega t)), c_i = a_i + b_i$

NOT the alternative $\sum_{i=-\infty}^{\infty} \tilde{c}_i$, tile $c_j = 1/2 a_j$
 $+ 1/2i b_j$

11.28 `fourier_predict`

expand a continuous fourier series at times t

11.29 `fourier_quadratic_interaction_coefficients`

11.30 `fourier_range`

approximate range of a continuous Fourier series with 2 components
 $\text{range}(y) = \max(y) - \min(y)$

11.31 `fourier_regress`

fit a continuous Fourier series to a set of sample points not
sampled
at equal intervals

11.32 `fourier_resampled_fit`

fits coefficients of a continuous Fourier transform,
but stores them as resampled values

11.33 `fourier_resampled_predict`

interpolates a continuous Fourier series that has been stored as
values
at their support points

11.34 `fourier_signed_square`

coefficients of the Fourier series of $|\cos a + \cos t|$ ($\cos a + \cos t$)
in general
 $\cos a$ is midrange
 $\cos t$ is tidal variation
c.f. Dronkers

11.35 fourier_transform

continuous fourier transformation of y
(not discrete fourier transformation dft/fft)

input:

b : data sampled at equal intervals
T : length of data in time or space, i.e. position of last
sample if
position of first sample is 0
T_max : maximum period to include

output :

A : fourier matrix
p : fourier transformation of b
tt : TODO

11.36 hyperbolic_fourier_box

11.37 idftmtx_man

inverse matrix for the discrete fourier transform in matlab style
with a limited number of columns, thus ignoring higher frequencies
keep 2nc+1 columns (mean and conj-complex pairs of nc frequencies)

11.38 laplace_2d_pwlinear

solution to the Laplacian in two dimensions for a finite
rectangular domain

with piecewise constant boundary conditions

linear system with 4 unknowns per frequency component

these are coefficients of s,c,sh,ch

(pu*(s + c) + qu*(s' + c'))*(shu + chu) = ru % upper bc

(pd*(s + c) + qd*(s' + c'))*(shd + chd) = rd % lower bc

((sl + cl)*(pl*(shl + chl) + ql*(shl' + chl')) = rl % left
bc

((sr + cr)*(pr*(shr + chr) + qr*(shr' + chr')) = rr % right
bc

least squares with piecewise integration

[x0,p,q,r] piecewise linear polynomials at the boundaries

11.39 nanfft

discrete fourier transform of a data series with gaps

11.40 peaks

peaks of the power spectrum of a discrete fourier transform

rule for peaks: there is no higher value left or right of the "peak"

until the signal drops to $p \cdot y_{\text{peak}}$, $p = 0.5$

works best, when spectrum has been smoothened

input :

f : frequency

y : absolute value of fourier transform (power spectrum)

L : length in space or time of series

output :

a0 : amplitude

s0 : standard deviation (error?) of amplitude

w0 : width of peak

lambda = wave length (period?)

pdx : index of peak

f : frequency (if not given as input)

11.41 roots_fourier

zeros of continuous fourier series

$$f = a_0 + \sum_{j=1}^n a_j \cos(j x) + b_j \sin(j x)$$

11.42 spectral_density

spectral density

11.43 test_complex_exp_product

11.44 test_fourier_filter

11.45 test_idftmtx

12 geometry/@Geometry

12.1 Geometry

12.2 arclength

arc length of a two dimensional curve

8th order accurate

does not require the segments length to vary smoothly

note: the curve can be considered parametric, e.g. $x = x(t)$, $y = y(t)$
and
and $t = t(s)$, but the error term contains derivatives of t ,
thus a non smooth t (strongly varying distance between points)
requires the scaling as done below

12.3 arclength_old

arc length of a two dimensional function

12.4 arclength_old2

arc length of a two dimensional function

12.5 base_point

base point (fusspunkt), i.e. point on a line with shortest distance
to another point

12.6 base_point_limited

base point (Fusspunkt) of a point on a line

12.7 centroid

centroid of a polygone

12.8 cosa_min_max

12.9 cross2

cross product in two dimensions

12.10 curvature

curvature of a function in two dimensions

12.11 ddot

sum of squares of cos of inner angles of triangle

12.12 distance

euclidan distance between two points

12.13 distance2

euclidean distance between two points
this function requires a and b of equal dimensions, or the least
the first pair or second pair to be a scalar

12.14 dot

dot product

12.15 edge_length

edge length

12.16 enclosed_angle

angle enclosed between two lines

12.17 enclosing_triangle

smallest enclosing equilateral triangle with bottom side parallel to
X axis

12.18 hexagon

coordinates of a hexagon, scaled and rotated

12.19 inPolygon

flag points contained in a polygon
much faster than matlab internal function

12.20 inTetra

flag points contained in tetrahedron

12.21 inTetra2

flag points contained in tetrahedron

12.22 inTriangle

flag points contained in triangle
function [flag, c] = inTriangle(P1,P2,P3,P0)

12.23 intersect

intersect between two lines

12.24 lineintersect

intersect of two lines

12.25 lineintersect1

intersect of two lines

12.26 minimum_distance_lines

minimum distance of two lines in three dimensions

12.27 mittenpunkt

mittenpunkt of a triangle

12.28 nagelpoint

nagelpoint of a triangle

12.29 onLine

12.30 orthocentre

orthocentre of triangle

12.31 plumb_line

12.32 poly_area

area of a polygon
function A = poly_area(x,y)

12.33 poly_edges

edges of a polygon

12.34 poly_set

associate point at arbitrary location with a polygon it is contained
in
and assign the value of the polygon to it

12.35 poly_width

width of polygon width holes by surface normals
holes / islands separated with NaN
order of points of outer boundary must be cw
order of points of holes must be ccw
note that this function does not give the true width for expanding
sections
use voronoi polygons for this

12.36 polyxpoly

intersections of two polygons

12.37 `project_to_curve`

closest point on a curve with respect to a point at distance to the curve

12.38 `quad_isconvex`

12.39 `random_disk`

draw random points on the unit disk

12.40 `random_simplex`

random point inside of a triangle

12.41 `sphere_volume`

volume of a sphere

12.42 `tetra_volume`

volume of a tetrahedron

12.43 `tobarycentric`

cartesian to barycentric coordinates

12.44 `tobarycentric1`

cartesian to barycentric coordinates

12.45 tobarycentric2

cartesian to barycentric coordinates

12.46 tobarycentric3

cartesian to barycentric coordinates

12.47 tri_angle

cos of angles of a triangle

12.48 tri_area

angle of a triangle

12.49 tri_centroid

centroid of a triangle

12.50 tri_distance_opposit_midpoint

distance between corner of a triangle and its opposing mid-point

12.51 tri_edge_length

edge length of a triangle

12.52 tri_edge_midpoint

mid point of a triangle

12.53 tri_excircle

excircle of a triangle

12.54 tri_height

height of a triangle

12.55 tri_incircle

incircle of a triangle

12.56 tri_isacute

flag acute triangles

12.57 tri_isobtuse

flag obtuse triangles

12.58 tri_semiperimeter

semiperimeter of a triangle

12.59 tri_side_length

edge length of triangle

13 geometry

13.1 Polygon

Simple 2D polygon class

Polygon properties:

x - x coordinates of polygon

y - y coordinates of polygon

nnodes - number of nodes in the polygon

Polygon methods:

in - checks whether given points lie inside, on the edge, or
outside of the polygon

area - returns the area of the polygon

centerline - computes the centerline of the river

iscw - check whether polygon is clockwise

reverse - reverse the order of the polygon

13.2 bounding_box

bounding box of X

13.3 curvature_1d

curvature of a sampled parametric curve in two dimensions

13.4 cvt

centroidal voronoi tessellation

13.5 deg_to_frac

degree, minutes and seconds to fractions

13.6 ellipse

n-points on an ellipse

13.7 ellipseX

x-coordinates of y-coordinates of an ellipse

13.8 ellipseY

13.9 first_intersect

get first intersection between lines in A and B

13.10 golden_ratio

golden ratio

13.11 hypot3

hypothenuse in 3D

13.12 meanangle

weighted mean of angles

13.13 meanangle2

mean angle

13.14 meanangle3

mean angle

13.15 meanangle4

mean angle

13.16 medianangle

median angle
angle, that has the smallest squared distance to all others

13.17 medianangle2

median angle

input
alpha : x*m, [rad] angle

output
ma : 1*m, [rad] median angle
sa : 1*m, [rad] standard error of median angle for uncorrelated
error

13.18 pilim

limit to +- pi

13.19 streamline_radius_of_curvature

streamline radius of curvature
simplifies when rotate to streamwise coordinates to $R = 1/dv/ds * u$

14 histogram/@Histogram

14.1 2x

14.2 Histogram

14.3 bimodes

14.4 cdf

14.5 cdfS

14.6 chi2test

14.7 cmoment

14.8 cmomentS

14.9 entropy

14.10 entropyS

14.11 iquantile

14.12 kstest

14.13 kurtosis

14.14 kurtosisS

14.15 mean

14.16 meanS

14.17 median

14.18 medianS

14.19 mode

14.20 modeS

14.21 moment

14.22 momentS

14.23 pdf

14.24 quantile

14.25 quantileS

14.26 setup

14.27 skewness

14.28 skewnessS

14.29 stairs

14.30 stairsS

14.31 std

14.32 stdS

14.33 var

14.34 varS

15 histogram

15.1 hist_man

15.2 histadapt

15.3 histconst

15.4 pdf_poly

15.5 plotcdf

15.6 test_histogram

16 linear-algebra

16.1 averaging_matrix_2

16.2 colnorm

norms of columns

16.3 condest_

estimation of the condition number

17 linear-algebra/coordinate-transformation

17.1 barycentric2cartesian

barycentric to cartesian coordinates

17.2 barycentric2cartesian3

convert barycentric to cartesian coordinates

17.3 cartesian2barycentric

cartesian to barycentric coordinates

17.4 cartesian_to_unit_triangle_basis

transform coordinates into unit triangle

17.5 ellipsoid2geoid

17.6 example_approximate_utm_conversion

17.7 latlon2utm

transform latitude and longitude to WGS84 UTM

17.8 latlon2utm_simple

17.9 lowrance_mercator_to_wgs84

convert lowrance coordinates to wgs84

based on spreadsheet by D Whitney King and Patty B at Lowrance

17.10 nmea2utm

convert nmea messages to utm coordinates

17.11 sn2xy

convert sn to xy coordinates

17.12 unit_triangle_to_cartesian

transform coordinates in unit triangle to cartesian coordinates

17.13 utm2latlon

convert wgs84 utm to latitude and longitude

17.14 xy2nt

project all points onto the cross section and assign them nz-coordinates

transform coordinate into N-T reference
rotate coordinate, so that cross section goes along x-axis
then x and y are n and t respectively scaled by width
N and T coordinates

17.15 xy2sn

convert cartesian to streamwise coordiantes

17.16 xy2sn_java

use java port for speed up

17.17 xy2sn_old

transform points from cartesian into streamwise coordinates

NOTE : prefer the java version, this has some problems with round off

18 linear-algebra

18.1 det2x2

2x2 matrix inverse of 2x2 matrices stacked along dim 3

18.2 det3x3

determinant of stacked 3x3 matrices

18.3 det4x4

determinant of stacked 4x4 matrices

18.4 diag2x2

diagonal of stacked 2x2 matrices

18.5 eig2x2

eigenvalues of stacked 2x2 matrices

19 linear-algebra/eigenvalue

19.1 eig_bisection

19.2 eig_inverse

19.3 eig_inverse_iteration

19.4 eig_power_iteration

20 linear-algebra/eigenvalue/jacobi-davidson

20.1 afun_jdm

20.2 davidson

20.3 jacobi_davidson

20.4 jacobi_davidson_qr

20.5 jacobi_davidson_qz

20.6 jacobi_davidson_simple

20.7 jdqr

```
% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal, A*V=W*R+Qschur*E, R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%   V*V=eye(j), Qschur'*V=0,
%   W=A*V, M=V'*W
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
```



```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%    $V^*V = \text{eye}(j)$ ,  $Qschur' * V = 0$ ,  $W' * W = \text{eye}(j)$ ,  $Qschur' * W = 0$ 
%    $(A * V - \tau * V) = W * R + Qschur * E$ ,  $E = Qschur' * (A * V - \tau * V)$ ,  $M = W' * V$ 
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur = [Rschur; zeros(1,k)], Qschur' * MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur,  $AV = A * V - \tau * V$ 
%    $V^*V = \text{eye}(j)$ ,  $W' * W = \text{eye}(j)$ ,  $Qschur' * V = 0$ ,  $Qschur' * W = 0$ ,
%    $(I - Qschur * Qschur') * AV = W * R$ ,  $M = W' * V$ ;  $R = W' * AV$ ;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur = [Rschur; zeros(1,k)], Qschur' * MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
  W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
  W=V*H; V(:,j+1)=[];R=R'*R; M=H(1:j,:)' ;
%===== ARNOLDI (for initializing spaces)
%=====
%===== END ARNOLDI
%=====

% not accurate enough M=Rw'\(M/Rv);
%===== COMPUTE SORTED JORDAN FORM
%=====

% compute vectors and matrices for skew projection
% solve preconditioned system
% 0 step of bicgstab eq. 1 step of bicgstab
% Then x is a multiple of b
% HIST=[0,1];
%   explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnm/snm]];
% sufficient accuracy. No need to update r,u
%   implicit preconditioning
% collect the updates for x in l-space

```

```

% but, do the orth to Q implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnrm/snrm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Q explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow, pause
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down
% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
r=r/rho; rho=1;
% HIST=rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
% HIST = rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow, pause
%----- compute schur form -----
A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults
%% 'v'

```

20.8 jdqr_sleijpen

```
% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal, A*V=W*R+Qschur*E, R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%   V*V=eye(j), Qschur'*V=0,
%   W=A*V, M=V'*W
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%   V*V=eye(j), Qschur'*V=0, W'*W=eye(j), Qschur'*W=0
%   (A*V-tau*V)=W*R+Qschur*E, E=Qschur'*(A*V-tau*V), M=W'*V
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur, AV=A*V-
%   tau*V
%   V*V=eye(j), W'*W=eye(j), Qschur'*V=0, Qschur'*W=0,
%   (I-Qschur*Qschur')*AV=W*R, M=W'*V; R=W'*AV;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```

```

% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
W=V*H; V(:,j+1)=[];R=R'*R; M=H(1:j,:)' ;
%===== ARNOLDI (for initializing spaces)
=====
%===== END ARNOLDI
=====
% not accurate enough M=Rw'\(M/Rv);
%===== COMPUTE SORTED JORDAN FORM
=====
% compute vectors and matrices for skew projection
% solve preconditioned system
% 0 step of bicgstab eq. 1 step of bicgstab
% Then x is a multiple of b
% HIST=[0,1];
% explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% implicit preconditioning
% collect the updates for x in l-space
% but, do the orth to Q implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Q explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow, pause
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down
% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow,% pause
r=r/rho; rho=1;
% HIST=rho;
% HIST=[HIST;rho];
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';

```

```

    plot(J,HIST(:,1),'*'); drawnow,% pause
% HIST = rho;
% HIST=[HIST;rho];
    HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
    plot(J,HIST(:,1),'*'); drawnow, pause
% HIST = rho;
% HIST=[HIST;rho];
    HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
    plot(J,HIST(:,1),'*'); drawnow, pause
%----- compute schur form -----
    A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults
%% 'v'

```

20.9 jdqr_vorst

```

% Read/set parameters
% Initiate global variables
% Return if eigenvalueproblem is trivial
% Initialize V, W:
%   V,W orthonormal, A*V=W*R+Qschur*E, R upper triangular
% The JD loop (Standard)
%   V orthogonal, V orthogonal to Qschur
%   V*V=eye(j), Qschur'*V=0,
%   W=A*V, M=V'*W
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
    Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   Both V and W orthonormal and orthogonal w.r.t. Qschur
%    $V^*V = \text{eye}(j)$ ,  $Qschur' * V = 0$ ,  $W' * W = \text{eye}(j)$ ,  $Qschur' * W = 0$ 
%    $(A * V - \tau * V) = W * R + Qschur * E$ ,  $E = Qschur' * (A * V - \tau * V)$ ,  $M = W' * V$ 
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   V W AV.
%   Both V and W orthonormal and orthogonal w.r.t. Qschur,  $AV = A * V - \tau * V$ 
%    $V^*V = \text{eye}(j)$ ,  $W' * W = \text{eye}(j)$ ,  $Qschur' * V = 0$ ,  $Qschur' * W = 0$ ,
%    $(I - Qschur * Qschur') * AV = W * R$ ,  $M = W' * V$ ;  $R = W' * AV$ ;
%
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
  Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```



```

% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
% The JD loop (Harmonic Ritz values)
%   W orthonormal, V and W orthogonal to Qschur,
%   W'*W=eye(j), Qschur'*V=0, Qschur'*W=0
%   W=(A*V-tau*V)-Qschur*E, E=Qschur'*(A*V-tau*V),
%   M=W'*V
% Compute approximate eigenpair and residual
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
%
%
%
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Check for convergence
% Expand the partial Schur form
Rschur=[Rschur;zeros(1,k)],Qschur'*MV(u)]; k=k+1;
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Expand preconditioned Schur matrix PinvQ
% Check for shrinking the search subspace
% Solve correction equation
% Expand the subspaces of the interaction matrix
W=V*Q; V=V(:,1:j)/R; E=E/R; R=eye(j); M=Q(1:j,:)' /R;
W=V*H; V(:,j+1)=[];R=R'*R; M=H(1:j,:)' ;
%===== ARNOLDI (for initializing spaces)
=====
%===== END ARNOLDI
=====

% not accurate enough M=Rw'\(M/Rv);
%===== COMPUTE SORTED JORDAN FORM
=====

% accepted separation between eigenvalues:
% no preconditioning
% solve left preconditioned system
% compute vectors and matrices for skew projection
% precondition and project r
% solve preconditioned system
% no preconditioning
% solve two-sided expl. precondition. system
% compute vectors and matrices for skew projection
% precondition and project r
% solve preconditioned system
% "unprecondition" solution
%%% u(:,j+1)=Atilde*u(:,j)

```

```

%%%% r(:,j+1)=Atilde*r(:,j)
%----- compute schur form -----
A*Q=Q*S, Q'*Q=eye(size(A));
% transform real schur form to complex schur form
%----- find order eigenvalues -----
%----- reorder schur form -----
%----- compute qz form -----
%----- sort eigenvalues -----
%----- sort qz form -----
% i>j, move ith eigenvalue to position j
% determine dimension
% defaults

```

20.10 jdqz

```

% Read/set parameters
% Return if eigenvalueproblem is trivial
% Initialize target, test space and interaction matrices
% V=RepGS(Qschur,V); [AV,BV]=MV(V); %%% more stability??
% W=RepGS(Zschur,eval(testspace)); %%% dangerous if sigma~lambda
% Solve the preconditioned correction equation
% Expand the subspaces and the interaction matrices
% Check for stagnation
% Solve projected eigenproblem
% Compute approximate eigenpair and residual
%=== an alternative, but less stable way of computing z =====
% display history
% save history
% check convergence
% EXPAND Schur form
% Expand preconditioned Schur matrix MinvZ=M\Zschur
% check for conjugate pair
% To detect whether another eigenpair is accurate enough
% restart if dim(V)> jmax
% Initialize target, test space and interaction matrices
% additional stabilisation. May not be needed
% V=RepGS(Zschur,V); [AV,BV]=MV(V);
% end add. stab.
% Solve the preconditioned correction equation
% expand the subspaces and the interaction matrices
% Check for stagnation
% compute approximate eigenpair
% Compute approximate eigenpair and residual
% display history
% save history
% check convergence
% expand Schur form

```

```

% ZastQ=Z'*Q0
% the final Qschur
% check for conjugate pair
% t perp Zschur, t in span(Q0,imag(q))
% To detect whether another eigenpair is accurate enough
% restart if dim(V)> jmax
%===== END JDQZ
=====
%=====
%===== PREPROCESSING
=====
%=====
%===== ARNOLDI (for initial spaces)
=====
%% then precondition=I and target = 0: apply Arnoldi with A
%===== END ARNOLDI
=====
%=====
%===== POSTPROCESSING
=====
%=====
%===== SORT QZ DECOMPOSITION INTERACTION MATRICES
=====
%===== COMPUTE SORTED JORDAN FORM
=====
%===== END JORDAN FORM
=====
%===== OUTPUT
=====
%=====
%===== UPDATE PRECONDITIONED SCHUR VECTORS
=====
%=====
%=====
%===== SOLVE CORRECTION EQUATION
=====
%=====
% solve preconditioned system
%=====

```

```

%===== LINEAR SOLVERS
=====
%=====

% [At,Bt]=MV(x); At=theta(2)*At-theta(1)*Bt;
% xtol=norm(r-At+Z*(Z'*At))/norm(r);
%===== Iterative methods
=====

% 0 step of bicgstab eq. 1 step of bicgstab
% Then x is a multiple of b
% HIST=[0,1];
% explicit preconditioning
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% implicit preconditioning
% collect the updates for x in l-space
% but, do the orth to Z implicitly
% compute norm in l-space
% HIST=[HIST;[nmv,rnorm/snorm]];
% sufficient accuracy. No need to update r,u
% Do the orth to Z explicitly
% In exact arithmetic not needed, but
% appears to be more stable.
% plot(HIST(:,1),log10(HIST(:,2)+eps),'*'), drawnow
% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
%=====

% 0 step of gmres eq. 1 step of gmres
% Then x is a multiple of b
HIST=1;
% Lucky break-down
HIST=[HIST;(gamma~=0)/sqrt(rho)];
% Lucky break-down
% solve in least square sense
HIST=log10(HIST+eps); J=[0:size(HIST,1)-1]';
plot(J,HIST(:,1),'*'); drawnow
%===== END SOLVE CORRECTION EQUATION
=====

%=====

%===== BASIC OPERATIONS
=====
%=====

y(1:5,1), pause
%===== COMPUTE r AND z
=====

```

```

% E*u=Q*sigma, sigma(1,1)>sigma(2,2)
%===== END computation r and z
=====
%=====

%===== Orthogonalisation
=====
%=====

%===== END Orthogonalisation
=====
%=====

%===== Sorts Schur form
=====
%=====

    kappa=max(norm(A,inf)/max(norm(B,inf),1.e-12),1);
    kappa=2^(round(log2(kappa)));
%----- compute the qz factorization -----
%----- scale the eigenvalues -----
%----- sort the eigenvalues -----
%----- swap the qz form -----
% repeat SwapQZ if angle is too small
%=====

%=====

% i>j, move ith eigenvalue to position j
% compute q s.t. C*q=(t(i,1)*S-s(i,1)*T)*q=0
C*P=Q*R
    check whether last but one diag. elt r nonzero
C*q
% end computation q
%===== END sort QZ decomposition interaction matrices
=====
%=====

%===== INITIALIZATION
=====
%=====

%=====

% defaults          %%%% search for 'xx' in fieldnames
%% 'ma'
%% 'sch'
%% 'to'
%% 'di'

```

```

% jmin=nselect+p0 %%%% 'jmi'
% jmax=jmin+p1 %%%% 'jma'
%% 'te'
%% 'pai'
%% 'av'
%% 'tr'
%% 'fix'
%% 'ns'
%% 'ch'
%% 'lso'
%% 'ls_m'
%% 'ls_t'
%% 'ls_e'
%% 'ty'
%% 'l_'
%% 'u_'
%% 'p_'
%% 'sca'
%% 'v0'
initiation
'standard'
'harmonic'
'searchspace'
%=====

% or Operator_Form=3 or Operator_Form=5???
%=====

%===== DISPLAY FUNCTIONS
=====
%=====

%=====

%=====

%=====

```

20.11 mfunc_jdm

20.12 mgs

20.13 minres_

20.14 mv_jacobi_davidson

21 linear-algebra

21.1 first

21.2 gershgorin_circle

range of eigenvalues determined by the gershgorin circle theorem

21.3 haussdorff

haussdorf dimension

box counting: count rectangles passed through by line (covered by polygon)

Koch snow flake 3:4 -> 1.2619

Kantor set 2:3, (4:9) -> 0.6309

quadrat 4:2, 9:3, 16:4 -> 2

21.4 eig2x2

reconstruct matrix from eigenvalue decomposition

21.5 inv2x2

2x2 inverse of stacked matrices

21.6 inv3x3

21.7 inv4x4

inverse of stacked 4x4 matrices

22 linear-algebra/lanczos

22.1 arnoldi

22.2 arnoldi_new

22.3 eigs_lanczos_man

22.4 lanczos

22.5 lanczos_

22.6 lanczos_biorthogonal

22.7 lanczos_biorthogonal_improved

22.8 `lanczos_ghep`

22.9 `mv_lanczos`

22.10 `reorthogonalise`

22.11 `test_lanczos`

23 `linear-algebra/linear-systems`

23.1 `gmres_man`

`break on convergence`

23.2 `minres_recycle`

24 `linear-algebra`

24.1 `lpmean`

`mean of pth-power of a`

24.2 `lpnorm`

`norm of lth-power of a`

24.3 matvec3

matrix-vector product of stacked matrices and vectors

24.4 max2d

maximum value and i-j index for matrix

24.5 mpoweri

approximation of A^p , where p is not integer by quadratic interpolation

24.6 mtimes2x2

24.7 mtimes3x3

product of stacked 3x3 matrices

24.8 nannorm

norm of a vector, skips nan-values

24.9 nanshift

shift vector, but set out of range values to NaN

24.10 nl

number rows (lines) of a matrix

analogue to unix nl command

24.11 normalise

normalise a vector or the columns of a matrix
note that the columns are independently normalised, and hence not
necessarily
orthogonal to each other use the gram schmidt algorithm for this (
qr or orth)

24.12 normalize1

normalize columns in x to [-1,1]

24.13 normrows

24.14 orth2

make matrix A orthogonal to B

24.15 orth_man

orthogonalize the columns of A

24.16 orthogonalise

make x orthogonal to Y

24.17 paddext

padd values to vactor
not suitable for noisy data
order = 0 : constant extrapolation (hold)
order = 1 : linear extrapolation

24.18 paddval1

padd values at end of x

24.19 paddval2

padd values to x

25 linear-algebra/polynomial

25.1 chebychev

chebycheff polynomials

25.2 piecewise_polynomial

evaluate piecewise polynomial

25.3 roots1

roots of linear functions

25.4 roots2

roots of quadratic function
 $c_1 x^2 + c_2 x + c_3 = 0$

25.5 roots2poly

25.6 roots3

25.7 roots4

25.8 test_roots4

25.9 vanderi_1d

vandermonde matrix of an integral

26 linear-algebra

26.1 randrot

random rotation matrix

26.2 right

get right column by shifting columns to left
extrapolate rightmost column

26.3 rot2

rotation matrix from angle

26.4 rot2dir

rotation matrix from direction vector

26.5 rot3

26.6 rotR

26.7 rownorm

26.8 simmilarity_matrix

26.9 spnorm

frobenius norm

26.10 spzeros

allocate a sparze matrix of zeros

26.11 test_roots3

26.12 transform_minmax

26.13 transpose3

transpose stacked 3x3 matrices

26.14 transposeall

27 logic

bitwise operations on integers

27.1 bitor_man

bitwise OR of the numbers of the columns of A

input:

A (positive integer)

28 master/plot

28.1 attach_boundary_value

28.2 cartesian_polar

28.3 img_vargrid

28.4 plot_basis_functions

28.5 plot_convergence

28.6 plot_dof

28.7 plot_eigenbar

28.8 `plot_error_estimation`

28.9 `plot_error_estimation_2`

28.10 `plot_error_fem`

28.11 `plot_fdm_kernel`

28.12 `plot_fdm_vs_fem`

28.13 `plot_fem_accuracy`

28.14 `plot_function_and_grid`

28.15 `plot_hat`

28.16 `plot_hydrogen_wf`

28.17 `plot_mesh`

28.18 `plot_mesh_2`

28.19 `plot_refine`

28.20 `plot_refine_3d`

28.21 `plot_runtime`

28.22 `plot_spectrum`

28.23 `plot_wavefunction`

29 `master/ported`

29.1 `assemble_2d_dphi_dphi`

29.2 `assemble_2d_phi_phi`

29.3 `assemble_3d_dphi_dphi`

29.4 assemble_3d_phi_phi

29.5 dV_2d_

29.6 derivative_2d

29.7 derivative_3d

29.8 element_neighbour_2d

29.9 prefetch_2d_

29.10 promote_2d_3_10

29.11 promote_2d_3_15

29.12 promote_2d_3_21

29.13 promote_2d_3_6

29.14 promote_3d_4_10

29.15 promote_3d_4_20

29.16 promote_3d_4_35

29.17 vander_2d

29.18 vander_3d

30 master/sandbox

30.1 adapt

30.2 assoc_laguerre

30.3 assoc_legendre

30.4 c23

31 master/sandbox/cg

31.1 cg

31.2 cg_coef_to_poly

31.3 errmat

31.4 lanczos

31.5 laplacian_2d

31.6 test_cg_eigs

31.7 test_lanczos

32 master/sandbox

32.1 condition_number_higher_order

32.2 confinement_dat

32.3 `convergence_2d_3d`

32.4 `convergence_matrix_powers`

32.5 `cut_out`

32.6 `derivative_2d`

32.7 `derivative_3d`

32.8 `dummy`

32.9 `eig_error`

32.10 `eigs_fix`

32.11 `energy_level`

32.12 `equalise`

32.13 example_int64

Basic operations

Matrix multiplication
Timing

33 master/sandbox/fem-matlab

33.1 boundary_circle

33.2 boundary_rectangle

33.3 geometry_circle_with_hole

33.4 geometry_rectangle

34 master/sandbox

34.1 fem_2d_estimate_error

34.2 fem_assemble_scratch

34.3 fem_s

34.4 fourier_h

34.5 grad_2d

34.6 grad_3d

34.7 gradient

34.8 harmonic_oscillator

34.9 hydrogen_2d_analytic

34.10 hydrogen_boxed

34.11 hydrogen_boxed_old

34.12 hydrogen_wave

% Hydrogen atom

34.13 hydrogen_wf

34.14 ichol_man

34.15 known_eigenvalue

34.16 kron_man

34.17 laguerre

34.18 laplacian_arbitrary_order_old

34.19 laplacian_convergence

34.20 laplacian_cut_out

34.21 laplacian_cylindrical

34.22 laplacian_non_uniform_old

34.23 `laplacian_polar`

34.24 `laplacian_simple`

34.25 `lderivative_3d`

34.26 `list_dat`

34.27 `matlab-horner`

34.28 `mesh_to_grid_2d_3`

34.29 `mg_mat`

34.30 `mv`

34.31 `orth2`

34.32 `partial_derivative_2d`

34.33 `partition_function`

34.34 `partition_function_old`

34.35 `poisson`

34.36 `poisson_fem`

34.37 `potential`

34.38 `powerc`

34.39 `quick_newihbour`

34.40 `radial`

34.41 `radial_convergence`

34.42 `radial_wafefunction`

34.43 refine_2d

34.44 refine_3d

34.45 relerr

34.46 restore_cw

34.47 runtime_bm

34.48 rydberg

34.49 s_old

34.50 snorm

34.51 spherical_harmonic

34.52 split_eig

34.53 sum1

34.54 sum3

35 master/sandbox/summation

35.1 acc

35.2 add

35.3 ape

35.4 mmul_accurately

35.5 sum_kahan

35.6 sum_pairwise

35.7 test_sum

36 master/sandbox

36.1 test_convergence_ill_conditioned

36.2 test_fem_1d

36.3 test_fem_2d

36.4 test_fem_3d

36.5 test_increase

36.6 test_lanczos_shift

36.7 test_ldl

36.8 test_power

36.9 trefethen_p8_fdm

36.10 wavefunc

36.11 xgrid

37 number-theory

37.1 ceiln

floor to leading n-digits

37.2 digitsb

number of digits with respect to specified base

37.3 floorn

floor to n-digits

37.4 iseven

true for even numbers in X

37.5 multichoosek

all combinations of length k from set values with repetitions
c.f. nchoosek, combinations without repetition

input :

x : scalar integer or vector of arbitrary numbers
k : length of subsets

output :

if x scalar : number of combinations
if x vector : the exact combinations

37.6 nchoosek_man

vectorised binomial coefficient
 $b = N! / K! (N-K)!$

37.7 pythagorean_triple

pythagorean triple

37.8 roundn

round to n digits

38 numerical-methods/differentiation

38.1 derivative1

first derivative on variable mesh
second order accurate

38.2 derivative2

second derivative on a variable mesh

39 numerical-methods/finite-difference

39.1 cdiff

differences of columns of X
degree = 1 : central first order differences
degree = 2 : central second order differences

39.2 `cdiffb`

differences of columns of X
degree = 1 : central first order differences
degree = 2 : central second order differences
TODO use difference matrix function for simplicity

39.3 `cmean`

single gaussian smoothing step with kernel $1/4*[1,2,1]$

39.4 `derivative_matrix_1_1d`

finite difference matrix of first derivative in one dimensions

39.5 `derivative_matrix_2_1d`

finite derivative matrix of second derivative in one dimension

39.6 `derivative_matrix_2d`

finite difference derivative matrix in two dimensions

39.7 `derivative_matrix_curvilinear`

derivative matrix on a curvilinear grid

39.8 `derivative_matrix_curvilinear_2`

derivative matrix on a two dimensional curvilinear grid
the grid has not necessarily to be orthogonal

39.9 difference_kernel

difference kernels for equispaced grids
c.f. Computing the Spectrum of the Confined Hydrogen Atom, Kastner,
2012

39.10 distmat

distance matrix for a 2 dimensional rectangular matrix

39.11 gradpde2d

objective function gradiend on two dimensional regular grid
numeric gradient for non-linear least squares optimisation
of a PDE on a rectangular grid
 $x_* = \min(f(x))$
 $f = (v(x) - v(x_*))^2 = f(x) + A \, dx + O(dx^2)$
 $a_{ij} = df_i/dx_j$

39.12 laplacian

39.13 laplacian_fdm

finite difference matrix of the laplacian
BC

39.14 left

left element of vector, leftmost column is extrapolated

39.15 lrmean

mean of the left and right element

40 numerical-methods/finite-difference/master

40.1 `fdm_adaptive_grid`

40.2 `fdm_adaptive_refinement_old`

40.3 `fdm_assemble_d1_2d`

40.4 `fdm_assemble_d2_2d`

40.5 `fdm_confinement`

40.6 `fdm_d_vargrid`

40.7 `fdm_h_unstructured`

40.8 `fdm_hydrogen_vargrid`

40.9 `fdm_mark_unstructured_2d`

40.10 `fdm_plot`

40.11 `fdm_plot_series`

40.12 `fdm_refine_2d`

40.13 `fdm_refine_3d`

40.14 `fdm_refine_unstructured_2d`

40.15 `fdm_schroedinger_2d`

40.16 `fdm_schroedinger_3d`

40.17 `relocate`

41 numerical-methods/finite-difference

41.1 `mid`

mid point between neighbouring vector elements

41.2 pwmid

segment end point to segment mid point transformation for regular 1
d grids

41.3 ratio

ratio of two subsequent values

41.4 steplength

step length of a vector if it were equispaced

41.5 swapoddeven

swap odd and even elements in a vector

41.6 test_derivative_matrix_2d

41.7 test_derivative_matrix_curvilinear

41.8 test_difference_kernel

42 numerical-methods/finite-element

42.1 Mesh_2d_java

42.2 Tree_2d.java

42.3 assemble_1d_dphi_dphi

42.4 assemble_1d_phi_phi

42.5 assemble_2d_dphi_dphi.java

42.6 assemble_2d_phi_phi.java

42.7 assemble_3d_dphi_dphi.java

42.8 assemble_3d_phi_phi.java

42.9 boundary_1d

42.10 boundary_2d

42.11 boundary_3d

42.12 `check_area_2d`

42.13 `circmesh`

42.14 `cropradius`

42.15 `display_2d`

42.16 `display_3d`

42.17 `distort`

42.18 `err_2d`

42.19 `estimate_err_2d_3`

42.20 `example_1d`

42.21 `example_2d`

42.22 `explode`

42.23 `fem_2d`

42.24 `fem_2d_heuristic_mesh`

42.25 `fem_get_2d_radial`

42.26 `fem_interpolation`

42.27 `fem_plot_1d`

42.28 `fem_plot_1d_series`

42.29 `fem_plot_2d`

42.30 `fem_plot_2d_series`

42.31 `fem_plot_3d`

42.32 fem_plot_3d_series

42.33 fem_plot_confine_series

42.34 fem_radial

adaptive grid
constant grid

42.35 flip_2d

42.36 get_mesh_arrays

42.37 hashkey

43 numerical-methods/finite-element/int

43.1 int_1d_gauss

43.2 int_1d_gauss_1

43.3 int_1d_gauss_2

43.4 int_1d_gauss_3

43.5 int_1d_gauss_4

43.6 int_1d_gauss_5

43.7 int_1d_gauss_6

43.8 int_1d_gauss_lobatto

43.9 int_1d_nc_2

43.10 int_1d_nc_3

43.11 int_1d_nc_4

43.12 int_1d_nc_5

43.13 int_1d_nc_6

43.14 int_1d_nc_7

43.15 int_1d_nc_7_hardy

43.16 int_2d_gauss_1

43.17 int_2d_gauss_12

43.18 int_2d_gauss_13

43.19 int_2d_gauss_16

43.20 int_2d_gauss_25

43.21 int_2d_gauss_3

43.22 int_2d_gauss_33

43.23 int_2d_gauss_6

43.24 int_2d_gauss_7

43.25 int_2d_gauss_9

43.26 int_2d_nc_10

43.27 int_2d_nc_15

43.28 int_2d_nc_21

43.29 int_2d_nc_3

43.30 int_2d_nc_6

43.31 int_3d_gauss_1

43.32 int_3d_gauss_11

43.33 int_3d_gauss_14

43.34 int_3d_gauss_15

43.35 int_3d_gauss_24

43.36 int_3d_gauss_4

43.37 int_3d_gauss_45

43.38 int_3d_gauss_5

43.39 int_3d_nc_11

43.40 int_3d_nc_4

43.41 int_3d_nc_6

43.42 int_3d_nc_8

44 numerical-methods/finite-element

44.1 interpolation_matrix

44.2 mark

44.3 mark_1d

44.4 mesh_1d_uniform

44.5 mesh_3d_uniform

44.6 mesh_interpolate

44.7 neighbour_1d

44.8 old

44.9 pdeeig_1d

44.10 pde eig_2d

44.11 pde eig_3d

44.12 polynomial_derivative_1d

44.13 potential_const

44.14 potential_coulomb

44.15 potential_harmonic_oscillator

44.16 project_circle

44.17 project_rectangle

44.18 promote_1d_2_3

44.19 promote_1d_2_4

44.20 promote_1d_2_5

44.21 promote_1d_2_6

44.22 quadrilaterate

44.23 recalculate_regularity_2d

44.24 refine_1d

44.25 refine_2d_21

44.26 refine_2d_structural

44.27 regularity_1d

44.28 regularity_2d

44.29 regularity_3d

```
{      T = [1 2 3 4];  
}
```

44.30 relocate_2d

44.31 test_circmesh

44.32 test_hermite

44.33 tri_assign_points

44.34 triangulation_uniform

44.35 vander_1d

van der Monde matrix

44.36 vanderd_1d

44.37 vanderi_1d

45 numerical-methods/finite-volume/@Advection

45.1 Advection

FVM treatment of the Advection equation

45.2 dot_advection

advection equation

46 numerical-methods/finite-volume/@Burgers

46.1 burgers_split

viscous Burgers' equation,
mixed analytic and numerical derivative in frequency space
by splitting scheme
 $u_t = -(0.5*u^2)_x + c*u_{xx}$

46.2 dot_burgers_fdm

viscous burgers' equation
 $u_t = -d/dx (1/2*u^2) + c d^2/dx^2 u_{xx}$

46.3 dot_burgers_fft

viscous Burgers' equation in frequency space
 $u_t + (0.5*u^2)_x = c*u_{xx}$

47 numerical-methods/finite-volume/@Finite_Volume

47.1 Finite_Volume

finite volume method for partial differential equations 1+1
dimensions
(time and space)

47.2 apply_bc

apply boundary conditions

47.3 solve

solve the the PDE by successively stepping in time
this is a trivial implmentation with constant step length
severity of diffusive error depends on dt/dx-ratio
stability depends on wave height
printf('Progress %2.1f%% %2.1fs\n',100*(t-Ti
(1))/(Ti(2)-Ti(1)),t_real);

47.4 step_split_strang

step in time, treat inhomogeneous part by Strang splitting
this scheme is not suitable for stationary solutions, for example
steady shallow water flow

47.5 step_unsplit

step in time, without splitting the inhomogeneous term

48 numerical-methods/finite-volume/@Flux_Limiter

48.1 Flux_Limiter

class of flux limiters

48.2 beam_warming

beam warming scheme
low resolution
note: works only if sign of eigenvalues point into the same
direction according to RL

48.3 fromm

fromme limiter
low res

48.4 lax_wendroff

lax wendroff scheme
second order accurate, but no tvd
this is effectively not a limiter
eq. 6.39 in randall, leveque

48.5 minmod

min-mod schock limiter

48.6 monotized_central

monotonized central flux limiter

48.7 muscl

muscl flux limiter

48.8 superbee

superbee limiter

48.9 upwind

godunov scheme
godunov, first order accurate

48.10 vanLeer

van Leer limiter

49 numerical-methods/finite-volume/@KDV

49.1 dot_kdv_fdm

korteweg de vries equation
 $u_t + (0.5*u^2)_x = c*u_{xxx}$

49.2 dot_kdv_fft

korteweg de vries equation
compute derivatives in frequency space
 $u_t + (0.5*u^2)_x = c*u_{xxx}$

49.3 kdv_split

korteweg de vries equation in frequency space,
derivative treated by splitting scheme

50 numerical-methods/finite-volume/@Reconstruct_Average_E

50.1 Reconstruct_Average_Evolve

Reconstruct Average Evolve Finite Volume Method for treatment of
1+1D pdes

McCronack Scheme

err = $O(dt^2) + O(dx^2)$, except as discontinuities

error:

```
h_xxx(3:end-2) = 1/dx^3*( -0.5*h(1:end-4) + h(2:end-3) - h(4:
    end-1) + 0.5*h(5:end) );
th = -1/6*dx^2*qh_.*(1 - (qh_*dt/dx).^2).*h_xxx;
```

50.2 advect_highres

single time step for the reconstruct evolve algorithm

50.3 advect_lowres

single time step
low resolution

51 numerical-methods/finite-volume

51.1 Godunov

Godunov, upwind method for systems of pdes

51.2 Lax_Friedrich

Lax-Friedrich-Method
for hyperbolic conservation laws
 $\text{err} = O(\Delta t) + O(\Delta x)$
 $|a \Delta t / \Delta x| < 1$

51.3 Measure

51.4 Roe

non linear roe solver for the SWE (randall, leveque 15.3.1)

The roe solver guarantess:

- A is diagonalisable with real eigenvalues (15.12)
- can be determined by a closed formula
- is an efficient replacement for true Rieman solver

51.5 fv_swe

wrapper for solving SWE

51.6 staggered_euler

forward euler method with staggered grid

51.7 staggered_grid

staggered grid approximation to the SWE

52 numerical-methods

52.1 grid2quad

extract rectangular elements of a structured grid
in form of an unstructured quad-mesh format

53 numerical-methods/integration

53.1 cumintL

cumulative integral from left to right

53.2 cumintR

cumulative integral from right to left

53.3 int_trapezoidal

integrate y along x with the trapezoidal rule

54 numerical-methods/interpolation/@Kriging

54.1 Kriging

```
class for Kriging interpolation
```

54.2 estimate_semivariance

```
estimate the parameter of the semivariance model for Kriging
interpolation
    % set up the regression matrix and solve for
    parameters
```

54.3 interpolate_

```
interpolate with Kriging method
```

```
this function may interpolate several quantities per coordinate,
using the same variogram, if the semivariance of the quantities
differs,
the user may prefer to estimate the semivariance and interpolate
each quantity
individually
```

```
Xs : source point coordinates
Vs : value at source points
Xt : target point coordinates
Vt : value at target points
E2t : squared interpolation error at target points
```

55 numerical-methods/interpolation/@RegularizedInterpolator1

55.1 RegularizedInterpolator1

```
class for regularized interpolation (Thikonov) on a 1D mesh
```

55.2 init

```
initialize the interpolator with a set of sampling points
```

56 numerical-methods/interpolation/@RegularizedInterpolator

56.1 RegularizedInterpolator2

```
class for regularized interpolation on an unstructures mesh (  
    interpolation)
```

56.2 init

```
initialize the interpolator with a set of point samples
```

57 numerical-methods/interpolation/@RegularizedInterpolator

57.1 RegularizedInterpolator3

```
class for regularized interpolation (Tikhonov) on a triangulation  
(unstructured mesh)
```

57.2 init

```
initialize the interpolator with a set of sampling points
```

58 numerical-methods/interpolation

58.1 IDW

```
spatial averaging by inverse distance weighting
```

58.2 IPoly

```
polynomial interpolation class
```


58.3 IRBM

```
interpolate by the radial basis function method
    fprintf(1,'Progress IRBM: %d%%\n',round(100*
        idx/size(Xi,1)));
```

58.4 ISparse

```
sparse interpolation class
```

58.5 Inn

```
nearest neighbour interpolation
```

58.6 Interpolator

```
interpolator super-class
    fprintf(1,'Progress: %f%% %fs\n',100*
        idx/size(Xt,1),t);
```

58.7 fixnan

```
fill nan-values in vector with gaps
```

58.8 idw1

```
spatial average by inverse distance weighting
```

58.9 idw2

```
spatial average by inverse distance weighting
```

58.10 inner2outer

linear interpolation of segment mid point to grid points at segment
ends
assumes equal grid spacing

58.11 inner2outer2

interpolate from element (segment) centres to edge points

58.12 interp1_limited

interpolate values, but not beyond a certain distance
this function is idempotent, i.e. it will not extrapolate over into
gaps
exceedint the limit and thus not spuriously extend the series when
called a second time on the same data

58.13 interp1_man

interpolate

58.14 interp1_save

make interpolation save to round off errors
the matlab internal interpolation suffers from rounding errors,
which
are unacceptable when values of X and Y are large (for example UTM
coordinates)
this normalization prevents this

58.15 interp1_slope

quadratic interpolation returning value and derivative(s)

58.16 `interp1_smooth`

58.17 `interp1_unique`

matlab fails to interpolate, when x values are not unique
this function makes the values unique before use

58.18 `interp2_man`

nearest neighbour interpolation in two dimensions

58.19 `interp_angle`

interpolate an angle

58.20 `interp_fourier`

interpolation by the fourier method

58.21 `interp_fourier_batch`

batch interpolation by the fourier interpolation

58.22 `interp_sn`

interpolate along streamwise coordinates
This gives similar result to setting aspect ratio for sN to
infinity,
but not quite, as the input point set is not dense (scale for sN to
infinity does not work)
 `sdx = sdx(sdx_);`

58.23 interp_sn2

interpolation in streamwise coordinates

58.24 interp_sn3

58.25 interp_sn_

58.26 limit_by_distance_1d

smooth subsequent values along a curve such that
 $v(x_0+dx) < v(x_0) + (ratio-1)*dx$
if v is the edge length in a resampled polygon, then $v_i/v_{(i+1)} <$
ratio
 $ratio^1 = \exp(a*1)$

58.27 resample1

interpolation along a parametric curve with variable step width

58.28 resample_d_min

resample a function

58.29 resample_vector

resample a track so that velocity vectors do not run into each
other

58.30 test_interp1_limited

59 numerical-methods

59.1 inverse_complex

60 numerical-methods/ode

60.1 bvp2_check_arguments

60.2 bvp2c

solve system of non-linear second order odes (in more than one variable)
as boundary value problems

odefun provides ode coefficients c:

$$c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4) \\ c_1 y'' + c_2 y' + c_3 y + c_4 = c_4$$

subject to the boundary conditions

bcfun provides v and p and optionally q, so that:

$$b_1 y + b_2 y' = f \\ q(x,1)*(p(x,1) y_l(x) + p(x,2) y_l'(x) \\ + q(x,2)*(p(x,1) y_r(x) + p(x,2) y_r'(x) = v(x)$$

where q weighs the waves travelling from left to right and right to left (default [1 1])

60.3 bvp2c2

solve second order boundary value problem via roots of the characteristic polynomial

input:

x : [nx1] discretized domain
n : number of vertices
nxc = n-1 : number of segments

```

bc : struct : boundary condition
    bc.p(1)*y(0) + bc.pd(2)*y'(0) = bc.val(1)
    bc.p(2)*y(L) + bc.pd(2)*y'(L) = bc.val(2)

```

output:

```

A : [2*nxc x 2*ns] discrctisation matrix
rhs : [2*nxc x 1] right hand size

```

```

y = A^-1 rhs

```

60.4 bvp2fdm

solve system of non-linear second order odes (in more than one variable)

as boundary value problems by the finite difference method

odefun provides ode coefficients c:

```

c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4)
c_1 y'' + c_2 y' + c_3 y + c_4 = 0

```

subject to the boundary conditions

bcfun provides v and p and optionally q, so that:

```

b_1 y + b_2 y' = f
q(x,1)*( p(x,1) y_l(x) + p(x,2) y_l'(x)
+ q(x,2)*( p(x,1) y_r(x) + p(x,2) y_r'(x) = v(x)

```

where q weighs the waves travelling from left to right and right to left (default [1 1])

60.5 bvp2wavetrain

solve second order boundary value problem by repeated integration

60.6 bvp2wavetwopass

two pass solution for the linearised wave equation

solve first for the wave number k, and then for y

60.7 ivp_euler_forward

solve intial value problem by the euler forward method

60.8 ivprk2

solve initial value problem by the two step runge kutta method

60.9 ode2_matrix

transformation matrix of second order ode
to left and right going wave

```
c = odefun(x)
c1 y'' + c2' y + c3 y == 0
y = y_p + y_m, left and right going wave
d/dx [y_p, y_m] = A*[y_m, y_p]
```

60.10 ode2characteristic

second order odes
transmitted and reflected wave

60.11 step_trapezoidal

single trapezoidal step

60.12 test_bvp2

61 numerical-methods/optimisation

61.1 armijo_stopping_criterion

armijo stopping criterion for optimizations

61.2 astar

astar path finding algorithm

61.3 binsearch

binary search on a line

61.4 bisection

bisection

61.5 box1

test objective function for optimisation routines

61.6 box2

61.7 cauchy

61.8 cauchy2

solve non-linear system by cuachy's method
slower than quadratic optimisation, but does not require a hessian
fun : objective function, returns
 f : scalar, objective function value
 g : nx1, gradient
x : nx1, initial position
opt : options

61.9 directional_derivative

directional (projected) derivative
d : derivative, highest first
p : series expansion around x0

61.10 dud

optimization by the dud algorithm

61.11 extreme3

extract maxima by quadratic approximation from sampled function `val`
(`t`)
intended to be called after `[mval, mid] = max(val)` for refinement
of
location and maximum

input
`t` : sampling time (uniformly spaced)
`v` : values at sampling times
output:
`tdx` : index where extremum should be computed
`t0` : location of the extremum
`val0` : value of extremum

$v'(dt0) = 0$ and $v''(dt0)$ determines type of extremum

61.12 extreme_quadratic

61.13 ftest

61.14 fzero_bisect

61.15 fzero_newton

61.16 grad

numerical gradient

61.17 hessian

numerical hessian

61.18 hessian_from_gradient

numerical hessian from gradient

61.19 hessian_projected

numerical hessian projected to one dimension

61.20 line_search

bisection routine

61.21 line_search2

bisection method

fun : objective funct
x0 : start value
f0 : objective function value at x0
g : gradient at x0
p : search direction from x0 (p = g for steepest descend)
h : initial step length (default 1)
lb : lower bound for x
up : upper bound for x

61.22 line_search_polynomial

```
polynomial line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

61.23 line_search_polynomial2

```
cubic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

61.24 line_search_quadratic

```
quadratic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

61.25 line_search_quadratic2

```
quadratic line search
```

61.26 line_search_wolfe

line search by wolfe method
c.f.: OPTIMIZATION THEORY AND METHODS - Nonlinear Programming, Sun,
Yuan

61.27 ls_bgfs

least squares by the bgfs method

61.28 ls_broyden

least squares by the broyden method
for rectangular / non symmetric systems
Numerical Optimization nokedal
Practical Methods of Optimization fletcher
c.f. gerber 1981
c.f. fletcher 1978 (more advanced, not used here)
c.f. Kelley 1999 ch. 4

BGFS:
Broyden 1965
Fletcher 1970
Goldfarb 1970
Shanno 1970

61.29 ls_generalized_secant

least squares by the secant method
Barnes, 1965
Wolfe, 1959
Fletcher 1980, 6.3
seber 2003
gerber

61.30 nlcg

non-linear conjugate gradient
input:
x : nx1 start vectort

opt : struct options
fdx : gradient constraint

61.31 nlls

non-linear least squares

61.32 picard

picard iteration

61.33 poly_extrema

extrema of a polynomial

61.34 quadratic_function

evaluate quadratic function in higher dimensions

61.35 quadratic_programming

optimize by quadratic programming

61.36 quadratic_step

single step of the quadratic programming

61.37 rosenbrock

rosenbrock test function

61.38 `sqrt_heron`

Heron's method for the square root

61.39 `test_directional_derivative`

61.40 `test_dud`

61.41 `test_fzero_newton`

61.42 `test_line_search_quadratic2`

61.43 `test_ls_generalized_secant`

61.44 `test_nlcg_6_order`

61.45 `test_nlls`

```
f = w*(p*abs(x-1).^4) + w*(1-p)*abs(x-1).^2;
```

62 numerical-methods/pde

62.1 `laplacian2d_fundamental_solution`

63 numerical-methods/piecewise-polynomials

63.1 Hermite1

hermite polynomial interpolation in 1d

63.2 hp2_fit

fit a hermite polynomial
coefficients are derivative free
x0 : left point of first segment
x1 : right point of last segment
n : number of segments
x : sample x-value
val : sample y-value
c : coefficients (values at points, no derivatives)

63.3 hp2_predict

prediction with pw hermite polynomial
c are values at support points

63.4 hp_predict

predict with piecewise hermite polynomial

63.5 hp_regress

fit piecewise hermite polynomial
coefficients are values and derivatives

63.6 lp_count

lagrangian basis for interpolation
count number of valid samples

63.7 `lp_predict`

lagrangian basis piecwie interpolation, predictor

63.8 `lp_regress`

63.9 `lp_regress_`

64 `regression/@PolyOLS`

64.1 `PolyOLS`

class for polynomial least squares

64.2 `coefftest`

64.3 `detrend`

detrending by polynomial regression

64.4 `fit`

fit a polynomial function
like `polyfit`, but returns parameter error estimates
TODO automatically activate `scaleflag`

64.5 `fit_`

fit a polynomial function

64.6 predict

predict polynomial function values

64.7 predict_

64.8 slope

slope by linear regression

65 regression/@PowerLS

65.1 PowerLS

class for power law regression

65.2 fit

fit a power law
like polyfit, but returns parameter error estimates

65.3 predict

predict with power law
S2 = diag((A*obj.C)*A');
L = Y - S;
U = Y + S;

65.4 predict_

66 regression/@Theil

66.1 Theil

Kendal-Theil-Sen robust regression

66.2 detrend

linear detrending of a set of samples by the Theil-Senn Slope

66.3 fit

fit slope and intercept to a set of sample with the Theil-Sen method

c : confidence interval $c = 2*ns*normcdf(1)$ for ns-sigma intervals

param : itercept and slope

P : confidence interval

66.4 predict

predict values and confidence intervals with the Theil-Sen method

66.5 slope

fit the slope with the Theil-Sen method

67 regression

linear and non-linear regression

67.1 Theil_Multivariate

extension of the Theil-Senn regression to higher dimensions by means of the Gauss-Seidel iteration

67.2 areg

regression using the pth-fraction of samples with smallest residual

67.3 ginireg

gini regression

67.4 hesssimplereg

hessian, gradient and objective function value of the simple
regression
 $\text{rhs} = p(1) + p(2) x + \text{eps}$

67.5 l1lin

solve $\|Ax - b\|_{L1}$ by means of linear programming

67.6 lsq_sparam

parameter covariance of the least squares regression

fun : model function for predtiction
b : sample values
 $f(p) = b$
p : parameter at point of evaluation (preferably optimum)

67.7 polyfitd

fit a polynomial of order n to a set of sampled values and sampled
values
of the derivative

x0 must contain at least for conditioning as otherwise the
intercept
cannot be determined

67.8 regression_method_of_moments

fit linear function $\|a \ b \ x = y\|_{L2}$ by the method of moments
 $y + \epsilon = \alpha + \beta x$

67.9 robustlinreg

fit a linear function by splitting the x-values at their median
 $(\text{med}(y_{\text{left}}) - \text{med}(y_{\text{right}})) / (\text{med}(x_{\text{left}}) - \text{med}(x_{\text{right}}))$
this approach performs poorly compared to the theil-senn operator

67.10 theil2

Theil senn-estimator for two dimensions (glm)

67.11 theil_generalised

generalization of the Theil-Senn operator to higher dimensions,
for arbitrary functions such as polynomials and multivariate
regression
either higher order polynomials or glm
c.f. "On theil's fitting method", Pegoraro, 1991

67.12 total_least_squares

total least squares

67.13 weighted_median_regression

weighted median regression
c.f. Scholz, 1978

68 set-theory

68.1 issubset

test if set B is subset of A in $O(n)$ -runtime

A : first set

B : second set

P : set of primes (auxiliary)

69 signal-processing

69.1 acf_effective_sample_size

effective sample size from acf

69.2 acf_genton

autocorrelation function

69.3 acfar1

Autocorrelation function of the finite AR1 process

$$\begin{aligned} a_k &= 1/(n-k) \sum x_{i+1} + (x_i + x_{i+k})\mu + \mu^2 \\ &= r^k + 1/n \sum_{ij} + 1/n \\ &\text{pause} \end{aligned}$$

69.4 acfar1_2

autocorrelation of the ar1 process

69.5 acfar2

impulse response of the ar2 process

69.6 acfar2_2

autocorrelation of the ar2 process
 $X_i + a_1 X_{i-1} + a_2 X_{i-2} = 0$

69.7 ar1_cutoff_frequency

69.8 ar1_effective_sample_size

effective sample size correction for autocorrelated series

69.9 ar1_mse_mu_single_sample

standard error of a single sample of an ar1 correlated process

69.10 ar1_mse_pop

variance of the population mean of a single realisation around zero

$$E[(\mu_N - 0)^2] = E[\mu_N^2]$$

69.11 ar1_mse_range

mean standard error of the mean of a range of values taken from an
ar1 process

69.12 ar1_spectrum

spectrum of the ar1 process

69.13 ar1_to_tikhonov

convert ar1 correlation to tikhonovs lambda

69.14 ar1_var_factor

variance correction factor for an autocorrelated finite process
n : [1 .. inf] population size
m : [1 .. n] samples size
rho : [-1 < rho < 1 (for convergence)] correlation of samples

69.15 ar1_var_factor_

variance of an autocorrelated finite process

69.16 ar1_var_range2

variance of sub sample starting at the end of the series
from the finite length first order autocorrelated process
$$s2 = 1/m^2 \sum_i^m \sum_j^m \rho^{-|i-j|}$$

69.17 ar1delay

approximate acf by the ar1 process
acf: autocovariance or autocorrelation function
nf : skip first samples (for mixed geometric-arithmetic series (ARMA))

69.18 ar1delay_old

autocorrelation of the residual

69.19 ar2conv

coefficients of the ar2 process determined from the two leading correlations
of the acf [1,r1,r2,...]

69.20 ar2dof

effective samples size for the ar2 process

69.21 ar2param

ar2 parameter estimation from first two terms of acf

```
acf = [1 a1 a2 ...]
```

69.22 asymwin

creates asymmetrical filter windows

filter will always have negative weights

69.23 autocorr_fft

autocorrelation function

69.24 bandpass

bandpass filter

69.25 bandpass2

bandpass filter

69.26 bartlett

Effective sample size factor for bartlett window

c.f. thiebaux

c.f spectral analysis-jenkins, eq. (6.3.27)

```
c = acf
```

note: results seems always to be 1 too low

T : reduction factor for dof

for ar1 with $a = \rho^k = \exp(-k/L)$, $T = 2L$

69.27 bartlett_spectrogram

bartlett spectrogramm
TODO sliding window

69.28 bin1d

bin values of v sampled at x into bins bounded by "edges"
apply function v to it

69.29 bin2d

bin values of V sampled at X and Y into the grid structured grid ex
,ey
apply function func to all values in the bin
func = mean : default
func = sum : non-normalized frequency histogram in 2D

69.30 binormrnd

generate two correlated normally distributed vectors

69.31 conv1_man

convolutions with padding

69.32 conv2_man

convolution in 2d

69.33 conv2z

69.34 conv30

convolve with rectangular window of length n
circular boundaries

69.35 conv_

convolution of a with b

69.36 conv_centered

convolve x with filter window f
when length of f is even, this guarantees a symmetric result (no
off by one
displacement) by making the length of f odd at first

69.37 convz

69.38 cosexpdelay

69.39 csmooth

smooth recursively with [1,2,1]/4 kernel

69.40 daniell_window

Daniell window for smoothing the power spectrum
c.f. Daniell 1946
Bloomfield 2000
meko 2015

69.41 danielle_window

danielle fourier window

69.42 db2neper

convert decibel to neper

69.43 db2power

power ratio from db

69.44 derive_danielle_weight

69.45 derive_limit_0_acfar

69.46 detect_peak

detect peaks in a vector
requires function value to fall to p*max before new value is
allowed

69.47 digital_low_pass_filter

design coefficients of a low pass filter with specified cut of
frequency
and sampling period
analogue low pass with pole at $s = -\omega_c = 1/\tau = 1/RC$
 $H_a = \tau/(\tau + s) = 1/(1 + \omega_c s)$

69.48 doublesum_ij

double sum of r^i

69.49 effective_sample_size_to_ar1

convert effective sample size to ar1 correlation

69.50 filt_hodges_lehman

69.51 filter1

filter along one dimension

69.52 filter2

filter columns of x (matlab does only support vector input)

69.53 filter_

invalidate values that exceed n-times the robust standard deviation

69.54 filteriir

filter adcp t-n data over time

v : nz,nt : values to be filtered

H : nt,1 : depth of ensemble

last : nt,1 : last bin above bottom that can be sampled without
side lobe interference

nf : scalar : number of reweighted iterations

when samples

- distance to bed is reference (advantageous for near-bed suspended
transport)

TODO for wash load: distance to surface is more relevant
interpolate depending on z

when depth changes, neighbouring indices do not correspond to same
relative position in the water column
relative position in the column (s-coordinate) smoothes values
near the bed: absolute distance to bed is chosen
near surface: absolute distance to surface is chosen
-> cubic transformation of index

faster and avoid aliasing (smoothing along z)
resample ensemble to same number of bins in S -> filter ->
resample back
use nonlinear transform z-s coordinates
-> resampling has to be local (Hi -> H-filtered)

filtered profile coordinates to sample coordinates
zf -> zi (special transform)
corresponding indices and fractions
filtration step (update of hf and vf)
sample coordinates to updated profile coordinates
(the inverse step is actually not necessary)
write filtered value

69.55 filterp

69.56 filterp1

fir filter with some fancy extras

69.57 filterstd

69.58 firls_man

design finite impulse response filter by the least squares method

69.59 flattopwin

the flat top window

69.60 frequency_response_boxcar

frequency response of a boxcar filter

69.61 freqz_boxcar

frequency response of a boxcar filter

69.62 gaussfilt1

filter data series with a gaussian window

69.63 hanchangewin

hanning window for change point detection

69.64 hanchangewin2

hanning window for change point detection

69.65 hanwin

hanning filter window

69.66 hanwin_

hanning filter window

69.67 highpass

high pass filter

69.68 kaiserwin

kaiser filter window

69.69 kalman

Kalman filter

69.70 lanczoswin

Lanczos window

69.71 last

lake tail, but for matrices

69.72 lowpass

low pass filter

69.73 lowpass2

design low pass filter with cutoff-frequency f1

69.74 lowpass_iir

iir-low pass

69.75 lowpass_iir_symmetric

two-sided iir low pass filter (for symmetry)

69.76 lowpassfilter2

low-pass filter of data

69.77 maxfilt1

69.78 meanfilt1

moving average filter with special treatment of the boundaries

69.79 medfilt1_man

moving median filter, supports columnwise operation

69.80 medfilt1_man2

moving median filter with special treatment of boundaries

69.81 medfilt1_padded

median filter with padding

69.82 medfilt1_reduced

median filter with padding

69.83 mid_term_single_sample

variance of single sample, mid term

69.84 minfilt1

69.85 mu2ar1

error variance of the mean of the finite length ar1 process

$(\mu)^2 = (\sum \text{epsi})^2 = \sum_i \sum_j \text{epsi}_i \text{epsi}_j = \sum_{ii}(\rho, n)/n^2$
this has the limit s^2 for $\rho \rightarrow 1$

69.86 mysmooth

69.87 nanautocorr

autocorrelation with nan-values

69.88 nanmedfilt1

medfilt1, skipping nans

69.89 neper2db

convert neper to db

69.90 peaks_man

peaks of a periodogram

69.91 polyfilt1

polynomial filter,
can be achieved by iteratively processing the data with
a mean (zero-order) filter

69.92 qmedfilt1

medfilt1, after fitting a quadratic polynomial

69.93 randar1

generate random ar1 process
e1 = randar1(sigma,p,n,m)

69.94 randar1_dual

draw random variables of two correlated ar1 processes

69.95 randar2

generate ar2 process

69.96 randarp

randomly generate the instance of an ar-p process

69.97 range_window

range of values within a certain range of indices (window)

69.98 rectwin

rectangular window

69.99 `recursive_sum`

69.100 `select_range`

69.101 `smooth1d_parametric`

smooth position of $p_0=x_0,y_0$ between $p_1=x_1,y_1$ and $p_2=x_2,y_2$,
so that distance to p_1 and p_2 becomes equal
and the chord length remains the same

69.102 `smooth2`

smooth vectos of X

69.103 `smooth_man`

69.104 `smooth_parametric`

smooth a parametric function given in x-y coordinates
`matvec2x2(R,[dxc;dyc])`

69.105 `smooth_parametric2`

parametrically smooth the curve

69.106 `smooth_with_splines`

69.107 smoothfft

filter with fast fourier transform

69.108 spectrogram

spectrogram

69.109 std_window

moving block standard deviation

69.110 sum_i_lag

sum of ar1 matrix with lag
 $\sum_{i=1}^n \rho^{|i-k|}$

69.111 sum_ii

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^n \rho^{|i-j|}$
this is for the variance, take square root for the standard
deviation factor

69.112 sum_ii_

69.113 sum_ij

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^m r^{|i-j|}$

69.114 sum_ij_

69.115 `sum_ij_partial_`

69.116 `sum_multivar`

`sum of matrix entries of bivariate ar1 process`

69.117 `test_acfar1`

69.118 `test_acfar1_2`

69.119 `test_acfar1_3`

69.120 `test_acfar1_4`

69.121 `test_acfar2`

69.122 `test_ar1_var_factor`

69.123 `test_ar1_var_factor_2`

69.124 `test_ar1_var_mu_single_sample`

69.125 test_ar1_var_pop

69.126 test_ar1_var_pop_1

69.127 test_ar1delay

69.128 test_bivariate_covariance_term

69.129 test_convexity

69.130 test_lanczoswin

69.131 test_madcorr

69.132 test_randar1

69.133 test_randar1_multivariate

69.134 test_randar2

69.135 test_sum_ij

69.136 test_sum_multivar

69.137 test_trifilt1

69.138 test_wautocorr

69.139 test_wavelet_transform

69.140 test_wordfilt

69.141 test_xar1_mid_term

69.142 tikhonov_to_ar1

convert coefficient of the tikhonov regularization to correlatioon
of the ar1 process

69.143 trapwin

trapezoidal filter window

69.144 `trifilt1`

filter with triangular window

69.145 `triwin`

triangular filter window

69.146 `triwin2`

triangular filter window

69.147 `varar1`

error variance of a single sample of a finite length ar1 process
with respect to the mean, averaged over the population

69.148 `welch_spectrogram`

welch spectrogram

69.149 `wfilt`

filter with window

69.150 `winbandpass`

filter with bandpass

69.151 `window_make_odd`

69.152 winfilt0

filter with window

69.153 winlength

window length for desired cutoff frequency
power at f_c is halved
 $H(wf) = 1/\sqrt{2} H(f)$
if the filter window were used as a low pass filter
note: the user should prefer a windowed ideal low pass filter
TODO, relate this to DOF

69.154 wmeanfilt

mean filter with window

69.155 wmedfilt

median filter with window

69.156 wordfilt

weighted order filter

69.157 wordfilt_edgeworth

weighed order filter

69.158 xar1

69.159 `xcorr_man`

cross correlation of two sampled ar1 processes

70 `sorting`

70.1 `sort2`

sort two numbers

70.2 `sort2d`

sort elements of matrix in X
returns row and column index of sorted values

71 `special-functions`

71.1 `bessel_sphere`

spherical Bessel function of the first kind

71.2 `digamma_man`

71.3 `hankel_sphere`

spherical Hankel function for the far field (incident plane wave)
first kind

71.4 hermite

probabilistic's hermite polynomial by recurrence relation

input :
n : order
x : value

output:
f : $H_n(x)$
df : $d/dx H_n(x)$

71.5 legendre_man

legendre polynomials

71.6 neumann_sphere

spherical Neumann function
Bessel function of the second kind

72 statistics

72.1 atan_s2

stadard deviation of the arcus tangens by means of taylor expansion

72.2 beta_mode_to_parameter

transform modes (mean and sd) to paramets of the beta function

72.3 coefficient_of_determination

72.4 conditional_expectation_normal

72.5 correlation_confidence_pearson

confidence intervals of the correlation coefficient
c.f. Fischer 1921

73 statistics/distributions

73.1 PDF

class for quasi-distributions from a set of sampling points

73.2 binorm_separation_coefficient

separation coefficient of a bimodal normal distribution

73.3 binormcdf

bio-modal gaussian distribution

73.4 binormfit

fit sum of two normal distribution to a histogram

73.5 binormpdf

73.6 edgeworth_cdf

edgeworth expansion of an unknown cumulative distribution
with mean μ , standard deviation σ , and third and fourth
cumulants
c.f. Rao 2010

73.7 edgeworth_pdf

probability density of and unknown distribution
with mean μ , standard deviation σ , and third and fourth
cumulants
c.f. Rao 2010

73.8 logn_mode2param

transform modes (μ, σ) to parameters of the log normal
distribution

73.9 logn_param2mode

transform parameters to mode (μ, σ) for the log normal
distribution

73.10 lognpdf_

log normal distribution called by modes rather than parameters

73.11 pdfsample

pdf from sample distribution
Note: better use kernel density estimates

73.12 t2cdf

Hotelling's T-squared cumulative distribution

73.13 t2inv

inverse of Hotelling's T-squared cumulative distribution

74 statistics

74.1 example_standard_error_of_sample_quantiles

74.2 f_var_finite

reduction of variance when sampling from a finite population
without replacement

74.3 gamma_mode_to_parameter

transform modes (μ, sd) to parameters of the gamma distribution

74.4 gaussfit3

74.5 gaussfit_quantile

74.6 hodges_lehmann_correlation

hodges_lehmann correlatoon coefficient
c.f. Shamos 1976
c.f. Bickel and Lehmann 1976
c.f. rousseeuw 1993
c.f. Shevlyakov 2011

74.7 hodges_lehmann_dispersion

dispersion determined by the hodges lehman method
asymptotic efficiency of dispersion estimates:
standard deviation: $E(s - \hat{s})/s = 2/\sqrt{2n} \sim 0.707/\sqrt{n}$
(100%)
hodges lehmann dispersion $E(s - \hat{s})/s = (\pi/3)^2 / (\sqrt{2n}) \sim$
 $0.775/\sqrt{n}$ (91%)

mad $E(s - \hat{s})/s \sim 1.17 s/\sqrt{n}$
 (60%)
 c.f. Shamos 1976
 c.f. Bickel and Lehmann 1976
 c.f. rousseeuw 1993
 nb: rousseeuw uses the 25th percentile, which is more efficient for
 small sample sizes

75 statistics/information-theory

75.1 akaike_information_criterion

akaike information criterion

 serr : rmse of model prediction
 n : effective sample size
 k : number of parameters

 c.f. akaike (1974)
 c.f. sugiura 1978

75.2 bayesian_information_criterion

bayesian information criterion

76 statistics

76.1 kurtncdf

76.2 kurtnpdf

76.3 kurtosis_bias_corrected

bias corrected kurtosis

76.4 limit

limit a by lower and upper bound

76.5 logfactorial

approximate log of the factorial

76.6 loglogpdf

76.7 lognfit_quantile

76.8 logskewcdf

76.9 logskewpdf

77 statistics/logu

77.1 lambertw_numeric

lambert-w function

77.2 logtrialtcdf

pdf of a logarithmic triangular distribution

77.3 logtrialtinv

inverse of the logarithmic triangular distribution

$$= (d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) - \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a)} - 1/\log(a)) c^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) + d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) + b \log(c) - d F \log(a) \log(c))) / (\log(a) - \log(b)))$$
$$x = (d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) - \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a)} - 1/\log(a)) c^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) + d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) + b \log(c) - d F \log(a) \log(c))) / (\log(a) - \log(b)))$$

77.4 logtrialtmean

mean of the logarithmic triangular distribution

77.5 logtrialtpdf

density of the logarithmic triangular distribution

77.6 logtrialtrnd

77.7 logtricdf

cumulative distribution of the logarithmic triangular distribution

77.8 logtriinv

invere of the logarithmic triangular distribution

77.9 logtrimean

mean of the logarithmic triangular distribution

77.10 logtripdf

probability density of the logarithmic triangular distribution

77.11 logtirnd

77.12 logucdf

probability density of the logarithmic uniform distribution

77.13 logucm

central moments of the log-uniform distribution

77.14 loguinv

inverse of the log-uniform distribution

77.15 logumean

mean of the log-uniform distribution

77.16 logupdf

pdf of the log uniform distribution

77.17 logurnd

random numbers following a log-uniform distribution

77.18 loguvar

variance of the log-uniform distribution

77.19 medlogu

median of the log-uniform distribution

77.20 test_logurnd

77.21 tricdf

cumulative distribution of the log-triangular distribution

77.22 triinv

inverse of the triangular distribution

77.23 trimedian

median of the triangular distribution

77.24 tripdf

probability density of the triangular distribution

77.25 trirnd

random numbers of the triangular distribution

78 statistics

78.1 maxnnormals

expected maximum of n normal variables
c.f. Wolperts
this is the median, not the mean of the maximum!
see median of gumbel

78.2 midrange

mid range of columns of X

78.3 minavg

solution of the minimum variance problem
minimise the variance of the weighted sum of n-independent
random variables with equal mean and individual variance

78.4 mode_man

79 statistics/moment-statistics

79.1 autocorr_man3

autocorrelation of the columns of X

79.2 autocorr_man4

autocorrelation for x if x is a vector, or individually for the columns of x if x is a matrix

c.f. box jenkins 2008 eq. 2.1.12

Note that it is faster to compute the acf in frequency space as done in the matlab internal function

79.3 autocorr_man5

autocorrelation of the columns of X

79.4 blockserr

estimate the standard error of potentially sequentially correlated data
by blocking
block length should be sufficiently larger than correlation length and sufficiently smaller than data length
this uses a sliding block approach, which reduces the variation of the error estimate

79.5 comoment

non-central higher order moments of the multivariate normal distribution

c.f. Moments and cumulants of the multivariate real and complex Gaussian distributions

note : there seem to be some typos in the original paper,
for x^4 c_{ii}^2 , the square seems to be missing

μ : $n \times 1$ mean vector

C : $n \times n$ covariance matrix

k : $n \times 1$ powers of variables in moments

79.6 corr_man

correlation of two vectors

79.7 cov_man

covariance matrix of two vectors

79.8 dof

mininum number of support points
for a polynomial of degree order in dim dimensions

79.9 edgeworth_quantile

inverse edgeworth expansion
c.f. cornis fisher 1937
c.f. Rao 2010
c.f. 2.50 in hall
CHERNOZHUKOV 3.3

79.10 effective_sample_size

effective sample size of the weighted mean of uncorrelated data
c.f. Kish

79.11 f_correlation

correction factor for standard error of the mean of n ar1-
correlated iid samples

79.12 f_finite

reduction factor of standard error for sampling from a finite
distribution
without replacement

79.13 lmean

mean of $x.^l$, not of abs

79.14 lmoment

l-moment of vector x

79.15 maskmean

mean of the masked values of X

79.16 masknanmean

79.17 mean1

mean of x

79.18 mean_man

mean and standard error of X

79.19 mse

mean squared error of residual vector res
this is de-facto the std for an unbiased residual

79.20 nanautocorr_man1

autocorrelation of a vector with nan-values

79.21 nanautocorr_man2

autocorrelation of a vector with nan-values

79.22 nanautocorr_man4

compute autocorrelation for x if x is a vector, or individually
for the
columns of x if x is a matrix
box jenkins 2008 eq. 2.1.12
TODO nan is problematic!
Note that it is faster to compute the acf in frequency space
as done in the matlab internal function

79.23 nancorr

(co)-correlation matrix when samples a NaN

79.24 nancumsum

cumulative sum, setting nan values to zero

79.25 nanlmean

mean of the l-th power of the absolute value of x

79.26 nanr2

coefficient of determination when samples are invalid

79.27 nanrms

root mean square value when sample contains nan-values

79.28 nanrmse

root mean square error from vector of residuals
this is de-facto the std for an unbiased residual

79.29 nanserr

standard error of x with respect to mean when x contains nan values

79.30 nanwmean

weighted mean
min_x sum w (x-mu)^2 => mu = sum(wx)/sum(w)
varargin can be dim
function [mu serr] = nanwmean(w,x)

79.31 nanwstd

weighed standard deviation

79.32 nanwvar

weighted variance of columns, corrected for degrees of freedom (bessel)

$$s^2 = \text{sum}(w*(x-\text{sum}(wx)/\text{sum}(w))^2)/\text{sum}(w)$$

79.33 nanxcorr

79.34 pearson

pearson correlation coefficient

79.35 pearson_to_kendall

conversion of pearson to kendall correlation coefficient
c.f. Kruskal 1958

79.36 pool_samples

pooled mean and standard deviation of several groups of different size, mean and standard deviation

79.37 qmean

trimmed mean

79.38 range_mean

79.39 rmse_

root mean square error computed from a residual vector
this is de-facto the std for an unbiased residual

79.40 serr

standard error of the mean of a set of uncorrelated samples

79.41 serr1

79.42 test_qskew

79.43 test_qstd_qskew_optimal_p

79.44 wautocorr

autocorrelation for x if x is a vector, or individually for the columns of x if x is a matrix
samples can be weighted

c.f. box jenkins 2008 eq. 2.1.12

c.f. autocorr_man4

Note that it is faster to compute the acf in frequency space as done in the matlab internal function

79.45 wcorr

correlation of two vectors when samples are weighted

79.46 wcov

covariance of two vectors when samples are weighted

79.47 wdof

effective degrees of freedom for weighted samples

79.48 wkurt

kurtosis with weighted samples

79.49 wmean

weighted mean

$\min_x \sum w (x - \mu)^2 \Rightarrow \mu = \sum(wx) / \sum(w)$

varargin can be dim

function [mu serr] = wmean(w,x)

79.50 wrms

weighted root mean square error

79.51 wserr

weighted root mean square error

79.52 wskew

skewness of a weighted set of samples

79.53 wstd

weighed standard deviation

79.54 wvar

weighted variance of columns, corrected for degrees of freedom (
bessel)
variance of the weighted sample mean of samples with same mean (but
not necessarily same variance)
 $s^2 = \text{sum } (w^2(x - \text{sum}(wx))^2)$

 $s2_mu$: error of mean, $s2_mu$: sd of prediction

80 statistics

80.1 nangeomean

80.2 nangeostd

geometric standard deviation ignoring nan-values

81 statistics/nonparametric-statistics

81.1 kernel1d

X : ouput x axis bins
xi : samples along x
m : number of bins in X
fun : kernel function
pdf : propability density of xi

81.2 kernel2d

kernel density estimate in two dimensions

82 statistics

82.1 normmoment

expected norm of $x.^n$, when values x in x are iid normal with mu
and sigma

82.2 normpdf2

pdf of the bivariate normal distribution

83 statistics/order-statistics

83.1 hodges_lehmann_location

hodges lehman location estimator

Asymptotic rms efficiency of location estimte:

mean: $1 \text{ s}/\sqrt{n}$
hodges lehman: $\sqrt{\pi/3} * s \sim 1.0233 \text{ s}/\sqrt{n}$
median: $\pi/2 \text{ s}/\sqrt{n} \sim 1.25 \text{ s} / \sqrt{n}$

83.2 kendall

kendall correlation coefficient

83.3 kendall_to_pearson

convert kendall rank correlation coefficient to the person product
moment
correlation coefficient

c.f. Kruska, 1985

83.4 mad2sd

transform median absolute deviation to standard deviation
for normal distributed values

83.5 madcorr

proxy correlation by median absolute deviation

83.6 median2_holder

83.7 median_ci

median and its confidence intervals under assumption of normality
 $se_me = \sqrt{1/2 \pi} \cdot 1.25331 \cdot sd/\sqrt{n}$

83.8 median_man

median and confidence intervals
c is a P value for the confidence interval,
default is 0.95 (2-sigma)
median of the columns of X

83.9 mediani

index of median, if median is not unique, any of the values is chosen

83.10 nanmadcorr

proxy correlation by median absolute deviation

83.11 nanwmedian

weighted median, skips nan-values

83.12 nanwquantile

weighted quantile, skips nan values

83.13 oja_median

two dimensional oja median

note: the multivariate median is not unique

oja 1983, for extension to multivariate function, see chaudhri

83.14 qkurtosis

kurtosis computed for quantiles

Note : this is a measurement of shape-tailedness and yields the same value for the

normal distribution as "kurtosis"

However, this is a separate statistic and hence requires different

methods for calculating P-values and hypothesis testing

83.15 qmoments

moments estimated from quantiles

83.16 qskew

skewness estimated from quantiles

Note : this is a measurement of shape-symmetry and yields the same value for the skew-normal distribution as "skewness"
However, this is an own statistic and hence requires different methods for calculating P-values and hypothesis testing

83.17 qskewq

skewness estimated by quantiles

83.18 qstdq

proxy standard deviation determined by quantiles

83.19 quantile1_optimisation

83.20 quantile2_breckling

quantile regression

83.21 quantile2_chaudhuri

quantile regression

83.22 quantile2_projected

quantile in two dimensions

83.23 quantile2_projected2

spatial quantile for chosen direction

83.24 quantile_envelope

83.25 quantile_regression_simple

simple quantile regression

83.26 ranking

ranking for spearman statistics

83.27 spatial_median

c.f. Oja 2008

is this the same as the oja simplex median (c.f. small 1990)?

83.28 spatial_quantile

spatial quantile

83.29 spatial_quantile2

spatial quantile

83.30 spatial_quantile3

spatial quantile

83.31 spatial_rank

unsigned rank

83.32 spatial_sign

spatial sign

83.33 spatial_signed_rank

signed rank

Note: this is only a true rank if X is normal with zero mean,
arbitrary variance

83.34 spearman

spearman's product moment coefficient

83.35 spearman_rank

83.36 spearman_to_pearson

conversion of spearman rank to person product moment correlation
coefficient

83.37 wmedian

weighted median

83.38 wquantile

weighted quantile

84 statistics

84.1 qstd

84.2 quantile_extrap

85 statistics/random-number-generation

85.1 laplacernd

random number of laplace distribution

85.2 randc

correlate to correlated standard normally distributed vectors

85.3 skewness2param

85.4 skewpdf_central_moments

85.5 skewrnd

random numbers of the skew normal distribution

85.6 skewrnd2

random numbers of the skew normal distribution

86 statistics

86.1 range

mid range

86.2 resample_with_replacement

87 statistics/resampling-statistics/@Jackknife

87.1 Jackknife

class for leave out 1 (delete 1) Jackknife estimates

note 1 : the 1-delete jackknife does not yield consistend estimates
for all functions,

in particular it will perform poorly on robust estimation
functions

this is overcome by the d-delete jackknife, where d has to
exceed the breakdown point

of the estimating function, for example \sqrt{n} for the
median

as this leads to unreasonably large number of repetitions,
bootstrap

is recommended for large sample cases (or blocking for
sequential data)

note 2 : as a linearisation, jackknife underestimates the error
variance in case of

dependence in the data

note 3 : studentisation and the leave out 1 jackknife are related

note 4 : the double 1 sample jackknife performs iferior to the d1
jackknife

87.2 estimated_STATIC

jackknife estimate of mean, bias and standard error
theta0 : estimate from all samples
thetad : set of estimates obtained by leaving out one data point
each
 last dimension of theta is assumed to be the jackknife
 dimension

87.3 matrix1_STATIC

matrix of estimation for leaving out two samples at a time

87.4 matrix2

matrix of estimations for jackknife with two samples left out

88 statistics/resampling-statistics

88.1 block_jackknife

88.2 jackknife_moments

moments determined by the jackknife

func : function of interest on the samples (e.g. mean)
A : parameter matrix
 columns : parameters
 rows : samples of the parameter sets
d : number of samples left out

88.3 moving_block_jackknife

blocked Jackknife for autocorrelated data
sliding block, statistically more efficient but computationally
expensive
note, number of blocks must be sufficiently large $h \sim \sqrt{n}$? $\ll n$

88.4 randblockterr

standard error of sequentially correlated data by blocking
block length should be sufficiently larger than correlation length
and sufficiently smaller than data length
this uses a sliding block approach, which reduces the variation of
the error estimate
TODO this does not work, randomly picking samples does not reveal
the correlation

88.5 resample

resample a vector and apply function to it

TODO, should be with replacement

n : number of samples
m : number of subsamples
cx : maximum number of combinations

89 statistics

89.1 scale_quantile_sd

scale factor for the standard deviation
of the asymptotic distribution of sample quantiles
(for normal distribution)
see cadwell, 1952

89.2 sd_sample_quantiles

89.3 skewpdf

skew-normal distribution
c.f. Azzalini 1985

89.4 trimmed_mean

trimmed mean

89.5 ttest2_man

two-sample t-test
here posix return value standard: h = 0 accepted, h = 1 failed
note: the matlab logic is inverse : h = 1 accepted, h = 0 failed
two sided univariate t-test

89.6 ttest_man

two-sample t-test
unequal sample size
equal variance

89.7 ttest_paired

paired t-test
unequal sample size
equal variance
more powerfull than unpaired test, as long as correlation between
x1 and x2 > 0

89.8 wgeomean

weighted geometric mean
function mu = wgeomean(w,x)

89.9 wgeovar

variance of the weighted geometric mean

89.10 wharmean

weighted harmonic mean

89.11 wharstd

89.12 wharvar

90 mathematics

mathematical functions of various kind

90.1 ternary_diagram

91 test/master

91.1 dat_test_lanczos_3d_k_20_n_40

91.2 poisson2d_blk

91.3 qr_implicit_givens_2

91.4 `spectral_derivative_2d`

91.5 `test_2d_eigensolver_hydrogen`

91.6 `test_2d_refine`

91.7 `test_3d_eigensolver_hydrogen`

91.8 `test_FEM`

91.9 `test_Mesh_3d`

91.10 `test_arnoldi`

91.11 `test_arpackc`

91.12 `test_assemble`

91.13 `test_assembly_performance`

91.14 test_bc_one_sided

91.15 test_compare_solvers

91.16 test_complete

91.17 test_convergence

91.18 test_convergence_b

91.19 test_df_2d

91.20 test_eig_algs

91.21 test_eig_inverse

91.22 test_eigs_lanczos

91.23 test_eigs_lanczos_1

91.24 test_eigs_lanczos_2

91.25 test_eigs_lanczos_performance

91.26 test_fdm

91.27 test_fdm_d_vargrid

91.28 test_fdm_spectral

91.29 test_fem

91.30 test_fem_1d

91.31 test_fem_1d_higher_order

91.32 test_fem_2d_adaptive

91.33 test_fem_2d_higher_order

91.34 test_fem_3d_higher_order

91.35 test_fem_3d_refine

91.36 test_fem_b

91.37 test_fem_derivative

91.38 test_fem_quadrature

91.39 test_final

91.40 test_fix_substitution

91.41 test_forward

91.42 test_get_sparse_arrays

91.43 test_harmonic_oscillator

91.44 test_high_order_fdm_periodic_bc

91.45 test_hydrogen_wf

91.46 test_ichol

91.47 test_interpolation

91.48 test_inverse_problem

91.49 test_it_vs_exact

91.50 test_jama

91.51 test_jd

91.52 test_jdqz

91.53 test_lanczos_2

91.54 test_lanczos_biorthogonal

91.55 test_laplacian

91.56 test_laplacian_non_uniform

91.57 test_laplacian_simple

91.58 test_mesh_2d_uniform

91.59 test_mesh_2d_uniform_2

91.60 test_mesh_circle

91.61 test_mesh_generation

91.62 test_mesh_interpolate

91.63 test_mg

91.64 test_minres_recycle

91.65 test_multigrid

91.66 test_nc

91.67 test_nonuniform_symmetric

91.68 test_pde

91.69 test_permutation

91.70 test_poison_fem

91.71 test_polar

91.72 test_potential

91.73 test_powers

91.74 test_precondition

91.75 test_project_rectangle

91.76 test_qr

91.77 test_quantum_well

91.78 test_radial_adaptive

91.79 test_radial_confinement

91.80 test_radial_fixes

91.81 test_refine_2d

91.82 test_refine_2d_b

91.83 test_refine_3d

91.84 `test_refine_structural`

91.85 `test_regularisation`

91.86 `test_round_off`

91.87 `test_schrödinger_potentials`

91.88 `test_uniform_mesh`

91.89 `test_vargrid`

92 `test`

92.1 `test_gaussfit3`

92.2 `test_mtimes3x3`

93 wavelet

93.1 continuous_wavelet_transform

continuous wavelet transform
follows "The Illustrated Wavelet Transform Handbook: Introductory
Theory and ..."

93.2 cwt_man

continuous fourier transform
as of time of implmentation, the matlab interal cwt is affected by
serious round-off errors and has issues with the scaling,
which is not the case here

93.3 example_wavelets

93.4 phasewrap

wrap the phase to +/- pi

93.5 test_cwt_man

93.6 test_phasewrap

93.7 test_wavelet

93.8 test_wavelet2

93.9 test_wavelet_analysis

93.10 test_wavelet_reconstruct

93.11 test_wtc

93.12 wavelet

wavelet windows

93.13 wavelet_reconstruct

inverses wavelet transform for single frequency
(reconstruction of time series)
n : window lengths in multiples of filter period $1/f_0$

93.14 wavelet_transform

wavelet transform for single frequency
n : window lengths in multiples of filter period $1/f_0$

94 mathematics

mathematical functions of various kind

94.1 wrapphase