

Manual for Package: mathematics

Revision 2:3M

Karl Kästner

October 14, 2019

Contents

1	complex-analysis	21
1.1	complex_exp_product_im_im	21
1.2	complex_exp_product_im_re	21
1.3	complex_exp_product_re_im	22
1.4	complex_exp_product_re_re	22
1.5	croots	23
1.6	root_complex	23
1.7	test_imroots	23
2	derivation	23
2.1	derive_acfar1	23
2.2	derive_ar2param	23
2.3	derive_arc_length	23
2.4	derive_fourier_power	23
2.5	derive_fourier_power_exp	24
2.6	derive_laplacian_curvilinear	24
2.7	derive_laplacian_fourier_pieewise_linear	24
2.8	derive_logtripdf	24
2.9	derive_smooth1d_parametric	24
2.10	simplify_atan	24
3	fourier/@STFT	24
3.1	STFT	24
3.2	itransform	25
3.3	stft_	25
3.4	stftmat	25
3.5	transform	25
4	fourier	25

4.1	amplitude_from_peak	25
4.2	dftmtx_man	26
4.3	example_fourier_window	26
4.4	fft_derivative	26
4.5	fft_man	26
4.6	fftsmooth	27
4.7	fix_fourier	27
4.8	fourier_axis	27
4.9	fourier_coefficient_piecewise_linear	28
4.10	fourier_coefficient_piecewise_linear_1	28
4.11	fourier_coefficient_ramp3	28
4.12	fourier_coefficient_ramp_pulse	28
4.13	fourier_coefficient_ramp_step	29
4.14	fourier_coefficient_square_pulse	29
4.15	fourier_derivative	29
4.16	fourier_expand	29
4.17	fourier_fit	29
4.18	fourier_interpolate	29
4.19	fourier_matrix	29
4.20	fourier_matrix2	30
4.21	fourier_matrix3	30
4.22	fourier_power	30
4.23	fourier_power_exp	30
4.24	fourier_predict	30
4.25	fourier_range	31
4.26	fourier_regress	31
4.27	fourier_resampled_fit	31
4.28	fourier_resampled_predict	31
4.29	fourier_signed_square	31
4.30	fourier_transform	31
4.31	hyperbolic_fourier_box	32
4.32	idftmtx_man	32
4.33	laplace_2d_pwlinear	32
4.34	nanfft	32
4.35	peaks	33
4.36	roots_fourier	33
4.37	spectral_density	33
4.38	test_complex_exp_product	33
4.39	test_idftmtx	33
5	geometry/@Geometry	34
5.1	Geometry	34
5.2	arclength	34
5.3	arclength_old	34

5.4	arclength_old2	34
5.5	base_point	34
5.6	base_point_limited	34
5.7	centroid	35
5.8	cross2	35
5.9	curvature	35
5.10	ddot	35
5.11	distance	35
5.12	distance2	35
5.13	dot	35
5.14	edge_length	35
5.15	enclosed_angle	36
5.16	enclosing_triangle	36
5.17	hexagon	36
5.18	inPolygon	36
5.19	inTetra	36
5.20	inTetra2	36
5.21	inTriangle	36
5.22	intersect	36
5.23	lineintersect	37
5.24	lineintersect1	37
5.25	minimum_distance_lines	37
5.26	mittenpunkt	37
5.27	nagelpoint	37
5.28	onLine	37
5.29	orthocentre	37
5.30	plumb_line	37
5.31	poly_area	38
5.32	poly_edges	38
5.33	poly_set	38
5.34	poly_width	38
5.35	polyxpoly	38
5.36	project_to_curve	38
5.37	random_disk	38
5.38	random_simplex	39
5.39	sphere_volume	39
5.40	tetra_volume	39
5.41	tobarycentric	39
5.42	tobarycentric1	39
5.43	tobarycentric2	39
5.44	tobarycentric3	39
5.45	tri_angle	39
5.46	tri_area	40
5.47	tri_centroid	40

5.48	tri_distance_opposit_midpoint	40
5.49	tri_edge_length	40
5.50	tri_edge_midpoint	40
5.51	tri_excircle	40
5.52	tri_height	40
5.53	tri_incircle	40
5.54	tri_isacute	41
5.55	tri_isobtuse	41
5.56	tri_semiperimeter	41
5.57	tri_side_length	41
6	geometry	41
6.1	Polygon	41
6.2	bounding_box	41
6.3	curvature_1d	42
6.4	cvt	42
6.5	deg_to_frac	42
6.6	ellipse	42
6.7	ellipseX	42
6.8	ellipseY	42
6.9	first_intersect	42
6.10	golden_ratio	42
6.11	hypot3	43
6.12	meanangle	43
6.13	meanangle2	43
6.14	meanangle3	43
6.15	meanangle4	43
6.16	medianangle	43
6.17	medianangle2	43
6.18	pilim	44
6.19	streamline_radius_of_curvature	44
7	linear-algebra	44
7.1	averaging_matrix_2	44
7.2	colnorm	44
7.3	condest_	44
8	linear-algebra/coordinate-transformation	44
8.1	barycentric2cartesian	44
8.2	barycentric2cartesian3	44
8.3	cartesian2barycentric	45
8.4	cartesian_to_unit_triangle_basis	45
8.5	example_approximate_utm_conversion	45
8.6	latlon2utm	45

8.7	latlon2utm_simple	45
8.8	lowrance_mercator_to_wgs84	45
8.9	nmea2utm	45
8.10	sn2xy	45
8.11	unit_triangle_to_cartesian	46
8.12	utm2latlon	46
8.13	xy2nt	46
8.14	xy2sn	46
8.15	xy2sn_java	46
8.16	xy2sn_old	46
9	linear-algebra	47
9.1	det2x2	47
9.2	det3x3	47
9.3	det4x4	47
9.4	diag2x2	47
9.5	eig2x2	47
9.6	first	47
9.7	gershgorin_circle	47
9.8	haussdorff	48
9.9	ieig2x2	48
9.10	inv2x2	48
9.11	inv3x3	48
9.12	inv4x4	48
9.13	lpmean	48
9.14	lpnorm	48
9.15	matvec3	49
9.16	max2d	49
9.17	mpoweri	49
9.18	mtimes2x2	49
9.19	mtimes3x3	49
9.20	nannorm	49
9.21	nanshift	49
9.22	nl	49
9.23	normalise	50
9.24	normalize1	50
9.25	normrows	50
9.26	orth2	50
9.27	orth_man	50
9.28	orthogonalise	50
9.29	paddext	50
9.30	paddval1	51
9.31	paddval2	51

10 linear-algebra/polynomial	51
10.1 chebychev	51
10.2 piecewise_polynomial	51
10.3 roots1	51
10.4 roots2	51
10.5 vanderi_1d	51
10.6 vandermonde	51
11 linear-algebra	52
11.1 randrot	52
11.2 right	52
11.3 rot2	52
11.4 rot2dir	52
11.5 rot3	52
11.6 rownorm	52
11.7 simmilarity_matrix	52
11.8 spnorm	52
11.9 spzeros	53
11.10 transpose3	53
11.11 transposeall	53
12 logic	53
12.1 bitor_man	53
13 number-theory	53
13.1 ceiln	53
13.2 digitsb	53
13.3 floorn	53
13.4 iseven	54
13.5 multichoosek	54
13.6 nchoosek_man	54
13.7 pythagorean_triple	54
13.8 roundn	54
14 numerical-methods/differentiation	54
14.1 derivative1	54
14.2 derivative2	55
15 numerical-methods/finite-difference	55
15.1 cdiff	55
15.2 cdiffb	55
15.3 cmean	55
15.4 derivative_matrix_1_1d	55
15.5 derivative_matrix_2_1d	55

15.6	derivative_matrix_2d	55
15.7	derivative_matrix_curvilinear	56
15.8	derivative_matrix_curvilinear_2	56
15.9	difference_kernel	56
15.10	distmat	56
15.11	gradpde2d	56
15.12	laplacian	56
15.13	laplacian_fdm	56
15.14	left	57
15.15	lrmean	57
15.16	mid	57
15.17	pwmid	57
15.18	ratio	57
15.19	steplength	57
15.20	swapoddeven	57
15.21	test_derivative_matrix_2d	57
15.22	test_derivative_matrix_curvilinear	58
15.23	test_difference_kernel	58
16	numerical-methods/finite-volume/@Advection	58
16.1	Advection	58
16.2	dot_advection	58
17	numerical-methods/finite-volume/@Burgers	58
17.1	burgers_split	58
17.2	dot_burgers_fdm	58
17.3	dot_burgers_fft	58
18	numerical-methods/finite-volume/@Finite_Volume	59
18.1	Finite_Volume	59
18.2	apply_bc	59
18.3	solve	59
18.4	step_split_strang	59
18.5	step_unsplit	59
19	numerical-methods/finite-volume/@Flux_Limiter	59
19.1	Flux_Limiter	59
19.2	beam_warming	60
19.3	fromm	60
19.4	lax_wendroff	60
19.5	minmod	60
19.6	monotized_central	60
19.7	muscl	60
19.8	superbee	60

19.9	upwind	61
19.10	vanLeer	61
20	numerical-methods/finite-volume/@KDV	61
20.1	dot_kdv_fdm	61
20.2	dot_kdv_fft	61
20.3	kdv_split	61
21	numerical-methods/finite-volume/@Reconstruct_Average_Evolve	61
21.1	Reconstruct_Average_Evolve	61
21.2	advect_highres	62
21.3	advect_lowres	62
22	numerical-methods/finite-volume	62
22.1	Godunov	62
22.2	Lax_Friedrich	62
22.3	Measure	62
22.4	Roe	62
22.5	fv_swe	63
22.6	staggered_euler	63
22.7	staggered_grid	63
23	numerical-methods	63
23.1	grid2quad	63
24	numerical-methods/integration	63
24.1	cumintL	63
24.2	cumintR	63
24.3	int_trapezoidal	63
25	numerical-methods/interpolation/@Kriging	64
25.1	Kriging	64
25.2	estimate_semivariance	64
25.3	interpolate_	64
26	numerical-methods/interpolation/@RegularizedInterpolator1	64
26.1	RegularizedInterpolator1	64
26.2	init	64
27	numerical-methods/interpolation/@RegularizedInterpolator2	65
27.1	RegularizedInterpolator2	65
27.2	init	65
28	numerical-methods/interpolation/@RegularizedInterpolator3	65
28.1	RegularizedInterpolator3	65

28.2	init	65
29	numerical-methods/interpolation	65
29.1	IDW	65
29.2	IPoly	65
29.3	IRBM	66
29.4	ISparse	66
29.5	Inn	66
29.6	Interpolator	66
29.7	fixnan	66
29.8	idw1	66
29.9	idw2	66
29.10	inner2outer	67
29.11	inner2outer2	67
29.12	interp1_limited	67
29.13	interp1_man	67
29.14	interp1_save	67
29.15	interp1_slope	67
29.16	interp1_smooth	68
29.17	interp1_unique	68
29.18	interp2_man	68
29.19	interp_angle	68
29.20	interp_fourier	68
29.21	interp_fourier_batch	68
29.22	interp_sn	68
29.23	interp_sn2	69
29.24	interp_sn3	69
29.25	interp_sn_	69
29.26	limit_by_distance_1d	69
29.27	resample1	69
29.28	resample_d_min	69
29.29	resample_vector	69
29.30	test_interp1_limited	69
30	numerical-methods	70
30.1	inverse_complex	70
31	numerical-methods/ode	70
31.1	bvp2_check_arguments	70
31.2	bvp2c	70
31.3	bvp2c2	70
31.4	bvp2fdm	71
31.5	bvp2wavetrain	71
31.6	bvp2wavetwopass	71

31.7	ivp_euler_forward	71
31.8	ivprk2	72
31.9	ode2_matrix	72
31.10	ode2characteristic	72
31.11	step_trapezoidal	72
31.12	test_bvp2	72
32	numerical-methods/optimisation	72
32.1	armijo_stopping_criterion	72
32.2	astar	72
32.3	binsearch	73
32.4	bisection	73
32.5	box1	73
32.6	box2	73
32.7	cauchy	73
32.8	cauchy2	73
32.9	directional_derivative	73
32.10	dud	74
32.11	extreme3	74
32.12	extreme_quadratic	74
32.13	ftest	74
32.14	grad	74
32.15	hessian	74
32.16	hessian_from_gradient	75
32.17	hessian_projected	75
32.18	line_search	75
32.19	line_search2	75
32.20	line_search_polynomial	75
32.21	line_search_polynomial2	76
32.22	line_search_quadratic	76
32.23	line_search_quadratic2	76
32.24	line_search_wolfe	76
32.25	ls_bgfs	76
32.26	ls_broyden	77
32.27	ls_generalized_secant	77
32.28	nleg	77
32.29	nlls	77
32.30	picard	77
32.31	poly_extrema	78
32.32	quadratic_function	78
32.33	quadratic_programming	78
32.34	quadratic_step	78
32.35	rosenbrock	78
32.36	sqrt_heron	78

32.37	test_directional_derivative	78
32.38	test_dud	78
32.39	test_line_search_quadratic2	78
32.40	test_ls_generalized_secant	79
32.41	test_nlcg_6_order	79
32.42	test_nlls	79
33	numerical-methods/piecewise-polynomials	79
33.1	Hermitel	79
33.2	hp2_fit	79
33.3	hp2_predict	79
33.4	hp_predict	79
33.5	hp_regress	80
33.6	lp_count	80
33.7	lp_predict	80
33.8	lp_regress	80
33.9	lp_regress_	80
34	regression/@PolyOLS	80
34.1	PolyOLS	80
34.2	coefftest	80
34.3	detrend	80
34.4	fit	81
34.5	fit_	81
34.6	predict	81
34.7	predict_	81
34.8	slope	81
35	regression/@PowerLS	81
35.1	PowerLS	81
35.2	fit	81
35.3	predict	82
35.4	predict_	82
36	regression/@Theil	82
36.1	Theil	82
36.2	detrend	82
36.3	fit	82
36.4	predict	82
36.5	slope	83
37	regression	83
37.1	Theil_Multivariate	83
37.2	areg	83

37.3	ginireg	83
37.4	hesssimplereg	83
37.5	l1lin	83
37.6	lsq_sparam	84
37.7	polyfitd	84
37.8	regression_method_of_moments	84
37.9	robustlinreg	84
37.10	theil2	84
37.11	theil_generalised	84
37.12	total_least_squares	85
37.13	weighted_median_regression	85
38	set-theory	85
38.1	issubset	85
39	signal-processing	85
39.1	acf_effective_sample_size	85
39.2	acf_genton	85
39.3	acfar1	86
39.4	acfar1_2	86
39.5	acfar2	86
39.6	acfar2_2	86
39.7	ar1_cutoff_frequency	86
39.8	ar1_effective_sample_size	86
39.9	ar1_mse_mu_single_sample	86
39.10	ar1_mse_pop	87
39.11	ar1_mse_range	87
39.12	ar1_spectrum	87
39.13	ar1_to_tikhonov	87
39.14	ar1_var_factor	87
39.15	ar1_var_factor_	87
39.16	ar1_var_range2	87
39.17	ar1delay	88
39.18	ar1delay_old	88
39.19	ar2conv	88
39.20	ar2dof	88
39.21	ar2param	88
39.22	asymwin	88
39.23	autocorr_fft	88
39.24	bandpass	89
39.25	bandpass2	89
39.26	bartlett	89
39.27	bartlett_spectrogram	89
39.28	bin1d	89

39.29	bin2d	89
39.30	binormrnd	90
39.31	conv1_man	90
39.32	conv2_man	90
39.33	conv2z	90
39.34	conv30	90
39.35	conv_	90
39.36	conv_centered	90
39.37	convz	90
39.38	cosexpdelay	91
39.39	csmooth	91
39.40	daniell_window	91
39.41	danielle_window	91
39.42	db2neper	91
39.43	db2power	91
39.44	derive_danielle_weight	91
39.45	derive_limit_0_acfar	91
39.46	detect_peak	92
39.47	digital_low_pass_filter	92
39.48	doublesum_ij	92
39.49	effective_sample_size_to_ar1	92
39.50	filt_hodges_lehman	92
39.51	filter1	92
39.52	filter2	92
39.53	filter_	93
39.54	filteriir	93
39.55	filterp	93
39.56	filterp1	94
39.57	filterstd	94
39.58	firls_man	94
39.59	flattopwin	94
39.60	frequency_response_boxcar	94
39.61	freqz_boxcar	94
39.62	gaussfilt1	94
39.63	hanchangewin	94
39.64	hanchangewin2	95
39.65	hanwin	95
39.66	hanwin_	95
39.67	highpass	95
39.68	kaiserwin	95
39.69	kalman	95
39.70	lanczoswin	95
39.71	last	95
39.72	lowpass	96

39.73	lowpass2	96
39.74	lowpass_iir	96
39.75	lowpass_iir_symmetric	96
39.76	lowpassfilter2	96
39.77	maxfilt1	96
39.78	meanfilt1	96
39.79	medfilt1_man	96
39.80	medfilt1_man2	97
39.81	medfilt1_padded	97
39.82	medfilt1_reduced	97
39.83	mid_term_single_sample	97
39.84	minfilt1	97
39.85	mu2ar1	97
39.86	nanautocorr	97
39.87	nanmedfilt1	97
39.88	neper2db	98
39.89	peaks_man	98
39.90	polyfilt1	98
39.91	qmedfilt1	98
39.92	randar1	98
39.93	randar1_dual	98
39.94	randar2	98
39.95	randarp	98
39.96	range_window	99
39.97	rectwin	99
39.98	recursive_sum	99
39.99	select_range	99
39.100	smooth1d_parametric	99
39.101	smooth2	99
39.102	smooth_man	99
39.103	smooth_parametric	99
39.104	smooth_parametric2	100
39.105	smoothfft	100
39.106	spectrogram	100
39.107	std_window	100
39.108	sum_i_lag	100
39.109	sum_ii	100
39.110	sum_ii_	100
39.111	sum_ij	100
39.112	sum_ij_	101
39.113	sum_ij_partial_	101
39.114	sum_multivar	101
39.115	test_acfar1	101
39.116	test_acfar1_2	101

39.117	test_acfar1_3	101
39.118	test_acfar1_4	101
39.119	test_acfar2	101
39.120	test_ar1_var_factor	101
39.121	test_ar1_var_factor_2	101
39.122	test_ar1_var_mu_single_sample	102
39.123	test_ar1_var_pop	102
39.124	test_ar1_var_pop_1	102
39.125	test_ar1delay	102
39.126	test_bivariate_covariance_term	102
39.127	test_convexity	102
39.128	test_lanczoswin	102
39.129	test_madcorr	102
39.130	test_randar1	102
39.131	test_randar1_multivariate	102
39.132	test_randar2	103
39.133	test_sum_ij	103
39.134	test_sum_multivar	103
39.135	test_trifilt1	103
39.136	test_wautocorr	103
39.137	test_wavelet_transform	103
39.138	test_wordfilt	103
39.139	test_xar1_mid_term	103
39.140	tikhonov_to_ar1	103
39.141	trapwin	104
39.142	trifilt1	104
39.143	triwin	104
39.144	triwin2	104
39.145	varar1	104
39.146	welch_spectrogram	104
39.147	wfilt	104
39.148	winbandpass	104
39.149	window_make_odd	105
39.150	winfilt0	105
39.151	winlength	105
39.152	wmeanfilt	105
39.153	wmedfilt	105
39.154	wordfilt	105
39.155	wordfilt_edgeworth	105
39.156	xar1	105
39.157	xcorr_man	106
40 sorting		106
40.1	sort2	106

40.2	sort2d	106
41	special-functions	106
41.1	bessel_sphere	106
41.2	hankel_sphere	106
41.3	hermite	106
41.4	legendre_man	107
41.5	neumann_sphere	107
42	statistics	107
42.1	atan_s2	107
42.2	beta_mode_to_parameter	107
42.3	correlation_confidence_pearson	107
43	statistics/distributions	107
43.1	PDF	107
43.2	binorm_separation_coefficient	107
43.3	binormcdf	108
43.4	binormfit	108
43.5	binormpdf	108
43.6	edgeworth_cdf	108
43.7	edgeworth_pdf	108
43.8	logn_mode2param	108
43.9	logn_param2mode	108
43.10	lognpdf_	109
43.11	pdfsample	109
43.12	t2cdf	109
43.13	t2inv	109
44	statistics	109
44.1	example_standard_error_of_sample_quantiles	109
44.2	f_var_finite	109
44.3	gamma_mode_to_parameter	109
44.4	hodges_lehmann_correlation	110
44.5	hodges_lehmann_dispersion	110
45	statistics/information-theory	110
45.1	akaike_information_criterion	110
45.2	bayesian_information_criterion	110
46	statistics	111
46.1	kurtncdf	111
46.2	kurtnpdf	111
46.3	kurtosis_bias_corrected	111
46.4	limit	111

46.5	logfactorial	111
46.6	loglogpdf	111
46.7	logskewcdf	111
46.8	logskewpdf	111
47	statistics/logu	112
47.1	lambertw_numeric	112
47.2	logtrialtcd	112
47.3	logtrialtinv	112
47.4	logtrialtmean	112
47.5	logtrialtpdf	112
47.6	logtrialtrnd	112
47.7	logtricdf	113
47.8	logtriinv	113
47.9	logtrimean	113
47.10	logtripdf	113
47.11	logtrirnd	113
47.12	logucdf	113
47.13	logucm	113
47.14	loguinv	113
47.15	logumean	114
47.16	logupdf	114
47.17	logurnd	114
47.18	loguvar	114
47.19	medlogu	114
47.20	test_logurnd	114
47.21	tricdf	114
47.22	triinv	114
47.23	trimedial	115
47.24	tripdf	115
47.25	trirnd	115
48	statistics	115
48.1	maxnnormals	115
48.2	midrange	115
48.3	minavg	115
48.4	mode_man	115
49	statistics/moment-statistics	116
49.1	autocorr_man3	116
49.2	autocorr_man4	116
49.3	autocorr_man5	116
49.4	blockserr	116
49.5	comoment	116

49.6	corr_man	117
49.7	cov_man	117
49.8	dof	117
49.9	edgeworth_quantile	117
49.10	effective_sample_size	117
49.11	f_correlation	117
49.12	f_finite	118
49.13	lmean	118
49.14	lmoment	118
49.15	maskmean	118
49.16	masknanmean	118
49.17	mean1	118
49.18	mean_man	118
49.19	mse	118
49.20	nanautocorr_man1	119
49.21	nanautocorr_man2	119
49.22	nanautocorr_man4	119
49.23	nancorr	119
49.24	nancumsum	119
49.25	nanlmean	119
49.26	nanr2	119
49.27	nanrms	120
49.28	nanrmse	120
49.29	nanserr	120
49.30	nanwmean	120
49.31	nanwstd	120
49.32	nanwvar	120
49.33	nanxcorr	120
49.34	pearson	121
49.35	pearson_to_kendall	121
49.36	pool_samples	121
49.37	qmean	121
49.38	range_mean	121
49.39	rmse	121
49.40	serr	121
49.41	serr1	121
49.42	test_kskew	122
49.43	test_qstd_kskew_optimal_p	122
49.44	wautocorr	122
49.45	wcorr	122
49.46	wcov	122
49.47	wdof	122
49.48	wkurt	122
49.49	wmean	123

49.50	wrms	123
49.51	wserr	123
49.52	wskew	123
49.53	wstd	123
49.54	wvar	123
50	statistics	124
50.1	nangeomean	124
50.2	nangeostd	124
51	statistics/nonparametric-statistics	124
51.1	kernel1d	124
51.2	kernel2d	124
52	statistics	124
52.1	normmoment	124
52.2	normpdf2	124
53	statistics/order-statistics	125
53.1	hodges_lehmann_location	125
53.2	kendall	125
53.3	kendall_to_pearson	125
53.4	mad2sd	125
53.5	madcorr	125
53.6	median2_holder	125
53.7	median_ci	126
53.8	median_man	126
53.9	mediani	126
53.10	nanmadcorr	126
53.11	nanwmedian	126
53.12	nanwquantile	126
53.13	oja_median	126
53.14	qkurtosis	127
53.15	qmoments	127
53.16	qskew	127
53.17	qskewq	127
53.18	qstdq	127
53.19	quantile1_optimisation	127
53.20	quantile2_breckling	128
53.21	quantile2_chaudhuri	128
53.22	quantile2_projected	128
53.23	quantile2_projected2	128
53.24	quantile_envelope	128
53.25	quantile_regression_simple	128

53.26	ranking	128
53.27	spatial_median	128
53.28	spatial_quantile	129
53.29	spatial_quantile2	129
53.30	spatial_quantile3	129
53.31	spatial_rank	129
53.32	spatial_sign	129
53.33	spatial_signed_rank	129
53.34	spearman	129
53.35	spearman_rank	129
53.36	spearman_to_pearson	130
53.37	wmedian	130
53.38	wquantile	130
54	statistics	130
54.1	qstd	130
55	statistics/random-number-generation	130
55.1	laplacernd	130
55.2	randc	130
55.3	skewrnd	130
55.4	skewrnd2	131
56	statistics	131
56.1	range	131
57	statistics/resampling-statistics/@Jackknife	131
57.1	Jackknife	131
57.2	estimated_STATIC	131
57.3	matrix1_STATIC	132
57.4	matrix2	132
58	statistics/resampling-statistics	132
58.1	block_jackknife	132
58.2	jackknife_moments	132
58.3	moving_block_jackknife	132
58.4	randblockserr	133
58.5	resample	133
59	statistics	133
59.1	scale_quantile_sd	133
59.2	skewpdf	133
59.3	trimmed_mean	133
59.4	ttest2_man	134
59.5	ttest_man	134

59.6	ttest_paired	134
59.7	wharmean	134
60	wavelet	134
60.1	contiuous_wavelet_transform	134
60.2	cwt_man	134
60.3	example_wavelets	135
60.4	phasewrap	135
60.5	test_cwt_man	135
60.6	test_phasewrap	135
60.7	test_wavelet	135
60.8	test_wavelet2	135
60.9	test_wavelet_analysis	135
60.10	test_wavelet_reconstruct	135
60.11	test_wtc	136
60.12	wavelet	136
60.13	wavelet_reconstruct	136
60.14	wavelet_transform	136
61	mathematics	136
61.1	wrapphase	136

1 complex-analysis

operations on complex numbers

1.1 complex_exp_product_im_im

product of the imaginary part of two complex exponentials

the product has two frequency components

```
input :
  c : complex amplitudes
  o : frequencies
output :
  cp : amplitude of the product
  op : frequencies of the product
```

1.2 complex_exp_product_im_re

product of the imaginary part of one and the real part of a second complex exponential

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

1.3 complex_exp_product_re_im

the product has two frequency components

product of the imaginary part of one and the real part of a second
complex exponential

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

1.4 complex_exp_product_re_re

product of the real part of two complex exponentials

$$\begin{aligned} \text{re}(c_1 \exp(i\omega_1 x)) * \text{re}(c_2 \exp(i\omega_2 x)) = \\ \frac{1}{2} * (\text{real}(c_1 * c_2 * \exp(i * (\omega_1 + \omega_2) * x)) \dots \\ + \text{real}(\text{conj}(c_1) * c_2 * \exp(i * (\omega_2 - \omega_1) * x))) \end{aligned}$$

the product has two frequency components

input :
 c : complex amplitudes
 o : frequencies
output :
 cp : amplitude of the product
 op : frequencies of the product

1.5 croots

nth-roots of a complex number

input:
c : complex number
n : order of root
 n must be rational, to obtain n solutions
 otherwise no finite set of solutions exists

r : roots of the complex number

1.6 root_complex

root of a complex number

1.7 test_imroots

2 derivation

derivation of several functions by means of symbolic computation

2.1 derive_acfar1

2.2 derive_ar2param

2.3 derive_arc_length

2.4 derive_fourier_power

2.5 `derive_fourier_power_exp`

2.6 `derive_laplacian_curvilinear`

2.7 `derive_laplacian_fourier_piecewise_linear`

2.8 `derive_logtripdf`

2.9 `derive_smooth1d_parametric`

2.10 `simplify_atan`

symbolic simplification of the arcus tangent

3 `fourier/@STFT`

3.1 `STFT`

class for short time fourier transform

Note: the interval T_i should be set to at least $2 \cdot \max(T)$, as
otherwise coefficients

tend to oscillate in the presence of noise

Note: for convenience, the independent variable is labeled as time
(t),

but the independent variable is arbitrary, so it works
likewise in space

3.2 itransform

inverse of the short time fourier transform

3.3 stft_

static wrapper for STFT

3.4 stftmat

transformation matrix for the short time fourier transform

3.5 transform

short time fourier transform

4 fourier

support and analysis functions both for the discrete (fast) fourier
transform (dft/fft)
and continuous fourier analysis (fourier series)

4.1 amplitude_from_peak

amplitude and standard deviation of the amplitude of a frequency
component

represented by a peak in the fourier domain

input :

h : peak height

w : peak width at half height

output:

a : amplitude in real space

s : standard deviation of the frequency (!)

4.2 dftmtx_man

fourier matrix in matlab style with a limited number of rows,
columns of higher frequencies are omitted

input :
n : number of samples
nr : number of columns

output :
F : fourier matrix

4.3 example_fourier_window

4.4 fft_derivative

derivative by fourier transform
exponential convergence for periodic functions
results in spurious oscillations for aperiodic functions

input:
x : data, sampled in equal intervals
k : order of the derivative

dx : kth-derivative of x

4.5 fft_man

fast fourier transform for complex input data

input:
F : data in real space

output :

F : fourier transformation of F

4.6 fftsmooth

smooth the fourier transform and determine upper and lower bound confidence intervals

input :
f :
sfunc : a smoothing function (for example fir convolution with rectangular window)
 returns filtered (mean) value and normalized fir window
nf : window length
nsigma : number of standard deviations for confidence intervals

output :
ff : filtered fourier transform
l : lower bound
u : upper bound

4.7 fix_fourier

fill gaps (missing data) by means of fourier extrapolation

fix periodic data series with fourier interpolation
longest gap should not exceed 1/2 of the shortest time span of interest (1/cutoff frequency)
note: this limit equals the position of first side lobe of the ft of a rectangular window with gap length

4.8 fourier_axis

return axis of frequencies and periods for the discrete fourier transform
as computed by fft (matlab-style)

input:
X : sample locations (equal interval)
L : length of samples
n : number of samples

output :
f : frequencies
T : periods
mask : mask for 1/2 of the fourier transform

(as both halves are complex conjugates)
N : frequency id

4.9 `fourier_coefficient_piecewise_linear`

fourier series coefficients of a piecewise linear function
(not coefficient of discrete fourier transform)
function can be discontinuous between intervals
scales domain length to 2π

input :
l,r : end points of piecewise linear function
lval, rval : values at end points
L : length of domain
n : number of samples/highest frequency

output :
a, b : coefficients for frequency components

4.10 `fourier_coefficient_piecewise_linear_1`

fourier series coefficients of a piecewise linear function
(not coefficient of discrete fourier transform)
function can be discontinuous between intervals
scales domain length to 2π

input :
X : end points of piecewise linear function
Y : values at end points

output :
ab : coefficients for frequency components

4.11 `fourier_coefficient_ramp3`

fourier series coefficient of a ramp

4.12 `fourier_coefficient_ramp_pulse`

fourier series coefficient of a ramp pulses

4.13 `fourier_coefficient_ramp_step`

fourier coefficient of a ramp-step

4.14 `fourier_coefficient_square_pulse`

fourier series coefficients of a square pulse

4.15 `fourier_derivative`

coefficients of the derivative of a fourier series
not of discrete fourier transform (fft)

4.16 `fourier_expand`

expand values of fourier series

4.17 `fourier_fit`

fit a fourier series to a set of sample points that are not spaced
in
equal intervals

4.18 `fourier_interpolate`

interpolate samples y sampled at moments (location) t to locations
 t_i

4.19 `fourier_matrix`

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

4.20 fourier_matrix2

transformation matrix for a continuous fourier series
(not for the discrete dft/fft)

4.21 fourier_matrix3

transformation matrix for the continous fourier transform
this is a matrix with $(2*n+1)$ real columns

4.22 fourier_power

powers of a continuous fourier series in sin/cos form

powers of $a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \phi))^p$
phase of first component assumed 0

frequencies higher than 2ω ignored in input
frequencies higher than 3ω not computed

4.23 fourier_power_exp

powers of the continuous fourier series
 $a^p = (u_r + u_1 \sin(\omega t) + u_2 \sin(\omega t + \phi))^p$
phase of first component assumed 0

higher orders than 2 ignored input
higher order than 3 not computed in output

$$y = a_0 + \sum (a_j \sin(j\omega t) + b_j \cos(j\omega t))$$
$$= \text{Real}(\sum_{i=0}^{\infty} c_i \exp(1i\omega t)), c_i = a_i + b_i$$

NOT the alternative $\sum_{i=-\infty}^{\infty} \tilde{c}_i$, tile $c_j = 1/2 a_j$
 $+ 1/2i b_j$

4.24 fourier_predict

expand a continous fourier series at times t

4.25 `fourier_range`

approximate range of a continuous Fourier series with 2 components
`range(y) = max(y) - min(y)`

4.26 `fourier_regress`

fit a continuous Fourier series to a set of sample points not
sampled
at equal intervals

4.27 `fourier_resampled_fit`

fits coefficients of a continuous Fourier transform,
but stores them as resampled values

4.28 `fourier_resampled_predict`

interpolates a continuous Fourier series that has been stored as
values
at their support points

4.29 `fourier_signed_square`

coefficients of the Fourier series of $|\cos a + \cos t|$ ($\cos a + \cos t$)
in general
 $\cos a$ is midrange
 $\cos t$ is tidal variation
c.f. Dronkers

4.30 `fourier_transform`

continuous Fourier transformation of y
(not discrete Fourier transformation dft/fft)

input:
 b : data sampled at equal intervals

`T` : length of data in time or space, i.e. position of last
sample if
position of first sample is 0
`T_max` : maximum period to include

output :

`A` : fourier matrix
`p` : fourier transformation of `b`
`tt` : TODO

4.31 hyperbolic_fourier_box

4.32 idftmtx_man

inverse matrix for the discrete fourier transform in matlab style
with a limited number of columns, thus ignoring higher frequencies
keep $2nc+1$ columns (mean and conj-complex pairs of nc frequencies)

4.33 laplace_2d_pwlinear

solution to the Laplacian in two dimensions for a finite
rectangular domain
with piecewise constant boundary conditions
linear system with 4 unknowns per frequency component
these are coefficients of `s,c,sh,ch`
 $(pu*(s + c) + qu*(s' + c'))*(shu + chu) = ru$ % upper bc
 $(pd*(s + c) + qd*(s' + c'))*(shd + chd) = rd$ % lower bc
 $((sl + cl)*(pl*(shl + chl) + ql*(shl' + chl'))) = rl$ % left
bc
 $((sr + cr)*(pr*(shr + chr) + qr*(shr' + chr'))) = rr$ % right
bc
least squares with piecewise integration
`[x0,p,q,r]` piecewise linear polynomials at the boundaries

4.34 nanfft

discrete fourier transform of a data series with gaps

4.35 peaks

peaks of the power spectrum of a discrete fourier transform

rule for peaks: there is no higher value left or right of the "peak"

until the signal drops to $p \cdot y_{\text{peak}}$, $p = 0.5$

works best, when spectrum has been smoothened

input :

f : frequency

y : absolute value of fourier transform (power spectrum)

L : length in space or time of series

output :

a0 : amplitude

s0 : standard deviation (error?) of amplitude

w0 : width of peak

lambda = wave length (period?)

pdx : index of peak

f : frequency (if not given as input)

4.36 roots_fourier

zeros of continuous fourier series

$$f = a_0 + \sum_{j=1}^n a_j \cos(j x) + b_j \sin(j x)$$

4.37 spectral_density

spectral density

4.38 test_complex_exp_product

4.39 test_idftmtx

5 geometry/@Geometry

5.1 Geometry

5.2 arclength

arc length of a two dimensional curve

8th order accurate

does not require the segments length to vary smoothly

note: the curve can be considered parametric, e.g. $x = x(t)$, $y = y(t)$
and

and $t = t(s)$, but the error term contains derivatives of t ,
thus a non smooth t (strongly varying distance between points)
requires the scaling as done below

5.3 arclength_old

arc length of a two dimensional function

5.4 arclength_old2

arc length of a two dimensional function

5.5 base_point

base point (fusspunkt), i.e. point on a line with shortest distance
to another point

5.6 base_point_limited

base point (Fusspunkt) of a point on a line

5.7 centroid

centroid pf a polygone

5.8 cross2

cross product in two dimensions

5.9 curvature

curvature of a function in two dimensions

5.10 ddot

sum of squares of cos of inner angles of triangle

5.11 distance

equclidan distance between two points

5.12 distance2

euclidean distance between two points
this function requires a and be of equal dimensions, or the least
the first pair or second pair to be a scalar

5.13 dot

dot product

5.14 edge_length

edge length

5.15 enclosed_angle

angle enclosed between two lines

5.16 enclosing_triangle

smallest enclosing equilateral triangle with bottom side parallel to
X axis

5.17 hexagon

coordinates of a hexagon, scaled and rotated

5.18 inPolygon

flag points contained in a polygon
much faster than matlab internal function

5.19 inTetra

flag points contained in tetrahedron

5.20 inTetra2

flag points contained in tetrahedron

5.21 inTriangle

flag points contained in triangle

5.22 intersect

intersect between two lines

5.23 lineintersect

intersect of two lines

5.24 lineintersect1

intersect of two lines

5.25 minimum_distance_lines

minimum distance of two lines in three dimensions

5.26 mittenpunkt

mittenpunkt of a triangle

5.27 nagelpoint

nagelpoint of a triangle

5.28 onLine

5.29 orthocentre

orthocentre of triangle

5.30 plumb_line

5.31 poly_area

area of a polygon

5.32 poly_edges

edges of a polygon

5.33 poly_set

associate point at arbitrary location with a polygon it is contained
in
and assign the value of the polygon to it

5.34 poly_width

width of polygon width holes by surface normals
holes / islands separated with NaN
order of points of outer boundary must be cw
order of points of holes must be ccw
note that this function does not give the true width for expanding
sections
use voronoi polygons for this

5.35 polyxpoly

intersections of two polygons

5.36 project_to_curve

closest point on a curve with respect to a point at distance to the
curve

5.37 random_disk

draw random points on the unit disk

5.38 random_simplex

random point inside of a triangle

5.39 sphere_volume

volume of a sphere

5.40 tetra_volume

volume of a tetrahedron

5.41 tobarycentric

cartesian to barycentric coordinates

5.42 tobarycentric1

cartesian to barycentric coordinates

5.43 tobarycentric2

cartesian to barycentric coordinates

5.44 tobarycentric3

cartesian to barycentric coordinates

5.45 tri_angle

cos of angles of a triangle

5.46 tri_area

angle of a triangle

5.47 tri_centroid

centroid of a triangle

5.48 tri_distance_opposit_midpoint

distance between corner of a triangle and its opposing mid-point

5.49 tri_edge_length

edge length of a triangle

5.50 tri_edge_midpoint

mid point of a triangle

5.51 tri_excircle

excircle of a triangle

5.52 tri_height

height of a triangle

5.53 tri_incircle

incircle of a triangle

5.54 tri_isacute

flag acute triangles

5.55 tri_isobtuse

flag obtuse triangles

5.56 tri_semiperimeter

semiperimeter of a triangle

5.57 tri_side_length

edge length of triangle

6 geometry

6.1 Polygon

Simple 2D polygon class

Polygon properties:

- x - x coordinates of polygon
- y - y coordinates of polygon
- nnodes - number of nodes in the polygon

Polygon methods:

- in - checks whether given points lie inside, on the edge, or outside of the polygon
- area - returns the area of the polygon
- centerline - computes the centerline of the river
- iscw - check whether polygon is clockwise
- reverse - reverse the order of the polygon

6.2 bounding_box

bounding box of X

6.3 curvature_1d

curvature of a sampled parametric curve in two dimensions

6.4 cvt

centroidal voronoi tessellation

6.5 deg_to_frac

degree, minutes and seconds to fractions

6.6 ellipse

n-points on an ellipse

6.7 ellipseX

x-coordinates of y-coordinates of an ellipse

6.8 ellipseY

6.9 first_intersect

get first intersection between lines in A and B

6.10 golden_ratio

golden ratio

6.11 hypot3

hypothenuse in 3D

6.12 meanangle

weighted mean of angles

6.13 meanangle2

mean angle

6.14 meanangle3

mean angle

6.15 meanangle4

mean angle

6.16 medianangle

median angle
angle, that has the smallest squared distance to all others

6.17 medianangle2

median angle

input
alpha : x*m, [rad] angle

ouput
ma : 1*m, [rad] median angle
sa : 1*m, [rad] standard error of median angle for uncorrelated
error

6.18 pilim

limit to $\pm \pi$

6.19 streamline_radius_of_curvature

streamline radius of curvature
simplifies when rotate to streamwise coordinates to $R = 1/dv/ds * u$

7 linear-algebra

7.1 averaging_matrix_2

7.2 colnorm

norms of columns

7.3 condest_

estimation of the condition number

8 linear-algebra/coordinate-transformation

8.1 barycentric2cartesian

barycentric to cartesian coordinates

8.2 barycentric2cartesian3

convert barycentric to cartesian coordinates

8.3 cartesian2barycentric

cartesian to barycentric coordinates

8.4 cartesian_to_unit_triangle_basis

transform coodinates into unit triangle

8.5 example_approximate_utm_conversion

8.6 latlon2utm

transform latitude and longitude to WGS84 UTM

8.7 latlon2utm_simple

8.8 lowrance_mercator_to_wgs84

convert lowrance coordinates to wgs84

based on spreadsheet by D Whitney King and Patty B at Lowrance

8.9 nmea2utm

convert nmea messages to utm coordinates

8.10 sn2xy

convert sn to xy coordinates

8.11 unit_triangle_to_cartesian

transform coordinates in unit triangle to cartesian coordinates

8.12 utm2latlon

convert wgs84 utm to latitude and longitude

8.13 xy2nt

project all points onto the cross section and assign them nz-coordinates

transform coordinate into N-T reference
rotate coordinate, so that cross section goes along x-axis
then x and y are n and t respectively scaled by width
N and T coordinates

8.14 xy2sn

convert cartesian to streamwise coordiantes

8.15 xy2sn_java

use java port for speed up

8.16 xy2sn_old

transform points from cartesian into streamwise coordinates

NOTE : prefer the java version, this has some problems with round off

9 linear-algebra

9.1 det2x2

2x2 matrix inverse of 2x2 matrices stacked along dim 3

9.2 det3x3

determinant of stacked 3x3 matrices

9.3 det4x4

determinant of stacked 4x4 matrices

9.4 diag2x2

diagonal of stacked 2x2 matrices

9.5 eig2x2

eigenvalues of stacked 2x2 matrices

9.6 first

9.7 gershgorin_circle

range of eigenvalues determined by the gershgorin circle theorem

9.8 haussdorff

haussdorf dimension

box counting: count rectangles passed through by line (covered by polygon)

Koch snow flake 3:4 -> 1.2619

Kantor set 2:3, (4:9) -> 0.6309

quadrat 4:2, 9:3, 16:4 -> 2

9.9 ieig2x2

reconstruct matrix from eigenvalue decomposition

9.10 inv2x2

2x2 inverse of stacked matrices

9.11 inv3x3

9.12 inv4x4

inverse of stacked 4x4 matrices

9.13 lpmean

mean of pth-power of a

9.14 lpnorm

norm of lth-power of a

9.15 `matvec3`

matrix-vector product of stacked matrices and vectors

9.16 `max2d`

maximum value and i-j index for matrix

9.17 `mpoweri`

approximation of A^p , where p is not integer by quadratic interpolation

9.18 `mtimes2x2`

9.19 `mtimes3x3`

product of stacked 3x3 matrices

9.20 `nannorm`

norm of a vector, skips nan-values

9.21 `nanshift`

shift vector, but set out of range values to NaN

9.22 `nl`

number rows (lines) of a matrix

analogue to unix `nl` command

9.23 normalise

normalise a vector or the columns of a matrix
note that the columns are independently normalised, and hence not
necessarily
orthogonal to each other use the gram schmidt algorithm for this (
qr or orth)

9.24 normalize1

normalize columns in x to [-1,1]

9.25 normrows

9.26 orth2

make matrix A orthogonal to B

9.27 orth_man

orthogonalize the columns of A

9.28 orthogonalise

make x orthogonal to Y

9.29 paddext

padd values to vactor
not suitable for noisy data
order = 0 : constant extrapolation (hold)
order = 1 : linear extrapolation

9.30 paddval1

padd values at end of x

9.31 paddval2

padd values to x

10 linear-algebra/polynomial

10.1 chebychev

chebycheff polynomials

10.2 piecewise_polynomial

evaluate piecewise polynomial

10.3 roots1

roots of linear functions

10.4 roots2

roots of quadratic function
 $c_1 x^2 + c_2 x + c_3 = 0$

10.5 vanderi_1d

vandermonde matrix of an integral

10.6 vandermonde

van der monde matrix

11 linear-algebra

11.1 randrot

random rotation matrix

11.2 right

get right column by shifting columns to left
extrapolate rightmost column

11.3 rot2

rotation matrix from angle

11.4 rot2dir

rotation matrix from direction vector

11.5 rot3

11.6 rownorm

11.7 simmilarity_matrix

11.8 spnorm

frobenius norm

11.9 spzeros

allocate a sparze matrix of zeros

11.10 transpose3

transpose stacked 3x3 matrices

11.11 transposeall

12 logic

bitwise operations on integers

12.1 bitor_man

bitwise OR of the numbers of the columns of A

input:
A (positive integer)

13 number-theory

13.1 ceiln

floor to leading n-digits

13.2 digitsb

number of digits with respect to specified base

13.3 floorn

floor to n-digits

13.4 iseven

true for even numbers in X

13.5 multichoosek

all combinations of length k from set values with repetitions
c.f. nchoosek, combinations without repetition

input :
 x : scalar integer or vector of arbitrary numbers
 k : length of subsets
output :
 if x scalar : number of combinations
 if x vector : the exact combinations

13.6 nchoosek_man

vectorised binomial coefficient
 $b = N! / K! (N-K)!$

13.7 pythagorean_triple

pythagorean triple

13.8 roundn

round to n digits

14 numerical-methods/differentiation

14.1 derivative1

first derivative on variable mesh
second order accurate

14.2 derivative2

second derivative on a variable mesh

15 numerical-methods/finite-difference

15.1 cdiff

differences of columns of X
degree = 1 : central first order differences
degree = 2 : central second order differences

15.2 cdiffb

differences of columns of X
degree = 1 : central first order differences
degree = 2 : central second order differences
TODO use difference matrix function for simplicity

15.3 cmean

single gaussian smoothing step with kernel $1/4*[1,2,1]$

15.4 derivative_matrix_1_1d

finite difference matrix of first derivative in one dimensions

15.5 derivative_matrix_2_1d

finite derivative matrix of second derivative in one dimension

15.6 derivative_matrix_2d

finite difference derivative matrix in two dimensions

15.7 derivative_matrix_curvilinear

derivative matrix on a curvilinear grid

15.8 derivative_matrix_curvilinear_2

derivative matrix on a two dimensional curvilinear grid
the grid has not necessarily to be orthogonal

15.9 difference_kernel

difference kernels for equispaced grids
c.f. Computing the Spectrum of the Confined Hydrogen Atom, Kastner,
2012

15.10 distmat

distance matrix for a 2 dimensional rectangular matrix

15.11 gradpde2d

objective function gradiend on two dimensional regular grid
numeric gradient for non-linear least squares optimisation
of a PDE on a rectangular grid
 $x_* = \min(f(x))$
 $f = (v(x) - v(x_*))^2 = f(x) + A \, dx + O(dx^2)$
 $a_{ij} = df_i/dx_j$

15.12 laplacian

15.13 laplacian_fdm

finite difference matrix of the laplacian
BC

15.14 left

left element of vector, leftmost column is extrapolated

15.15 lrmean

mean of the left and right element

15.16 mid

mid point between neighbouring vector elements

15.17 pwmid

segment end point to segment mid point transformation for regular 1
d grids

15.18 ratio

ratio of two subsequent values

15.19 steplength

step length of a vector if it were equispaced

15.20 swapoddeven

swap odd and even elements in a vector

15.21 test_derivative_matrix_2d

15.22 test_derivative_matrix_curvilinear

15.23 test_difference_kernel

16 numerical-methods/finite-volume/@Advection

16.1 Advection

FVM treatment of the Advection equation

16.2 dot_advection

advection equation

17 numerical-methods/finite-volume/@Burgers

17.1 burgers_split

viscous Burgers' equation,
mixed analytic and numerical derivative in frequency space
by splitting scheme
 $u_t = -(0.5*u^2)_x + c*u_{xx}$

17.2 dot_burgers_fdm

viscous burgers' equation
 $u_t = -d/dx (1/2*u^2) + c d^2/dx^2 u_{xx}$

17.3 dot_burgers_fft

viscous Burgers' equation in frequency space
 $u_t + (0.5*u^2)_x = c*u_{xx}$

18 numerical-methods/finite-volume/@Finite_Volume

18.1 Finite_Volume

finite volume method for partial differential equations 1+1
dimensions
(time and space)

18.2 apply_bc

apply boundary conditions

18.3 solve

solve the the PDE by successively stepping in time
this is a trivial implmentation with constant step length
severity of diffusive error depends on dt/dx-ratio
stability depends on wave height

```
printf('Progress %2.1f%% %2.1fs\n',100*(t-Ti
(1))/(Ti(2)-Ti(1)),t_real);
```

18.4 step_split_strang

step in time, treat inhomogeneous part by Strang splitting
this scheme is not suitable for stationary solutions, for example
steady shallow water flow

18.5 step_unsplit

step in time, without splitting the inhomogeneous term

19 numerical-methods/finite-volume/@Flux_Limiter

19.1 Flux_Limiter

class of flux limiters

19.2 beam_warming

beam warming scheme
low resolution
note: works only if sign of eigenvalues point into the same
direction according to RL

19.3 fromm

fromme limiter
low res

19.4 lax_wendroff

lax wendroff scheme
second order accurate, but no tvd
this is effectively not a limiter
eq. 6.39 in randall, leveque

19.5 minmod

min-mod schock limiter

19.6 monotized_central

monotonized central flux limiter

19.7 muscl

muscl flux limiter

19.8 superbee

superbee limiter

19.9 upwind

godunov scheme
godunov, first order accurate

19.10 vanLeer

van Leer limiter

20 numerical-methods/finite-volume/@KDV

20.1 dot_kdv_fdm

korteweg de vries equation
 $u_t + (0.5u^2)_x = c u_{xxx}$

20.2 dot_kdv_fft

korteweg de vries equation
compute derivatives in frequency space
 $u_t + (0.5u^2)_x = c u_{xxx}$

20.3 kdv_split

korteweg de vries equation in frequency space,
derivative treated by splitting scheme

21 numerical-methods/finite-volume/@Reconstruct_Average_Evolve

21.1 Reconstruct_Average_Evolve

Reconstruct Average Evolve Finite Volume Method for treatment of
1+1D pdes

McCronack Scheme
 $\text{err} = O(\Delta t^2) + O(\Delta x^2)$, except as discontinuities
error:

```

h_xxx(3:end-2) = 1/dx^3*( -0.5*h(1:end-4) + h(2:end-3) - h(4:
    end-1) + 0.5*h(5:end) );
th = -1/6*dx^2*qh_.*(1 - (qh_*dt/dx).^2).*h_xxx;

```

21.2 advect_highres

single time step for the reconstruct evolve algorithm

21.3 advect_lowress

single time step
low resolution

22 numerical-methods/finite-volume

22.1 Godunov

Godunov, upwind method for systems of pdes

22.2 Lax_Friedrich

Lax-Friedrich-Method
for hyperbolic conservation laws
 $\text{err} = O(\text{dt}) + O(\text{dx})$
 $|a \text{ dt/dx}| < 1$

22.3 Measure

22.4 Roe

non linear roe solver for the SWE (randall, leveque 15.3.1)

The roe solver guarantess:

- A is diagonalisable with real eigenvalues (15.12)
- can be determined by a closed formula
- is an efficient replacement for true Rieman solver

22.5 fv_swe

wrapper for solving SWE

22.6 staggered_euler

forward euler method with staggered grid

22.7 staggered_grid

staggered grid approximation to the SWE

23 numerical-methods

23.1 grid2quad

extract rectangular elements of a structured grid
in form of an unstructured quad-mesh format

24 numerical-methods/integration

24.1 cumintL

cumulative integral from left to right

24.2 cumintR

cumulative integral from right to left

24.3 int_trapezoidal

integrate y along x with the trapezoidal rule

25 numerical-methods/interpolation/@Kriging

25.1 Kriging

```
class for Kriging interpolation
```

25.2 estimate_semivariance

```
estimate the parameter of the semivariance model for Kriging
interpolation
    % set up the regression matrix and solve for
    parameters
```

25.3 interpolate_

```
interpolate with Krieking method

this function may interpolate several quantities per coordinate,
using the same variogram, if the semivariance of the quantities
differs,
the user may prefer to estimate the semivariance and interpolate
each quantity
individually

Xs : source point coordinates
Vs : value at source points
Xt : targe point coordinates
Vt : value at target points
E2t : squared interpolation error at target points
```

26 numerical-methods/interpolation/@RegularizedInterpolator

26.1 RegularizedInterpolator1

```
class for regularized interpolation (Thikonov) on a 1D mesh
```

26.2 init

```
initialize the interpolator with a set of sampling points
```


27 numerical-methods/interpolation/@RegularizedInterpolator2

27.1 RegularizedInterpolator2

class for regularized interpolation on an unstructures mesh (
interpolation)

27.2 init

initialize the interpolator with a set of point samples

28 numerical-methods/interpolation/@RegularizedInterpolator3

28.1 RegularizedInterpolator3

class for regularized interpolation (Tikhonov) on a triangulation
(unstructured mesh)

28.2 init

initialize the interpolator with a set of sampling points

29 numerical-methods/interpolation

29.1 IDW

spatial averaging by inverse distance weighting

29.2 IPoly

polynomial interpolation class

29.3 IRBM

```
interpolate by the radial basis function method
    fprintf(1,'Progress IRBM: %d%%\n',round(100*
        idx/size(Xi,1)));
```

29.4 ISparse

```
sparse interpolation class
```

29.5 Inn

```
nearest neighbour interpolation
```

29.6 Interpolator

```
interpolator super-class
    fprintf(1,'Progress: %f%% %fs\n',100*
        idx/size(Xt,1),t);
```

29.7 fixnan

```
fill nan-values in vector with gaps
```

29.8 idw1

```
spatial average by inverse distance weighting
```

29.9 idw2

```
spatial average by inverse distance weighting
```

29.10 inner2outer

linear interpolation of segment mid point to grid points at segment
ends
assumes equal grid spacing

29.11 inner2outer2

interpolate from element (segment) centres to edge points

29.12 interp1_limited

interpolate values, but not beyond a certain distance
this function is idempotent, i.e. it will not extrapolate over into
gaps
exceedint the limit and thus not spuriously extend the series when
called a second time on the same data

29.13 interp1_man

interpolate

29.14 interp1_save

make interpolation save to round off errors
the matlab internal interpolation suffers from rounding errors,
which
are unacceptable when values of X and Y are large (for example UTM
coordinates)
this normalization prevents this

29.15 interp1_slope

quadratic interpolation returning value and derivative(s)

29.16 interp1_smooth

29.17 interp1_unique

matlab fails to interpolate, when x values are not unique
this function makes the values unique before use

29.18 interp2_man

nearest neighbour interpolation in two dimensions

29.19 interp_angle

interpolate an angle

29.20 interp_fourier

interpolation by the fourier method

29.21 interp_fourier_batch

batch interpolation by the fourier interpolation

29.22 interp_sn

interpolate along streamwise coordinates
This gives similar result to setting aspect ratio for sN to
infinity,
but not quite, as the input point set is not dense (scale for sN to
infinity does not work)
 sdx = sdx(sdx_);

29.23 interp_sn2

interpolation in streamwise coordinates

29.24 interp_sn3

29.25 interp_sn_

29.26 limit_by_distance_1d

smooth subsequent values along a curve such that
 $v(x_0+dx) < v(x_0) + (\text{ratio}-1)*dx$
if v is the edge length in a resampled polygon, then $v_i/v_{(i+1)} <$
ratio
 $\text{ratio}^1 = \exp(a*1)$

29.27 resample1

interpolation along a parametric curve with variable step width

29.28 resample_d_min

resample a function

29.29 resample_vector

resample a track so that velocity vectors do not run into each other

29.30 test_interp1_limited

30 numerical-methods

30.1 inverse_complex

31 numerical-methods/ode

31.1 bvp2_check_arguments

31.2 bvp2c

solve system of non-linear second order odes (in more than one variable)
as boundary value problems

odefun provides ode coefficients c:
$$c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4)$$
$$c_1 y'' + c_2 y' + c_3 y + c_4 = c_4$$

subject to the boundary conditions
bcfun provides v and p and optionally q, so that:

$$b_1 y + b_2 y' = f$$
$$q(x,1)*(p(x,1) y_l(x) + p(x,2) y_l'(x)) + q(x,2)*(p(x,1) y_r(x) + p(x,2) y_r'(x)) = v(x)$$

where q weighs the waves travelling from left to right and right to left (default [1 1])

31.3 bvp2c2

solve second order boundary value problem via roots of the characteristic polynomial

input:

x : [nx1] discretized domain
n : number of vertices
nxc = n-1 : number of segments

```

bc : struct : boundary condition
    bc.p(1)*y(0) + bc.pd(2)*y'(0) = bc.val(1)
    bc.p(2)*y(L) + bc.pd(2)*y'(L) = bc.val(2)

```

output:

```

A : [2*nxc x 2*ns] discrctisation matrix
rhs : [2*nxc x 1] right hand size

```

```

y = A^-1 rhs

```

31.4 bvp2fdm

solve system of non-linear second order odes (in more than one variable)

as boundary value problems by the finite difference method

odefun provides ode coefficients c:

```

c(x,1) y''(x) + c(x,2) y'(x) + c(x,3) y = c(x,4)
c_1 y'' + c_2 y' + c_3 y + c_4 = 0

```

subject to the boundary conditions

bcfun provides v and p and optionally q, so that:

```

b_1 y + b_2 y' = f
q(x,1)*( p(x,1) y_l(x) + p(x,2) y_l'(x)
+ q(x,2)*( p(x,1) y_r(x) + p(x,2) y_r'(x) = v(x)

```

where q weighs the waves travelling from left to right and right to left (default [1 1])

31.5 bvp2wavetrain

solve second order boundary value problem by repeated integration

31.6 bvp2wavetwopass

two pass solution for the linearised wave equation

solve first for the wave number k, and then for y

31.7 ivp_euler_forward

solve intial value problem by the euler forward method

31.8 ivprk2

solve initial value problem by the two step runge kutta method

31.9 ode2_matrix

transformation matrix of second order ode
to left and right going wave

```
c = odefun(x)
c1 y'' + c2' y + c3 y == 0
y = y_p + y_m, left and right going wave
d/dx [y_p, y_m] = A*[y_m, y_p]
```

31.10 ode2characteristic

second order odes
transmitted and reflected wave

31.11 step_trapezoidal

single trapezoidal step

31.12 test_bvp2

32 numerical-methods/optimisation

32.1 armijo_stopping_criterion

armijo stopping criterion for optimizations

32.2 astar

astar path finding algorithm

32.3 binsearch

binary search on a line

32.4 bisection

bisection

32.5 box1

test objective function for optimisation routines

32.6 box2

32.7 cauchy

32.8 cauchy2

solve non-linear system by cuachy's method
slower than quadratic optimisation, but does not require a hessian
fun : objective function, returns
 f : scalar, objective function value
 g : nx1, gradient
x : nx1, initial position
opt : options

32.9 directional_derivative

directional (projected) derivative
d : derivative, highest first
p : series expansion around x0

32.10 dud

optimization by the dud algorithm

32.11 extreme3

extract maxima by quadratic approximation from sampled function `val`
(`t`)
intended to be called after `[mval, mid] = max(val)` for refinement
of
location and maximum

input
`t` : sampling time (uniformly spaced)
`v` : values at sampling times
output:
`tdx` : index where extremum should be computed
`t0` : location of the extremum
`val0` : value of extremum

$v'(dt0) = 0$ and $v''(dt0)$ determines type of extremum

32.12 extreme_quadratic

32.13 ftest

32.14 grad

numerical gradient

32.15 hessian

numerical hessian

32.16 hessian_from_gradient

numerical hessian from gradient

32.17 hessian_projected

numerical hessian projected to one dimension

32.18 line_search

bisection routine

32.19 line_search2

bisection method

fun : objective funct
x0 : start value
f0 : objective function value at x0
g : gradient at x0
p : search direction from x0 (p = g for steepest descend)
h : initial step length (default 1)
lb : lower bound for x
up : upper bound for x

32.20 line_search_polynomial

polynomial line search
fun : objective funct
x0 : start value
f0 : objective function value at x0
g : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h : initial step length (default 1)
lb : lower bound for x
up : upper bound for x

32.21 line_search_polynomial2

```
cubic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

32.22 line_search_quadratic

```
quadratic line search
fun : objective funct
x0  : start value
f0  : objective function value at x0
g   : gradient at x0
dir : search direction from x0 (p = g for steepest descend)
h   : initial step length (default 1)
lb  : lower bound for x
up  : upper bound for x
```

32.23 line_search_quadratic2

```
quadratic line search
```

32.24 line_search_wolfe

```
line search by wolfe method
c.f.: OPTIMIZATION THEORY AND METHODS - Nonlinear Programming, Sun,
      Yuan
```

32.25 ls_bgfs

```
least squares by the bgfs method
```

32.26 ls_broyden

least squares by the broyden method
for rectangular / non symmetric systems
Numerical Optimization nodedal
Practical Methods of Optimization fletcher
c.f. gerber 1981
c.f. fletcher 1978 (more advanced, not used here)
c.f. Kelley 1999 ch. 4

BGFS:
Broyden 1965
Fletcher 1970
Goldfarb 1970
Shanno 1970

32.27 ls_generalized_secant

least squares by the secant method
Barnes, 1965
Wolfe, 1959
Fletcher 1980, 6.3
seber 2003
gerber

32.28 nlcg

non-linear conjugate gradient
input:
x : nx1 start vectort
opt : struct options
fdx : gradient constraint

32.29 nlls

non-linear least squares

32.30 picard

picard iteration

32.31 poly_extrema

extrema of a polynomial

32.32 quadratic_function

evaluate quadratic function in higher dimensions

32.33 quadratic_programming

optimize by quadratic programming

32.34 quadratic_step

single step of the quadratic programming

32.35 rosenbrock

rosenbrock test function

32.36 sqrt_heron

Heron's method for the square root

32.37 test_directional_derivative

32.38 test_dud

32.39 test_line_search_quadratic2

32.40 test_ls_generalized_secant

32.41 test_nlcg_6_order

32.42 test_nlls

```
f = w'*(p*abs(x-1).^4) + w'*(1-p)*abs(x-1).^2;
```

33 numerical-methods/piecewise-polynomials

33.1 Hermite1

hermite polynomial interpolation in 1d

33.2 hp2_fit

fit a hermite polynomial
coefficients are derivative free
x0 : left point of first segment
x1 : right point of last segment
n : number of segments
x : sample x-value
val : sample y-value
c : coefficients (values at points, no derivatives)

33.3 hp2_predict

prediction with pw hermite polynomial
c are values at support points

33.4 hp_predict

predict with piecewise hermite polynomial

33.5 hp_regress

fit piecewise hermite polynomial
coefficients are values and derivatives

33.6 lp_count

lagrangian basis for interpolation
count number of valid samples

33.7 lp_predict

lagrangian basis piecwie interpolation, predicor

33.8 lp_regress

33.9 lp_regress_

34 regression/@PolyOLS

34.1 PolyOLS

class for polynomial least squares

34.2 coefftest

34.3 detrend

detrending by polynomial regression

34.4 fit

fit a polynomial function
like polyfit, but returns parameter error estimates

34.5 fit_

fit a polynomial function

34.6 predict

predict polynomial function values

34.7 predict_

34.8 slope

slope by linear regression

35 regression/@PowerLS

35.1 PowerLS

class for power law regression

35.2 fit

fit a power law
like polyfit, but returns parameter error estimates

35.3 predict

```
predict with power law
    S2 = diag((A*obj.C)*A');
    L  = Y - S;
    U  = Y + S;
```

35.4 predict_

36 regression/@Theil

36.1 Theil

Kendal-Theil-Sen robust regression

36.2 detrend

linear detrending of a set of samples by the Theil-Senn Slope

36.3 fit

fit slope and intercept to a set of sample with the Theil-Sen method

```
c      : confidence interval c = 2*ns*normcdf(1) for ns-sigma
        intervals
param : itercept and slope
P      : confidence interval
```

36.4 predict

predict values and confidence intervals with the Theil-Sen method

36.5 slope

fit the slope with the Theil-Sen method

37 regression

linear and non-linear regression

37.1 Theil_Multivariate

extension of the Theil-Senn regression to higher dimensions by means of the Gauss-Seidel iteration

37.2 areg

regression using the pth-fraction of samples with smallest residual

37.3 ginireg

gini regression

37.4 hesssimplereg

hessian, gradient and objective function value of the simple regression
 $\text{rhs} = p(1) + p(2) x + \text{eps}$

37.5 l1lin

solve $\|Ax - b\|_{L1}$ by means of linear programming

37.6 lsq_sparam

parameter covariance of the least squares regression

fun : model function for prediction
b : sample values
 $f(p) = b$
p : parameter at point of evaluation (preferably optimum)

37.7 polyfitd

fit a polynomial of order n to a set of sampled values and sampled values of the derivative

x0 must contain at least for conditioning as otherwise the intercept cannot be determined

37.8 regression_method_of_moments

fit linear function $\|a \ b \ x = y\|_{L2}$ by the method of moments
 $y + \text{eps} = \text{alpha} + \text{beta} * x$

37.9 robustlinreg

fit a linear function by splitting the x-values at their median
 $(\text{med}(y_{\text{left}}) - \text{med}(y_{\text{right}})) / (\text{med}(x_{\text{left}}) - \text{med}(x_{\text{right}}))$
this approach performs poorly compared to the theil-senn operator

37.10 theil2

Theil senn-estimator for two dimensions (glm)

37.11 theil_generalised

generalization of the Theil-Senn operator to higher dimensions,
for arbitrary functions such as polynomials and multivariate
regression
either higher order polynomials or glm
c.f. "On theil's fitting method", Pegoraro, 1991

37.12 total_least_squares

total least squares

37.13 weighted_median_regression

weighted median regression
c.f. Scholz, 1978

38 set-theory

38.1 issubset

test if set B is subset of A in $O(n)$ -runtime

A : first set
B : second set
P : set of primes (auxiliary)

39 signal-processing

39.1 acf_effective_sample_size

effective sample size from acf

39.2 acf_genton

autocorrelation function

39.3 acfar1

Autocorrelation function of the finite AR1 process

$$\begin{aligned} a_k &= 1/(n-k) \sum x_{i-k} x_i + (x_i + x_{i+k})\mu + \mu^2 \\ &= r^k + 1/n \sum_{i=k+1}^n x_i + 1/n \\ &\quad \text{pause} \end{aligned}$$

39.4 acfar1_2

autocorrelation of the ar1 process

39.5 acfar2

impulse response of the ar2 process

39.6 acfar2_2

autocorrelation of the ar2 process
 $X_i + a_1 X_{i-1} + a_2 X_{i-2} = 0$

39.7 ar1_cutoff_frequency

39.8 ar1_effective_sample_size

effective sample size correction for autocorrelated series

39.9 ar1_mse_mu_single_sample

standard error of a single sample of an ar1 correlated process

39.10 ar1_mse_pop

variance of the population mean of a single realisation around zero

$$E[(\mu_N - 0)^2] = E[\mu_N^2]$$

39.11 ar1_mse_range

mean standard error of the mean of a range of values taken from an ar1 process

39.12 ar1_spectrum

spectrum of the ar1 process

39.13 ar1_to_tikhonov

convert ar1 correlation to tikhonovs lambda

39.14 ar1_var_factor

variance correction factor for an autocorrelated finite process

n : [1 .. inf] population size

m : [1 .. n] samples size

rho : [-1 < rho < 1 (for convergence)] correlation of samples

39.15 ar1_var_factor_

variance of an autocorrelated finite process

39.16 ar1_var_range2

variance of sub sample starting at the end of the series

from the finite length first order autocorrelated process

$$s2 = 1/m^2 \sum_i \sum_j \rho^{|i-j|}$$

39.17 ar1delay

approximate acf by the ar1 process
acf: autocovariance or autocorrelation function
nf : skip first samples (for mixed geometric-arithmetic series (ARMA))

39.18 ar1delay_old

autocorrelation of the residual

39.19 ar2conv

coefficients of the ar2 process determined from the two leading correlations
of the acf [1,r1,r2,...]

39.20 ar2dof

effective samples size for the ar2 process

39.21 ar2param

ar2 parameter estimation from first two terms of acf
acf = [1 a1 a2 ...]

39.22 asymwin

creates asymmetrical filter windows
filter will always have negative weights

39.23 autocorr_fft

autocorrelation function

39.24 bandpass

bandpass filter

39.25 bandpass2

bandpass filter

39.26 bartlett

Effective sample size factor for bartlett window
c.f. thiebaux
c.f spectral analysis-jenkins, eq. (6.3.27)
 $c = acf$
note: results seams always to be 1 tac too low
T : reduction factor for dof
for ar1 with $a = \rho^k = \exp(-k/L)$, $T = 2L$

39.27 bartlett_spectrogram

bartlet spectrogramm
TODO sliding window

39.28 bin1d

bin values of v sampled at x into bins bounded by "edges"
apply function v to it

39.29 bin2d

bin values of V sampled at X and Y into the grid structured grid ex
,ey
apply function func to all walues in the bin
func = mean : default
func = sum : non-normalized frequency histogram in 2D

39.30 binormrnd

generate two correlated normally distributed vectors

39.31 conv1_man

convolutions with padding

39.32 conv2_man

convolution in 2d

39.33 conv2z

39.34 conv30

convolve with rectangular window of length n
circular boundaries

39.35 conv_

convolution of a with b

39.36 conv_centered

convolve x with filter window f
when length of f is even, this guarantees a symmetric result (no
off by on
displacement) by making the length of f odd at first

39.37 convz

39.38 cosexpdelay

39.39 csmooth

smooth recursively with $[1,2,1]/4$ kernel

39.40 daniell_window

Daniell window for smoothing the power spectrum
c.f. Daniell 1946
Bloomfield 2000
meko 2015

39.41 danielle_window

danielle fourier window

39.42 db2neper

convert decibel to neper

39.43 db2power

power ratio from db

39.44 derive_danielle_weight

39.45 derive_limit_0_acfar

39.46 detect_peak

detect peaks in a vector
requires function value to fall to $p \cdot \max$ before new value is
allowed

39.47 digital_low_pass_filter

design coefficients of a low pass filter with specified cut of
frequency
and sampling period
analogue low pass with pole at $s = -\omega_c = 1/\tau = 1/RC$
 $H_a = \tau / (\tau + s) = 1 / (1 + \omega_c \cdot s)$

39.48 doublesum_ij

double sum of r^i

39.49 effective_sample_size_to_ar1

convert effective sample size to ar1 correlation

39.50 flt_hodges_lehman

39.51 filter1

filter along one dimension

39.52 filter2

filter columns of x (matlab does only support vector input)

39.53 filter_

invalidate values that exceed n-times the robust standard deviation

39.54 filteriir

filter adcp t-n data over time

v : nz,nt : values to be filtered

H : nt,1 : depth of ensemble

last : nt,1 : last bin above bottom that can be sampled without
side lobe interference

nf : scalar : number of reweighted iterations

when samples

- distance to bed is reference (advantageous for near-bed suspended
transport)

TODO for wash load: distance to surface is more relevant
interpolate depending on z

when depth changes, neighbouring indices do not correspond to same
relative position in the water column

relative position in the column (s-coordinate) smoothes values

near the bed: absolute distance to bed is chosen

near surface: absolute distance to surface is chosen

-> cubic transformation of index

faster and avoid aliasing (smoothing along z)

resample ensemble to same number of bins in S -> filter ->

resample back

use nonlinear transform z-s coordinates

-> resampling has to be local (Hi -> H-filtered)

filtered profile coordinates to sample coordinates

zf -> zi (special transform)

corresponding indices and fractions

filtration step (update of hf and vf)

sample coordinates to updated profile coordinates

(the inverse step is actually not necessary)

write filtered value

39.55 filterp

39.56 filterp1

fir filter with some fancy extras

39.57 filterstd

39.58 firls_man

design finite impulse response filter by the least squares method

39.59 flattopwin

the flat top window

39.60 frequency_response_boxcar

frequency response of a boxcar filter

39.61 freqz_boxcar

frequency response of a boxcar filter

39.62 gaussfilt1

filter data series with a gaussian window

39.63 hanchangewin

hanning window for change point detection

39.64 hanchangewin2

nanning window for chage point detection

39.65 hanwin

hanning filter window

39.66 hanwin_

hanning filter window

39.67 highpass

high pass filter

39.68 kaiserwin

kaiser filter window

39.69 kalman

Kalman filter

39.70 lanczoswin

Lanczos window

39.71 last

lake tail, but for matrices

39.72 lowpass

low pass filter

39.73 lowpass2

design low pass filter with cutoff-frequency f1

39.74 lowpass_iir

iir-low pass

39.75 lowpass_iir_symmetric

two-sided iir low pass filter (for symmetry)

39.76 lowpassfilter2

low-pass filter of data

39.77 maxfilt1

39.78 meanfilt1

moving average filter with special treatment of the boundaries

39.79 medfilt1_man

moving median filter, supports columnwise operation

39.80 medfilt1_man2

moving median filter with special treatment of boundaries

39.81 medfilt1_padded

median filter with padding

39.82 medfilt1_reduced

median filter with padding

39.83 mid_term_single_sample

variance of single sample, mid term

39.84 minfilt1

39.85 mu2ar1

error variance of the mean of the finite length ar1 process

$(\mu)^2 = (\sum \text{epsi})^2 = \sum_i \sum_j \text{epsi}_i \text{epsi}_j = \sum_{ii}(\rho, n)/n^2$
this has the limit s^2 for $\rho \rightarrow 1$

39.86 nanautocorr

autocorrelation with nan-values

39.87 nanmedfilt1

medfilt1, skipping nans

39.88 neper2db

convert neper to db

39.89 peaks_man

peaks of a periodogram

39.90 polyfilt1

polynomial filter,
can be achieved by iteratively processing the data with
a mean (zero-order) filter

39.91 qmedfilt1

medfilt1, after fitting a quadratic polynomial

39.92 randar1

generate random ar1 process
e1 = randar1(sigma,p,n,m)

39.93 randar1_dual

draw random variables of two correlated ar1 processes

39.94 randar2

generate ar2 process

39.95 randarp

randomly generate the instance of an ar-p process

39.96 range_window

range of values within a certain range of indices (window)

39.97 rectwin

rectangular window

39.98 recursive_sum

39.99 select_range

39.100 smooth1d_parametric

smooth position of $p_0=x_0, y_0$ between $p_1=x_1, y_1$ and $p_2=x_2, y_2$,
so that distance to p_1 and p_2 becomes equal
and the chord length remains the same

39.101 smooth2

smooth vectos of X

39.102 smooth_man

39.103 smooth_parametric

smooth a parametric function given in x-y coordinates
`matvec2x2(R, [dxc;dyc])`

39.104 smooth_parametric2

parametrically smooth the curve

39.105 smoothfft

filter with fast fourier transform

39.106 spectrogram

spectrogram

39.107 std_window

moving block standard deviation

39.108 sum_i_lag

sum of ar1 matrix with lag
 $\sum_{i=1}^n \rho^{|i-k|}$

39.109 sum_ii

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^n \rho^{|i-j|}$
this is for the variance, take square root for the standard
deviation factor

39.110 sum_ii_

39.111 sum_ij

sum of ar1 matrix
 $\sum_{i=1}^n \sum_{j=1}^m r^{|i-j|}$

39.112 `sum_ij_`

39.113 `sum_ij_partial_`

39.114 `sum_multivar`

`sum of matrix entries of bivariate ar1 process`

39.115 `test_acfar1`

39.116 `test_acfar1_2`

39.117 `test_acfar1_3`

39.118 `test_acfar1_4`

39.119 `test_acfar2`

39.120 `test_ar1_var_factor`

39.121 `test_ar1_var_factor_2`

39.122 test_ar1_var_mu_single_sample

39.123 test_ar1_var_pop

39.124 test_ar1_var_pop_1

39.125 test_ar1delay

39.126 test_bivariate_covariance_term

39.127 test_convexity

39.128 test_lanczoswin

39.129 test_madcorr

39.130 test_randar1

39.131 test_randar1_multivariate

39.132 test_randar2

39.133 test_sum_ij

39.134 test_sum_multivar

39.135 test_trifilt1

39.136 test_wautocorr

39.137 test_wavelet_transform

39.138 test_wordfilt

39.139 test_xar1_mid_term

39.140 tikhonov_to_ar1

convert coefficient of the tikhonov regularization to correlatioon
of the ar1 process

39.141 trapwin

trapezoidal filter window

39.142 trifilt1

filter with triangular window

39.143 triwin

triangular filter window

39.144 triwin2

triangular filter window

39.145 varar1

error variance of a single sample of a finite length ar1 process
with respect to the mean, averaged over the population

39.146 welch_spectrogram

welch spectrogram

39.147 wfilt

filter with window

39.148 winbandpass

filter with bandpass

39.149 `window_make_odd`

39.150 `winfilt0`

filter with window

39.151 `winlength`

window length for desired cutoff frequency
power at f_c is halved
 $H(wf) = 1/\sqrt{2} H(f)$
if the filter window were used as a low pass filter
note: the user should prefer a windowed ideal low pass filter
TODO, relate this to DOF

39.152 `wmeanfilt`

mean filter with window

39.153 `wmedfilt`

median filter with window

39.154 `wordfilt`

weighted order filter

39.155 `wordfilt_edgeworth`

weighed order filter

39.156 `xar1`

39.157 `xcorr_man`

cross correlation of two sampled ar1 processes

40 `sorting`

40.1 `sort2`

sort two numbers

40.2 `sort2d`

sort elements of matrix in X
returns row and column index of sorted values

41 `special-functions`

41.1 `bessel_sphere`

spherical Bessel function of the first kind

41.2 `hankel_sphere`

spherical Hankel function for the far field (incident plane wave)
first kind

41.3 `hermite`

probabilistic's hermite polynomial by recurrence relation

input :
n : order
x : value

output:
f : $H_n(x)$
df : $d/dx H_n(x)$

41.4 legendre_man

legendre polynomials

41.5 neumann_sphere

spherical Neumann function
Bessel function of the second kind

42 statistics

42.1 atan_s2

stadard deviation of the arcus tangens by means of taylor expansion

42.2 beta_mode_to_parameter

transform modes (mean and sd) to paramets of the beta function

42.3 correlation_confidence_pearson

confience intervals of the correlation coefficient
c.f. Fischer 1921

43 statistics/distributions

43.1 PDF

class for quasi-distributions from a set of sampling points

43.2 binorm_separation_coefficient

separation coefficient of a bimodal normal distribution

43.3 binormcdf

bio-modal gaussian distribution

43.4 binormfit

fit sum of to normal distribution to a histogram

43.5 binormpdf

43.6 edgeworth_cdf

edgeworth expansion of an unknown cumulative distribution
with mean μ , standard deviation σ , and third and fourth
cumulants
c.f. Rao 2010

43.7 edgeworth_pdf

probability density of and unknown distribution
with mean μ , standard deviation σ , and third and fourth
cumulants
c.f. Rao 2010

43.8 logn_mode2param

transform modes (μ, σ) to parameters of the log normal
distribution

43.9 logn_param2mode

transform parameters to mode (μ, σ) for the log normal
distribution

43.10 lognpdf_

log normal distribution called by modes rather than parameters

43.11 pdfsample

pdf from sample distribution

Note: better use kernel density estimates

43.12 t2cdf

Hotelling's T-squared cumulative distribution

43.13 t2inv

inverse of Hotelling's T-squared cumulative distribution

44 statistics

44.1 example_standard_error_of_sample_quantiles

44.2 f_var_finite

reduction of variance when sampling from a finite population
without replacement

44.3 gamma_mode_to_parameter

transform modes (mu,sd) to parameters of the gamma distribution

44.4 hedges_lehmann_correlation

hedges_lehmann correlatoon coefficient
c.f. Shamos 1976
c.f. Bickel and Lehmann 1976
c.f. rousseeuw 1993
c.f. Shevlyakov 2011

44.5 hedges_lehmann_dispersion

dispersion determined by the hedges lehman method
asymptotic efficiency of dispersion estimates:
standard deviation: $E(s - \hat{s})/s = 2/\sqrt{2n} \sim 0.707/\sqrt{n}$
(100%)
hedges lehmann dispersion $E(s - \hat{s})/s = (\pi/3)^2 / (\sqrt{2n}) \sim$
 $0.775/\sqrt{n}$ (91%)
mad $E(s - \hat{s})/s \sim 1.17 s/\sqrt{n}$
(60%)
c.f. Shamos 1976
c.f. Bickel and Lehmann 1976
c.f. rousseeuw 1993
nb: rousseeuw uses the 25th percentile, which is more efficient for
small sample sizes

45 statistics/information-theory

45.1 akaike_information_criterion

akaike information criterion

serr : rmse of model prediction
n : effective sample size
k : number of parameters

c.f. akaike (1974)
c.f. sugiura 1978

45.2 bayesian_information_criterion

bayesian information criterion

46 statistics

46.1 kurtncdf

46.2 kurtnpdf

46.3 kurtosis_bias_corrected

bias corrected kurtosis

46.4 limit

limit a by lower and upper bound

46.5 logfactorial

approximate log of the factorial

46.6 loglogpdf

46.7 logskewcdf

46.8 logskewpdf

47 statistics/logu

47.1 lambertw_numeric

lambert-w function

47.2 logtrialtcdf

pdf of a logarithmic triangular distribution

47.3 logtrialtinv

inverse of the logarithmic triangular distribution

$$= (d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) - \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a) - 1/\log(a)}) c)^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) + d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) + b \log(c) - d F \log(a) \log(c))) / (\log(a) - \log(b)))$$
$$x = (d F \log(a) \log(b) + a \log(b) - b \log(a) - d F \log(a) \log(c) - a \log(c) + d F \log(b) \log(c) + b \log(c) - d F \log^2(b)) / ((\log(a) - \log(b)) W((a^{-1/(\log(a) - \log(b))}) (b^{-\log(c)/\log(a) - 1/\log(a)}) c)^{-\log(a)/(\log(a) - \log(b))}) (-d F \log^2(b) + a \log(b) + d F \log(a) \log(b) + d F \log(c) \log(b) - b \log(a) - a \log(c) + b \log(c) - d F \log(a) \log(c))) / (\log(a) - \log(b)))$$

47.4 logtrialtmean

mean of the logarithmic triangular distribution

47.5 logtrialtpdf

density of the logarithmic triangular distribution

47.6 logtrialtrnd

47.7 logtricdf

cumulative distribution of the logarithmic triangular distribution

47.8 logtriinv

inverse of the logarithmic triangular distribution

47.9 logtrimean

mean of the logarithmic triangular distribution

47.10 logtripdf

probability density of the logarithmic triangular distribution

47.11 logtirnd

47.12 logucdf

probability density of the logarithmic uniform distribution

47.13 logucm

central moments of the log-uniform distribution

47.14 loguinv

inverse of the log-uniform distribution

47.15 logumean

mean of the log-uniform distribution

47.16 logupdf

pdf of the log uniform distribution

47.17 logurnd

random numbers following a log-uniform distribution

47.18 loguvar

variance of the log-uniform distribution

47.19 medlogu

median of the log-uniform distribution

47.20 test_logurnd

47.21 tricdf

cumulative distribution of the log-triangular distribution

47.22 triinv

inverse of the triangular distribution

47.23 trimedian

median of the triangular distribution

47.24 tripdf

probability density of the triangular distribution

47.25 trirnd

random numbers of the triangular distribution

48 statistics

48.1 maxnnormals

expected maximum of n normal variables
c.f. Wolperts
this is the median, not the mean of the maximum!
see median of gumbel

48.2 midrange

mid range of columns of X

48.3 minavg

solution of the minimum variance problem
minimise the variance of the weighted sum of n-independent
random variables with equal mean and individual variance

48.4 mode_man

49 statistics/moment-statistics

49.1 autocorr_man3

autocorrelation of the columns of X

49.2 autocorr_man4

autocorrelation for x if x is a vector, or individually for the columns of x if x is a matrix

c.f. box jenkins 2008 eq. 2.1.12

Note that it is faster to compute the acf in frequency space as done in the matlab internal function

49.3 autocorr_man5

autocorrelation of the columns of X

49.4 blockserr

estimate the standard error of potentially sequentially correlated data
by blocking
block length should be sufficiently larger than correlation length and sufficiently smaller than data length
this uses a sliding block approach, which reduces the variation of the error estimate

49.5 comoment

non-central higher order moments of the multivariate normal distribution

c.f. Moments and cumulants of the multivariate real and complex Gaussian distributions

note : there seem to be some typos in the original paper,
for x^4 c_{ii}^2 , the square seems to be missing

mu : nx1 mean vector
C : nxn covariance matrix
k : nx1 powers of variables in moments

49.6 corr_man

correlation of two vectors

49.7 cov_man

covariance matrix of two vectors

49.8 dof

mininum number of support points
for a polynomial of degree order in dim dimensions

49.9 edgeworth_quantile

inverse edgeworth expansion
c.f. cornis fisher 1937
c.f. Rao 2010
c.f. 2.50 in hall
CHERNOZHUKOV 3.3

49.10 effective_sample_size

effective sample size of the weighted mean of uncorrelated data
c.f. Kish

49.11 f_correlation

correction factor for standard error of the mean of n ar1-
correlated iid samples

49.12 `f_finite`

reduction factor of standard error for sampling from a finite
distribution
without replacement

49.13 `lmean`

mean of $x.^l$, not of `abs`

49.14 `lmoment`

l-moment of vector `x`

49.15 `maskmean`

mean of the masked values of `X`

49.16 `masknanmean`

49.17 `mean1`

mean of `x`

49.18 `mean_man`

mean and standard error of `X`

49.19 `mse`

mean squared error of residual vector `res`
this is de-facto the `std` for an unbiased residual

49.20 nanautocorr_man1

autocorrelation of a vector with nan-values

49.21 nanautocorr_man2

autocorrelation of a vector with nan-values

49.22 nanautocorr_man4

compute autocorrelation for x if x is a vector, or individually
for the
columns of x if x is a matrix
box jenkins 2008 eq. 2.1.12
TODO nan is problematic!
Note that it is faster to compute the acf in frequency space
as done in the matlab internal function

49.23 nancorr

(co)-correlation matrix when samples a NaN

49.24 nancumsum

cumulative sum, setting nan values to zero

49.25 nanlmean

mean of the l-th power of the absolute value of x

49.26 nanr2

coefficient of determination when samples are invalid

49.27 nanrms

root mean square value when sample contains nan-values

49.28 nanrmse

root mean square error from vector of residuals
this is de-facto the std for an unbiased residual

49.29 nanserr

standard error of x with respect to mean when x contains nan values

49.30 nanwmean

weighted mean
 $\min_x \sum w (x - \mu)^2 \Rightarrow \mu = \frac{\sum wx}{\sum w}$
varargin can be dim
function [mu serr] = nanwmean(w,x)

49.31 nanwstd

weighed standard deviation

49.32 nanwvar

weighted variance of columns, corrected for degrees of freedom (bessel)

$s^2 = \frac{\sum w (x - \frac{\sum wx}{\sum w})^2}{\sum w}$

49.33 nanxcorr

49.34 pearson

pearson correlation coefficient

49.35 pearson_to_kendall

conversion of pearson to kendall correlation coefficient
c.f. Kruskal 1958

49.36 pool_samples

pooled mean and standard deviation of several groups of different
size, mean and standard deviation

49.37 qmean

trimmed mean

49.38 range_mean

49.39 rmse

root mean square error computed from a residual vector
this is de-facto the std for an unbiased residual

49.40 serr

standard error of the mean of a set of uncorrelated samples

49.41 serr1

49.42 test_qskew

49.43 test_qstd_qskew_optimal_p

49.44 wautocorr

autocorrelation for x if x is a vector, or individually for the columns of x if x is a matrix
samples can be weighted

c.f. box_jenkins 2008 eq. 2.1.12

c.f. autocorr_man4

Note that it is faster to compute the acf in frequency space as done in the matlab internal function

49.45 wcorr

correlation of two vectors when samples are weighted

49.46 wcov

covariance of two vectors when samples are weighted

49.47 wdof

effective degrees of freedom for weighted samples

49.48 wkurt

kurtosis with weighted samples

49.49 wmean

weighted mean

$\min_x \sum w (x - \mu)^2 \Rightarrow \mu = \frac{\sum wx}{\sum w}$

varargin can be dim

function [mu serr] = wmean(w,x)

49.50 wrms

weighted root mean square error

49.51 wserr

weighted root mean square error

49.52 wskew

skewness of a weighted set of samples

49.53 wstd

weighed standard deviation

49.54 wvar

weighted variance of columns, corrected for degrees of freedom (bessel)

variance of the weighted sample mean of samples with same mean (but not necessarily same variance)

$s^2 = \frac{\sum w^2 (x - \sum wx)^2}{\sum w}$

s2_mu : error of mean, s2_mu : sd of prediction

50 statistics

50.1 nangeomean

50.2 nangeostd

geometric standard deviation ignoring nan-values

51 statistics/nonparametric-statistics

51.1 kernel1d

X : ouput x axis bins
xi : samples along x
m : number of bins in X
fun : kernel function
pdf : propability density of xi

51.2 kernel2d

kernel density estimate in two dimensions

52 statistics

52.1 normmmoment

expected norm of $x.^n$, when values x in x are iid normal with mu
and sigma

52.2 normpdf2

pdf of the bivariate normal distribution

53 statistics/order-statistics

53.1 hodges_lehmann_location

hodges lehman location estimator

Asymptotic rms efficiency of location estimate:

mean: $1 \text{ s}/\sqrt{n}$

hodges lehman: $\sqrt{\pi/3} * s \sim 1.0233 \text{ s}/\sqrt{n}$

median: $\pi/2 \text{ s}/\sqrt{n} \sim 1.25 \text{ s} / \sqrt{n}$

53.2 kendall

kendall correlation coefficient

53.3 kendall_to_pearson

convert kendall rank correlation coefficient to the person product
moment
correlation coefficient

c.f. Kruska, 1985

53.4 mad2sd

transform median absolute deviation to standard deviation
for normal distributed values

53.5 madcorr

proxy correlation by median absolute deviation

53.6 median2_holder

53.7 median_ci

median and its confidence intervals under assumption of normality
`se_me = sqrt(1/2 pi) 1.25331 * sd/sqrt(n)`

53.8 median_man

median and confidence intervals
`c` is a P value for the confidence interval,
default is 0.95 (2-sigma)
median of the columns of `X`

53.9 mediani

index of median, if median is not unique, any of the values is
chosen

53.10 nanmadcorr

proxy correlation by median absolute deviation

53.11 nanwmedian

weighted median, skips nan-values

53.12 nanwquantile

weighted quantile, skips nan values

53.13 oja_median

two dimensional oja median
note: the multivariate median is not unique

oja 1983, for extension to multivariate function, see chaudhri

53.14 qkurtosis

kurtosis computed for quantiles

Note : this is a measurement of shape-tailedness and yields the same value for the normal distribution as "kurtosis"
However, this is a separate statistic and hence requires different methods for calculating P-values and hypothesis testing

53.15 qmoments

moments estimated from quantiles

53.16 qskew

skewness estimated from quantiles

Note : this is a measurement of shape-symmetry and yields the same value for the skew-normal distribution as "skewness"
However, this is an own statistic and hence requires different methods for calculating P-values and hypothesis testing

53.17 qskewq

skewness estimated by quantiles

53.18 qstdq

proxy standard deviation determined by quantiles

53.19 quantile1_optimisation

53.20 quantile2_breckling

quantile regression

53.21 quantile2_chaudhuri

quantile regression

53.22 quantile2_projected

quantile in two dimensions

53.23 quantile2_projected2

spatial quantile for chosen direction

53.24 quantile_envelope

53.25 quantile_regression_simple

simple quantile regression

53.26 ranking

ranking for spearman statistics

53.27 spatial_median

c.f. Oja 2008
is this the same as the oja simplex median (c.f. small 1990)?

53.28 spatial_quantile

spatial quantile

53.29 spatial_quantile2

spatial quantile

53.30 spatial_quantile3

spatial quantile

53.31 spatial_rank

unsigned rank

53.32 spatial_sign

spatial sign

53.33 spatial_signed_rank

signed rank

Note: this is only a true rank if X is normal with zero mean,
arbitrary variance

53.34 spearman

spearman's product moment coefficient

53.35 spearman_rank

53.36 spearman_to_pearson

conversion of spearman rank to person product moment correlation coefficient

53.37 wmedian

weighted median

53.38 wquantile

weighted quantile

54 statistics

54.1 qstd

55 statistics/random-number-generation

55.1 laplacernd

random number of laplace distribution

55.2 randc

correlate to correlated standard normally distributed vectors

55.3 skewrnd

random numbers of the skew normal distribution

55.4 skewrnd2

random numbers of the skew normal distribution

56 statistics

56.1 range

mid range

57 statistics/resampling-statistics/@Jackknife

57.1 Jackknife

class for leave out 1 (delete 1) Jackknife estimates

note 1 : the 1-delete jackknife does not yield consistend estimates
for all functions,

in particular it will perform poorly on robust estimation
functions

this is overcome by the d-delete jackknife, where d has to
exceed the breakdown point

of the estimating function, for example \sqrt{n} for the
median

as this leads to unreasonably large number of repetitions,
bootstrap

is recommended for large sample cases (or blocking for
sequential data)

note 2 : as a linearisation, jackknife underestimates the error
variance in case of

dependence in the data

note 3 : studentisation and the leave out 1 jackknife are related

note 4 : the double 1 sample jackknife performs iferior to the d1
jackknife

57.2 estimated_STATIC

jackknife estimate of mean, bias and standard error

theta0 : estimate from all samples

thetad : set of estimates obtained by leaving out one data point
each

last dimension of theta is assumed to be the jackknife
dimension

57.3 matrix1_STATIC

matrix of estimation for leaving out two samples at a time

57.4 matrix2

matrix of estimations for jackknife with two samples left out

58 statistics/resampling-statistics

58.1 block_jackknife

58.2 jackknife_moments

moments determined by the jackknife

func : function of interest on the samples (e.g. mean)
A : parameter matrix
 columns : parameters
 rows : samples of the parameter sets
d : number of samples left out

58.3 moving_block_jackknife

blocked Jackknife for autocorrelated data
sliding block, statistically more efficient but computationally
expensive
note, number of blocks must be sufficiently large $h \sim \sqrt{n}$? $\ll n$

58.4 randblockterr

standard error of sequentially correlated data by blocking
block length should be sufficiently larger than correlation length
and sufficiently smaller than data length
this uses a sliding block approach, which reduces the variation of
the error estimate
TODO this does not work, randomly picking samples does not reveal
the correlation

58.5 resample

resample a vector and apply function to it

TODO, should be with replacement

n : number of samples
m : number of subsamples
cx : maximum number of combinations

59 statistics

59.1 scale_quantile_sd

scale factor for the standard deviation
of the asymptotic distribution of sample quantiles
(for normal distribution)
see cadwell, 1952

59.2 skewpdf

skew-normal distribution
c.f. Azzalini 1985

59.3 trimmed_mean

trimmed mean

59.4 ttest2_man

two-sample t-test
here posix return value standard: h = 0 accepted, h = 1 failed
note: the matlab logic is inverse : h = 1 accepted, h = 0 failed
two sided univariate t-test

59.5 ttest_man

two-sample t-test
unequal sample size
equal variance

59.6 ttest_paired

paired t-test
unequal sample size
equal variance
more powerfull than unpaired test, as long as correlation between
x1 and x2 > 0

59.7 wharmean

weighted harmonic mean

60 wavelet

60.1 continuous_wavelet_transform

continuous wavelet transform
follows "The Illustrated Wavelet Transform Handbook: Introductory
Theory and ..."

60.2 cwt_man

continuous fourier transform
as of time of implmentation, the matlab interal cwt is affected by
serious round-off errors and has issues with the scaling,
which is not the case here

60.3 example_wavelets

60.4 phasewrap

wrap the phase to +/- pi

60.5 test_cwt_man

60.6 test_phasewrap

60.7 test_wavelet

60.8 test_wavelet2

60.9 test_wavelet_analysis

60.10 test_wavelet_reconstruct

60.11 test_wtc

60.12 wavelet

wavelet windows

60.13 wavelet_reconstruct

inverses wavelet transform for single frequency
(reconstruction of time series)
n : window lengths in multiples of filter period $1/f_0$

60.14 wavelet_transform

wavelet transform for single frequency
n : window lengths in multiples of filter period $1/f_0$

61 mathematics

mathematical functions of various kind

61.1 wrapphase