

Computational Physics – Lecture 19

(27 April 2020)

Recap

In the previous class, we learnt how to convolve and deconvolve two functions numerically.

Today's plan

In today's class, we want to learn something called spectral analysis of functions.

Introduction

An important type of mathematical analysis enabled by Fourier transforms is power spectral estimation.

Introduction

An important type of mathematical analysis enabled by Fourier transforms is power spectral estimation.

This has many experimental applications: optimal filtering, noise handling, signal processing, speech recognition, sonar/radar, etc.

Introduction

An important type of mathematical analysis enabled by Fourier transforms is power spectral estimation.

This has many experimental applications: optimal filtering, noise handling, signal processing, speech recognition, sonar/radar, etc.

Power spectral estimation also has theoretical applications in study of stochastic quantities, e.g., in statistical mechanics, astrophysics, and cosmology.

Introduction

Usually, by the word *spectrum*, we mean a relationship between the value of some parameter and frequency.

Introduction

Usually, by the word *spectrum*, we mean a relationship between the value of some parameter and frequency.

We see spectra of light intensity in optics.

Introduction

Usually, by the word *spectrum*, we mean a relationship between the value of some parameter and frequency.

We see spectra of light intensity in optics.

Every physical phenomenon – electromagnetic, thermal, mechanical, etc. – has an associated spectrum.

Introduction

Usually, by the word *spectrum*, we mean a relationship between the value of some parameter and frequency.

We see spectra of light intensity in optics.

Every physical phenomenon – electromagnetic, thermal, mechanical, etc. – has an associated spectrum.

In Fourier analysis, we split a function into frequency components. So Fourier transforms give a natural way to generalise the notion of a spectrum to arbitrary functions.

Parseval's theorem

Parseval's theorem says that

$$\int_{-\infty}^{\infty} dx f_1(x)[f_2(x)]^* = \int_{-\infty}^{\infty} dk \tilde{f}_1(k)[\tilde{f}_2(k)]^*. \quad (1)$$

This is because

$$\int_{-\infty}^{\infty} dx f_1(x)[f_2(x)]^* \quad (2)$$

$$= \int_{-\infty}^{\infty} dx \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk \cdot \tilde{f}(k) \cdot \exp(ikx) \right] [f_2(x)]^* \quad (3)$$

$$= \int_{-\infty}^{\infty} dx \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk \cdot \tilde{f}_1(k) \cdot \exp(ikx) \right] [f_2(x)]^* \quad (4)$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk \cdot \tilde{f}_1(k) \left[\int_{-\infty}^{\infty} dx \exp(ikx)[f_2(x)]^* \right] \quad (5)$$

$$= \int_{-\infty}^{\infty} dk \cdot \tilde{f}_1(k)[\tilde{f}_2(k)]^* \quad (6)$$

$$(7)$$

Power spectrum

So now if $f_1 = f_2$, we have

$$\int_{-\infty}^{\infty} dx |f_1(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_1(k)|^2. \quad (8)$$

Power spectrum

So now if $f_1 = f_2$, we have

$$\int_{-\infty}^{\infty} dx |f_1(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_1(k)|^2. \quad (8)$$

(We used this result in our previous discussion on optimal filters.)

Power spectrum

So now if $f_1 = f_2$, we have

$$\int_{-\infty}^{\infty} dx |f_1(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_1(k)|^2. \quad (8)$$

(We used this result in our previous discussion on optimal filters.)

The function $|\tilde{f}_1(k)|^2$, which is a function of k , is called the *power spectrum* or the *power spectral density* of $f(x)$.

Power spectrum

So now if $f_1 = f_2$, we have

$$\int_{-\infty}^{\infty} dx |f_1(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_1(k)|^2. \quad (8)$$

(We used this result in our previous discussion on optimal filters.)

The function $|\tilde{f}_1(k)|^2$, which is a function of k , is called the *power spectrum* or the *power spectral density* of $f(x)$.

The *average power* over a scale X is defined as

$$p_{\text{avg}} = \frac{1}{X} \int_0^X dx |f^2(x)|. \quad (9)$$

Random functions

The notion of power spectrum is particularly useful when $f(x)$ describes a stochastic process.

Random functions

The notion of power spectrum is particularly useful when $f(x)$ describes a stochastic process.

For example, $f(x)$ could describe experimental noise, with x being time.

Random functions

The notion of power spectrum is particularly useful when $f(x)$ describes a stochastic process.

For example, $f(x)$ could describe experimental noise, with x being time.

Or, $f(x)$ could describe a fundamentally stochastic quantity such as a quantum field or the cosmological matter distribution.

Random functions

The notion of power spectrum is particularly useful when $f(x)$ describes a stochastic process.

For example, $f(x)$ could describe experimental noise, with x being time.

Or, $f(x)$ could describe a fundamentally stochastic quantity such as a quantum field or the cosmological matter distribution.

In such cases $f(x)$ is a random variable with a certain probability distribution.

Random functions

But then the quantity

$$\tilde{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx \cdot f(x) \cdot \exp(-ikx)$$

is also a random variable.

Random functions

But then the quantity

$$\tilde{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx \cdot f(x) \cdot \exp(-ikx)$$

is also a random variable.

The problem is that for such stochastic processes the integral

$$\int_{-\infty}^{\infty} dx |f(x)|$$

does not converge.

Random functions

But then the quantity

$$\tilde{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx \cdot f(x) \cdot \exp(-ikx)$$

is also a random variable.

The problem is that for such stochastic processes the integral

$$\int_{-\infty}^{\infty} dx |f(x)|$$

does not converge.

So their Fourier transform does not exist. (Dirichlet condition.)

Random functions

But then the quantity

$$\tilde{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx \cdot f(x) \cdot \exp(-ikx)$$

is also a random variable.

The problem is that for such stochastic processes the integral

$$\int_{-\infty}^{\infty} dx |f(x)|$$

does not converge.

So their Fourier transform does not exist. (Dirichlet condition.)

“If the signal has non-zero power, it has infinite energy. If it has finite energy, it has zero average power.”

Random functions

So we have here a class of function for which the Fourier integral does not converge but they have finite average power.

Random functions

So we have here a class of function for which the Fourier integral does not converge but they have finite average power.

Unfortunately, such functions turn out to be important in physics (noise, fundamental stochastic fields).

Random functions

So we have here a class of function for which the Fourier integral does not converge but they have finite average power.

Unfortunately, such functions turn out to be important in physics (noise, fundamental stochastic fields).

In such cases, we study *statistics in Fourier space*.

Power spectrum of a random function

We can begin by defining a truncated function

$$f_X(x) = \begin{cases} f(x) & \text{if } |x| \leq X \\ 0, & \text{otherwise,} \end{cases} \quad (10)$$

so that

$$f(x) = \lim_{X \rightarrow \infty} f_X(x).$$

No $f_X(x)$ satisfies the Dirichlet condition and a Fourier transform exists.

Power spectrum of a random function

Now we know from Parseval's theorem that

$$\int_{-\infty}^{\infty} dx |f_X(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (11)$$

Power spectrum of a random function

Now we know from Parseval's theorem that

$$\int_{-\infty}^{\infty} dx |f_X(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (11)$$

If we divide both sides by $2X$ we get

$$\frac{1}{2X} \int_{-\infty}^{\infty} dx |f_X(x)|^2 = \frac{1}{2X} \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (12)$$

Power spectrum of a random function

Now we know from Parseval's theorem that

$$\int_{-\infty}^{\infty} dx |f_X(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (11)$$

If we divide both sides by $2X$ we get

$$\frac{1}{2X} \int_{-\infty}^{\infty} dx |f_X(x)|^2 = \frac{1}{2X} \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (12)$$

The LHS is now the average power in the original function over the scale $2X$.

Power spectrum of a random function

Now we know from Parseval's theorem that

$$\int_{-\infty}^{\infty} dx |f_X(x)|^2 = \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (11)$$

If we divide both sides by $2X$ we get

$$\frac{1}{2X} \int_{-\infty}^{\infty} dx |f_X(x)|^2 = \frac{1}{2X} \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2. \quad (12)$$

The LHS is now the average power in the original function over the scale $2X$.

We still cannot take the limit $X \rightarrow \infty$ because the Fourier transform does not exist in that limit.

Power spectrum of a random function

But we can take the expectation value.

$$E \left\{ \frac{1}{2X} \int_{-\infty}^{\infty} dx |f_X(x)|^2 \right\} = E \left\{ \frac{1}{2X} \int_{-\infty}^{\infty} dk |\tilde{f}_X(k)|^2 \right\}. \quad (13)$$

which gives us

$$\frac{1}{2X} \int_{-\infty}^{\infty} dx E \{ |f_X(x)|^2 \} = \frac{1}{2X} \int_{-\infty}^{\infty} dk E \left\{ |\tilde{f}_X(k)| \right\}^2. \quad (14)$$

or

$$\lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-\infty}^{\infty} dx \bar{f}^2(x) = \lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-\infty}^{\infty} dk E \left\{ |\tilde{f}_X(k)| \right\}^2. \quad (15)$$

which gives

$$\langle \bar{f}^2(x) \rangle = \int_{-\infty}^{\infty} dk \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}. \quad (16)$$

(Bar: ensemble average; Angle brackets: time average)

Power spectrum of a random function

Time average of the ensemble rms value is just the rms value so

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}. \quad (17)$$

Power spectrum of a random function

Time average of the ensemble rms value is just the rms value so

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}. \quad (17)$$

The quantity

$$P(k) = \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}$$

is the power spectral density for random function $f(X)$.

Power spectrum of a random function

Time average of the ensemble rms value is just the rms value so

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}. \quad (17)$$

The quantity

$$P(k) = \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}$$

is the power spectral density for random function $f(X)$.

Remember that

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk P(k)$$

so the integral of the power spectrum is the variance of the field!

Power spectrum of a random function

Time average of the ensemble rms value is just the rms value so

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}. \quad (17)$$

The quantity

$$P(k) = \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)| \right\}^2}{2X}$$

is the power spectral density for random function $f(X)$.

Remember that

$$\bar{f}^2(x) = \int_{-\infty}^{\infty} dk P(k)$$

so the integral of the power spectrum is the variance of the field!

We have discovered a simple connection between the power spectrum and statistics.

Wiener-Khinchin Theorem

There is another interesting property of the power spectrum of a random function.

Wiener-Khinchin Theorem

There is another interesting property of the power spectrum of a random function.

We have

$$P(k) = \lim_{X \rightarrow \infty} \frac{E \left\{ |\tilde{f}_X(k)|^2 \right\}}{2X} \quad (18)$$

$$= \lim_{X \rightarrow \infty} \frac{1}{2X} E \left\{ \left[\int_{-\infty}^{\infty} dx_1 f_X(x_1) \exp(-ikx_1) \right] \right. \quad (19)$$

$$\left. \left[\int_{-\infty}^{\infty} dx_2 f_X(x_2) \exp(ikx_2) \right] \right\} \quad (20)$$

where we have used the fact that

$$|\tilde{f}_X(k)|^2 = f_X(k) f_X(-k)$$

Wiener-Khinchin Theorem

The above will give us

$$P(k) = \lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dx_1 \{ E [f_X(x_1) f_X(x_2)] \\ \times \exp (ik(x_2 - x_1)) \} \quad (21)$$

Wiener-Khinchin Theorem

The above will give us

$$P(k) = \lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dx_1 \{ E [f_X(x_1) f_X(x_2)] \\ \times \exp (ik(x_2 - x_1)) \} \quad (21)$$

The quantity $E [f_X(x_1) f_X(x_2)]$ is called the *correlation function*.

$$E [f_X(x_1) f_X(x_2)] = \begin{cases} R(x_1, x_2), & \text{if } |x_1|, |x_2| < X \\ 0, & \text{otherwise,} \end{cases} \quad (22)$$

Wiener-Khinchin Theorem

The above will give us

$$P(k) = \lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dx_1 \{ E[f_X(x_1)f_X(x_2)] \\ \times \exp(ik(x_2 - x_1)) \} \quad (21)$$

The quantity $E[f_X(x_1)f_X(x_2)]$ is called the *correlation function*.

$$E[f_X(x_1)f_X(x_2)] = \begin{cases} R(x_1, x_2), & \text{if } |x_1|, |x_2| < X \\ 0, & \text{otherwise,} \end{cases} \quad (22)$$

Wiener-Khinchin Theorem

If we write $x_2 - x_1 = \xi$ and $dx_2 = d\xi$, we get

$$P(k) = \int_{-\infty}^{\infty} d\xi \left[\lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-X}^X dx_1 R(x_1, x_1 + \xi) \right. \\ \left. \times \exp(-ik\xi) \right]. \quad (23)$$

So the power spectrum is the Fourier transform of the time-averaged correlation function!

Wiener-Khinchin Theorem

If we write $x_2 - x_1 = \xi$ and $dx_2 = d\xi$, we get

$$P(k) = \int_{-\infty}^{\infty} d\xi \left[\lim_{X \rightarrow \infty} \frac{1}{2X} \int_{-X}^X dx_1 R(x_1, x_1 + \xi) \right. \\ \left. \times \exp(-ik\xi) \right]. \quad (23)$$

So the power spectrum is the Fourier transform of the time-averaged correlation function!

Wiener-Khinchin Theorem

For isotropic or stationary processes, the correlation function is independent of x so

$$\langle R(x_1, x_1 + \xi) \rangle = R(\xi), \quad (24)$$

and we get

$$\tilde{P}(k) = \int_{-\infty}^{\infty} dx \cdot R(x) \cdot \exp(-ikx) \quad (25)$$

and

$$R(x) = \int_{-\infty}^{\infty} dk \cdot \tilde{P}(k) \cdot \exp(ikx). \quad (26)$$

Wiener-Khinchin Theorem

For isotropic or stationary processes, the correlation function is independent of x so

$$\langle R(x_1, x_1 + \xi) \rangle = R(\xi), \quad (24)$$

and we get

$$\tilde{P}(k) = \int_{-\infty}^{\infty} dx \cdot R(x) \cdot \exp(-ikx) \quad (25)$$

and

$$R(x) = \int_{-\infty}^{\infty} dk \cdot \tilde{P}(k) \cdot \exp(ikx). \quad (26)$$

This is called the Wiener-Khinchin Theorem.

Computation

Our task is now to compute the power spectrum using DFT.

Computation

Our task is now to compute the power spectrum using DFT.

Because we know how to calculate the FT using the DFT, discretising the power spectrum of $f(x)$ is easy.

Computation

Our task is now to compute the power spectrum using DFT.

Because we know how to calculate the FT using the DFT, discretising the power spectrum of $f(x)$ is easy.

But if $f(x)$ is a random function, we now know that the power spectrum measures statistics of the distribution from which $f(x)$ is derived.

Computation

Our task is now to compute the power spectrum using DFT.

Because we know how to calculate the FT using the DFT, discretising the power spectrum of $f(x)$ is easy.

But if $f(x)$ is a random function, we now know that the power spectrum measures statistics of the distribution from which $f(x)$ is derived.

How well does the DFT-based computation approximate those statistics?

Estimators

Note that we are now trying to compute statistics of the population from that of a sample.

Estimators

Note that we are now trying to compute statistics of the population from that of a sample.

This is done by means of *estimators*.

Estimators

Note that we are now trying to compute statistics of the population from that of a sample.

This is done by means of *estimators*.

For example

$$\hat{m} = \frac{1}{n} \sum_0^{n-1} x_n \quad (27)$$

is an estimator for the mean.

Good and bad estimators

An estimator is good, or consistent, if

- ▶ its expectation value is equal to the true value of the statistic that it is measuring
- ▶ its variance is small

Good and bad estimators

An estimator is good, or consistent, if

- ▶ its expectation value is equal to the true value of the statistic that it is measuring
- ▶ its variance is small

For example, the above estimator for the mean is consistent.

But the estimator

$$S^2 = \frac{1}{n} \sum (x_i - \bar{x})^2 \quad (28)$$

is not a consistent estimator for variance. (It is biased.) The estimator

$$S^2 = \frac{1}{n-1} \sum (x_i - \bar{x})^2 \quad (29)$$

is consistent.

Periodogram

We can start by defining a power spectrum estimator as

$$P_n(k_q) = \frac{1}{n} |\tilde{f}(k_q)|^2 \quad (30)$$

Periodogram

We can start by defining a power spectrum estimator as

$$P_n(k_q) = \frac{1}{n} |\tilde{f}(k_q)|^2 \quad (30)$$

This most straightforward way of writing the discrete power spectrum is called the *periodogram*.

Periodogram

We can start by defining a power spectrum estimator as

$$P_n(k_q) = \frac{1}{n} |\tilde{f}(k_q)|^2 \quad (30)$$

This most straightforward way of writing the discrete power spectrum is called the *periodogram*.

Suppose we also discretise the correlation function as

$$R_n(p) = \frac{1}{n} \sum f_r \cdot f_{r+p} \quad (31)$$

Periodogram

Then we have

$$\begin{aligned}P_n(q) &= \sum R_n(p) \cdot \exp\left(\frac{-i2\pi qp}{n}\right) \\&= \sum \left[\frac{1}{n} \sum f_r \cdot f_{r+p} \right] \cdot \exp\left(\frac{-i2\pi qp}{n}\right) \\&= \frac{1}{n} \left[\sum f_n \exp\left(\frac{i2\pi qr}{n}\right) \right] \cdot \left[\sum f_{r+p} \exp\left(\frac{-i2\pi q(r+p)}{n}\right) \right] \\&= \frac{1}{n} |\tilde{f}(k_q)|^2\end{aligned}$$

Periodogram

Is our estimator unbiased?

Periodogram

Is our estimator unbiased?

The expectation value is

$$E [P_n(q)] = E \left[\sum R_n(p) \cdot \exp \frac{-i2\pi qp}{n} \right] \quad (32)$$

$$= \sum E [R_n(p)] \cdot \exp \frac{-i2\pi qp}{n} \quad (33)$$

Periodogram

Is our estimator unbiased?

The expectation value is

$$E [P_n(q)] = E \left[\sum R_n(p) \cdot \exp \frac{-i2\pi qp}{n} \right] \quad (32)$$

$$= \sum E [R_n(p)] \cdot \exp \frac{-i2\pi qp}{n} \quad (33)$$

What is the expectation value of $R_n(p)$?

Periodogram

Is our estimator unbiased?

The expectation value is

$$E [P_n(q)] = E \left[\sum R_n(p) \cdot \exp \frac{-i2\pi qp}{n} \right] \quad (32)$$

$$= \sum E [R_n(p)] \cdot \exp \frac{-i2\pi qp}{n} \quad (33)$$

What is the expectation value of $R_n(p)$?

It is given by

$$E [R_n(p)] = \left(1 - \frac{|p|}{n} \right) R(p), \quad (34)$$

where $R(p)$ is the true correlation function $E [f(x_r)f(x_r + p\Delta)]$

Periodogram

Let us denote

$$a(p) = \left(1 - \frac{|p|}{n}\right) \quad (35)$$

then we have

$$E [P_n(q)] = \sum R(p) a(p) \exp \frac{-i2\pi qp}{n} \quad (36)$$

Periodogram

Let us denote

$$a(p) = \left(1 - \frac{|p|}{n}\right) \quad (35)$$

then we have

$$E[P_n(q)] = \sum R(p)a(p) \exp \frac{-i2\pi qp}{n} \quad (36)$$

Using the discrete convolution theorem, this means that

$$E[P_n(q)] = \sum P(s)A(q-s) \quad (37)$$

where A is the Fourier transform of a and is of the form

$$A(k) \propto \left[\frac{\sin 2\pi kn/2}{\sin 2\pi k/2} \right]^2 \quad (38)$$

So the estimator is biased but becomes asymptotically unbiased for $n \rightarrow \infty$.

Periodogram

But the variance of $P_n(q)$ can be shown to give $P^2(q)$

$$\lim_{n \rightarrow \infty} \text{var} [P_n(q)] = P^2(q), \quad (39)$$

so our estimator, the periodogram, is not consistent.

Periodogram

But the variance of $P_n(q)$ can be shown to give $P^2(q)$

$$\lim_{n \rightarrow \infty} \text{var} [P_n(q)] = P^2(q), \quad (39)$$

so our estimator, the periodogram, is not consistent.

Many solutions to this problem have been proposed.

Periodogram

But the variance of $P_n(q)$ can be shown to give $P^2(q)$

$$\lim_{n \rightarrow \infty} \text{var} [P_n(q)] = P^2(q), \quad (39)$$

so our estimator, the periodogram, is not consistent.

Many solutions to this problem have been proposed.

One such solution is the Bartlett method.

Periodogram

But the variance of $P_n(q)$ can be shown to give $P^2(q)$

$$\lim_{n \rightarrow \infty} \text{var} [P_n(q)] = P^2(q), \quad (39)$$

so our estimator, the periodogram, is not consistent.

Many solutions to this problem have been proposed.

One such solution is the Bartlett method.

Note that x_1, x_2, \dots, x_l are uncorrelated random variables with mean $E(X)$ and variance σ^2 then the quantity

$$\frac{x_1 + x_2 + \dots + x_l}{l} \quad (40)$$

has mean $E(X)$ and variance σ^2/l .

Bartlett Method

We can now split our sample of n points into l segments of m points each and average the periodograms of each segment.

Bartlett Method

We can now split our sample of n points into l segments of m points each and average the periodograms of each segment.

For the averaged periodogram the bias will go to zero and the variance will go to zero as $l \rightarrow \infty$.

Bartlett Method

We can now split our sample of n points into l segments of m points each and average the periodograms of each segment.

For the averaged periodogram the bias will go to zero and the variance will go to zero as $l \rightarrow \infty$.

This is therefore a consistent method of deriving the power spectrum.

Higher-dimensional FT

We should mention for completeness that we have discussed on one-dimensional FTs so far.

Higher-dimensional FT

We should mention for completeness that we have discussed on one-dimensional FTs so far.

Higher-dimensional FTs are straitforward to define. For example

$$\tilde{f}(k) = \frac{1}{(\sqrt{2\pi})^2} \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdot f(x_1, x_2) \cdot \exp(-i(k_1 x_1 + k_2 x_2))$$

and its discrete version

$$\tilde{f}_{qr} = \frac{1}{\sqrt{mn}} \sum \sum f_{ps} \exp \left(-i2\pi \left[\frac{pq}{n} + \frac{rs}{m} \right] \right).$$

Higher-dimensional FT

We should mention for completeness that we have discussed on one-dimensional FTs so far.

Higher-dimensional FTs are straightforward to define. For example


$$\tilde{f}(k) = \frac{1}{(\sqrt{2\pi})^2} \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdot f(x_1, x_2) \cdot \exp(-i(k_1 x_1 + k_2 x_2))$$

and its discrete version

$$\tilde{f}_{qr} = \frac{1}{\sqrt{mn}} \sum \sum f_{ps} \exp \left(-i2\pi \left[\frac{pq}{n} + \frac{rs}{m} \right] \right).$$


Coding higher-dimensional DFT is hard! But libraries are available.

2d power spectrum




A poisson-distributed field.

2d power spectrum



2d power spectrum of the previous field.

3d power spectrum



3d power spectrum of the matter distribution in the universe.

What we have learnt so far

- ▶ Fourier transform, Fourier series, and Discrete Fourier transform
- ▶ Fast Fourier Transform algorithm
- ▶ DFT as a means to compute the FT
- ▶ Convolution and deconvolution of functions
- ▶ Optimal filtering
- ▶ Power spectra and statistics
- ▶ Higher-dimensional FTs