

Computational Physics – Lecture 12

(18 March 2020)

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

1. You are welcome to turn on your video. But keep it on if you have low bandwidth.

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

1. You are welcome to turn on your video. But keep it on if you have low bandwidth.
2. Mute Keep your microphone unless you are speaking to the class.

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

1. You are welcome to turn on your video. But keep it on if you have low bandwidth.
2. Mute Keep your microphone unless you are speaking to the class.
3. If you would like to say something, click on the “Raise Hand” button. (Find it now!)

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

1. You are welcome to turn on your video. But keep it on if you have low bandwidth.
2. Mute Keep your microphone unless you are speaking to the class.
3. If you would like to say something, click on the “Raise Hand” button. (Find it now!)
4. You can use the chat window also.

Housekeeping

We are having a class over Zoom for the first time, so some rules are necessary:

1. You are welcome to turn on your video. But keep it on if you have low bandwidth.
2. Mute Keep your microphone unless you are speaking to the class.
3. If you would like to say something, click on the “Raise Hand” button. (Find it now!)
4. You can use the chat window also.
5. I am recording this lecture, so you can catch up if you end up having to leave due to a technical problem.

Where are we?

We have been studying numerical techniques to solve ordinary differential equations.

Where are we?

We have been studying numerical techniques to solve ordinary differential equations.

Here is what we have learnt so far:

Where are we?

We have been studying numerical techniques to solve ordinary differential equations.

Here is what we have learnt so far:

1. Euler's Method
2. Error analysis and optimum step size for Euler's Method
3. Taylor and Runge-Kutta Methods
4. Multi-dimensional differential equations
5. Simultaneous differential equations
6. Higher-order differential equations
7. Adaptive step-size control

Today's plan

Today we want to learn about a case in which the Euler or Runge-Kutta fail catastrophically.

Today's plan

Today we want to learn about a case in which the Euler or Runge-Kutta fail catastrophically.

Worse, in such cases it may be impossible to use adaptive step-size control.

Today's plan

Today we want to learn about a case in which the Euler or Runge-Kutta fail catastrophically.

Worse, in such cases it may be impossible to use adaptive step-size control.

We want to see what these cases are, why they behave like this, and what can be done about them.

An example

Consider the initial value problem

$$y' = \lambda y, \tag{1}$$

where $\lambda < 0$ and $y(0) = \alpha$.

An example

Consider the initial value problem

$$y' = \lambda y, \tag{1}$$

where $\lambda < 0$ and $y(0) = \alpha$.

In this case, we happen to know the solution analytically.

An example

Consider the initial value problem

$$y' = \lambda y, \tag{1}$$

where $\lambda < 0$ and $y(0) = \alpha$.

In this case, we happen to know the solution analytically.

The solution is

$$y(t) = \alpha e^{\lambda t}, \tag{2}$$

which is a decreasing function of t because λ is negative.

An example

Consider the initial value problem

$$y' = \lambda y, \tag{1}$$

where $\lambda < 0$ and $y(0) = \alpha$.

In this case, we happen to know the solution analytically.

The solution is

$$y(t) = \alpha e^{\lambda t}, \tag{2}$$

which is a decreasing function of t because λ is negative.

But we are interested in getting this solution numerically.

An example

Consider the initial value problem

$$y' = \lambda y, \tag{1}$$

where $\lambda < 0$ and $y(0) = \alpha$.

In this case, we happen to know the solution analytically.

The solution is

$$y(t) = \alpha e^{\lambda t}, \tag{2}$$

which is a decreasing function of t because λ is negative.

But we are interested in getting this solution numerically.

(That is, we are interested in getting an *accurate numerical approximation* to this exact solution.)

Solve example problem using Euler's Method

Let us apply Euler's Method to our given initial value problem with mesh points

$$t_j = jh \quad j = 0, 1, 2, \dots, N, \quad (3)$$

where as usual

$$h = \frac{b - a}{N} \quad (4)$$

is the step size. (In our case a is 0 and b is some arbitrary positive real number.)

Solve example problem using Euler's Method

Let us apply Euler's Method to our given initial value problem with mesh points

$$t_j = jh \quad j = 0, 1, 2, \dots, N, \quad (3)$$

where as usual

$$h = \frac{b - a}{N} \quad (4)$$

is the step size. (In our case a is 0 and b is some arbitrary positive real number.)

What is the numerical approximation to the solution $y(t)$ according to this method?

Solve example problem using Euler's Method

Well, as we have studied before, the solution according to Euler's Method is

$$w_0 = \alpha \tag{5}$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \tag{6}$$

Solve example problem using Euler's Method

Well, as we have studied before, the solution according to Euler's Method is

$$w_0 = \alpha \quad (5)$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \quad (6)$$

which, for our example, becomes

$$w_0 = \alpha \quad (7)$$

$$w_{j+1} = w_j + h \cdot \lambda w_j, \quad (8)$$

Solve example problem using Euler's Method

Well, as we have studied before, the solution according to Euler's Method is

$$w_0 = \alpha \quad (5)$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \quad (6)$$

which, for our example, becomes

$$w_0 = \alpha \quad (7)$$

$$w_{j+1} = w_j + h \cdot \lambda w_j, \quad (8)$$

or, written differently, we have

$$w_0 = \alpha \quad (9)$$

$$w_{j+1} = (1 + h\lambda)w_j. \quad (10)$$

Solve example problem using Euler's Method

So, with Euler's Method and our example initial value problem, we have

$$w_{j+1} = (1 + h\lambda)^{j+1}w_0. \quad (11)$$

Solve example problem using Euler's Method

So, with Euler's Method and our example initial value problem, we have

$$w_{j+1} = (1 + h\lambda)^{j+1}w_0. \quad (11)$$

That is,

$$w_{j+1} = (1 + h\lambda)^{j+1}\alpha, \quad (12)$$

because $w_0 = \alpha$.

Solve example problem using Euler's Method

So, with Euler's Method and our example initial value problem, we have

$$w_{j+1} = (1 + h\lambda)^{j+1}w_0. \quad (11)$$

That is,

$$w_{j+1} = (1 + h\lambda)^{j+1}\alpha, \quad (12)$$

because $w_0 = \alpha$.

Is this an accurate approximation to the true solution?

Solve example problem using Euler's Method

So, with Euler's Method and our example initial value problem, we have

$$w_{j+1} = (1 + h\lambda)^{j+1}w_0. \quad (11)$$

That is,

$$w_{j+1} = (1 + h\lambda)^{j+1}\alpha, \quad (12)$$

because $w_0 = \alpha$.

Is this an accurate approximation to the true solution?

We can answer than question by calculating the error. (We can do this for this example problem because we know the true solution.)

Compute the error in our application of Euler's Method

Recall that we define the absolute error as

$$|y(t_j) - w_j|. \quad (13)$$

Compute the error in our application of Euler's Method

Recall that we define the absolute error as

$$|y(t_j) - w_j|. \quad (13)$$

In our case, this is

$$|e^{jh\lambda} - (1 + h\lambda)^j| |\alpha|, \quad (14)$$

or, in other words,

$$|(e^{h\lambda})^j - (1 + h\lambda)^j| |\alpha| \quad (15)$$

Compute the error in our application of Euler's Method

Recall that we define the absolute error as

$$|y(t_j) - w_j|. \quad (13)$$

In our case, this is

$$|e^{jh\lambda} - (1 + h\lambda)^j| |\alpha|, \quad (14)$$

or, in other words,

$$|(e^{h\lambda})^j - (1 + h\lambda)^j| |\alpha| \quad (15)$$

So our Euler solution is a good approximation if the value $(1 + h\lambda)^j$ is close to the value of $(e^{h\lambda})^j$.

Compute the error in our application of Euler's Method

Recall that we define the absolute error as

$$|y(t_j) - w_j|. \quad (13)$$

In our case, this is

$$|e^{jh\lambda} - (1 + h\lambda)^j| |\alpha|, \quad (14)$$

or, in other words,

$$|(e^{h\lambda})^j - (1 + h\lambda)^j| |\alpha| \quad (15)$$

So our Euler solution is a good approximation if the value $(1 + h\lambda)^j$ is close to the value of $(e^{h\lambda})^j$. Is this the case?

Since $\lambda < 0$, the quantity $(e^{h\lambda})^j$ goes to zero as j increases.

Since $\lambda < 0$, the quantity $(e^{h\lambda})^j$ goes to zero as j increases.

But the quantity $(1 + h\lambda)^j$ does not go to zero as j increases unless $|1 + h\lambda| < 1$.

Since $\lambda < 0$, the quantity $(e^{h\lambda})^j$ goes to zero as j increases.

But the quantity $(1 + h\lambda)^j$ does not go to zero as j increases unless $|1 + h\lambda| < 1$.

So we will get a reasonable solution only if $-2 < h\lambda < 0$ or in other words if

$$h < \frac{2}{|\lambda|}. \quad (16)$$

Since $\lambda < 0$, the quantity $(e^{h\lambda})^j$ goes to zero as j increases.


But the quantity $(1 + h\lambda)^j$ does not go to zero as j increases unless $|1 + h\lambda| < 1$.

So we will get a reasonable solution only if $-2 < h\lambda < 0$ or in other words if


$$h < \frac{2}{|\lambda|}. \quad (16)$$

For other values of the step size h , Euler's Method will fail for this problem.


Demo: $y' = -10y$ with $y(0) = 1$




Demo: $y' = -10y$ with $y(0) = 1$



Demo: $y' = -10y$ with $y(0) = 1$



Demo: $y' = -10y$ with $y(0) = 1$



Try other methods?

You might think that if Euler's Method has this problem then we should use some other method, such as our favourite, the fourth-order Runge-Kutta method.

Try other methods?

You might think that if Euler's Method has this problem then we should use some other method, such as our favourite, the fourth-order Runge-Kutta method.

But the problem occurs these methods too.

Try other methods?

You might think that if Euler's Method has this problem then we should use some other method, such as our favourite, the fourth-order Runge-Kutta method.

But the problem occurs these methods too.

In general, you get an expression of the form

$$w_{j+1} = Q(h\lambda) \cdot w_j, \quad (17)$$

where Q is some function, which took the form $1 + h\lambda$ in the special case of Euler's Method.

Try other methods?

You might think that if Euler's Method has this problem then we should use some other method, such as our favourite, the fourth-order Runge-Kutta method.

But the problem occurs these methods too.

In general, you get an expression of the form

$$w_{j+1} = Q(h\lambda) \cdot w_j, \quad (17)$$

where Q is some function, which took the form $1 + h\lambda$ in the special case of Euler's Method.

The method is then accurate only if the value of $(Q(h\lambda))^j$ is close to the value of $(e^{h\lambda})^j$.

Try other methods?

You might think that if Euler's Method has this problem then we should use some other method, such as our favourite, the fourth-order Runge-Kutta method.

But the problem occurs these methods too.

In general, you get an expression of the form

$$w_{j+1} = Q(h\lambda) \cdot w_j, \quad (17)$$

where Q is some function, which took the form $1 + h\lambda$ in the special case of Euler's Method.

The method is then accurate only if the value of $(Q(h\lambda))^j$ is close to the value of $(e^{h\lambda})^j$.

The error will grow without bounds if $|Q(h\lambda)| > 1$.

Try other methods?

For an n th-order Taylor Method, for example, you can see that our condition for accuracy becomes

$$\left| 1 + h\lambda + \frac{1}{2}h^2\lambda^2 + \cdots + \frac{1}{n!}h^n\lambda^n \right| < 1. \quad (18)$$

Try other methods?

For an n th-order Taylor Method, for example, you can see that our condition for accuracy becomes

$$\left| 1 + h\lambda + \frac{1}{2}h^2\lambda^2 + \cdots + \frac{1}{n!}h^n\lambda^n \right| < 1. \quad (18)$$

This immediately tells you the accuracy condition for the fourth-order Runge-Kutta Method if you remember that the fourth-order Runge-Kutta Method is essentially a fourth-order Taylor Method.

Try other methods?

For an n th-order Taylor Method, for example, you can see that our condition for accuracy becomes

$$\left| 1 + h\lambda + \frac{1}{2}h^2\lambda^2 + \cdots + \frac{1}{n!}h^n\lambda^n \right| < 1. \quad (18)$$

This immediately tells you the accuracy condition for the fourth-order Runge-Kutta Method if you remember that the fourth-order Runge-Kutta Method is essentially a fourth-order Taylor Method.

So how do we come up with a method that stays accurate at reasonable values of the step-size h ?

A trick: Implicit integration

We can refer to the set $h < 2/|\lambda|$ as the **stability region** of Euler's Method.

A trick: Implicit integration

We can refer to the set $h < 2/|\lambda|$ as the **stability region** of Euler's Method.

So our job now is to come up with methods that have a large stability region.

A trick: Implicit integration

We can refer to the set $h < 2/|\lambda|$ as the **stability region** of Euler's Method.

So our job now is to come up with methods that have a large stability region.

We do this by means of a trick called **implicit integration** or **backward integration**.

Backward (or Implicit) Euler Method

We remember that the Euler method is given by

$$w_0 = \alpha \tag{19}$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \tag{20}$$

Backward (or Implicit) Euler Method

We remember that the Euler method is given by

$$w_0 = \alpha \tag{19}$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \tag{20}$$

This is actually the **Forward**, or **Explicit**, Euler Method.

Backward (or Implicit) Euler Method

We remember that the Euler method is given by

$$w_0 = \alpha \quad (19)$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \quad (20)$$

This is actually the **Forward**, or **Explicit**, Euler Method.

The *Backward*, or *Implicit*, Euler Method is given by

$$w_0 = \alpha \quad (21)$$

$$w_{j+1} = w_j + h \cdot f(w_{j+1}, t_{j+1}), \quad (22)$$

Backward (or Implicit) Euler Method

We remember that the Euler method is given by

$$w_0 = \alpha \quad (19)$$

$$w_{j+1} = w_j + h \cdot f(w_j, t_j), \quad (20)$$

This is actually the **Forward**, or **Explicit**, Euler Method.

The *Backward*, or *Implicit*, Euler Method is given by

$$w_0 = \alpha \quad (21)$$

$$w_{j+1} = w_j + h \cdot f(w_{j+1}, t_{j+1}), \quad (22)$$

This small change makes a big difference. It transforms our Euler method from being catastrophically inaccurate to being stable for any value of step size h .

Backward (or Implicit) Euler Method

Why is this method called implicit?

Backward (or Implicit) Euler Method

Why is this method called implicit?

Because the right-hand side is now unknown:

$$w_0 = \alpha \tag{23}$$

$$w_{j+1} = w_j + h \cdot f(w_{j+1}, t_{j+1}), \tag{24}$$

so we will need to solve for w_{j+1} implicitly.

Backward (or Implicit) Euler Method

Why is this method called implicit?

Because the right-hand side is now unknown:

$$w_0 = \alpha \quad (23)$$

$$w_{j+1} = w_j + h \cdot f(w_{j+1}, t_{j+1}), \quad (24)$$

so we will need to solve for w_{j+1} implicitly.

For our example problem, $y' = \lambda y$ with $\lambda < 0$, the implicit Euler method will give

$$w_0 = \alpha \quad (25)$$

$$w_{j+1} = w_j + h \cdot \lambda w_{j+1}, \quad (26)$$

Backward (or Implicit) Euler Method

Why is this method called implicit?

Because the right-hand side is now unknown:

$$w_0 = \alpha \quad (23)$$

$$w_{j+1} = w_j + h \cdot f(w_{j+1}, t_{j+1}), \quad (24)$$

so we will need to solve for w_{j+1} implicitly.

For our example problem, $y' = \lambda y$ with $\lambda < 0$, the implicit Euler method will give

$$w_0 = \alpha \quad (25)$$

$$w_{j+1} = w_j + h \cdot \lambda w_{j+1}, \quad (26)$$

Luckily, solving for w_{j+1} is easy in this case.

Backward (or Implicit) Euler Method

The solution is

$$w_{j+1} = \frac{w_j}{1 - h\lambda} \quad (27)$$

or

$$w_{j+1} = \left(\frac{\alpha}{1 - h\lambda} \right)^{j+1}. \quad (28)$$

Backward (or Implicit) Euler Method

The solution is


$$w_{j+1} = \frac{w_j}{1 - h\lambda} \quad (27)$$

or


$$w_{j+1} = \left(\frac{\alpha}{1 - h\lambda} \right)^{j+1}. \quad (28)$$

Now we see that that for $\lambda < 0$, the term $(1 - h\lambda)^{j+1}$ will be < 1 as long as $|1 - h\lambda| > 1$ so the error will now not grow catastrophically. We have expanded our region of stability by a large margin.

Demo: $y' = -10y$ with $y(0) = 1$



Demo: $y' = -10y$ with $y(0) = 1$



Backward (or Implicit) Euler Method

We can similarly develop an implicit version of the fourth-order Runge-Kutta method.

Backward (or Implicit) Euler Method

We can similarly develop an implicit version of the fourth-order Runge-Kutta method.

If the differential equation is not linear, solving for w_{j+1} is hard. We will then have to use a root-finding method such as Newton-Raphson.

Why is all this important?

We have seen that implicit methods did better than explicit methods for our example initial value problem.

Why is all this important?

We have seen that implicit methods did better than explicit methods for our example initial value problem.

But so what? Is all this relevant beyond our little example?

Why is all this important?

We have seen that implicit methods did better than explicit methods for our example initial value problem.

But so what? Is all this relevant beyond our little example?

The answer is unfortunately yes. Implicit methods are useful in solving stiff differential equations, which unfortunately are quite common in physics.

What is a stiff differential equation?

Consider the system of equations

$$u_1' = 9u_1 + 24u_2 + 5 \cos t - 13 \sin t, \quad (29)$$

$$u_2' = -24u_1 - 51u_2 - 9 \cos t + \frac{1}{3} \sin t, \quad (30)$$

with $u_1(0) = 4/3$ and $u_2(0) = 2/3$.

What is a stiff differential equation?

Consider the system of equations

$$u_1' = 9u_1 + 24u_2 + 5 \cos t - 13 \sin t, \quad (29)$$

$$u_2' = -24u_1 - 51u_2 - 9 \cos t + \frac{1}{3} \sin t, \quad (30)$$

with $u_1(0) = 4/3$ and $u_2(0) = 2/3$.

The unique solution to this system of equations is

$$u_1(t) = 2e^{-3t} - e^{-39t} + \frac{1}{3} \cos t \quad (31)$$

$$u_2(t) = -e^{-3t} + 2e^{-39t} - \frac{1}{3} \cos t. \quad (32)$$

What is a stiff differential equation?

We see that the solution has a decaying part or a *transient*, namely e^{-39t} . The solution want to move closer to

$$u_1(t) = \frac{1}{3} \cos t \quad (33)$$

$$u_2(t) = -\frac{1}{3} \cos t, \quad (34)$$

as soon as possible. That is why it is **stiff**.

What is a stiff differential equation?

t	$u_1(t)$ (exact)	$u_1(t)$ (RK4 $h = 0.1$)
0.1	1.793061	-2.645169
0.2	1.423901	-18.45158
0.3	1.131575	-87.47221
0.4	0.9094086	-934.0722
0.5	0.7387877	-1760.016
...

What is a stiff differential equation?

In general, in a higher-order initial value problem, the unique solution can have such transient components.

What is a stiff differential equation?

In general, in a higher-order initial value problem, the unique solution can have such transient components.

This is the case, e.g., for equations describing an equilibrium between different processes (such as chemical reactions).

Exercise

Solve $y' = -30y$ with $0 \leq t \leq 1.5$ and $y(0) = \frac{1}{3}$

- ▶ Using Forward Euler with $h = 0.1$
- ▶ Using Backward Euler with $h = 0.1$
- ▶ Using Forward fourth-order Runge Kutta with $h = 0.1$

Make a plot showing these solutions and the exact solution.

Remember!

1. Get good sleep and nutrition
2. Keep yourself busy with work but avoid stress and anxiety
3. Maintain good hygiene