Divorce-induced health decline and the impact on mortality risk*

Kar Man Tan† Pavel Grigoriev‡

November 8, 2026

Abstract

Divorce is linked to elevated mortality. However, the causal mechanisms underlying this relationship are complex and remain insufficiently identified. Health is one such mechanism and is the focus of this paper. Using linked administrative data from the German Pension Insurance Fund (DRV), we examine whether divorce increases the risk of an initial musculoskeletal rehabilitation, and how much of the divorce-mortality gap runs through this channel. Rehabilitation is administered by the DRV and is granted only once a musculoskeletal condition has become chronic and threatens the ability to work. Our outcome measures the crossing of an institutional threshold rather than the onset of disease. We build annual risk sets of individuals experiencing a first (observable) divorce and individuals who have not yet divorced in that year and then use propensity score matching to generate comparable matches. Results: Divorce effect on cumulative incidence of musculoskeletal rehab, divorce effect on mortality, results from decomposition

Keywords: divorce, health, rehabilitation, mortality

JEL Classification: H75, I12, J12

*The authors are grateful to the DFG (German Research Foundation) for financial support (Project number: 544381616).

†Corresponding author: karman.tan@bib.bund.de. Federal Institute for Population Research (BiB), Friedrich-Ebert-Allee 4, 65185 Wiesbaden, Germany

‡Federal Institute for Population Research (BiB)

1 Introduction

The link between divorce and mortality has been widely discussed in social epidemiological studies. Register-based studies have used Finnish data (Metsä-Simola and Martikainen, 2013), Norwegian data (Berntsen and Kravdal, 2012) and English/Welsh data (Grundy and Tomassini, 2010). Until recently, Germany has been a difficult setting for this question. Brockmann and Klein (2004) is the only study to the best of our knowledge which explicitly analyzes the link between divorce and mortality although the dataset used in the study is the SOEP[1] which is not an administrative dataset. Since 2011, it is possible to link different administrative datasets from the German Pension Insurance Fund (DRV) using a unique identifier, allowing us to access full income biographies, divorce information[2], rehabilitation history and death. The availability of this dataset led to the conception of the project *Divorce and Diagnosis (DiDi): Register-based Analysis on the Role of Divorce in Health and Employment Dynamics*[3], which constitutes our study.

Our study focuses on the interaction between divorce, health and mortality. While there can be a direct link between divorce and mortality if divorce results in suicide (Kposowa, 2000; Stevenson and Wolfers, 2006) or homicide, causal evidence remains scarce when the link is indirect. Hence, we chose health as it is one possible indirect link tying together divorce and mortality. There can, however, still be threats to identification as health is not only a consequence but also a cause of divorce (Torvik et al., 2015). Furthermore, the individuals who choose to divorce may not necessarily be random. Many of the register-based studies make use of hazard models to estimate excess mortality risk after divorce and controls for sociodemographic and socioeconomic variables. For Germany, Brockmann and Klein (2004) account for health by using the self-reported health measure from the German SOEP and control for variables associated with marital status to find that divorced men have 51% higher mortality than married men while there is no significant effect for women. Leopold (2018) presents a very careful multi-outcome treatment of the gendered consequences of divorce over time, but it is not designed to control for the health that occurs before from the health consequences which follow the divorce. In our paper, we control for pre-existing health trends and other factors which occur before the divorce by using propensity score matching to answer the following questions

RQ1: Does divorce raise the risk of musculoskeletal rehabilitation[4] among divorced and non-divorced individuals?

RQ2: Is some of the excess mortality caused by divorce occurring because divorce caused additional musculoskeletal deterioration?

For context, the DRV only grants rehabilitation once the musculoskeletal condition is chronic to the point that it threatens working ability (Deutsche Rentenversicherung Bund, 2005). We do not argue that divorce causes musculoskeletal disease but rather whether divorce moves individuals who carry a latent vulnerability across the legal

[1] German Socioeconomic Panel

[2] This information is available due to the law of splitting pension points.

[3] This project is a collaboration between the Federal Institute for Population Research, Hertie School, Charité - Universitätsmedizin Berlin and the DRV.

[4] Musculoskeletal diseases are the most common cause of lost work time in Germany.

threshold where rehabilitation becomes necessary. Please see Section 2.3 for a more detailed discussion.

Results... maybe policy implications?

Section 2 provides a literature review, Section 3 introduces the data and methodology, Section 4 presents the results, Section 5 discusses the implications and Section 6 concludes.

2 Literature review

2.1 Link between divorce and mortality

The link between divorce and mortality risk is widely documented in sociology and epidemiology. In a meta-analysis, Shor et al. (2012) run meta-regressions on 625 mortality risk estimates over 104 studies and find that the mean hazard ratio was 1.30 for divorced individuals. The hazard ratio is slightly higher for men at 1.37 and lower for women at 1.22, although the difference between genders decrease with age. However, heterogeneity on the effect on death exists depending on the time since divorce. Metsä-Simola and Martikainen (2013) find that excess mortality among Finnish men, tied to accidental, violent or alcohol-related deaths, is highest immediately after divorce and declines over time. On the other hand, Berntsen and Kravdal (2012) finds that excess mortality among Norwegian men increases with time since divorce and remarriage can change the magnitude of the effect. In Germany, the evidence is limited. Brockmann and Klein (2004) use event-history models which account for marital transitions and duration in each marital status. They find evidence of selection into marriage and that mortality risk is lower when the marriage duration is longer.

Most of the previous studies establish a robust association, however, causal evidence remains scarce as divorce is not random. Sbarra and Nietert (2009) follow 1376 individuals in the Charleston Heart Study for forty years and show that the definition of the exposure determines the size of the effect. Individuals who are continuously separated or divorced have a higher hazard ratio (1.66) compared to individuals who have ever been separated or divorced (0.98). Wiborg and Gugushvili (2025) initially find raised hazard ratios for divorced men (1.59) and women (1.47) using Norwegian data. After using twin-pairs to account for genetic and early developmental and environmental factors, the relationship between divorce and mortality disappears.

2.2 Divorce and health

The effect of divorce on health is difficult to isolate due to the possibility of reverse causality. Joung et al. (1998), using Dutch data, find that married individuals who had more subjective health complaints were 1.5 times more likely to divorce, while individuals who reported multiple chronic conditions were twice as likely to divorce. Torvik et al. (2015) corroborates Joung et al. (1998)'s findings with Norwegian data. Poor health behaviors including smoking, drinking, and lack of exercise in addition to mental distress predict subsequent divorce. Metsä-Simola and Martikainen (2013), with Finnish data, find that psychotropic medication use among individuals who divorce is already elevated five years before hand and peaks at around six to nine months just

before divorce. Wade and Pevalin (2004) find worse mental health among those divorcing or divorced than among the married using British data. Generally, poor health tends to precede divorce.

To deal with health selection, Kohn and Averett (2014) estimates a dynamic model that accounts for the persistence of health and how it interacts with relationship status using 18 waves of the British Household Panel Survey. The result is that divorce has a marginally significant effect on the health of men younger than 45. Sbarra et al. (2014) also finds only a marginal effect of divorce on health. After matching individuals on divorce probability, Sbarra et al. (2014) finds that only individuals who were already depressed before divorce experienced major depressive episodes following divorce. With respect to mental health, Björkenstam et al. (2013) find that divorced individuals have thrice the risk of psychiatric inpatient care compared with married individuals in Sweden. Meulman et al. (2023) find similar results in the Netherlands where divorced individuals had higher health expenditures dedicated to mental health care. In terms of physical health, the evidence is heterogeneous. Zulkarnain and Korenman (2019) find that women who divorce at older ages tend to have more physical changes than men. Leopold (2018), on the other hand, find that men are more vulnerable in the short run in terms of well-being although the gender differences narrow in the medium run. Pellón-Elexpuru et al. (2024), in a meta-analysis across 94 studies find that divorced individuals report worse self-rated physical conditions with higher complaints of musculoskeletal, cardiovascular and cerebrovascular conditions although the effect is not significant.

2.3 Divorce and musculoskeletal disease

Evidence on the pathway from divorce to musculoskeletal impairment is rather limited and ambiguous. However, we chose to focus on musculoskeletal diseases because they are a major source of disability and work interruption in Germany[5]. The DRV Reha-Atlas (Deutsche Rentenversicherung Bund, 2024) reports that, of all completed medical rehabilitations, orthopedic conditions are the most common, accounting for 35.6% of rehabilitations among women and 35.5% among men. Brenscheidt et al. (2026) reports that musculoskeletal diseases account for 21% of certified incapacity days in 2024, which is the largest diagnostic group.

Sjøgren et al. (2009), using Danish data, find that chronic pain is more common among individuals who are divorced, separated or widowed, where musculoskeletal diseases accounted for 66.8% of the cause of chronic pain. However, Choi and Wagner (2026), using the Health and Retirement Survey find that individuals who transition from marriage to divorce tend to have a lower probability of high-impact pain. On the other hand, Cimas et al. (2026) using SHARE[6] data, find that there are gender differences. While divorced women have a higher incidence of chronic musculoskeletal pain, men have a lower prevalence. Meanwhile, Andorsen et al. (2017) finds that there is no significant effect between divorce and musculoskeletal complaints using Norwegian data.

While the impact of divorce on musculoskeletal diseases is ambiguous, the evidence on

[5]Please see the section on Rehabilitation system in Germany in the Appendix.

[6]Survey of Health, Aging and Retirement in Europe

the relationship between social support and the consequences of musculoskeletal diseases is more homogeneous. Prang et al. (2015) finds that individuals who have support from family friends and neighbors tend to have less persistent pain associated with musculoskeletal diseases, while widowed, separated or divorced individuals, in the absence of a network, tend to have a higher probability of persistent pain. Woods (2005) corroborates this and finds that a lack of support is associated with higher musculoskeletal morbidity, absence from work and subsequently, failure to return to work. Reese et al. (2010) finds that the quality of marriages matter, i.e. individuals with arthritis in problematic marriages tend to have more pain complaints and disability compared to individuals in well-adjusted marriages. This could signify that a decline in the relationship quality prior to divorce leads to poorer support, hence more complaints of pain. Couch et al. (2015) and Tamborini et al. (2016) find that divorce is associated with subsequent health threatening the ability to work, especially among individuals who do not remarry, although they do not differentiate by types of diseases.

2.4 Musculoskeletal disease and death

Musculoskeletal conditions are rarely direct causes of death in the way that cardiovascular diseases are. They often operate indirectly, for example through inflammation affecting the cardiovascular disease risk. Rheumatoid arthritis and chronic inflammatory activity can contribute to oxidative stress, dysfunction of blood vessels, arterial build-up and other cardiovascular risks (England et al., 2018). Cardiovascular mortality among people with rheumatoid arthritis is about 60% higher than the general population (Meulman et al., 2009). For non-inflammatory conditions, musculoskeletal conditions causes pain and restricted movement, which then makes people avoid physical activity. This reduces cardiovascular health and fitness, which then potentially increases the risk of comorbidities. Nüesch et al. (2011) finds that English patients with knee or hip osteoarthritis tend to have higher all-cause and cardiovascular mortality. Hawker et al. (2014) finds a similar result with Canadian patients.

Jansson, Mittendorfer-Rutz, and Alexanderson (2012) is more related to our study. Using Swedish register data, the authors compared individuals who were absent from work due to musculoskeletal diseases with people who were absent from work due to other diseases and people who were not absent from work. They find that among three categories of musculoskeletal diseases, the all-cause morbidity was 1.3 to 1.5 times higher. Given that musculoskeletal diseases account for the highest share of rehabilitations facilitated by the DRV, it is important that rehabilitation can encourage return to work and by extension, reduce the all-cause mortality attributed to musculoskeletal diseases. Bethge et al. (2019) finds that work-related medical rehabilitation tends to increase the return to work rate by 5.8 percentage points and shortens the median time to return by 9.46 days compared to conventional medical rehabilitation. In our study, the first completed DRV-facilitated musculoskeletal rehabilitation should be interpreted as an indicator of serious, work-relevant impairment and also receipt of potentially beneficial treatment.

3 Data and methodology

3.1 Data

The Research Data Center of the DRV allows linkages of different datasets starting from the reporting year 2011 through a unique identifier (Forschungsdatenzentrum Rentenversicherung, 2025). The datasets we use are AKVS[7], LTVS[8], RTBN[9], RTWF[10], RSD[11] and VA[12]. The time period covered by the data is 2011 to 2023. All estimates use the 20% sample extract, which contains 17,559,407 individuals and 185,518,738 person-years.

The AKVS, LTVS and RTBN are merged by the DRV to form the BASE dataset. The AKVS dataset contains individuals who are still actively contributing to the pension fund hence it generally represents individuals who are employed. The LTVS dataset contains individuals who have paused pension fund contributions but have not yet retired (nor drawn from their pension fund). It may include individuals who are unemployed, undergoing further education, on parental leave, etc. The RTBN dataset contains individuals who are drawing from their pension fund, i.e. retired individuals. The merged dataset, BASE, allows us to map out the full life-course coverage of individuals. If we only considered only RTBN, the sample is biased towards individuals who have contributed and lived long enough to draw a pension, ignoring those individuals who have undergone rehabilitation or died before drawing a pension. Considering only AKVS ignores rehabilitation efforts and gray divorces. To round up the dataset, it is also important to consider LTVS which details what happens between employment spells and retirement.

We then supplement the merged dataset with the RTWF, RSD and the VA. The RTWF details pension terminations due to deaths. If the death information from the merged dataset is inconsistent with the information in the RTWF, the individuals (cases) are dropped. This affects 102 individuals. Addtionally, individuals can have duplicate rows arising from BASE or RTWF. This affects 10,345,764 person-years with 11,297,317 excess rows. We drop the excess rows with more missing values. The RSD dataset details the completed rehabilitations and include information such as the broad diagnosis group, the type of rehabilitation benefits applied for, start and end date of rehabilitation. The RSD dataset also includes individuals with withdrawn, transferred or rejected applications. We only include individuals whose applications were successful and who completed the rehabilitation spell. Furthermore, about 5% of individuals undergo three or more rehabilitation spells per year. We also drop these individuals. We then supplement the merged dataset by including the additional rehabilitation information for the remaining individuals in the RSD dataset. The VA dataset includes information related to divorce and only contains divorced individuals. From the VA, we can obtain the date of divorce, pension supplements and deductions which result from divorce. Note that the VA only records divorces which have become final and where pension rights were equalized (See appendix on Updating the evolution of $\mathbf{L}_{i,t}$).

[7] Active Insured Persons
[8] Latent Insured Persons
[9] Insured Persons' Pension Portfolio
[10] Termination of Pensions due to Death
[11] Completed Rehabilitation in the Insurance History
[12] Pension Equalization Statistics

3.2 Matching

Since divorce is not randomly assigned, divorced and non-divorced individuals may differ systematically before divorce occurs. We therefore use 1:1 matching without replacement to construct a comparison group of individuals who are similar with respect to observed pre-treatment characteristics. The causal interpretation relies on conditional exchangeability (Rosenbaum and Rubin, 1983). This means that, conditional on the observed characteristics used in matching, individuals who divorce and comparable individuals who do not divorce are assumed to have had similar potential health trajectories in the absence of divorce. In other words, it is assumed that there are no remaining unobserved factors that jointly affect the probability of divorce and the subsequent health outcomes. However, the administrative data may not fully capture factors such as relationship conflict or social support. Some of these factors may already show up in the data before divorce, for instance in earnings, employment or rehabilitation use, which we use below to examine whether divorced individuals and their controls followed different paths before divorce.

We define treatment as the first observed divorce from the earliest observed marriage spell. The year of divorce is fixed as t_0. We include divorces observed between 2012 and 2018, so that each cohort can be observed for at least five years after divorce. For each treatment year, we form a separate within-year risk set. For treated individuals whose first observed divorce occurs in t_0, eligible controls are individuals who have not yet experienced a divorce by that year. Not yet divorced means that no divorce from the VA has been recorded in the panel by t_0. Controls may divorce later. We do not restrict the control group to individuals who never divorce because this would define control eligibility using information that occurs after t_0. Since the panel starts in 2011, to avoid individual whose divorces occured before 2011 entering the control pool, we exclude controls with a recorded end of a first marriage in or before t_0. Individuals must be alive at t_0 and must have an observed record in $t_0 - 1$. We exclude individuals with a musculoskeletal rehabilitation up to and including t_0. They may previously have received rehabilitation for other conditions. The resulting sample therefore consists of individuals who are at risk of their first observed musculoskeletal rehabilitation after t_0.

We require treated and control individuals to be matched exactly on `gender`, 5-year `age_band` and whether any rehabilitation has occurred by $t_0 - 1$, i.e. `ever_had_any_rehab_to_date`. Then, we estimate a logistic propensity score model separately for each treatment year. The propensity score represents the estimated probability of divorce in t_0, conditional on observed characteristics measured before divorce at $t_0 - 1$. The numeric characteristics we include are `age`, `income`, `pension_income`, `cumulative_rehabs_to_date`, `cumulative_mental_rehabs_to_date`, `years_since_last_rehab`, `rehab_starts_in_year`, `full_time_contribution_periods`[13], `part_time_contribution_periods`[14], `regular_employment`[15], `unemployment`[16], `marginal_employment`[17], `care_leave`[18] and `number_of_children`. The binary characteristics we include are

[13]Periods during which pension contributions were recorded for full-time work.
[14]Periods during which pension contributions were recorded for part-time work
[15]Days in reporting year spent in regular employment
[16]Days in reporting year spent unemployed
[17]Days in reporting year for mini- or midi-jobs
[18]Days in reporting year spent in caring capacity, *Pflegezeit*

ever_had_mental_rehab_to_date, unsuccessful_rehab_application_this_year, missing_full_time_contribution, missing_part_time_contribution and child_born_in_year. The categorical variables included are federal_state, broad_occupational_category[19], income_source[20], education[21] and youngest_child_age_group[22].

The numeric variables are standardized. Missing numeric and binary values are filled with the median and most common value respectively, along with an indicator for missingness. Categories with fewer than ten observations in a treatment year are pooled together in a separate category for 'rare' categories.

We also deliberately leave marital status out of the matching. Using marital status would select many future divorcees as controls, since the marital status is partially constructed using the VA which only records divorces: individuals coded as married are disproportionately those whose marriage later ends in divorce. In an earlier run with a 4% sample, 32% of individuals who were selected as controls were eventually divorced by 2018. Among the matched controls, 8.4% (10,717 out of 127,395) have marital status coded as divorced in $t_0 - 1$, 6.4% are coded as married, and 83.3% have no information on marital status. The comparison is therefore between individuals who divorce in t_0 and individuals who do not divorce in t_0, most of whose marital status is unknown, rather than individuals known to remain married.

We treat missingness in earnings, pension income and contribution periods as potentially informative. In particular, observed earnings with missing pension income generally indicate active employment, whereas observed pension income with missing earnings generally indicates pension receipt. We retain the observed earnings and pension values and include a categorical measure describing the joint observability of the two income sources. We also include separate indicators for missing contribution-period information. This allows the matching model to use both the recorded values and the information contained in their missingness patterns.

Within each treatment year and exact-match group, we use optimal matching (Rosenbaum, 1989). The matching procedure first maximizes the number of treated individuals who can be matched within the caliper (maximum absolute propensity score difference of 0.01) and then chooses the combination of pairs that minimizes the total propensity score distance. Since treatment years are matched in chronological order, an individual who was used as a control in an earlier year is removed from the pool of controls of later years. This individual, however, can still enter a later risk set as a treated individual if they experienced divorce. To accommodate the large annual control pools, the propensity score model is fitted using all treated individuals and, where necessary, a reproducibly selected sample of controls, with a nominal estimation sample

[19]Classification of Occupations 1988 created by the German Federal Employment Agency (Bundesinstitut für Berufsbildung (BIBB), n.d.). We used the first digit which signifies occupational area. For values with only two digits, we code the values as zero. What about 555 (disabled) and 666 (rehabilitants)? Why are these classified as occupations?

[20]This variable has the following categories regarding the source of income: income only from working, income only from pensions, income from earnings and pensions, no income from earnings or pensions. The latter two categories are uncommon.

[21]Defined as low, medium and high where medium indicates that the highest education level is a Bachelor's degree, and high a Master's degree

[22]The age groups are 0-2, 3-5, 6-9, 10-17, 18+.

target of 500,000 observations[23]. The resulting propensity scores are corrected for the control-sampling fraction by adding the logarithm of the fraction to each individual's predicted log-odds (King and Zeng, 2001), and the fitted model is then used to calculate propensity scores for every eligible treated individual and control. The computational limit therefore affects how the propensity model is estimated but does not remove controls from the matching pool. Similarly, the optimal assignment in very large exact-match groups is solved on a reduced set of candidate pairs.

We assess covariate balance before and after matching using standardized mean differences. Balance is assessed separately for continuous variables, binary variables, each level of categorical variables and indicators of missingness. An absolute standardized mean difference below 0.10 is treated as evidence of adequate balance on the corresponding observed characteristic (Austin, 2009). If residual imbalance above this threshold remains after matching, the affected variables are additionally included in another specification of the outcome model (see Identification section). This adjustment is intended to address limited remaining imbalance and is not used as a substitute for unsuccessful matching.

Matching on $t_0 - 1$ alone makes divorced individuals and their controls alike in that year only, which may be a potential threat to the conditional exchangeability assumption. Individuals who look alike at $t_0 - 1$ may nevertheless have reached that point along different paths (i.e. different pre-divorce trends), and if those paths continue after t_0, part of the post-divorce contrast in rehabilitation would reflect a pre-existing trend rather than divorce. We first conducted an event study from $t_0 - 3$ to $t_0 + 5$ by taking the difference in values of unsuccessful rehabilitation applications, earnings, pension income, missingness of pension income, days in the year spent employed, unemployed, marginally employed, caretaking and prior rehabilitation activity between the treated and control in each matched pair and conducting a Wald test (Wald, 1943) to determine whether the differences across time are significant for each variable. We regress each within-pair difference on dummies for each year relative to t_0, with $t_0 - 1$ as the reference year and standard errors clustered by pair, and test jointly whether the coefficients for $t_0 - 3$ and $t_0 - 2$ are zero. For earnings, rehabilitation starts and unsuccessful applications, the coefficients are significantly different from zero, showing that divorced individuals and their controls were not on the same path before $t_0 - 1$.

We therefore repeat the matching exercise with three lags, i.e. $t_0 - 3$, $t_0 - 2$ and $t_0 - 1$ instead of only $t_0 - 1$, adding earnings, pension income, income source, unsuccessful applications, rehabilitation starts and days spent in regular employment, unemployment, marginal employment and care at $t_0 - 2$ and $t_0 - 3$ to the lag-1 variables. Because the panel starts in 2011, requiring three observed pre-divorce years restricts this specification to divorces in 2014–2018. Matching on earlier years leads to differences in pension, income missingness of pension income and unsuccessful applications to not be jointly significant (Table 9). The Wald test, however, rejects for earnings, rehabilitation starts and marginal employment days. For earnings, the remaining diffrencs are below 1% of mean earnings. Because $t_0 - 2$ and $t_0 - 3$ now enter the matching, the remaining coefficients show how closely the matched histories agree rather than providing an independent test. We use the lag-3 specification as our preferred specification

[23] Please see the section on Dealing with a large sample of controls in the Appendix

Matching on the earlier years roughly halves the pre-divorce differences in earnings, and the differences in pension income, missingness of pension income and unsuccessful applications are no longer significant (Table 9). The Wald test still rejects for earnings and rehabilitation starts. For earnings, the remaining differences are below 1% of mean earnings. For rehabilitation starts, most of the remaining difference reflects $t_0 - 1$, the one pre-divorce year in which rehabilitation starts are not a matching variable: the gap is -0.07 and 0.16 per 100 people in $t_0 - 3$ and $t_0 - 2$ but 0.36 in $t_0 - 1$. Because these years now enter the matching, the remaining coefficients show how closely the matched histories agree rather than providing an independent test. We use the lag-3 specification as our preferred specification.

One caveat with matching on unemployment days is that similar mean day counts can coexist with different shares of individuals experiencing unemployment. In the lag-3 specification, mean number of days spent unemployed in the pooled matched sample was almost identical $t_0 - 1$ among divorced and controls (47.2 vs 48.0 days), but 21.8% of women who were about to divorce spent some days in unemployment in $t_0 - 1$ compared to only 16.7% of controls. In $t_0 - 3$ and $t_0 - 2$, the shares were similar though (14.2% vs 13.8% and 15.4% vs 14.1% respectively), suggesting that there is an early consequence of separation for women. This pattern is consistent with labor market changes preceding divorce, but does not establish that divorce caused them.

Using the matched sample, we estimate the effect of divorce on first entry into musculoskeletal rehabilitation and all-cause mortality. Musculoskeletal rehabilitation is examined over a 5-year follow-up period, with death treated as a competing event. Mortality is then followed for *up to* ten years where permitted by the end of the observation period in 2023.

3.3 Identification

For identification, we focus on the average treatment effect among the matched treated individuals.

3.3.1 RQ1: Musculoskeletal rehabilitation

We follow each matched individual from t_0+1 to t_0+5 and estimate a discrete-time hazard model:

$$\begin{aligned} \lambda_{i,\tau}^{\text{rehab}} &= \Pr(R_{i\tau} = 1 | \text{no musculoskeletal rehab before } \tau, \text{ alive at } \tau) \\ &= \Lambda_{\text{rehab}}\left(\alpha_\tau + \beta D_i + \mathbf{x}_i'\gamma\right) \end{aligned} \tag{1}$$

where $R_{i,\tau}$ is a dummy indicating individual i's first entry into musculoskeletal rehabilitation in time in follow-up year $t_0 + \tau$ where $\tau \in \{1, \ldots, 5\}$, α_τ are fixed effects, D_i is a dummy indicating divorce at t_0 and $\mathbf{x}_i$ are other variables measured at $t_0 - 1$ that we add depending on the specification. We estimate three specifications for Equation 1. In the first specification, we omit $\mathbf{x}_i$ and rely only on matching to deal with selection bias. In the second specification, we include variables whose absolute standardized mean difference after matching exceed 0.10 into $\mathbf{x}_i$. Standard errors are clustered at the level of the matched pair. In the third specification, $\mathbf{x}_i$ is instead replaced by $\mathbf{x}_{i,\tau-1}$, which is

the value of each variable in the year before each follow-up year. By conditioning on variables in $t_0 + \tau - 1$, we remove some of the variation attributed to divorce. However, we include the specification for completeness and to identify whether other post-divorce factors can affect musculoskeletal rehabilitation. Furthermore, since matching is exact on gender, we estimate all three specifications separately for men and women.

The hazard odds ratio from Equation 1 is difficult to interpret on its own as it does not show how often rehabilitation occurs, hence we also estimate the cumulative incidence of a first musculoskeletal rehabilitation over the 5-year follow up period using a discrete Aalen-Johansen estimator[24] (Aalen and Johansen, 1978). We report four values at the the 5-year horizon, 1) cumulative rehabilitation risk among treated individuals, 2) cumulative rehabilitation risk among controls, 3) risk difference per 1000 individuals and 4) the corresponding risk ratio. Confidence intervals are obtained by resampling whole matched pairs 200 times.

3.3.2 RQ2: Mortality

We follow each individual from $t_0 + 1$ until $t_0 + 10$ or 2023, the last year of the sample, whichever occurs first and estimate a discrete-time hazard model similar to Equation 1:

$$\begin{aligned} \lambda_{i,\tau}^{\text{death}} &= \Pr(Y_{i,\tau} = 1 | \text{alive at beginning of follow-up year } \tau) \\ &= \Lambda_{\text{death}}\left(\alpha_\tau + \beta D_i + \mathbf{x}_i'\gamma\right) \end{aligned} \tag{2}$$

where $Y_{i,\tau}$ is a dummy indicating that individual i dies in follow-up year $t_0 + \tau$, α_τ are time fixed effects, D_i is a dummy indicating divorce at t_0 and $\mathbf{x}_i$ are other variables measured prior to t_0. Just as in Equation 1, we estimate two specifications for Equation 2 by varying the inclusion of variables in $\mathbf{x}_i$ based on absolute standardized mean differences after matching. We complement the pooled hazard odds ratio with cumulative mortality risks using the Kaplan-Meier estimator (Kaplan and Meier, 1958). We again report cumulative mortality risk among treated and controls, the risk difference per 1000 individuals and the corresponding risk ratio at the 5-year horizon[25]. Just like for RQ1, the lag-3 specification is our preferred specification.

In addition to the straightforward analysis of divorce on mortality, we run a mediation analysis to determine how much of the mortality difference can be attributed to 'early'

[24]This estimator recognizes that individuals who die can no longer enter rehabilitation. For example, suppose there are 100 people where 10 enter rehabilitation in $t_0 + 1$, 20 die before rehabilitation and 70 are still alive and have not entered rehabilitation. Suppose in $t_0 + 2$, 7 people out of the remaining 70 enter rehabilitation. If we treat death as censoring rather than as a competing event, the proportion surviving up to $t_0 + 1$ is $1 - \frac{10}{100} = 0.9$, and up to $t_0 + 2$ is $1 - \frac{7}{70} = 0.9$, $\therefore$ the cumulative rehabilitation probability is $1 - (0.9 \times 0.9) = 0.19$, which is an overestimation. With the Aalen-Johansen estimator, we consider the proportion which actually entered rehabilitation by each follow-up year before dying. In $t_0 + 1$, the proportion entering rehabilitation is 0.1, however, after $t_0 + 1$, the proportion remaining alive without having entered rehabilitation is $1 - \frac{10+20}{100} = 0.7$. In $t_0 + 2$, $\frac{7}{70} = 0.1$ is the share of people who enter rehabilitation, but since only 70% of the original population was alive without having entered rehabilitation at the beginning of $t_0 + 2$, the contribution to the incidence is $0.7 \times 0.1 = 0.07$, hence the cumulative rehabilitation probability is 0.17

[25]The hazard model uses up to ten years of follow-up although later follow-up years are available only for earlier divorce cohorts.

musculoskeletal rehabilitation as result of divorce. A single binary mediator measured one or two years after divorce is insufficient for this question since it can ignore deterioration that occurs later in the follow-up window, such as economic and rehabilitation histories that change after divorce. For example, divorce can potentially change earnings which can subsequently have the ability to affect the probability of entering musculoskeletal rehabilitation and dying.

Let $D_i \in \{0,1\}$ be divorce status fixed at t_0, $\mathbf{z}_i$ the available pre-treatment matching variables for the lag-3 specification, $H = 5$ the fixed mortality and mediation horizon, $\mathbf{L}_{i,t}$ the post-divorce history (`income`, `pension_income`, `unsuccessful_rehab_application_this_year`, `non_ms_rehab`[26], `income_source`, and the days in a year spent in `regular_employment`, `unemployment`, `marginal_employment` and `care_leave`) in time t and $M_{i,t-1} \in 0,1$ is a dummy indicating whether individual i entered musculoskeletal rehabilitation by the end of time $t-1$. We use information in time $t-1$ to predict information in time t. We cannot use information from time t since we are unsure whether the dependent or independent variables are recorded first. We use weighted maximum likelihood to account for individuals actually observed in the data in a given year.

We estimate the probability of an individual i having a record for the year $t_o + t$ given that they are alive:

$$\Pr(R_{i,t} = 1 | D_i, \mathbf{z}_i, \mathbf{L}_{i,t-1}) = \Lambda_{\text{obs}}(\alpha_t + \beta D_i + \mathbf{z}_i'\psi + \mathbf{L}_{i,t-1}'\psi_L + \mu M_{i,t-1}) \tag{3}$$

We then predict $\hat{p}_{\text{obs},i}(u)$, i.e. the probability that individual i has a record in time u given their information up until $u-1$ and inverse this to compute weight:

$$\hat{\omega}_{i,t} = \min\left(\prod_{u=1}^{t} \frac{1}{\max\left(\hat{p}_{\text{obs},i}(u), \underline{p}\right)}, \bar{\omega} \right) \tag{4}$$

We set the maximum weight, $\bar{\omega} = 20$, to prevent one trajectory from having too much influence, and the minimum observation probability to $\underline{p} = 0.02$. We then estimate the parameters of a model on the probability of first musculoskeletal rehabilitation in year t given that the individual did not experience musculoskeletal rehab in $t-1$ and other information in $t-1$[27]:

$$\begin{aligned} \lambda_{i,t}^{\text{rehab}} &= \Pr(M_{i,t} = 1 | M_{i,t-1} = 0, D_i, \mathbf{z}_i, \mathbf{L}_{i,t-1}) \\ &= \Lambda_{\text{rehab}}(\kappa_t + \delta D_i + \mathbf{z_i}'\eta + \mathbf{L}_{i,t-1}'\eta_L) \end{aligned} \tag{5}$$

Similarly, we estimate the parameters for a model on the probability of death in year t given that an individual was alive at the end of $t-1$[28] is given by:

[26] A dummy indicating that an individual has a non-musculoskeletal rehabilitation that year.

[27] Weighed version: $\ell_{\text{rehab}} = \sum_{i,t} \hat{\omega}_{i,t-1} \left[M_{i,t} \log \lambda_{i,t}^{\text{rehab}} + (1 - M_{i,t}) \log \left(1 - \lambda_{i,t}^{\text{rehab}}\right) \right]$

[28] Weighted version: $\ell_{\text{death}} = \sum_{i,t} \hat{\omega}_{i,t-1} \left[Y_{i,t} \log \lambda_{i,t}^{\text{death}} + (1 - Y_{i,t}) \log \left(1 - \lambda_{i,t}^{\text{death}}\right) \right]$

$$\begin{aligned}\lambda_{i,t}^{\text{death}} &= \Pr(Y_{i,t}=1|\text{alive at the end of } t-1) \\ &= \Lambda_{\text{death}}\left(\alpha_t + \theta_D D_i + \theta_M M_{i,t} + \theta_{DM} D_i M_{i,t} + \mathbf{z}_i'\rho + \mathbf{L}_{i,t-1}'\rho_L\right)\end{aligned} \tag{6}$$

The rehabilitation and death models use $\hat{\omega}_{i,t-1}$, with $\hat{\omega}_{i,0} = 1$, to account for the availability of the earlier records needed to construct the predictors. The models for the post-divorce history, $\mathbf{L}_{i,t}$, use $\hat{\omega}_{i,t}$ because fitting these models also requires the values recorded in year t. Once fitted, the history models are used to simulate how these variables evolve under each hypothetical divorce status.

In the BASE and RSD datasets, the observation can be missing for $\mathbf{L}_{i,t}$ and $M_{i,t}$. The gaps are often not random. For example, a curiosity of the BASE dataset is that records often stop in the year before death. Suppose an individual has observations for 2015 and 2016 and died in 2017. If we stopped at 2016, which is the last observation, we would end up discarding the known death in 2017. We retain the death year information, but, since 2017 is missing, we will not know whether that individual started rehabilitation. When jointly estimating the models in Equations 5 and 6 for such cases, we attach a 50% weight to each rehabilitation possibility. In other words, suppose the rehabilitation probability estimated from Equation 5 is 20% and then subsequently, when this probability is fed into Equation 6, the death probability is 1%, meaning that the probability of rehabilitation and then death is $80\% \times 1\% = 0.8\%$. Conversely, when the probability of no rehabilitation, 80%, is fed into Equation 6, the death probability is then 5% meaning that the probability of no rehabilitation then death is $20\% \times 5\% = 1\%$. We then weigh these two probabilities by 50% $\implies 0.5 \times 0.8\% + 0.5 \times 1\% = 0.9\%$. We assume that the reason for records stopping is unrelated to rehabilitation and death[29], e.g. the records do not stop because of deteriorating health.

Since $\mathbf{L}_{i,t}$ is itself affected by divorce and evolves annually, we cannot hold it fixed at its observed values when generating the hypothetical divorce paths needed for the mediation decomposition[30]. Hence, we need to model the evolution of $\mathbf{L}_{i,t}$ among individuals who survive, i.e. observations where $Y_{i,t} = 0$, so that we can simulate the counterfactual histories under the hypothetical divorce status for our mediation analysis (Robins, 1986; Lin et al., 2017). We fit nine separate models on the survivors, weighted by the same $\hat{\omega}_{i,t}$ (Lin et al., 2017) used in the weighted versions of Equations 5 and 6. We use a log-scale linear regressions for `income` and `pension_income`, logit for `unsuccessful_rehab_application_this_year` and `non_ms_rehab`, and multinomial logit for `income_source`. For the four employment day related counts, we use a two part model: a logit when there is at least one day and a linear regression for the number of days. These nine models condition on divorce status D_i, the covariates used for matching $\mathbf{z}_i$, $\mathbf{L}_{i,t-1}$, current mediator $M_{i,t}$, prior mediator $M_{i,t-1}$ and follow-up-year fixed effects. The two-part models for `income` and `pension_income` additionally include the current `income_source`, the models for the four employment-day counts additionally

[29] As a sensitivity analysis, when an observation is missing in the year of death, we fill up the row with last observed values. This has 100 bootstrap replicates.

[30] Suppose an individual earns 30,000 in 2013, divorces in 2014 and then earns 22,000 in 2015. Divorce could have caused the earnings to drop to 22,000 if the individual switched jobs. To determine the individual's rehabilitation or death risk if they had not divorced, we cannot plug in 22,000 because 22,000 is already corrupted by divorce. We need a clean number uncorrupted by divorce.

include the current `income_source`, `income`, `pension_income` and the day counts simulated before them, and the logit models include all of these, with the model for `non_ms_rehab` also including the current `unsuccessful_rehab_application_this_year`. Please see the section on Updating the evolution of $\mathbf{L}_{i,t}$ in the Appendix.

Once the parameters for these nine equations are estimated, we simulate the counterfactual histories, one year at a time using the following process.

1. Rehab probability (uses $\mathbf{L}_{i,t-1}$)→ draw→ $M_{i,t}$
    - Since the simulation starts in the first follow-up year, t_0+1, we initialize using the individual's true observed history in t_0-1, i.e[31], $\mathbf{L}_{i,t_0-1}$, and estimate the probability of rehabilitation, $\lambda^{\text{rehab}}_{i,t_0+1}$. Since we need to feed M_{i,t_0+1} into the next step and not $\lambda^{\text{rehab}}_{i,t_0+1}$, we draw a random number from a uniform distribution $U(0,1)$ and if it is smaller than $\lambda^{\text{rehab}}_{i,t_0-1}$, then the value of M_{i,t_0+1} is set at 1 and 0 otherwise.
2. Death probability (uses $\mathbf{L}_{i,t-1}$, $M_{i,t}$) → if individual i survives, continue
3. History-update (uses $\mathbf{z}_i$, D_i, $\mathbf{L}_{i,t-1}$, $M_{i,t-1}$, $M_{i,t}$) → produces $\mathbf{L}_{i,t}$
    - We use $\mathbf{L}_{i,t_0-1}$, M_{i,t_0}(= 0 since no one in the sample has had a musculoskeletal rehabilition by t_0) and M_{i,t_0+1} to simulate $\mathbf{L}_{i,t_0+1}$.
4. Hand $\mathbf{L}_{i,t}$ forward, repeat for $t+1$

True divorce status D_i is fixed at baseline and can simply be plugged into Equation 5 for rehabilitation. In order to find the counterfactual effect, we can simply plug in the opposite value since there are only two divorce statuses and each status has a direct opposite. Hence, in step 1, we obtained the rehabilition probability and thus, predicted rehabilitation occurrence at time t, $M_{i,t}(d_i)$ for an individual with divorce status d_i. For individual i whose plugged in divorce status is $d_i = 1$, denote the rehabilitation occurrence in time t as $M_{i,t}(1)$ and when $d_i = 0$, $M_{i,t}(0)$. An individual's rehabilitation timing is given by $M_i(1)$ and $M_i(0)$[32].We then obtain the sets, $M(1)$ and $M(0)$, which contain the rehabilitation timing for all matched treated individuals.

Subsequently, when we cross $d_i \in \{0,1\}$ with the resulting $M_i(d_i)$ and $M_i(d_i')$ where $d_i \neq d_i'$, we yield four different possibilities $\in \{(0,M_i(0)),(0,M_i(1)),(1,M_i(0)),(1,M_i(1))\}$. We might be tempted to plug in these four possibilities into Equations 5, 6 and the equations for updating $\mathbf{L}_{i,t}$ (in the Appendix) but $M_i(d_i)$ is not a fixed characteristic but the output of a year-by-year stochastic process. In other words, since $M_i(d_i)$ is based on *at least one random* draw under a hypothetical divorce status, there is no direct opposite like in the case with divorce status. Furthermore, we have two dimensions, d_i and $M_i(d_i)$, where we need a counterfactual but it is difficult to defend *individual* counterfactuals for $(0,M_i(1))$ and $(1,M_i(0))$. Think of this like a natural experiment. If we only wanted a clean effect of divorce, we can externally impose divorce on some individuals. The individuals who we forced to divorce would experience the rehabilitation timing of someone with a divorced status, likewise for the non-divorced. If

[31]We do not use the history in t_0 to initialize since it gives both treated and controls the same starting point before simulating counterfactuals. At t_0, the treated may already be affected by divorce.

[32]For example, if $M_i(d_i) = t+2$, then $M_{i,t'}(d_i) = 0$ when $0 < t' < t+2$ and $M_{i,t+2}(d_i) = 1$.

we wanted a clean direct effect of divorce and an indirect effect attributable to rehabilitation, we not only need to force someone to divorce but also experience the rehabilitation timing of a non-divorced person at the same time and vice versa, i.e. we need the off-diagonal or counterfactuals. This is not possible with a natural experiment as we cannot impose both dimensions simultaneously. For example, suppose we want to plug in the $(0, M_i(1))$ combination. Suppose the probability of rehabilitation under divorce is 80% and under non-divorce 5%. This means the timing of $M_i(1)$ is likely to be earlier than that of $M_i(0)$. Hence, plugging in $d_i = 0$ together with $M_i(1)$ (which is, to put it crudely, 'contaminated' with individual i's identity when $d_i = 1$) into Equation 6 would be like imposing both dimensions simultaneously. This argument also holds for the $(1, M_i(0))$ combination. To manage this 'contamination', we sever the link between individual i and their estimated rehabilitation timing by attaching another individual's rehabilitation timing. While $(0, M_i(0))$ and $(1, M_i(1))$ do not have this 'contamination' problem, we also sever the link between the individual and estimated rehabilitation timing for consistency.

To do this, we make use of the distribution of $M(1)$ and $M(0)$. This would then be similar to imposing an external rehabilitation timing. Before step 2, we replace individual i's value of $M_i(d_i)$ with individual j's $M_j(d_j)$ where $d_i = d_j$. This means that when both individuals i and j have an identical plugged in divorce status, individual i will have individual j's rehabilitation timing. This second-best approach helps us identify what the average death risk for individuals with a particular plugged in divorce status looks like under a random redistribution of rehabilitation timing[33].

Suppose individual i received individual j's value when $d_i = 0$, and individual k's value when $d_i = 1$. This means that individual i's hypothetical rehabilitation timings going into step 2 take the values of $M_j(0)$ and $M_k(1)$ respectively. We can then plug in the four combinations, $\in \{(0, M_j(0)), (0, M_k(1)), (1, M_j(0)), (1, M_k(1))\}$, into Equation 6 in step 2 for individual i in time t. We then update $\mathbf{L}_{i,t}$ and repeat all four steps until we reach five years.

Since individual i's $M_{i,t}$ depends on a random draw which subsequently affects $\mathbf{L}_{i,t}$, we re-run the entire process 25 times with different draws for the rehabilitation occurrence to average out simulation randomness. Let $M_{i,r}(d_i)$ and $Y_{i,r}(d_i, M(d_i'))$ denote individual i's simulated rehabilitation and death indicators by the horizon of five years from t_0 and $R = 25$ denote the number of Monte Carlo repeats. The Monte Carlo average for individual i is given by:

$$\begin{aligned} \hat{M}_i(d_i) &= \frac{1}{R} \sum_{r=1}^{R} M_{i,r}(d_i) \\ \hat{Y}_i(d_i, M(d_i')) &= \frac{1}{R} \sum_{r=1}^{R} Y_{i,r}(d_i, M(d_i')) \end{aligned} \tag{7}$$

[33] One caveat is that, it may be possible that individual j's rehabilitation timing is unlikely to be produced given individual i's rehabilitation probability, for example, if individual j's probability of rehabilitation in year 1 is high (e.g., 90%) while individual i's is low (e.g., 10%). This is called a positivity or overlap concern (VanderWeele et al., 2014; Petersen et al., 2012). To deal with overlap concern, we only draw from individuals in the same gender × 5-year age band × any prior rehab group.

We then average across the matched treated individuals. Let $\mathcal{T}$ denote the set of N_M matched treated individuals:

$$\begin{aligned} \hat{M}(d) &= \frac{1}{N_M} \sum_{i \in \mathcal{T}} \hat{M}_i(d_i) \\ \hat{Y}(d, M(d')) &= \frac{1}{N_M} \sum_{i \in \mathcal{T}} \hat{Y}_i(d_i, M(d'_i)) \end{aligned} \tag{8}$$

In Equation 8, we plug in the four different combinations to obtain *population* averages of the rehabilitation risk under the hypothetical divorce status, d and mortality risk under the hypothetical divorce d combined with the rehabilitation timing of divorce status d' where $d, d' \in \{0, 1\}$. We can then decompose into the direct effect and indirect effect:

$$\begin{aligned} \widehat{\text{Total effect}} &= \hat{Y}(1, M(1)) - \hat{Y}(0, M(0)) \\ \widehat{\text{Direct effect}} &= \hat{Y}(1, M(0)) - \hat{Y}(0, M(0)) \\ \widehat{\text{Indirect effect}} &= \hat{Y}(1, M(1)) - \hat{Y}(1, M(0)) \end{aligned} \tag{9}$$

The total effect is simply answering what the difference in mortality risk is between divorced and non-divorced individuals within five years. The direct effect captures part of the difference in mortality risk related to divorce but unrelated to rehabilitation, i.e. given that rehabilitation timing is held constant, how does divorce (potentially operating through financial strain, loneliness, stress) affect mortality risk. The indirect effect answers how much of the difference in mortality risk is attributed to divorce shifting individuals onto the rehabilitation timing distribution typical of divorced individuals. We calculated interventional direct and indirect effects (VanderWeele et al., 2014) which is different from natural direct and indirect effects (Pearl, 2001). A natural indirect effect answers what would happen to individual i's mortality risk if their rehabilitation timing shifted from what it looks like when individual i is non-divorced to what it would look like if individual i is divorced while holding divorce status constant at $d = 1$. In contrast, an interventional indirect effect answers what would happen to average mortality risk if the population's rehabilitation timing shifts from the distribution typical of non-divorced individuals to a distribution typical of divorced individuals while holding divorce status constant at $d = 1$.

Before computing the decomposition, we check whether the fitted models reproduce the observed data. We simulate each group under its own divorce status and compare the simulated yearly cumulative incidence of musculoskeletal rehabilitation, cumulative mortality, and mean earnings and pension income with the observed values. The reason we do this is because in an earlier version of the code, due to the runaway earnings, the simulated musculoskeletal rehab came out at 30% against the true 4.6-8.6% observed. For mortality, if the absolute diffence exceeds 10 per 1,000, then we do not proceed with the decomposition so far. So awaiting fixes. I will not put this part in the final draft because I will probably tweak the simulation parameters and tolerances until the conditions are met so that decomposition can proceed.

Finally, we then bootstrap the matched pairs 100 times (Efron and Tibshirani, 1993) with replacement and repeat the entire procedure. We do this to obtain the 95% confidence interval for the total, direct and indirect effects.

4 Results

4.1 Summary statistics

	Before matching			After matching		
Variable (measured in $t_0 - 1$ unless stated)	Divorced	Control	SMD	Divorced	Control	SMD
Panel A: Continuous variables						
Age (years)	45.96	50.97	−0.340	45.96	45.97	−0.001
Earnings (euros)	22,511	14,107	0.412	22,511	23,709	−0.057
Earnings, $t_0 - 2$ (euros)	21,538	13,630	0.396	21,538	22,777	−0.060
Earnings, $t_0 - 3$ (euros)	20,679	13,119	0.388	20,679	22,036	−0.067
Pension income (euros)	70.56	230.56	−0.438	70.56	63.50	0.027
Pension income, $t_0 - 2$ (euros)	61.82	214.94	−0.440	61.82	55.30	0.027
Pension income, $t_0 - 3$ (euros)	54.45	200.10	−0.440	54.45	48.32	0.028
Full-time contribution periods	22.05	79.12	−0.435	22.05	20.67	0.016
Part-time contribution periods	3.29	8.56	−0.297	3.29	3.00	0.020
Cumulative rehabilitations to date	0.151	0.035	0.320	0.151	0.150	0.003
Cumulative mental health rehabilitations to date	0.064	0.009	0.250	0.064	0.058	0.020
Years since last rehabilitation[a]	1.59	1.99	−0.244	1.59	1.73	−0.087
Rehabilitation starts, $t_0 - 2$	0.036	0.007	0.180	0.036	0.033	0.016
Rehabilitation starts, $t_0 - 3$	0.029	0.007	0.147	0.029	0.030	−0.007
Days in regular employment[b]	258.20	248.79	0.060	258.20	264.06	−0.039
Days unemployed[b]	47.16	34.74	0.121	47.16	48.04	-0.008
Days in marginal employment[b]	50.28	47.23	0.027	50.28	49.30	0.009
Days in care leave[b]	3.29	3.14	0.005	3.29	3.22	0.002
Number of children	0.083	0.050	0.090	0.083	0.102	−0.043
Panel B: Binary variables (%)						
Ever had a mental health rehabilitation	5.31	0.78	0.266	5.31	4.76	0.025
Unsuccessful rehabilitation application	2.64	0.43	0.180	2.64	2.24	0.026
Unsuccessful rehabilitation application, $t_0 - 2$	2.39	0.44	0.166	2.39	2.02	0.025
Unsuccessful rehabilitation application, $t_0 - 3$	1.98	0.44	0.141	1.98	1.76	0.016
Full-time contribution data missing	91.66	71.57	0.537	91.66	92.61	−0.035
Part-time contribution data missing	91.66	71.57	0.537	91.66	92.61	−0.035
Regular employment days missing[b]	14.34	38.15	−0.562	14.34	13.14	0.035
Unemployment days missing[b]	13.62	35.82	−0.532	13.62	12.84	0.023
Marginal employment days missing[b]	13.52	35.72	−0.533	13.52	12.76	0.023
Care leave days missing[b]	13.25	35.60	−0.539	13.25	12.45	0.024
Child born in year	0.04	0.15	−0.035	0.04	0.09	−0.020
Panel C: Categorical variables[c]	Levels	Before			After	
Income source	4	0.663			0.035	
Income source, $t_0 - 2$	4	0.614			0.038	
Income source, $t_0 - 3$	4	0.600			0.046	
Federal state[d]	16/17	0.253			0.025	
Occupation category[d]	10/11	0.448			0.024	
Education[d]	3/4	0.203			0.033	
Age group of youngest child	6	0.087			0.050	
Age band	16	0.337			0.000	
Sex	2	0.018			0.000	
Any rehabilitation by $t_0 - 1$	2	0.350			0.000	

Table 1: Covariate balance before and after matching, lag-3 specification, 2014–2018

Notes: 84,461 matched pairs. SMD is the standardized mean difference, divorced minus control. Earnings and pension income are set to zero when the record for that year is unobserved for matching convenience.

[a]Defined only for those with a prior rehabilitation (88.3% of divorced vs 97.2% before matching missing prior rehabilitation).

[b]Employment related days are missing for some person-years. Missing records are indicated by 'Regular employment/Unemployment/Marginal employment/Care leave days missing' in Panel B. Missing day counts are filled with the median before matching.

[c] We calculate SMD for each category separately and report the largest SMD across all levels

[d] Where two levels are shown, this indicates that missing values are counted as its own category.

Table 1 compares the treated (divorced) and controls before and after matching. Before matching, the treated are younger than the controls (46.0 vs 51.0 years), there are more currently employed individuals than pensioners, and more have had prior rehabilitation, including mental health rehabilitation. After matching, no variable has an SMD exceeding 0.10. The average SMD (from the 73 continuous variables, binary variables and each category) falls from 0.239 before matching to 0.022 after matching. The variables with the largest remaining difference after matching are years since last rehabilitation (-0.087) and earnings at $t_0 - 3$ (-0.067). Before matching, earnings are observed for 86.75% of treated and 57.24% of controls and after matching 86.75% and 87.38% respectively. Pension income is observed for 8.34% of trated and 28.43% of controls before matching, and after matching 8.34% and 7.39% respectively. Among those with observed earnings, the mean earnings are €25,950 for treated and €24,646 for controls before matching and €27,132 after matching. Similarly, among those with observed pension income, the mean is €846 for treated and for controls before and after matching, the means are €811.0 and €860 respectively. The mean full-time contribution periods is 264.3 for treated and respectively 278.3 and 279.9 for controls before and after matching. The missingness rate for full-time and part-time contribution periods is not only numerically identical but matches pension income missingness exactly, which is consistent with all three variables tracking the same employment vs status. For a more detailed breakdown, please see the section on 6 in the Appendix.

4.2 RQ1: Musculoskeletal rehabilitation

	Lag-3, 2014–2018	Lag-1, 2014–2018	Lag-1, 2012–2018
Panel A: Men			
Hazard model: odds ratio			
Divorce[a,*]	2.043 [1.929, 2.165]	2.084 [1.966, 2.208]	1.968 [1.879, 2.061]
Divorce[b,*]	1.993 [1.878, 2.114]	2.042 [1.924, 2.168]	1.930 [1.841, 2.024]
Cumulative incidence at five years			
Divorced (%)	8.30 [8.06, 8.57]	8.29 [8.05, 8.54]	8.42 [8.24, 8.61]
Controls (%)	4.16 [3.94, 4.32]	4.08 [3.91, 4.29]	4.38 [4.22, 4.54]
Risk difference (per 1,000)	41.4 [39.0, 44.5]	42.1 [38.7, 45.1]	40.4 [37.8, 42.7]
Risk ratio[c]	1.995	2.033	1.922
Matched pairs	41,430	41,661	62,562
Person-years	399,168	401,491	602,609
First MS rehabilitations	5,161	5,154	8,008
Panel B: Women			
Hazard model: odds ratio			
Divorce[a,*]	2.303 [2.175, 2.438]	2.297 [2.170, 2.431]	2.195 [2.097, 2.298]
Divorce[b,*]	2.196 [2.070, 2.330]	2.206 [2.080, 2.340]	2.099 [2.002, 2.201]
Cumulative incidence at five years			
Divorced (%)	8.81 [8.58, 9.08]	8.77 [8.52, 9.09]	8.77 [8.56, 8.95]
Controls (%)	3.93 [3.72, 4.09]	3.93 [3.77, 4.10]	4.11 [3.98, 4.25]
Risk difference (per 1,000)	48.7 [46.0, 51.9]	48.4 [45.4, 51.9]	46.6 [44.3, 49.0]
Risk ratio[c]	2.239	2.234	2.136
Matched pairs	43,031	43,445	64,833
Person-years	417,755	421,723	629,336
First MS rehabilitations	5,481	5,515	8,345

Table 2: Effect of divorce on first musculoskeletal rehabilitation entry, separately for men and women, current run.

Notes: Brackets contain 95% confidence intervals. The SMD-adjusted specification is identical to the matching-only specification and is therefore omitted. Risk intervals use 200 matched-pair bootstrap replicates; odds-ratio intervals use pair-clustered standard errors. Risk difference is treated minus control. Cumulative incidence accounts for death as a competing event.

[*] All reported divorce odds ratios have $p < 0.001$.

[a] Specification with only divorce and follow-up year dummies.

[b] Specification with full controls at $t-1$ for each follow-up year t with missingness indicators. These controls may be affected by divorce, so this specification provides a separate conditional comparison.

[c] Only point estimates are reported because I forgot to save the sex-specific risk-ratio CIs.

Panel A in Table 2 reports reports the discrete-time hazard model in Equation 1 among men, and Panel B reports the corresponding estimates for women, with death as a competing event. In the lag-3 specification, the matching-only odds ratios are 2.043 ($p < 0.001$) for men and 2.303 ($p < 0.001$) for women, indicating approximately 104% and 130% higher annual odds of entering rehabilitation, respectively, over the five-year follow-up. With previous-year controls, the odds ratios decrease slightly to 1.993 ($p < 0.001$) for men and 2.196 ($p < 0.001$) for women. At five years, cumulative incidence among men is 8.30% for divorcees and 4.16% for controls, a difference of 41.4

rehabilitation entries per 1,000 people ($p < 0.05$). Among women, the corresponding risks are 8.81% and 3.93%, a difference of 48.7 entries per 1,000 ($p < 0.05$).




Figure 1: Cumulative incidence of first musculoskeletal rehabilitation

Notes: 41,430 matched pairs for men and 43,031 for women. 5-year risk difference is 41.4 (95% CI [38.96, 44.48]) and 48.7 (95% CI [45.97, 51.88]) per 1,000 for men and women respectively. Women minus men is 7.3 per 1,000 (95% CI [2.84, 12.19]). Confidence intervals use 200 bootstrap replications. Death is a competing event.

Figure 1 displays the 5-year cumulative effect of divorce on starting a musculoskeletal rehabilitation. For men, the 5-year cumulative risk difference is 41.14 per 1,000 between divorced and controls. For women, it is 48.7 per 1,000. The constructed confidence intervals exclude zero. The overall incidences among divorced are 8.3% and 8.8% for men and women respectively, compared with 4.1% and 3.9% among controls.

One possible interpretation is that divorce affects the transition from latent or previously untreated musculoskeletal problems into formal rehabilitation differently for men and women. Biological differences in susceptibility to musculoskeletal conditions may shape the underlying health problems on which these post-divorce processes operate. Wolf et al. (2015) find that women appear more biologically susceptible to particular degenerative and joint conditions due to differences in anatomy, hormones and biomechanics. However, certain injury patterns are more common in men, and when men develop osteoporotic hip fractures, outcomes may be worse. Hence, they cannot by themselves explain why the divorce-control contrast is slightly higher for women.

To explore possible explanations involving the transition into formal rehabilitation, we next examine associations with age, previous applications and recorded economic circumstances. In the specification with annually updated $t-1$ covariates and missingness indicators, each additional year of age within a five-year age band is associated with 2.9% higher annual odds of first musculoskeletal rehabilitation entry (OR 1.029; $p < 0.001$). Although pre-divorce application histories enter the matching, the annually updated model also accounts for subsequent applications. Within the matched sample, an unsuccessful rehabilitation application in $t-1$ is associated with

higher odds of entry at time t (OR 4.230; $p < 0.001$). Higher recorded earnings are also positively associated with entry (OR 1.393; $p < 0.001$), whereas higher recorded pension income is negatively associated (OR 0.573; $p < 0.001$). The income odds ratios refer to a one-standard-deviation increase in log-transformed amounts. These conditional associations provide information about the rehabilitation entry process, although their role in generating the divorce–control difference remains to be established.

There are further tests I want to conduct to shed more light on the mechanism beween divorce and rehabilitation entry. Eventually, I would substitute rehabilitation incidence with rehabilitation duration, `mcdams_succeed_1`.

There are three hypotheses:

1. In a failing marriage, individuals might not have time to seek out rehabilitation for existing complaints because they are dealing with the stress of a difficult relationship and maybe still exerting effort to try to preserve the marriage. Once divorce occurs, the individual may finally have free time to prioritize seeking rehabilitation.
    - Run extra analysis using rehabilitation application dates. Among people who already had musculoskeletal issues BEFORE divorce (but we don't observe them in the data since we excluded individuals with prior musculoskeletal rehabilitation), does divorce make them more likely to apply for their first rehabilitation?
    - Check rehabilitation application patterns before divorce (we already used unsuccessful applications in matching and in the event study, we showed that both had parallel trends prior to t_0 although there is a tiny significant difference in unsuccessful applications between divorced and controls per year - difference in level not trend). There shouldn't be a huge difference between divorced and controls prior to t_0. If there is a divergence after t_0, i.e. applications from divorced increase while those for controls remain, then that can provide support for the hypothesis that individuals finally have time to seek help.
    - Link first musculoskeletal rehabilitation starts from $t_0 + 1$ to their original application dates. Estimate how many additional starts per 1,000 matched individuals came from applications filed before, in the same month as, or after the end of marriage month. If much of the excess comes from applications already filed before the marriage end month, we can examine whether applications were awaiting a decision or had already been approved and waiting to start but it doesn't really help for this hypothesis.
    - For divorced, use the court decision date as second timing marker.
2. Even in failing marriages, partners are usually still obligated to perform physical tasks, like lifting, chores, etc. After divorce, however, individuals are often on their own, and have no one to rely on for the physically challenging tasks. Initially, divorced individuals might be able to bear the brunt, but over time, bodies might give in increasing likelihood of musculoskeletal rehabilitation.
    - With the existing data, we can plot the annual first musculoskeletal rehabilitation starts in each follow-up year since we already have the table

somewhere. There are about 8 to 10 starts in each follow-up year, so this first point might not support the hypothesis.

- As a proxy of whether an individual generally needs help performing physical tasks, we can look at periods of incapacity before divorce. If a period of incapacity existed, this can signify that the individual is likelier to delegate physically demanding tasks when a partner is present. See point 4 too.
- We should also check whether the individuals receiving a musculoskeletal rehab in years 3 to 5 are the same people who also underwent a musculoskeletal rehabilitation in years 1 to 2. Then we can recalculate the incidence instead and see if the same share of divorced and control undergo their first musculoskeletal rehabilitation. We have this in the existing data. Even if you count for the unique individuals, divorced individuals still are twice as likely to experience musculoskeletal rehabilitation start.
- We should also look at total musculoskeletal rehabilitation days per 1,000 divorced and control individuals including AND excluding repeat episodes over 5 years or in each follow-up year.

3. In some marriages, both partners have unequal expectations regarding the standard of cleanliness that needs to be maintained. In failing marriages, through pettiness or cold war, one partner may have to constantly pick up after the other. After a divorce, it might be easier for the individual who has higher standards of cleanliness to maintain the standard.
    - This can't really be tested unless we use SOEP.
    - Maybe we can look at children and see whether the musculoskeletal entries differ among people with children and people without and whether that difference is constant among divorce and controls
4. Losing a partner's financial support makes restoring work capacity a lot more valuable. Since the individual is now depending on their own paycheck, they can't afford to NOT seek help because if they were to become disabled, they no onger have anyone to rely on.
    - Record work impairment, `mcauan` (1: unable to work, 2: able to work, 0: legacy code or missing), at first entry. This can tell us whether the extra starts are attributed to people who are already suffering from some sort of work incapacity.
    - We can also test whether individuals who were already working before the split are more likely to apply for first musculoskeletal rehabilitation. If applications or rehabilitations rise among individuals who were already working, then this can point towards an individual's desire to protect existing earnings.
    - Check the point above for each income quintile. The urgency to protect existing earnings might differ for different income groups.
5. Divorce could change access to DRV rehabilitation. There are two ways to qualify for DRV rehab, i.e. either have an accumulated 15-year contribution history

(regardless of when the contributions are made) OR 6 months of qualifying contributions in the previous 2 years.

- With pension equalization, an individual can have additional records added to contribution history which then pushes them across the threshold for qualification.

4.3 RQ2: Mortality

	Lag-3, 2014–2018	Lag-1, 2014–2018	Lag-1, 2012–2018
Panel A: Men			
Hazard model: odds ratio			
Divorce[a]	1.284 [1.217, 1.355]	1.284 [1.217, 1.354]	1.251 [1.201, 1.303]
Divorce[b]	1.211 [1.139, 1.287]	1.204 [1.133, 1.279]	1.158 [1.105, 1.214]
Cumulative mortality at five years			
Divorced (%)	3.49 [3.32, 3.63]	3.48 [3.32, 3.66]	3.21 [3.09, 3.37]
Controls (%)	3.12 [2.96, 3.29]	3.06 [2.92, 3.21]	2.99 [2.86, 3.11]
Risk difference (per 1,000)	3.6 [1.6, 5.9]	4.2 [1.9, 6.8]	2.2 [0.7, 4.3]
Risk ratio[c]	1.116	1.138	1.075
Matched pairs	41,430	41,661	62,562
Person-years (full follow-up)	580,506	583,645	990,325
Deaths (full follow-up)	4,933	4,952	8,267
Panel B: Women			
Hazard model: odds ratio			
Divorce[a]	1.171 [1.076, 1.274]	1.142 [1.050, 1.241]	1.199 [1.124, 1.279]
Divorce[b]	1.050 [0.954, 1.156]	1.038 [0.944, 1.140]	1.074 [0.998, 1.155]
Cumulative mortality at five years			
Divorced (%)	1.25 [1.13, 1.35]	1.25 [1.13, 1.36]	1.18 [1.08, 1.28]
Controls (%)	1.33 [1.22, 1.43]	1.34 [1.23, 1.46]	1.24 [1.17, 1.33]
Risk difference (per 1,000)	−0.8 [−2.2, 0.6]	−0.9 [−2.5, 0.5]	−0.6 [−1.8, 0.5]
Risk ratio[c]	0.937	0.930	0.948
Matched pairs	43,031	43,445	64,833
Person-years (full follow-up)	610,804	616,512	1,039,740
Deaths (full follow-up)	2,050	2,093	3,528

Table 3: Effect of divorce on all-cause mortality, separately for men and women.

Notes: Brackets contain 95% confidence intervals. The SMD-adjusted specification is identical to the matching-only specification and is therefore omitted. Risk and risk-difference intervals use 200 matched-pair bootstrap replicates; odds-ratio intervals use pair-clustered standard errors. Cumulative mortality is estimated as one minus Kaplan–Meier survival. Risk differences are divorced minus control, expressed per 1,000 people. Differences and ratios are calculated using unrounded risk estimates.

[a]Specification with only divorce and follow-up year dummies.

[b]Specification with full controls measured at $t-1$ for each follow-up year t, with missingness indicators. These controls may be affected by divorce, so this specification provides a separate conditional comparison.

[c]Only point estimates are reported because sex-specific risk-ratio confidence intervals were not saved in the exported outputs.

Panel A in Table 3 reports mortality estimates for men, and Panel B reports the corresponding estimates for women. In the preferred lag-3 specification, the

matching-only odds ratios are 1.284 ($p < 0.001$) for men and 1.171 ($p < 0.001$) for women, indicating approximately 28% and 17% higher annual odds of death, respectively, over the available follow-up. With previous-year controls, the odds ratio is 1.211 ($p < 0.001$) for men and 1.050 (insignificant) for women. At five years, cumulative mortality among men is 3.49% for divorcees and 3.12% for controls, a difference of 3.6 deaths per 1,000 people ($p < 0.05$). Among women, the corresponding risks are 1.25% and 1.33%, a difference of −0.8 deaths per 1,000 (insignificant). The pooled hazard estimates summarize a longer follow-up period than the five-year risk contrasts and therefore need not have the same direction when mortality differences change over time.




Figure 2: Five-year cumulative all-cause mortality among divorcees and matched controls, by sex

Notes: 41,430 matched pairs for men and 43,031 for women. The five-year risk difference, divorced minus control, is 3.63 per 1,000 for men (95% CI [1.60, 5.91]) and −0.84 per 1,000 for women (95% CI [−2.23, 0.56]). The difference between the female and male risk differences is −4.47 per 1,000 (95% CI [−6.95, −1.96]). Confidence intervals use 200 matched-pair bootstrap replications. Cumulative mortality is estimated as one minus Kaplan–Meier survival. Shaded areas show pointwise 95% confidence intervals for each risk curve.

The difference in mortality rate is higher for controls in the first two follow-up years. The overall mortality rate is 1.49 per 1,000 and 3.58 per 1,000 for divorced and controls respectively in the first follow-up year. In the second follow-up year, the overall mortality

rate is 3.41 and 4.06 per 1,000 for divorced and controls. This difference reverses sign from the third follow-up year onwards with the divorced mortality rate at 5.06, 6.26 and 7.25 per 1,000 in follow-up years 3, 4 and 5. For controls the mortality rate is 4.42, 4.33 and 5.72 per 1,000. This is true for both genders, although most of this effect is driven by men. The later excess mortality offsets the earlier mortality deficit before the cumulative differnce becomes positive. For the pooled sample, this occurs by year 5, for men year 4. For women, the later excess does not fully offset the earlier deficit within five years. The 5-year cumulative difference is 3.63 deaths per 1,000 among men ($p < 0.001$), compared with -0.84 among women (insignificant). The women-minus-men contrast is -4.47 per 1,000 ($p < 0.001$), indicating a significantly larger 5-year mortality difference among men.

DONE How much more likely are divorced people to die during the follow up period

DONE Compare lag1 and lag3 spec mortality risk

- Cumulative mortality over 5 years
- Total, direct and indirect effects plus 95% confidence interval.
- How often is it the case that the borrowed rehab timing is completely off?
- Total, direct and indirect effects for men vs women.

5 Discussion

6 Conclusion

References

Aalen, Odd O. and Søren Johansen (1978). "An Empirical Transition Matrix for Non-Homogeneous Markov Chains Based on Censored Observations". In: *Scandinavian Journal of Statistics* 5.3, pp. 141–150.

Andorsen, Ole Fredrik et al. (2017). "A prospective cohort study on risk factors of musculoskeletal complaints (pain and/or stiffness) in a general population: The Tromsø study". In: *PLOS ONE* 12.7, e0181417. DOI: 10.1371/journal.pone.0181417.

Austin, Peter C. (2009). "Balance Diagnostics for Comparing the Distribution of Baseline Covariates between Treatment Groups in Propensity-Score Matched Samples". In: *Statistics in Medicine* 28.25, pp. 3083–3107. DOI: 10.1002/sim.3697.

Berntsen, Kjersti Norgård and Øystein Kravdal (2012). "The relationship between mortality and time since divorce, widowhood or remarriage in Norway". In: *Social Science & Medicine* 75.12, pp. 2267–2274.

Bethge, Matthias et al. (2019). "Effects of Nationwide Implementation of Work-Related Medical Rehabilitation in Germany: Propensity Score Matched Analysis". In: *Occupational and Environmental Medicine* 76.12, pp. 913–919. DOI: 10.1136/oemed-2019-106065. URL: https://doi.org/10.1136/oemed-2019-106065.

Björkenstam, Emma et al. (2013). “Risk of new psychiatric episodes in the year following divorce in midlife: Cause or selection? A nationwide register-based study of 703,960 individuals”. In: *International Journal of Social Psychiatry* 59.8, pp. 801–804.

Brenscheidt, S. et al. (2026). *Arbeitswelt im Wandel: Zahlen – Daten – Fakten (2026)*. baua: Praxis 3315. Dortmund: Bundesanstalt für Arbeitsschutz und Arbeitsmedizin, p. 84. DOI: 10.21934/baua:praxis20260304. URL: https://www.baua.de/DE/Angebote/Publikationen/Praxis/A122 (visited on 08/09/2026).

Brockmann, Hilke and Thomas Klein (2004). “Love and death in Germany: The marital biography and its effect on mortality”. In: *Journal of Marriage and Family* 66.3, pp. 567–581.

Bundesinstitut für Berufsbildung (BIBB) (n.d.). *German Classification of Occupations 1988 (KldB 88)*. Accessed 1 November 2026. URL: https://metadaten.bibb.de/en/classification/detail/5.

Bundesrepublik Deutschland (1988). *Sozialgesetzbuch Fünftes Buch (SGB V)—Gesetzliche Krankenversicherung, §§ 47–49 and 51*. German. https://www.gesetze-im-internet.de/sgb_5/. Current consolidated version; accessed 9 August 2026.

Bundesrepublik Deutschland (1989). *Sozialgesetzbuch Sechstes Buch (SGB VI)—Gesetzliche Rentenversicherung, §§ 10 and 11*. German. https://www.gesetze-im-internet.de/sgb_6/. Current consolidated version; accessed 9 August 2026.

Bundesrepublik Deutschland (1994). *Gesetz über die Zahlung des Arbeitsentgelts an Feiertagen und im Krankheitsfall (Entgeltfortzahlungsgesetz, EntgFG), §§ 3 and 5*. German. https://www.gesetze-im-internet.de/entgfg/. Current consolidated version; accessed 9 August 2026.

Choi, Seung-Won Emily and Brandon Wagner (2026). “Marital status and high-impact pain among middle-aged and older adults: A population-based longitudinal study in the United States”. In: *The Journal of Pain* 41, p. 106208. DOI: 10.1016/j.jpain.2026.106208.

Cimas, Marta et al. (2018). “Chronic musculoskeletal pain in European older adults: Cross-national and gender differences”. In: *European Journal of Pain* 22.2, pp. 333–345. DOI: 10.1002/ejp.1123.

Couch, Kenneth A., Christopher R. Tamborini, and Gayle L. Reznik (2015). “The long-term health implications of marital disruption: Divorce, work limits, and Social Security disability benefits among men”. In: *Demography* 52.5, pp. 1487–1512. DOI: 10.1007/s13524-015-0424-z.

Deutsche Rentenversicherung (n.d.[a]). *Allgemeine medizinische Reha*. German. https://www.deutsche-rentenversicherung.de/DRV/DE/Reha/Medizinische-Reha/Allgemeine-med-Reha/allgemeine-med-reha_node.html. Accessed 9 August 2026.

Deutsche Rentenversicherung (n.d.[b]). *Übergangsgeld*. German. https://www.deutsche-rentenversicherung.de/DRV/DE/Reha/Warum-Reha/uebergangsgeld.html. Accessed 9 August 2026.

Deutsche Rentenversicherung Bund (2005a). *Kriterien zur Rehabilitationsbedürftigkeit für Leistungen zur medizinischen Rehabilitation bei Erkrankungen des Stütz- und Bewegungsapparates aus der Sicht der Deutschen Rentenversicherung Bund*. Tech. rep. Stand 18.3.2005/1.10.2005. Deutsche Rentenversicherung Bund. URL: https://www.deutsche-rentenversicherung.de/SharedDocs/Downloads/DE/

Experten/infos_fuer_aerzte/begutachtung/leitlinien_rehabeduerftigkeit_stuetz_bewegungsapparat_kurzfassung_pdf.pdf.

Deutsche Rentenversicherung Bund (2005b). *Kriterien zur Rehabilitationsbedürftigkeit für Leistungen zur medizinischen Rehabilitation bei Erkrankungen des Stütz- und Bewegungsapparates aus der Sicht der Deutschen Rentenversicherung Bund*. German. Guideline, short version. Status 18 March/1 October 2005; accessed 9 August 2026. Deutsche Rentenversicherung Bund. URL: https://www.deutsche-rentenversicherung.de/SharedDocs/Downloads/DE/Experten/infos_fuer_aerzte/begutachtung/leitlinien_rehabeduerftigkeit_stuetz_bewegungsapparat_kurzfassung_pdf.pdf.

Deutsche Rentenversicherung Bund (2024). *Reha-Atlas 2024: Die Teilhabeleistungen der Deutschen Rentenversicherung in Zahlen, Fakten und Trends*. Reha-Atlas. Reports rehabilitation activity and expenditure for 2023. Deutsche Rentenversicherung Bund. URL: https://www.deutsche-rentenversicherung.de/SharedDocs/Downloads/DE/Statistiken-und-Berichte/Rehaatlas/2024/rehaatlas-2024-download.html (visited on 08/09/2026).

Deutscher Bundestag (Apr. 3, 2009). *Gesetz über den Versorgungsausgleich (Versorgungsausgleichsgesetz – VersAusglG)*. Article 1 of the Gesetz zur Strukturreform des Versorgungsausgleichs (VAStrRefG), BGBl. I, p. 700. In force since 1 September 2009. Last amended by Article 1 of the Act of 12 May 2021, BGBl. I, p. 1085. Short-marriage exclusion: §3(3); de minimis exclusion: §18; contractual waiver: §§6–8. URL: https://www.gesetze-im-internet.de/versausglg/.

Efron, Bradley and Robert J. Tibshirani (1993). *An Introduction to the Bootstrap*. Vol. 57. Monographs on Statistics and Applied Probability. New York: Chapman and Hall/CRC. ISBN: 978-0-412-04231-7. DOI: 10.1007/978-1-4899-4541-9.

England, Bryant R. et al. (2018). "Increased Cardiovascular Risk in Rheumatoid Arthritis: Mechanisms and Implications". In: *BMJ* 361, k1036. DOI: 10.1136/bmj.k1036. URL: https://doi.org/10.1136/bmj.k1036.

Grundy, Emily MD and Cecilia Tomassini (2010). "Marital history, health and mortality among older men and women in England and Wales". In: *BMC Public Health* 10.1, p. 554.

Hawker, Gillian A. et al. (2014). "All-Cause Mortality and Serious Cardiovascular Events in People with Hip and Knee Osteoarthritis: A Population Based Cohort Study". In: *PLOS ONE* 9.3, e91286. DOI: 10.1371/journal.pone.0091286. URL: https://doi.org/10.1371/journal.pone.0091286.

Jansson, Catarina, Ellenor Mittendorfer-Rutz, and Kristina Alexanderson (2012). "Sickness Absence because of Musculoskeletal Diagnoses and Risk of All-Cause and Cause-Specific Mortality: A Nationwide Swedish Cohort Study". In: *Pain* 153.5, pp. 998–1005. DOI: 10.1016/j.pain.2012.01.028. URL: https://doi.org/10.1016/j.pain.2012.01.028.

Joung, Inez M. A. et al. (1998). "A longitudinal study of health selection in marital transitions". In: *Social Science & Medicine* 46.3, pp. 425–435. DOI: 10.1016/S0277-9536(97)00186-X.

Kaplan, E. L. and Paul Meier (1958). "Nonparametric Estimation from Incomplete Observations". In: *Journal of the American Statistical Association* 53.282, pp. 457–481. DOI: 10.1080/01621459.1958.10501452.

King, Gary and Langche Zeng (2001). “Logistic Regression in Rare Events Data”. In: *Political Analysis* 9.2, pp. 137–163. DOI: 10.1093/oxfordjournals.pan.a004868.

Koch, Elisabeth, ed. (2025). *Münchener Kommentar zum Bürgerlichen Gesetzbuch, Band 9: Familienrecht I, §§ 1297–1588, VersAusglG, GewSchG, LPartG*. 10th ed. Munich: C.H. Beck. ISBN: 978-3-406-81029-9.

Kohn, Jennifer L. and Susan L. Averett (2014). “The effect of relationship status on health with dynamic health and persistent relationships”. In: *Journal of Health Economics* 36, pp. 69–83. DOI: 10.1016/j.jhealeco.2014.03.010.

Kposowa, Augustine J (2000). “Marital status and suicide in the National Longitudinal Mortality Study”. In: *Journal of Epidemiology & Community Health* 54.4, pp. 254–261.

Leopold, Thomas (2018). “Gender differences in the consequences of divorce: A study of multiple outcomes”. In: *Demography* 55.3, pp. 769–797.

Lin, Sheng-Hsuan et al. (2017a). “Mediation analysis for a survival outcome with time-varying exposures, mediators, and confounders”. In: *Statistics in Medicine* 36.26, pp. 4153–4166. DOI: 10.1002/sim.7426.

Lin, Sheng-Hsuan et al. (2017b). “Parametric Mediational g-Formula Approach to Mediation Analysis with Time-varying Exposures, Mediators, and Confounders”. In: *Epidemiology* 28.2, pp. 266–274. DOI: 10.1097/EDE.0000000000000609.

Metsä-Simola, Niina and Pekka Martikainen (2013a). “Divorce and changes in the prevalence of psychotropic medication use: A register-based longitudinal study among middle-aged Finns”. In: *Social Science & Medicine* 94, pp. 71–80. DOI: 10.1016/j.socscimed.2013.06.027.

Metsä-Simola, Niina and Pekka Martikainen (2013b). “The short-term and long-term effects of divorce on mortality risk in a large Finnish cohort, 1990–2003”. In: *Population Studies* 67.1, pp. 97–110.

Meulman, Iris et al. (2023). “Estimating healthcare expenditures after becoming divorced or widowed using propensity score matching”. In: *The European Journal of Health Economics* 24.7, pp. 1047–1060. DOI: 10.1007/s10198-022-01532-z.

Meune, Christophe et al. (2009). “Trends in Cardiovascular Mortality in Patients with Rheumatoid Arthritis over 50 Years: A Systematic Review and Meta-Analysis of Cohort Studies”. In: *Rheumatology* 48.10, pp. 1309–1313. DOI: 10.1093/rheumatology/kep252. URL: https://doi.org/10.1093/rheumatology/kep252.

Nüesch, Eveline et al. (2011). “All Cause and Disease Specific Mortality in Patients with Knee or Hip Osteoarthritis: Population Based Cohort Study”. In: *BMJ* 342, p. d1165. DOI: 10.1136/bmj.d1165. URL: https://doi.org/10.1136/bmj.d1165.

Pearl, Judea (2001). “Direct and Indirect Effects”. In: *Proceedings of the Seventeenth Conference on Uncertainty in Artificial Intelligence*. UAI’01. San Francisco, CA: Morgan Kaufmann Publishers Inc., pp. 411–420.

Pellón-Elexpuru, Inés et al. (2024). “Divorce and physical health: A three-level meta-analysis”. In: *Social Science & Medicine* 352, p. 117005. DOI: 10.1016/j.socscimed.2024.117005.

Petersen, Maya L. et al. (2012). “Diagnosing and Responding to Violations in the Positivity Assumption”. In: *Statistical Methods in Medical Research* 21.1, pp. 31–54. DOI: 10.1177/0962280210386207.

Prang, Khic-Houy, Janneke Berecki-Gisolf, and Sharon Newnam (2015). “Recovery from musculoskeletal injury: The role of social support following a transport accident”. In: *Health and Quality of Life Outcomes* 13, p. 97. DOI: 10.1186/s12955-015-0291-8.

Reese, Jennifer Barsky et al. (2010). “Pain and functioning of rheumatoid arthritis patients based on marital status: Is a distressed marriage preferable to no marriage?” In: *The Journal of Pain* 11.10, pp. 958–964. DOI: 10.1016/j.jpain.2010.01.003.

Robins, James M. (1986). “A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect”. In: *Mathematical Modelling* 7.9-12, pp. 1393–1512. DOI: 10.1016/0270-0255(86)90088-6.

Rosenbaum, Paul R. (1989). “Optimal Matching for Observational Studies”. In: *Journal of the American Statistical Association* 84.408, pp. 1024–1032. DOI: 10.2307/2290079.

Rosenbaum, Paul R and Donald B Rubin (1983). “The central role of the propensity score in observational studies for causal effects”. In: *Biometrika* 70.1, pp. 41–55.

Sbarra, David A and Paul J Nietert (2009). “Divorce and death: Forty years of the Charleston Heart Study”. In: *Psychological Science* 20.1, pp. 107–113.

Sbarra, David A et al. (2014). “Marital dissolution and major depression in midlife: A propensity score analysis”. In: *Clinical Psychological Science* 2.3, pp. 249–257. DOI: 10.1177/2167702613498727.

Shor, Eran et al. (2012). “Meta-analysis of marital dissolution and mortality: Reevaluating the intersection of gender and age”. In: *Social Science & Medicine* 75.1, pp. 46–59. DOI: 10.1016/j.socscimed.2012.03.010.

Sjøgren, Per et al. (2009). “Epidemiology of chronic pain in Denmark: An update”. In: *European Journal of Pain* 13.3, pp. 287–292. DOI: 10.1016/j.ejpain.2008.04.007.

Stevenson, Betsey and Justin Wolfers (2006). “Bargaining in the Shadow of the Law: Divorce Laws and Family Distress”. In: *The Quarterly Journal of Economics* 121.1, pp. 267–288. DOI: 10.1093/qje/121.1.267.

Tamborini, Christopher R., Gayle L. Reznik, and Kenneth A. Couch (2016). “Work disability among women: The role of divorce in a retrospective cohort study”. In: *Journal of Health and Social Behavior* 57.1, pp. 98–117. DOI: 10.1177/0022146515627849.

Torvik, Fartein Ask et al. (2015). “Health, health behaviors, and health dissimilarities predict divorce: Results from the HUNT study”. In: *BMC Psychology* 3.1, p. 13.

VanderWeele, Tyler J., Stijn Vansteelandt, and James M. Robins (2014). “Effect Decomposition in the Presence of an Exposure-Induced Mediator-Outcome Confounder”. In: *Epidemiology* 25.2, pp. 300–306. DOI: 10.1097/EDE.0000000000000034.

Wade, Terrance J and David J Pevalin (2004). “Marital transitions and mental health”. In: *Journal of Health and Social Behavior* 45.2, pp. 155–170.

Wald, Abraham (1943). “Tests of Statistical Hypotheses Concerning Several Parameters When the Number of Observations Is Large”. In: *Transactions of the American Mathematical Society* 54.3, pp. 426–482. DOI: 10.1090/S0002-9947-1943-0012401-3.

Wiborg, Øyvind Nicolay and Alexi Gugushvili (2025). “Does partnership predict mortality? Evidence from a twin fixed effects study design”. In: *SSM – Population Health* 30, p. 101805. DOI: 10.1016/j.ssmph.2025.101805.

Wolf, Jennifer Moriatis et al. (2015). “Male and female differences in musculoskeletal disease”. In: *JAAOS-Journal of the American Academy of Orthopaedic Surgeons* 23.6, pp. 339–347.

Woods, Valerie (2005). “Work-related musculoskeletal health and social support”. In: *Occupational Medicine* 55.3, pp. 177–189. DOI: 10.1093/occmed/kqi085.

Zulkarnain, Alice and Sanders Korenman (2019). "Divorce and health in middle and older ages". In: *Review of Economics of the Household* 17.4, pp. 1081–1106. DOI: 10.1007/s11150-018-9435-z.

Appendix

Rehabilitation system in Germany

For an individual covered by statutory health insurance, musculoskeletal pain first enters the health insurance system through sickness absence rather than through the DRV. If pain makes an employee unable to work, the individual must immediately inform the employer of the incapacity and the expected duration. If the incapacity lasts longer than three calendar days, a medical certificate needs to be acquired from the individual's general practitioner, although the employer may require certification from the first day. The employer continues to pay the normal salary for up to six weeks for incapacity caused by the same illness (Bundesrepublik Deutschland, 1994). After the six weeks, if the illness continues, the statutory health insurer steps in and pays *Krankengeld*, which is usually about 70% of covered gross income up to a maximum of 90% of net income. For the same illness, this benefit is limited to 78 weeks within a 3-year period, counted from the first day of incapacity, meaning that the time of illness paid by the employer is included in the 78 weeks (Bundesrepublik Deutschland, 1988). Medical rehabilitation facilitated by the DRV requires a separate application and assessment of whether earning capacity is substantially endangered or already reduced, if rehabilitation is expected to prevent or improve that reduction (Bundesrepublik Deutschland, 1989; Deutsche Rentenversicherung, n.d.). This application is usually independent of the six weeks and *Krankengeld* although the health insurer can allow the individual ten weeks to submit a rehabilitation application indicating that earning capacity is threatened (Bundesrepublik Deutschland, 1988). For musculoskeletal conditions, the DRV assessment focuses on documented functional limitations and their consequences for work and social participation. Chronic pain, psychosocial circumstances surrounding incapacity, comorbidity, and active cooperation are considerations in this assessment rather than four cumulative eligibility conditions. Isolated pain is insufficient to establish rehabilitation need for certain hip, knee, and shoulder conditions (Deutsche Rentenversicherung Bund, 2005). Approved medical rehabilitation is normally provided on an inpatient or full-day outpatient basis and usually lasts approximately three weeks. The employer continues wages during rehabilitation if entitlement remains, otherwise, the DRV may pay *Ubergangsgeld*, normally 68% of previous net earnings or 75% for an insured person with a qualifying child, while entitlement to *Krankengeld* is suspended (Deutsche Rentenversicherung; Bundesrepublik Deutschland, n.d. 1988).

Dealing with a large sample of controls

Suppose in 2015, there are 50,000 individuals who are divorced and 2 million eligible controls. Using all 2 million controls to estimate the logit model would cause memory issues and take a long time to run. So we imposed a cap of $450,000(500,000 - 50,000)$ eligible controls which are then randomly selected, although the same seed is used for reproducibility. If there are less than 450,000 eligible controls, we use all of them to

estimate the parameters for the logit model. Once we have estimated the logit model, we estimate the propensity score for everyone, not only the 450,000 controls and 50,000 treated. Then we match based on propensity scores and end up with a treated-control pair, i.e. in 2015, we would have 50,000 treated and 50,000 controls.

The controls used for fitting are selected by a seeded hash of the individual id, so that the same individuals are selected in every run. Since the fitting sample contains all treated individuals but only a fraction, f, of the controls, this inflates the log-odds of divorce by $\log(\frac{1}{f})$ for everyone. To rectify this, we add $\log f$ to each individual's predicted log-odds before converting it to a probability (King and Zeng, 2001).

The matching step faces a similar problem. With optimal matching, we need a distance for each treated and control pair in an exact-match group, and there can me millions of such pairs. In an earlier run on a 4% sample, divorced individuals tend to have a scores of between 0.12 to 0.19, while controls have scores around 0.001. Hence, divorced individuals may have no controls nearby. In optimal matching, these divorced individuals may be left unmatched to keep the overall distance low. We impose a 'penalty distance' which is the product of the number of unmatched individuals multiplied by 0.01. With such a penalty, then the algorithm is less likely to leave divorced individuals unmatched. If the number of pairs is too large for a dense solver, we only keep the n nearest controls where n is the number of divorced individuals in that exact-match group.

Updating the evolution of $\mathbf{L}_{i,t}$

We first estimate the parameters of the model for `income_source`. Let `income_source` be denoted by $S_{i,t}$:

$$\Pr(S_{i,t} = k|\cdot) = \frac{\exp\left(c_k + \delta_k D_i + \mathbf{z}_i'\eta_k + \mathbf{L}_{i,t-1}'\eta_{L,k} + \mu_k M_{i,t-1} + \zeta_k M_{i,t} + \tau_{k,t}\right)}{\sum_{k'}\exp(\cdot)}$$

where c_k is the constant and $\tau_{k,t}$ is the time fixed effect at year t for category k.

Then, we estimate the parameters for the models on `income` and `pension_income`. Let $A_{i,t}^v$ represent `income` and `pension_income`. The first part is a logit for whether the amount is positive:

$$\begin{aligned}\Pr\left(A_{i,t}^v \mid \cdot\right) = \Lambda\Big(&a_v + b_v D_i + \mathbf{z}_i' g_v + \mathbf{L}_{i,t-1}' g_{L,v} + m_v M_{i,t-1} + n_v M_{i,t} \\ &+ \tau_{v,t}^0 + \phi_v' \mathbb{1}\left[\widetilde{S}_{i,t}\right]\Big)\end{aligned}$$

The second part is a linear regression of the log amount among individuals with a positive `income` or `pension_income`:

$$\log(1 + A_{i,t}^v) = c_v + \beta_v D_i + \mathbf{z}_i'\gamma_v + \mathbf{L}_{i,t-1}'\gamma_{L,v} + \mu_v M_{i,t-1} + \zeta_v M_{i,t} + \tau_{v,t} + \phi_v' \mathbb{1}[\tilde{S}_{i,t}] + \varepsilon_{i,t}^v$$

where v indicates whether we are estimating `income` or `pension_income`, a_v and c_v are constants, $\tilde{S}_{i,t}$ was the previously estimated `income_source`, $\tau^0_{v,t}$ and $\tau_{v,t}$ are time fixed effects. In the simulation, `income` is drawn only if $\tilde{S}_{i,t}$ has the values 'earnings only' or 'earnings and pension', while `pension_income` is drawn when $\tilde{S}_{i,t}$ has the values 'pension only' or 'earnings and pension', otherwise the amount is zero. If the first part leads to a positive amount, then we add a random error within the estimated residual standard deviation. This is to introduce some variation in case the simulated individual shares identical characteristics with others. We also cap the result at the largest observed log amount.

Next, we estimate the models for the four employment days related counts $E^e_{i,t}$ where e indicates whether we are estimating the days in the year for `regular_employment`, `unemployment`, `marginal_employment` or `care_leave`. This is also the order in which we estimate and update. Each count again has two parts. The first part is a logit for $E^e_{i,t} > 0$:

$$\begin{aligned}\Pr\left(E^e_{i,t} > 0 \mid \cdot\right) = \Lambda\Big(&a_e + b_e D_i + \mathbf{z}_i' g_e + \mathbf{L}'_{i,t-1} g_{L,e} + m_e M_{i,t-1} + n_e M_{i,t} \\ &+ \tau^0_{e,t} + \phi_e' \mathbb{1}\left[\widetilde{S}_{i,t}\right] + \tilde{\mathbf{A}}'_{i,t} \pi^0_e + \tilde{\mathbf{E}}^{<e'}_{i,t} \theta^0_e\Big)\end{aligned}$$

where $\tilde{\mathbf{A}}_{i,t}$ contains $\log(1+\tilde{A}^v_{i,t})$ for the simulated `income` and `pension_income` and $\tilde{\mathbf{E}}'^{<e}_{i,t}$ contains $\log(1+\tilde{E}^{e'}_{i,t})$ for the day counts simulated before e. For the second part, we transform the number of days into a share of the largest observed count $\bar{E}^e \leq 366$, $q^e_{i,t} = (E^e_{i,t} - 0.5)/\bar{E}^e \in (0,1)$, where subtracting half a day keeps the share strictly between 0 and 1. Among individuals with at least one day, we estimate:

$$\begin{aligned}\log\left(\frac{q^e_{i,t}}{1-q^e_{i,t}}\right) = &c_e + \beta_e D_i + \mathbf{z}_i' \gamma_e + \mathbf{L}'_{i,t-1} \gamma_{L,e} + \mu_e M_{i,t-1} + \zeta_e M_{i,t} \\ &+ \tau^0_{e,t} + \phi_e' \mathbb{1}\left[\widetilde{S}_{i,t}\right] + \tilde{\mathbf{A}}'_{i,t} \pi^0_e + \tilde{\mathbf{E}}^{<e'}_{i,t} \theta_e + \epsilon^e_{i,t}\end{aligned}$$

as a first step in avoiding impossible values. We then obtain the mean $\hat{\mu}^e_{i,t}$ and residual standard deviation, $\hat{\sigma}_e = \left(\Sigma_{i,t}\hat{\omega}_{i,t}(\hat{\epsilon}^e_{i,t})^2 / \Sigma_{i,t}\hat{\omega}_{i,t}\right)^{0.5}$. The number of days is then:

$$\tilde{E}^e_{i,t} = \min\left\{\max\left\{\text{round}\left(\bar{E}^e \Lambda\left(\hat{\mu}^e_{i,t} + \hat{\sigma}_e u^e_{i,t}\right) + 0.5\right), 1\right\}, \bar{E}^e\right\}, \qquad u^e_{i,t} \sim N(0,1)$$

We finally estimate the parameters for the model on the binary indicators `non_ms_rehab` and `unsuccessful_rehab_application_this_year`:

$$\begin{aligned}\Pr\left(B^1_{i,t} = 1 \mid \cdot\right) = \Lambda\Big(&\alpha_1 + \theta_1 D_i + \mathbf{z}_i' \rho_1 + \mathbf{L}'_{i,t-1} \rho_{L,1} + \mu_1 M_{i,t-1} + \nu_1 M_{i,t} \\ &+ \tau_{1,t} + \widetilde{\mathbf{A}}'_{i,t} \pi_1 + \mathbb{1}\left[\widetilde{S}_{i,t} = 1\right] \pi_{S,1}\Big)\end{aligned}$$

$$\Pr\left(B^2_{i,t}=1 \mid \cdot,\, B^1_{i,t}\right) = \Lambda\Big(\alpha_2 + \theta_2 D_i + \mathbf{z}_i'\rho_2 + \mathbf{L}'_{i,t-1}\rho_{L,2} + \mu_2 M_{i,t-1} + \nu_2 M_{i,t} + \tau_{2,t} + \widetilde{\mathbf{A}}'_{i,t}\pi_2 + \mathbb{1}\left[\widetilde{S}_{i,t}\right]\pi_{S,2} + \pi_B \widetilde{B}^1_{i,t}\Big)$$

where $B^1_{i,t}$ is a dummy representing `unsuccessful_rehab_application_this_year` and is estimated first, $B^2_{i,t}$ is a a dummy representing `non_ms_rehab` which is estimated second hence also contains the predicted $\widetilde{B}^1_{i,t}$ and $\widetilde{\mathbf{A}}_{i,t}$ is a vector containing the drawn level ($\log(1+x)$ not log) of the simulated current amounts of (`income`, `pension_income`, and days in `employment`, `unemployment`, `marginal_employment` and `care_leave`) which we just estimated in the regressions above.

Descriptive summary

Year	Individuals	Divorces	% of panel[a]	All rehab	MS rehab	Deaths
2011	13,829,551	23,364	0.169	219,182	95,087	31,808
2012	13,857,384	22,640	0.163	228,922	100,297	169,528
2013	13,890,621	21,730	0.156	226,893	107,153	174,345
2014	13,971,071	21,200	0.152	233,652	112,715	170,579
2015	14,066,307	20,216	0.144	233,004	111,258	179,268
2016	14,173,132	18,958	0.134	229,302	107,608	178,304
2017	14,271,784	18,167	0.127	227,455	106,209	184,061
2018	14,427,112	17,152	0.119	233,189	108,016	189,902
2019	14,525,854	16,616	0.114	244,078	115,975	186,690
2020	14,520,197	15,527	0.107	198,377	94,086	198,307
2021	14,555,740	13,885	0.095	214,640	101,387	205,950
2022	14,682,308	11,618	0.079	225,418	105,466	216,538
2023	14,747,677	9,350	0.063	224,790	102,330	226,362
Total	17,559,407	230,423		2,938,902	1,367,587	2,311,642
Person-years	185,518,738					

Table 4: Overall data description

Notes: Each individual is counted once per year. Divorces are counted in the year of the first divorce we observe. Of the 230,423 individuals who divorced, 226,855 divorces are from the first recorded marriage. Rehabilitation counts are successful, completed rehabilitation spells. MS stands for musculoskeletal. MS counts codes 20, 53 and 56 (Table 8). [a] Divorces as a percentage of individuals

The number of divorces fell steadily from 23,364 divorces in 2011 (0.169% of individuals observed in 2011) to 9,350 divorces in 2023 (0.063% of individuals in 2023). The national divorce rate also fell during this period although at a slower rate. The reason for this difference may be attributed to who gets counted in the VA dataset. The VA dataset explicitly excludes marriages which lasted three years or less or if the individuals divorcing waived the pension equalization through a prenuptial agreement or divorce settlement. Furthermore, if the income of the individuals divorcing are roughly equal, then the family court allows pension equalization to be skipped (Deutscher Bundestag, 2009; Koch, 2025).

Category		Codes	Events	Unique persons	Share (%)
1	Musculoskeletal	20 / 53, 56	1,367,587	797,759	46.5
2	Mental health	30 / 55	446,382	343,054	15.2
3	Neoplasms[a]	26 / 48	360,033	≤283,584	12.3
4	Nervous system[a]	28 / 52	155,632	≤112,451	5.3
5	Cardiovascular[b]	22	90,534	75,206	3.1
6	Digestive & metabolic[b]	23	82,370	68,791	2.8
7	Skin[a]	25 / 57	77,322	≤61,093	2.6
8	Other conditions[b]	31	67,891	53,149	2.3
9	Endocrine[b]	24	58,945	49,335	2.0
10	Respiratory[b]	21	40,945	34,765	1.4
	Top-10 total		2,747,641		93.5
	All groups		2,938,902	1,754,448	100.0

Table 5: Top ten rehabilitation diagnosis groups

Notes: The DRV reclassified rehabilitation approvals from ICD-10-based diagnosis groups (codes 20–33) to treatment-based indicator groups (codes 40–59) on 1 January 2021; a small number of spells already carry the new codes in 2018–2020. Combined groups show old / new codes; event counts are summed across both systems.

[a]Mapping inferred from year-by-year volume patterns; code 52 (neurology) may additionally include stroke cases previously under code 22 (cardiovascular).

[b]Old-system code only since it is possible that the indicator groups system redistributes these diagnostic chapters across multiple treatment-based codes (principally code 51, internal medicine), preventing one-to-one mapping. These counts therefore end in 2021–2023 with the last old-code spells.

Unique-person counts for MS and mental health are counted directly across their harmonized codes. Other combined groups retain upper bounds (≤), since the same person may appear in both coding systems. In the current run, mental health variables count codes 30 and 55. The pre-divorce histories used for matching end before the new code was in regular use.

Across the 20% sample, 1,754,448 individuals have at least one completed rehabilitation (about 10.0% of all individuals in the sample). There are 2,938,902 rehabilitations covered. In any given year, about 1.3% of the observed population starts a rehabilitation. The most common rehabilitation type is for musculoskeletal related conditions, covering 46.5% off all rehabilitations, followed mental health rehabilitation, covering 15.2% of all rehabilitations.

Table 6: Descriptive statistics of the matching variables in the matched sample, lag-3 specification, 2014–2018

Variable ($t_0 - 1$ unless stated)	N	Mean	Median	SD	Min	Max	Conditional mean
Panel A: Continuous variables							
Age (years)	168,922	45.97	46.00	9.85	19.00	93.00	45.97
Earnings (euros)	168,922	23,110	19,970	21,206	0	105,088	26,542.91
Earnings, $t_0 - 2$ (euros)	168,922	22,157	18,820	20,785	0	91,800	25,677.61
Earnings, $t_0 - 3$ (euros)	168,922	21,358	17,867	20,351	0	95,271	24,645.46
Pension income (euros)	168,922	67.03	0.00	257.21	0.00	2,986.34	852.38
Pension income, $t_0 - 2$ (euros)	168,922	58.56	0.00	238.42	0.00	2,886.77	832.58
Pension income, $t_0 - 3$ (euros)	168,922	51.39	0.00	221.27	0.00	2,839.35	812.45
Full-time contribution periods (months)	168,922	21.36	0.00	88.19	0.00	610.00	271.64
Part-time contribution periods (months)	168,922	3.14	0.00	14.36	0.00	441.00	39.95
Cumulative rehabilitations to date	168,922	0.150	0.000	0.455	0.000	7.000	1.284
Cumulative mental health rehabilitations to date	168,922	0.061	0.000	0.281	0.000	5.000	1.206
Years since last rehabilitation	19,766	1.66	1.00	1.58	0.00	6.00	2.370
Rehabilitation starts, $t_0 - 2$	168,922	0.035	0.000	0.203	0.000	2.000	1.131
Rehabilitation starts, $t_0 - 3$	168,922	0.030	0.000	0.191	0.000	2.000	1.144
Regular employment (days)	145,713	261.15	365.00	152.24	0.00	366.00	330.10
Unemployed (days)	146,572	47.60	0.00	110.92	0.00	366.00	236.54
Marginal employment (days)	146,720	49.79	0.00	114.45	0.00	366.00	248.65
Care leave (*Pflegezeit*, days)	147,216	3.25	0.00	32.77	0.00	366.00	288.12
Number of children	168,922	0.093	0.000	0.441	0.000	3.000	1.912
Panel B: Binary variables (% with the characteristic)							
Ever had a mental health rehabilitation	168,922	5.03					
Unsuccessful rehabilitation application	168,922	2.44					
Unsuccessful rehabilitation application, $t_0 - 2$	168,922	2.21					
Unsuccessful rehabilitation application, $t_0 - 3$	168,922	1.87					
Full-time contribution period missing	168,922	92.14					
Part-time contribution period missing	168,922	92.14					
Child born in year	168,922	0.06					

Notes: For earnings, pension income and contribution periods, the conditional means shown are means among recorded values: zeros introduced for missing records are excluded, but any genuinely recorded zeros are retained. Their means excluding every zero have not yet been established. For all other continuous variables, conditional means exclude every zero. Employment-day summaries use available observations, including genuine zeros; their conditional means use only observations with positive days.

The preferred lag-3 sample contains 84,461 matched pairs. In the year before divorce, individuals are 46.0 years old on average. Mean earnings are €23,110 and median earnings are €19,970; among observations with recorded earnings, the mean is €26,543. Mean earnings increase from €21,358 three years before divorce to €23,110 one year before divorce. Pension income has a median of zero, with a mean of €67.03 across all observations and €852.38 among observations with recorded pension income. Note that earnings, pension income and contribution periods include zeros assigned when the underlying variable is unobserved. Employment-day statistics use available observations, including genuine zeros, and exclude missing day counts. For earnings, pension income and contribution periods, the conditional means are calculated among observations with recorded values, retaining any genuinely recorded zeros. For the remaining continuous variables, the conditional means exclude every zero. Previous rehabilitation is recorded for 11.7% of matched observations, while 5.03% have a history of mental health rehabilitation. Years since last rehabilitation is defined only for those with a prior rehabilitation. Among observations with available employment information, regular employment averages 261.15 days and unemployment averages 47.60 days. Among observations with positive days, the corresponding means are 330.10

and 236.54 days. These statistics describe the pooled treated and control sample.

	Treated		Controls (year slots)	
Restriction	Kept	Excluded	Kept	Excluded
Panel A: lag-3 specification, 2014–2018 (preferred)				
First divorce in 2014–2018	95,693	—	70,909,406[a]	—
Not yet divorced at t_0	—	—	70,280,109	629,297
No first-marriage end in or before t_0	—	—	66,597,324	3,682,785
Divorce ends first-observed marriage	94,577	1,116	—	—
Alive and no MS rehab starting in t_0	92,754	1,823	66,043,471	553,853
Has a record in $t_0 - 1$	92,178	576	64,312,108	1,731,363
Has a valid age in $t_0 - 1$	92,178	0	64,310,753	1,355
No MS rehab before t_0	85,107	7,071	62,986,747	1,324,006
Has records in $t_0 - 2$ and $t_0 - 3$	84,461	646	59,961,604	3,025,143
Matched pairs	84,461	—	84,461	—
Panel B: lag-1 specification, 2012–2018				
First divorce in 2012–2018	140,063	—	98,657,411[a]	—
Not yet divorced at t_0	—	—	97,914,601	742,810
No first-marriage end in or before t_0	—	—	92,679,605	5,234,996
Divorce ends first-observed marriage	138,946	1,117	—	—
Alive and no MS rehab starting in t_0	136,332	2,614	91,913,444	766,161
Has a record in $t_0 - 1$	135,574	758	89,679,572	2,233,872
Has a valid age in $t_0 - 1$	135,574	0	89,678,216	1,356
No MS rehab before t_0	127,396	8,178	88,183,324	1,494,892
Matched pairs	127,395	—	127,395	—

Table 7: Specification samples

Notes: Treated individuals enter in exactly one year, so the treated columns count people. Controls can be eligible in several years, so the control columns count year slots. Hence, the 59,961,604 eligible control slots in Panel A belong to 13,138,278 individuals, and the 88,183,324 in Panel B to 14,363,580. The exclusion of controls with a recorded first-marriage end in or before t_0 is shown separately. Survival at t_0 and absence of an MS rehabilitation starting in t_0 are combined into one reporting row. Iintermediate counts are not reported. [a]Person-years in the treatment years.

To build the matches for the lag-3 specification, we use first observed divorces from 2014 to 2018 so that each cohort has at least five years of follow-up. In this window, there are 95,693 divorces. After imposing restrictions, we are left with 84,461 individuals. See Table 7 above. For the lag-1 specification, we use first observed divorces from 2012 to 2018. Similarly, after imposing restrictions and matching, we are left with 127,395 divorces out of 140,063.

t_0	First divorces	Eligible control slots	Matched pairs	Mean gap ($\times 10^{-6}$)	Maximum gap	Mean score (pre-match, $\times 10^{-3}$)	
						Treated	Controls
Panel A: lag-3 specification, 2014–2018							
2014	21,200	12,052,653	19,495	1.026	0.003244	3.49	1.63
2015	20,216	11,999,378	18,232	1.385	0.009110	3.31	1.53
2016	18,958	11,951,822	16,676	1.056	0.003119	3.12	1.40
2017	18,167	11,951,191	15,605	0.417	0.000662	2.76	1.31
2018	17,152	12,006,560	14,453	1.830	0.008302	2.71	1.21
Total	95,693		84,461	1.135	0.009110		
Panel B: lag-1 specification, 2012–2018							
2012	22,640	12,623,077	21,754	0.592	0.002776	3.20	1.74
2013	21,730	12,573,500	20,535	1.059	0.008991	3.12	1.64
2014	21,200	12,545,036	19,611	0.472	0.000544	3.21	1.57
2015	20,216	12,545,945	18,373	0.702	0.002985	2.99	1.47
2016	18,958	12,569,959	16,791	1.025	0.009391	2.85	1.34
2017	18,167	12,626,407	15,739	0.476	0.001860	2.55	1.25
2018	17,152	12,699,400	14,592	0.385	0.000971	2.52	1.16
Total	140,063		127,395	0.684	0.009391		

Table 8: Matching by treatment year

Notes: First divorces are all first observed divorces in the year, before any restriction. Eligible control slots are counted before controls used in earlier treatment years are removed from the pool available for matching. Matching proceeds chronologically without control reuse. Mean gap is the average absolute difference in propensity scores within matched pairs; the caliper is 0.01. Maximum gaps are reported on the original propensity-score scale. Total-row mean gaps are averages across all matched pairs. The final two columns report mean propensity scores in the eligible pools before matching, after the sampling correction.

Table 8 shows the matching results. Of the 84,461 eligible treated individuals for the lag-3 specification (and 127,395 for the lag-1 specification), the match rate is 100% (and respectively 99.99%). No control is used twice as a control and every treated individual has at least one control in their exact-match group (ie. their gender × 5-year age band × prior rehabilitation group). Although controls used in earlier years are removed from the pool available for later matching, this does not lead to a decline in matching success. In the lag-3 specification, eligible control slots decrease slightly from 12.05 million in 2014 to 12.01 million in 2018, although the change is not monotonic. These counts describe the eligible pools before earlier control use is removed. It is also possible for individuals to appear first as a control and then as a treated if that individual divorces eventually.

Across the lag-3 matched sample, 603 individuals out of 168,319 unique individuals appear as both treated and control (about 0.36%). There is a median gap of two years between being a control and being a treated. The matched sample also has close to an even split on gender with 41,430 male and 43,031 female pairs (49.1% vs 50.9%). 11.70% of individuals in the lag-3 specification and 9.35% in the lag-1 specification have had some rehabilitation before t_0. Across all years, the mean absolute gap between propensity scores within a pair is 1.135×10^{-6} in the lag-3 specification and 0.684×10^{-6} in the lag-1 specification. The maximum gaps are 0.009110 and 0.009391 respectively, both below the 0.01 caliper. The propensity scores are low because divorce is relatively rare: in the preferred specification, annual mean scores range from approximately 2.7 to 3.5 per 1,000 among divorcees and 1.2 to 1.6 per 1,000 among controls before matching.

Year relative to t_0	Earnings (euros)	Pension income (euros)	Rehabilitation starts (per 100 people)	Unsuccessful applications (pp)	Pension income missing (pp)	Regular employment (days)	Unemployed (days)	Marginal employment (days)	Care leave (days)
−3	−159.2	−0.93	−0.50	−0.18	0.07	1.33	−0.75	2.20	0.13
	(40.5)	(0.49)	(0.13)	(0.09)	(0.05)	(0.68)	(0.47)	(0.56)	(0.15)
−2	−41.4	−0.55	−0.05	−0.03	0.03	0.97	−0.38	1.67	0.10
	(30.2)	(0.39)	(0.14)	(0.10)	(0.04)	(0.56)	(0.41)	(0.45)	(0.12)
−1	0.0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	—	—	—	—	—	—	—	—	—
0	357.3	−1.69	1.59	0.83	0.18	2.72	13.87	−5.32	−0.51
	(32.5)	(0.48)	(0.12)	(0.09)	(0.05)	(0.54)	(0.42)	(0.44)	(0.13)
1	562.0	−1.94	2.29	0.92	0.17	4.71	15.52	−7.67	−1.08
	(46.0)	(0.71)	(0.14)	(0.09)	(0.07)	(0.69)	(0.50)	(0.55)	(0.19)
2	679.1	0.54	2.57	0.78	−0.09	3.96	14.43	−8.25	−1.16
	(54.7)	(0.86)	(0.14)	(0.09)	(0.08)	(0.76)	(0.54)	(0.61)	(0.23)
3	644.8	2.46	2.74	0.76	−0.29	2.19	14.45	−7.62	−1.23
	(61.9)	(0.99)	(0.14)	(0.09)	(0.09)	(0.81)	(0.56)	(0.65)	(0.27)
4	617.7	2.19	2.91	0.85	−0.30	1.53	13.11	−7.09	−1.41
	(68.9)	(1.18)	(0.15)	(0.09)	(0.10)	(0.84)	(0.58)	(0.68)	(0.30)
5	634.7	4.07	2.91	0.85	−0.45	0.03	11.00	−6.54	−1.41
	(75.5)	(1.35)	(0.15)	(0.09)	(0.11)	(0.88)	(0.58)	(0.70)	(0.33)
Joint Wald statistic	18.34	3.62	17.41	4.50	1.91	4.35	2.54	18.21	0.88
p-value	< 0.001	0.164	< 0.001	0.105	0.385	0.113	0.281	< 0.001	0.644
Pairs, t_0-3	84,461	84,461	84,461	84,461	84,461	68,802	69,759	69,770	70,142
Pairs, t_0-2	84,461	84,461	84,461	84,461	84,461	68,928	69,743	69,833	70,230
Pairs, t_0-1	84,461	84,461	84,461	84,461	84,461	70,976	71,501	71,649	72,134
Pairs, t_0+5	81,443	81,443	81,443	81,443	81,443	60,924	61,454	61,846	62,041

Table 9: Differences between matched pairs relative to t_0-1, from t_0-3 to t_0+5 (lag-3 specification, 2014–2018)

Notes: Each entry is the mean within-pair difference between divorcees and controls in that year, minus the reference-year mean difference at t_0-1. Standard errors in parentheses are clustered on the matched pair. The joint Wald tests concern the coefficients for t_0-3 and t_0-2 (two degrees of freedom). Outcome-specific observed-pair counts appear in the final rows. Employment-day counts are included only when both members of a pair have usable values; missing day counts are not filled. The composition of observed pairs can therefore change across years. Employment-day means are not participation shares. Reference-year zeros do not imply that divorcees and controls have identical levels in t_0-1. Because the pre-divorce values also enter matching, these rows describe the remaining differences in the histories used to select controls.

The estimates show how the difference between divorcees and their matched controls changes relative to the year before divorce. The earnings gap is €159.2 lower at t_0-3 and €41.4 lower at t_0-2 than at t_0-1, indicating that divorcees' earnings were already rising relative to controls before the recorded divorce. The rehabilitation-start gap also increases towards divorce. Joint tests reject unchanged pre-divorce gaps for earnings, rehabilitation starts and marginal-employment days (all $p < 0.001$), while providing no statistically significant evidence of changes in pension income, unsuccessful applications, pension-income missingness, regular-employment days, unemployment days or care-leave days.

After divorce, the earnings and rehabilitation gaps increase further: by t_0+5, they are respectively €634.7 and 2.91 rehabilitation starts per 100 people above their differences at t_0-1. The unemployment-day gap is 15.52 days above its reference-year difference in t_0+1 and remains 11.00 days above it in t_0+5. The regular-employment gap initially increases by 4.71 days in t_0+1, but is only 0.03 days above its reference-year difference by t_0+5. At that endpoint, the marginal-employment and care-leave gaps are respectively 6.54 and 1.41 days below their reference-year differences.

These are changes in the gaps between the groups, rather than their absolute

differences in each year. The employment estimates concern recorded day counts, not the shares of individuals with any days. Although the lag-3 specification incorporates three years of pre-divorce information into matching, some differences in pre-divorce trajectories remain and must be considered when interpreting the subsequent estimates. Changes in the composition of pairs with available employment information may also contribute to the patterns during follow-up.

	Divorced		Controls	
What happened	*n*	%	*n*	%
Neither recorded rehabilitation nor death	75,319	89.18	79,178	93.75
Died before any recorded MSK rehabilitation	1,915	2.27	1,867	2.21
Started MSK rehabilitation, survived	7,160	8.48	3,393	4.02
Started MSK rehabilitation, then died	67	0.08	23	0.03
Total matched assignments	84,461	100.00	84,461	100.00

Table 10: Recorded outcomes in the five years after t_0, lag-3 specification, 2014–2018

Notes: Outcomes cover $t_0 + 1$ through $t_0 + 5$. All percentages use the full matched denominator of 84,461 assignments in each group. The four outcome categories describe assignments entering follow-up. The 23 matched controls absent from the follow-up risk sets are reported separately and are not assigned an outcome during this window. Individuals appearing as both treated and control can contribute to both groups. "Survived" means no recorded death through $t_0 + 5$. Deaths come from the recorded year of death. Rehabilitation and death recorded in the same calendar year are classified as rehabilitation followed by death; their order within that year is not observed. Deaths after rehabilitation are calculated as all deaths minus deaths before rehabilitation. Missing annual records during follow-up contribute no recorded rehabilitation start. Percentages may not sum to 100 because of rounding.

In the five years after t_0, most individuals have neither a recorded musculoskeletal rehabilitation nor a recorded death: 89.18% among divorcees and 93.75% among controls entering follow-up. First MSK rehabilitation is recorded for 7,160 divorcees (8.48%) and 3,393 controls (4.02%). Death before any recorded MSK rehabilitation occurs among 2.27% of divorcees and 2.21% of controls. A further 67 divorcees and 23 controls have both a recorded rehabilitation and a subsequent death, including cases assigned this order because both events occur in the same calendar year.