

## More Applications of The Pumping Lemma

### The Pumping Lemma:

For infinite context-free language  $L$

there exists an integer  $m$  such that

for any string  $w \in L$ ,  $|w| \geq m$

we can write  $w = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

and it must be:

$$uv^i xy^i z \in L, \quad \text{for all } i \geq 0$$

## Non-context free languages

$$\{a^n b^n c^n : n \geq 0\} \quad \{vv : v \in \{a,b\}^*\}$$

## Context-free languages

$$\{a^n b^n : n \geq 0\} \quad \{ww^R : w \in \{a,b\}^*\}$$

**Theorem:** The language

$$L = \{vv : v \in \{a,b\}^*\}$$

is **not** context free

**Proof:** Use the Pumping Lemma  
for context-free languages

$$L = \{vv : v \in \{a,b\}^*\}$$

Assume for contradiction that  $L$   
is context-free

Since  $L$  is context-free and infinite  
we can apply the pumping lemma

$$L = \{vv : v \in \{a,b\}^*\}$$

Pumping Lemma gives a magic number  $m$   
such that:

Pick any string of  $L$  with length at least  $m$

we pick:  $a^m b^m a^m b^m \in L$

$$L = \{vv : v \in \{a,b\}^*\}$$

We can write:  $a^m b^m a^m b^m = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

Pumping Lemma says:

$$uv^i xy^i z \in L \quad \text{for all } i \geq 0$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$


We examine all the possible locations of string  $vxy$  in  $a^m b^m a^m b^m$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $vxy$  is within the first  $a^m$

$$v = a^{k_1} \quad y = a^{k_2} \quad k_1 + k_2 \geq 1$$




$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $vxy$  is within the first  $a^m$

$$v = a^{k_1} \quad y = a^{k_2} \quad k_1 + k_2 \geq 1$$



$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $vxy$  is within the first  $a^m$

$$a^{m+k_1+k_2} b^m a^m b^m = uv^2 xy^2 z \notin L$$

$$k_1 + k_2 \geq 1$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 1:**  $vxy$  is within the first  $a^m$

$$a^{m+k_1+k_2} b^m a^m b^m = uv^2 xy^2 z \notin L$$

However, from Pumping Lemma:  $uv^2 xy^2 z \in L$


**Contradiction!!!**

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:**  $v$  is in the first  $a^m$   
 $y$  is in the first  $b^m$

$$v = a^{k_1} \quad y = b^{k_2} \quad k_1 + k_2 \geq 1$$




$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:**  $v$  is in the first  $a^m$   
 $y$  is in the first  $b^m$

$$v = a^{k_1} \quad y = b^{k_2} \quad k_1 + k_2 \geq 1$$



$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:**  $v$  is in the first  $a^m$   
 $y$  is in the first  $b^m$

$$a^{m+k_1} b^{m+k_2} a^m b^m = uv^2 xy^2 z \notin L$$

$$k_1 + k_2 \geq 1$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 2:**  $v$  is in the first  $a^m$   
 $y$  is in the first  $b^m$

$$a^{m+k_1} b^{m+k_2} a^m b^m = uv^2 xy^2 z \notin L$$

However, from Pumping Lemma:  $uv^2 xy^2 z \in L$

**Contradiction!!!**




$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:**  $v$  overlaps the first  $a^m b^m$   
 $y$  is in the first  $b^m$

$$v = a^{k_1} b^{k_2} \quad y = b^{k_3} \quad k_1, k_2 \geq 1$$




$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:**  $v$  overlaps the first  $a^m b^m$   
 $y$  is in the first  $b^m$

$$v = a^{k_1} b^{k_2} \quad y = b^{k_3} \quad k_1, k_2 \geq 1$$



$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:**  $v$  overlaps the first  $a^m b^m$   
 $y$  is in the first  $b^m$

$$a^m b^{k_2} a^{k_1} b^{m+k_3} a^m b^m = uv^2 xy^2 z \notin L$$

$$k_1, k_2 \geq 1$$

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 3:**  $v$  overlaps the first  $a^m b^m$   
 $y$  is in the first  $b^m$

$$a^m b^{k_2} a^{k_1} b^{k_3} a^m b^m = uv^2 xy^2 z \notin L$$

However, from Pumping Lemma:  $uv^2 xy^2 z \in L$


**Contradiction!!!**

$$L = \{vv : v \in \{a,b\}^*\}$$

$$a^m b^m a^m b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

**Case 4:**  $v$  in the first  $a^m$   
 $y$  Overlaps the first  $a^m b^m$

Analysis is similar to case 3



**Other cases:**  $vxy$  is within  $a^m b^m a^m b^m$

or

$a^m b^m a^m b^m$

or

$a^m b^m a^m b^m$

Analysis is similar to case 1:

$a^m b^m a^m b^m$

More cases:  $vxy$  overlaps  $a^m \boxed{b^m} a^m b^m$

or

$a^m b^m \boxed{a^m} b^m$

Analysis is similar to cases 2,3,4:

$\boxed{a^m b^m} a^m b^m$

There are no other cases to consider

Since  $|vxy| \leq m$ , it is impossible

$vxy$  to overlap:

$a^m \boxed{b^m a^m} b^m$

nor

$a^m \boxed{b^m a^m b^m}$

nor

$\boxed{a^m b^m a^m} b^m$

In all cases we obtained a **contradiction**

**Therefore:** The original assumption that

$$L = \{vv : v \in \{a,b\}^*\}$$

is context-free must be wrong

**Conclusion:**  $L$  is not context-free

Non-context free languages

$$\{a^n b^n c^n : n \geq 0\}$$

$$\{ww : w \in \{a,b\}^*\}$$

$$\{a^{n!} : n \geq 0\}$$

Context-free languages

$$\{a^n b^n : n \geq 0\}$$

$$\{ww^R : w \in \{a,b\}^*\}$$

**Theorem:** The language

$$L = \{a^{n!} : n \geq 0\}$$

is **not** context free

**Proof:** Use the Pumping Lemma  
for context-free languages

$$L = \{a^{n!} : n \geq 0\}$$

Assume for **contradiction** that  $L$   
is context-free

Since  $L$  is context-free and infinite  
we can apply the pumping lemma

$$L = \{a^{n!} : n \geq 0\}$$

Pumping Lemma gives a magic number  $m$   
such that:

Pick any string of  $L$  with length at least  $m$

we pick:  $a^{m!} \in L$

$$L = \{a^{n!} : n \geq 0\}$$

We can write:  $a^{m!} = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

Pumping Lemma says:

$$uv^i xy^i z \in L \quad \text{for all } i \geq 0$$

$$L = \{a^{n!} : n \geq 0\}$$


$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

We examine all the possible locations of string  $vxy$  in  $a^{m!}$

There is only one case to consider

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$




$$v = a^{k_1} \quad y = a^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$



$$L = \{a^{n!} : n \geq 0\}$$


$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$



$$v = a^{k_1} \quad y = a^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$



$$k = k_1 + k_2$$

$$v = a^{k_1} \quad y = a^{k_2} \quad 1 \leq k \leq m$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

$$a^{m!+k} = uv^2xy^2z$$

$$1 \leq k \leq m$$


Since  $1 \leq k \leq m$ , for  $m \geq 2$  we have:

$$m!+k \leq m!+m$$

$$< m!+m!m$$

$$= m!(1+m)$$

$$= (m+1)!$$



$$m! < m!+k < (m+1)!$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

$$m! < m! + k < (m+1)!$$



$$a^{m!+k} = uv^2xy^2z \notin L$$

$$L = \{a^{n!} : n \geq 0\}$$

$$a^{m!} = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

However, from Pumping Lemma:  $uv^2xy^2z \in L$

$$a^{m!+k} = uv^2xy^2z \notin L$$

**Contradiction!!!**

We obtained a **contradiction**

**Therefore:** The original assumption that  
 $L = \{a^{n!} : n \geq 0\}$   
is context-free must be wrong

**Conclusion:**  $L$  is not context-free

## Non-context free languages

$$\{a^n b^n c^n : n \geq 0\} \quad \{ww : w \in \{a, b\}^*\}$$

$$\{a^{n^2} b^n : n \geq 0\} \quad \{a^{n!} : n \geq 0\}$$

## Context-free languages

$$\{a^n b^n : n \geq 0\} \quad \{ww^R : w \in \{a, b\}^*\}$$

**Theorem:** The language

$$L = \{a^{n^2} b^n : n \geq 0\}$$

is **not** context free

**Proof:** Use the Pumping Lemma  
for context-free languages

$$L = \{a^{n^2} b^n : n \geq 0\}$$

Assume for **contradiction** that  $L$   
is context-free

Since  $L$  is context-free and infinite  
we can apply the pumping lemma

$$L = \{a^{n^2} b^n : n \geq 0\}$$

Pumping Lemma gives a magic number  $m$   
such that:

Pick any string of  $L$  with length at least  $m$

we pick:  $a^{m^2} b^m \in L$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

We can write:  $a^{m^2} b^m = uvxyz$

with lengths  $|vxy| \leq m$  and  $|vy| \geq 1$

Pumping Lemma says:

$$uv^i xy^i z \in L \quad \text{for all } i \geq 0$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$


We examine all the possible locations

of string  $vxy$  in  $a^{m^2} b^m$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$


Most complicated case:  $v$  is in  $a^m$   
 $y$  is in  $b^m$



$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

$$v = a^{k_1} \quad y = b^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$




$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Most complicated sub-case:  $k_1 \neq 0$  and  $k_2 \neq 0$

$$v = a^{k_1} \quad y = b^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$






$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Most complicated sub-case:  $k_1 \neq 0$  and  $k_2 \neq 0$

$$v = a^{k_1} \quad y = b^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$



$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Most complicated sub-case:  $k_1 \neq 0$  and  $k_2 \neq 0$

$$v = a^{k_1} \quad y = b^{k_2} \quad 1 \leq k_1 + k_2 \leq m$$

$$a^{m^2-k_1} b^{m-k_2} = uv^0xy^0z$$

$$k_1 \neq 0 \text{ and } k_2 \neq 0 \quad 1 \leq k_1 + k_2 \leq m$$



$$(m - k_2)^2 \leq (m - 1)^2 \text{ (since } k_2 \geq 1)$$

$$= m^2 - 2m + 1$$

$$< m^2 - k_1 \text{ (since } k_1 \leq m)$$



$$m^2 - k_1 \neq (m - k_2)^2$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

$$m^2 - k_1 \neq (m - k_2)^2$$



$$a^{m^2 - k_1} b^{m - k_2} = uv^0 xy^0 z \notin L$$

$$L = \{a^{n^2} b^n : n \geq 0\}$$

$$a^{m^2} b^m = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

However, from Pumping Lemma:  $uv^0xy^0z \in L$

$$a^{m^2-k_1} b^{m-k_2} = uv^0xy^0z \notin L$$

Contradiction!!!

When we examine the rest of the cases  
we also obtain a contradiction

In all cases we obtained a **contradiction**

**Therefore:** The original assumption that

$$L = \{a^{n^2} b^n : n \geq 0\}$$

is context-free must be wrong

**Conclusion:**  $L$  is not context-free