


Ćwiczenie M7

Przyrodzy:

Fotokomórka

Wzrost AXIS



1. Zależność t^2 od h $m = 99,8 \pm 0,1 \text{ g}$

	$h=20\text{cm}$	$h=30\text{cm}$	$h=40\text{cm}$	$h=50\text{cm}$
	$t [s]$	$t [s]$	$t [s]$	$t [s]$
1.	3,782	5,040	5,883	6,577
2.	3,944	4,892	5,860	6,646
3.	3,838	4,904	5,850	6,616
4.	3,944	4,999	5,873	6,491
5.	3,890	4,833	5,868	6,586
6.	3,831	4,892	5,843	6,534
7.	3,308	4,997	5,885	6,506
8.	3,886	4,978	5,884	6,431
9.	3,888	4,929	5,911	6,400
10.	3,953	4,986	5,823	6,567

Zależność t^2 od d_w^2

masa odważników na nitce: $109,4 \pm 0,1 \text{ g}$

masa każdego z odważników na wahadle: $123,8 \pm 0,3 \text{ g}$

szerokość walcu: $10,0 \text{ mm}$ średnica walcu: $45,9 \text{ mm}$

Odległość krainców:	$99,9 \text{ mm}$	$178,5 \text{ mm}$	$258,6 \text{ mm}$	$339,0 \text{ mm}$
	$t [\text{s}]$	$t [\text{s}]$	$t [\text{s}]$	$t [\text{s}]$
1.	3,764	4,143	4,903	5,595
2.	3,653	4,072	4,887	5,493
3.	3,834	4,198	4,667	5,438
4.	3,664	4,112	4,793	5,647
5.	3,712	4,182	4,685	5,536
6.	3,745	4,281	4,751	5,461
7.	3,683	4,123	4,776	5,568
8.	3,739	4,050	4,818	5,769
9.	3,691	4,180	4,798	5,448
10.	3,650	4,250	4,773	5,568

Zależność t^2 od $\frac{1}{m}$

	$m = 99,8 \text{ g}$	$m = 119,7 \text{ g}$	$m = 139,6 \text{ g}$	$m = 159,6 \text{ g}$
	$t [\text{s}]$	$t [\text{s}]$	$t [\text{s}]$	$t [\text{s}]$
1.	5,040	4,472	4,255	3,925
2.	4,892	4,480	4,273	4,002
3.	4,904	4,500	4,218	3,848
4.	4,999	4,592	4,218	3,916
5.	4,833	4,413	4,237	3,835
6.	4,892	4,601	4,021	3,961
7.	4,997	4,582	4,176	3,903
8.	4,978	4,632	4,105	3,973
9.	4,929	4,572	4,269	3,857
10	4,986	4,492	4,227	3,982