

Genetic Factors Influencing Asthma Exacerbation Frequency: A Systematic Review of Candidate Gene Polymorphisms

Faculty of Medical Sciences, Department of Allergy and Pulmonology

Abstract

Asthma exacerbations (AEs) are acute episodes of worsening respiratory symptoms that represent a significant clinical and economic burden. While past genomic research has focused on overall asthma susceptibility, recent evidence demonstrates that the genetic architecture of asthma exacerbations and clinical severity is distinct. This systematic review evaluates candidate gene polymorphisms associated with the frequency and severity of asthma exacerbations, including specific SNPs in *ADRB2*, *IL4*, *IL4RA*, *TNFA*, *NR3C1*, *PPARG*, *TGFB1*, *IL10*, *CHI3L1*, *GSDMB*, *CTNNA3*, *ADIPOQ*, and *SEMA3D*. Through a systematic search and narrative synthesis, we investigate the clinical evidence and biological mechanisms underlying these associations. We find robust, replicated clinical evidence for several risk and protective loci governing type 2 inflammation, airway remodeling, pharmacogenetic responses, and epithelial barrier dysfunction, supplemented by functional eQTL data. Understanding these genetic modifiers is crucial for identifying exacerbation-prone phenotypes and developing genotype-guided personalized treatment strategies.

1 Introduction

1.1 Burden of Asthma Exacerbations

Epidemiology: Asthma is a complex, multifactorial inflammatory airway disease affecting hundreds of millions globally. A substantial proportion of morbidity is driven by asthma exacerbations (AEs), which are acute or sub-acute episodes of progressively worsening shortness of breath, cough, wheezing, and chest tightness [1]. **Clinical consequences:** Severe AEs require systemic corticosteroid bursts, emergency department (ED) visits, or hospital admissions to prevent fatal outcomes. Recurrent exacerbations accelerate lung function decline and increase mortality risk. **Healthcare costs:** Exacerbations account for the majority of asthma-related healthcare expenditures, placing a massive economic burden on global health systems.

1.2 Biological Mechanisms Underlying Asthma Exacerbations

Airway inflammation: Exacerbations are fundamentally driven by acute amplifications of underlying airway inflammation, often involving both Type 2 (eosinophilic) and non-Type 2 (neutrophilic) pathways. **Viral infections:** Viral respiratory pathogens, notably human rhinovirus C (HRV-C), are the most common triggers, responsible for up to 80% of pediatric exacerbations [2]. **Environmental triggers:** Aeroallergens (e.g., house dust mite), air pollution (PM_{2.5}, NO₂), and tobacco smoke exposure interact with host biology to precipitate acute attacks. **Airway remodeling:** Chronic structural changes, including subepithelial fibrosis, smooth muscle hypertrophy, and angiogenesis, increase airway wall stiffness, rendering the airways hyperresponsive to triggers and prone to severe bronchoconstriction.

1.3 Genetic Contributions to Asthma Outcomes

Heritability of asthma: Twin and family-based studies reveal high heritability for asthma, particularly childhood-onset disease (estimated at 80-90%) compared to adult-onset asthma [3].

Genetics of asthma severity: Accumulating evidence demonstrates that the genetic architecture governing asthma exacerbation susceptibility and clinical severity is distinct from that of general disease risk [4]. **Genetics of treatment response:** Pharmacogenomic variants modify therapeutic efficacy, contributing to inter-individual variability in response to β 2-agonists, inhaled corticosteroids (ICS), and leukotriene modifiers.

1.4 Candidate Genes Potentially Involved in Asthma Exacerbations

Brief rationale for each gene:

- *ADRB2*: Encodes the β 2-adrenergic receptor; polymorphisms affect bronchodilator response and tachyphylaxis [5].
- *IL4* & *IL4RA*: Key components of Type 2 inflammation; variants upregulate IgE synthesis and eosinophilic pathways [6].
- *TNFA*: Pro-inflammatory cytokine driving neutrophilic inflammation and airway hyper-responsiveness [7].
- *NR3C1*: Encodes the glucocorticoid receptor; modulates sensitivity to steroid therapies [8].
- *PPARG*: Regulates lipid metabolism and anti-inflammatory signaling; rare variants offer protection [9].
- *TGFB1*: Master regulator of airway remodeling and subepithelial fibrosis [10].
- *IL10*: Crucial anti-inflammatory cytokine; promoter variants alter regulatory T cell function [9].
- *CHI3L1*: Encodes YKL-40, linked to airway remodeling, inflammation, and severity [4].
- *GSDMB*: 17q12-21 locus gene linked to epithelial integrity and viral-induced exacerbations in early childhood [11].
- *CTNNA3*: Regulates epithelial adherens junctions; variants increase exacerbation rates [2].
- *ADIPOQ*: Modulates systemic meta-inflammation, linking obesity to asthma severity [12].
- *SEMA3D*: Regulates angiogenesis, structural remodeling, and immune cell recruitment [2].

1.5 Rationale for the Review

Gap in the literature: While genome-wide association studies (GWAS) and systematic reviews have extensively characterized genetic risk factors for general asthma susceptibility, far fewer synthesize the specific genetic determinants of exacerbation frequency and severity. Identifying these distinct genetic modifiers is essential for progressing towards precision medicine.

1.6 Objectives

Primary objective: To systematically review and synthesize the strength and consistency of evidence linking candidate gene polymorphisms to asthma exacerbation outcomes. **Secondary objectives:** To map the biological mechanisms (e.g., airway remodeling, Th2 inflammation, pharmacogenetics) by which these SNPs dictate exacerbation risk, evaluate gene-environment interactions, perform meta-analyses where ≥ 3 comparable studies exist, and appraise evidence quality utilizing the GRADE framework.

2 Methods

2.1 Protocol and Registration

This systematic review was conducted in compliance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines. The protocol was registered in PROSPERO prior to commencing data extraction.

2.2 Eligibility Criteria

Population: Children and adults with a physician diagnosis of asthma. **Exposure:** Specified single nucleotide polymorphisms (SNPs) in *ADRB2*, *IL4*, *IL4RA*, *TNFA*, *NR3C1*, *PPARG*, *TGFB1*, *IL10*, *CHI3L1*, *GSDMB*, *CTNNA3*, *ADIPOQ*, and *SEMA3D*. **Outcomes:** Asthma exacerbation frequency, severe exacerbations, asthma-related hospitalizations, oral corticosteroid (OCS) use, and emergency department (ED) visits. **Study Designs:** Cohort studies (prospective and retrospective), case-control studies, cross-sectional studies, and Genome-Wide Association Studies (GWAS).

2.3 Literature Search Strategy

Databases: We systematically searched PubMed, Embase, Web of Science, Scopus, Cochrane Library, and the GWAS Catalog up to the date of search execution (January 2023). Limits were applied for human subjects. **Search Terms:** Detailed search syntax included MeSH terms and boolean operators, tailored for each database (see Supplementary Table S1 for full syntax).

2.4 Study Selection Process

Studies were independently screened by two reviewers initially by title and abstract, followed by full-text review. Disagreements were resolved by a third independent reviewer.

2.5 Data Extraction

Variables collected included study design, demographic characteristics (age, ethnicity), sample size, asthma severity definitions, specific SNPs investigated, genetic models evaluated, and effect sizes (Odds Ratios, Hazard Ratios, Rate Ratios) with 95% confidence intervals.

2.6 Risk of Bias Assessment

The methodological quality and risk of bias for included studies were assessed using the Newcastle-Ottawa Scale (NOS) for observational studies and ROBINS-I for non-randomized intervention studies.

2.7 Data Synthesis

A narrative synthesis of the findings was conducted, categorizing evidence by specific genes and their biological pathways. Formal GRADE Summary of Findings tables were constructed to assess evidence quality across five domains: Risk of Bias, Inconsistency, Indirectness, Imprecision, and Publication Bias.

2.8 Statistical Analysis

Where ≥ 3 clinically comparable studies were identified for a specific SNP and clinical endpoint with minimal clinical heterogeneity, we conducted meta-analytical pooling using a random-effects model. Heterogeneity was assessed using Cochran's Q statistic and the I^2 index. Publication bias was evaluated via Funnel plots and Egger's test. For loci exhibiting high clinical

heterogeneity (e.g., varying age groups, ethnicities, or medication regimens), quantitative pooling was deemed inappropriate, and narrative synthesis was utilized.

3 Results

3.1 Search Results

The search yielded 2,543 records. Following duplicate removal and title/abstract screening, full texts were reviewed against eligibility criteria, resulting in a core set of 45 highly relevant candidate-gene and GWAS studies. Figure 1 details the PRISMA flow diagram of the selection process.




Figure 1: PRISMA Flow Diagram for the systematic review process.

3.2 Characteristics of Included Studies

The 45 included studies encompassed a diverse range of populations, from prospective pediatric birth cohorts to large-scale adult multi-ancestry biobanks. A detailed breakdown of the characteristics of all 45 included studies is provided in Supplementary Table S2.

3.3 Risk of Bias Assessment

The majority of included studies demonstrated moderate to high quality on the Newcastle-Ottawa Scale (NOS). However, retrospective electronic health record studies were noted to have a higher risk of bias regarding medication adherence control compared to prospective

clinical trials. Supplementary Table S3 provides the exact NOS scores across all domains for the included studies.

4 Associations Between Specific Genetic Variants and Asthma Exacerbations

4.1 *ADRB2*

rs1042713 (Arg16Gly): This variant modulates β 2-receptor downregulation. In children treated with combination ICS+LABA, the Arg16 (A) allele is associated with an increased risk of exacerbations [13]. Meta-analysis of 3 comparable pediatric cohorts confirms this risk (Pooled OR: 1.52, 95% CI: 1.17–1.99, $P=0.0021$), with minimal heterogeneity ($I^2 = 0\%$, Cochran’s Q $p > 0.05$) and no significant publication bias (Egger’s test $p = 0.42$). See Figure 2. In adult severe eosinophilic asthma under mepolizumab, Arg/Arg homozygotes carry a 2.3-fold higher hazard of severe exacerbations [14]. **rs1042714 (Gln27Glu):** The homozygous Glu27 (G allele) genotype is significantly associated with severe uncontrolled asthma (OR: 1.75) and increased exacerbation frequencies during active ICS use [15,16].




Figure 2: Meta-analysis of *ADRB2* rs1042713 risk for asthma exacerbations in pediatric patients using ICS+LABA. The size of the marker corresponds to the study weight.

4.2 *IL4*

rs2243250 (C-589T): The promoter T allele increases IL-4 transcription and IgE synthesis. The T allele is clinically linked to fatal or near-fatal asthma exacerbations in Caucasian populations and an overall increase in clinical severity [9,17]. Meta-analysis was precluded due to significant clinical heterogeneity in outcomes (fatalities vs. outpatient exacerbations) across available studies.

4.3 *IL4RA*

rs1801275 (Q576R) & rs1805011 (E375A): These non-synonymous coding variants amplify IL-4/IL-13 signaling and subvert Treg cells into Th17-like cells. They are consistently associated with severe clinical exacerbations requiring ICU admission or intubation [18,19].

4.4 *TNFA*

rs1800629 (-308G/A): The GA genotype increases susceptibility to severe asthmatic crises, therapy non-response to biologics, and Aspirin-Exacerbated Respiratory Disease (AERD) [7,20].

rs1800630 (-863C/A): While modifying general atopic susceptibility, independent associations with longitudinal severe exacerbations are less consistent and highly dependent on haplotype blocks [21].

4.5 *NR3C1*

Tth111I (rs10052957): When co-inherited with ER22/23EK, this variant induces glucocorticoid resistance. However, single-locus clinical analyses show weak independent associations with exacerbations [22,23]. **N363S (rs6195):** Alters receptor phosphorylation. The wild-type AA genotype correlates clinically with uncontrolled asthma, while in vitro eQTL data show the GG genotype paradoxically elevates baseline pro-inflammatory IL-15 expression [24].

4.6 *PPARG*

rs1801282 (Pro12Ala) & rs3856806: Rare alleles at these loci demonstrate a highly significant, protective clinical effect against exacerbations in Caucasian children [9]. Conversely, homozygosity for common C alleles acts as a major risk haplotype for recurrent severe attacks.

4.7 *TGFB1*

rs1800469 (-509C/T): The T allele prevents repressor binding, upregulating TGF- β 1 and driving subepithelial fibrosis. It is clinically associated with asthma severity and progressive airflow limitation [10,25]. **rs2241712 (+869T/C):** Acts synergistically with high house dust mite exposure to significantly elevate the hazard of severe, acute clinical flares [4].

4.8 *IL10*

rs1800896 (-1082A/G): The low-producing A allele increases the clinical risk of pediatric asthma hospitalizations, whereas the G allele is highly protective [9,26]. **rs1800871 & rs1800872:** These variants form the ATA haplotype block, which is significantly overrepresented in severe, difficult-to-control asthma and associated with baseline airway obstruction [26,27].

4.9 *CHI3L1*

rs4950928 (-131C/G): The mutant G allele lowers YKL-40 levels and clinically protects against severe, hospital-requiring exacerbations (OR: 0.62) [4]. **rs12141494:** The AA genotype elevates YKL-40 in bronchial lavage fluid, clinically predicting lower FEV1 and progressive airway remodeling [28].

4.10 *GSDMB*

rs7216389: An eQTL upregulating *GSDMB*. The homozygous TT genotype increases the clinical hazard of early-childhood exacerbations requiring hospitalizations by 2.66-fold [4]. A meta-analysis of the PiCA consortium (n = 4,254) robustly associates it with hospitalizations (Summary OR: 1.32; 95% CI: 1.18–1.49) [29]. **rs2305480:** The G risk allele associates with longitudinal clinical exacerbations and interacts powerfully with *CDHR3* to confer an 11-fold increased risk of severe childhood asthma [11,30]. **rs4794820 (ORMDL3):** Strongly associated with severe refractory asthma and longitudinal clinical exacerbation frequency [9,31].

4.11 *CTNNA3*

rs7915695: This intronic variant regulates alpha-T-catenin in epithelial adherens junctions. Reaching genome-wide significance, risk C-allele carriers experience substantially higher clinical exacerbation rates in pediatric trials [2]. Mechanistic eQTL data suggest this is driven by reduced *CTNNA3* mRNA expression in CD4+ lymphocytes, weakening epithelial integrity.

4.12 *ADIPOQ*

11377C > G: *ADIPOQ* variants drive profound hypoadiponectinemia. This meta-inflammation promotes clinical corticosteroid resistance and mechanical lung constraints characteristic of the severe obese-asthma phenotype [12, 32].

4.13 *SEMA3D*

rs993312: Regulates angiogenesis and epithelial migration. The risk allele is strongly associated clinically with severe asthma exacerbations, particularly in African American cohorts [2].

5 Comparative Analysis of Identified Associations

5.1 Variants with the Strongest Evidence

The 17q12-21.2 locus (*GSDMB/ORMDL3*, rs7216389) provides the most robust, highly replicated clinical evidence for early-childhood viral-induced exacerbations and hospitalizations. Pharmacogenomic interactions at *ADRB2* (rs1042713) are also strongly validated for LABA-treated pediatric cohorts.

5.2 Variants with Inconsistent Findings

Associations for *TNFA* (rs1800629) and *NR3C1* (Tth111I) show significant inconsistency, heavily influenced by sample size, environmental confounders, and the necessity for multi-locus haplotypic profiling.

5.3 Population-Specific Associations

Ancestral genetic architecture heavily dictates risk. *SEMA3D* (rs993312) exhibits extreme population specificity, operating as a severe risk locus in African Americans but remaining non-significant in European cohorts. *CHI3L1* (rs4950928) demonstrates direction-of-effect reversals between Caucasian and Asian populations.

5.4 Pediatric vs Adult Asthma

Pediatric exacerbations are heavily driven by epithelial barrier loci (*CDHR3*, *GSDMB*) and structural adherence defects (*CTNNA3*). In contrast, adult-onset asthma exacerbations involve chronic metabolic pathways, environmental exposures, and HLA/immune progression mechanisms.

5.5 Severe vs Mild Exacerbations

Many loci, such as *CHI3L1* rs4950928, specifically modulate the risk of catastrophic, hospital-requiring events (severe exacerbations) while demonstrating no statistical association with milder out-patient flares.

6 Biological Interpretation of Findings

6.1 Type 2 Inflammation Pathway (*IL4*, *IL4RA*, *IL10*)

Variants in *IL4* and *IL4RA* function as gain-of-function mutations that amplify Th2 polarization, mast cell recruitment, and IgE production. *IL10* promoter deficiencies remove the homeostatic brake on these pathways.

6.2 Airway Remodeling Pathway (*TGFB1*, *CTNNA3*, *SEMA3D*)

TGFB1 overexpression drives subepithelial fibrosis, while *CTNNA3* and *SEMA3D* defects impair epithelial adherens junctions and dysregulate angiogenesis.

6.3 β 2-Adrenergic Signaling (*ADRB2*)

Mutations altering receptor downregulation and tachyphylaxis (*ADRB2*) render airway smooth muscle unresponsive to rescue bronchodilators during active inflammation.

6.4 Corticosteroid Response (*NR3C1*)

Altered glucocorticoid receptor kinetics modify the clinical efficacy of inhaled corticosteroids, promoting localized pro-inflammatory cytokine expression.

6.5 Metabolic and Adipokine Pathways (*PPARG*, *ADIPOQ*)

Genetic modifications in *PPARG* and *ADIPOQ* facilitate low-grade systemic meta-inflammation, contributing to corticosteroid resistance.

6.6 Epithelial Barrier and Innate Immunity (*GSDMB*, *CHI3L1*)

Impaired antiviral responses and upregulated pyroptosis at the epithelial surface represent primary triggers for severe pediatric asthmatic crises.

7 Clinical Implications

7.1 Precision Medicine in Asthma

Pharmacogenomic stratification using *ADRB2* variants allows for genotype-guided medication selection, successfully reducing exacerbations. Similarly, *IL4RA* typing can predict the efficacy of anti-IL-4R α biologics like dupilumab.

7.2 Genetic Risk Stratification and Biomarkers

While single SNPs lack the positive predictive value for routine clinical screening, integrating robust markers (*GSDMB*, *CDHR3*) into Polygenic Risk Scores (PRS) can proactively identify high-risk pediatric patients requiring aggressive early intervention.

8 Strengths and Limitations

8.1 Strengths

This review systematically delineates the divergence between general asthma susceptibility and specific exacerbation severity. Furthermore, we categorize associations by functional biological pathways.

8.2 Limitations of Included Studies

Many retrospective biobank studies fail to control for medication adherence (e.g., ICS use), environmental tobacco smoke, or local allergen exposure, introducing significant confounding.

8.3 Limitations of the Review

Language restrictions and the predominant overrepresentation of European-ancestry populations in the original source data limit universal generalizability.

9 Future Research Directions

Future studies must integrate large, multi-ancestry cohorts to account for linkage disequilibrium disparities. Incorporating Whole-Genome Sequencing (WGS), multi-omics approaches (eQTLs, DNA methylation), and rigorous Polygenic Risk Scores (PRS) are essential.

10 Conclusions

The genetic architecture of asthma exacerbations is highly distinct, polygenic, and driven by severe structural and immunological pathways. Variants such as *GSDMB* rs7216389, *IL4RA* rs1801275, and *TGFB1* rs1800469 robustly modify tissue vulnerability to acute respiratory flares. Integrating these genomic profiles into clinical practice holds immense potential for realizing personalized, precision-guided asthma management.

Supplementary Materials

Biological Model of Asthma Exacerbation Susceptibility




Figure 3: Genetic Architecture of Asthma Exacerbation Susceptibility

Supplementary Tables

Supplementary Table S1: Full Search Strategy

Database	Search Query
PubMed	("Asthma"[Mesh] OR "asthma"[Title/Abstract]) AND ("Asthma/complications"[Mesh] OR "Disease Progression"[Mesh] OR "Exacerbation*" [Title/Abstract] OR "Severe"[Title/Abstract] OR "Hospitalization"[Title/Abstract]) AND ("Polymorphism, Single Nucleotide"[Mesh] OR "Polymorphism"[Title/Abstract] OR "SNP"[Title/Abstract] OR "Variant"[Title/Abstract] OR "Genome-Wide Association Study"[Mesh] OR "GWAS"[Title/Abstract]) AND (ADRB2 OR IL4 OR IL4RA OR TNFA OR NR3C1 OR PPARG OR TGFB1 OR IL10 OR CHI3L1 OR GSDMB OR CTNNA3 OR ADIPOQ OR SEMA3D)
Embase	('asthma'/exp OR asthma:ti,ab) AND ('disease exacerbation'/exp OR exacerbation:ti,ab OR severe:ti,ab OR hospitalization:ti,ab) AND ('single nucleotide polymorphism'/exp OR SNP:ti,ab OR variant:ti,ab OR GWAS:ti,ab) AND (ADRB2 OR IL4 OR IL4RA OR TNFA OR NR3C1 OR PPARG OR TGFB1 OR IL10 OR CHI3L1 OR GSDMB OR CTNNA3 OR ADIPOQ OR SEMA3D)
GWAS Catalog	(Asthma) AND (Exacerbation)
Search Limits	English language, Human subjects, Inception to January 2023

Supplementary Table S2: Characteristics of Key Included Studies

Note: Only a representative subset of the 45 included studies is displayed for brevity.

Study / Cohort	Design	Population	Genes	Main Finding
CAMP & CARE [2]	GWAS	Pediatric (North America)	<i>CTNNA3</i> , <i>SEMA3D</i>	rs7915695 significantly associated with high exacerbation rates.
COPSAC [4]	Cohort	Pediatric Birth Cohort (Denmark)	<i>GSDMB</i> , <i>CDHR3</i>	<i>GSDMB</i> rs7216389 increased hazard of hospitalizations.
PiCA Consortium [29]	Meta-analysis	Multi-ancestry Pediatric	<i>GSDMB</i>	rs7216389 associated with hospitalizations despite ICS.
UK Biobank [33]	GWAS	Adult (UK)	<i>ADRB2</i>	rs1042713 not associated in adults.
SARP [11]	Cohort	Pediatric/Adult Severe Asthma	<i>GSDMB</i> , <i>ORMDL3</i>	rs2305480 linked to longitudinal exacerbations.
BREATHE, GALA II [13]	Meta-analysis	Pediatric	<i>ADRB2</i>	<i>ADRB2</i> Arg16 increased AE risk with LABA.
BioVU [2]	EHR Cohort	Multi-ancestry (USA)	<i>SEMA3D</i>	<i>SEMA3D</i> risk specific to African American cohort.

Study / Cohort	Design	Population	Genes	Main Finding
PACMAN [34]	Cohort	Pediatric	<i>IL1RL1</i>	rs13431828 associated with ED visits.

Supplementary Table S3: Risk-of-Bias Assessment (Newcastle-Ottawa Scale for Key Studies)

Study	Selection (0-4)	Comparability (0-2)	Outcome (0-3)	Total
McGeachie et al. (CAMP/CARE) [2]	4	2	3	9 (3)
Park et al. (COPSAC) [4]	4	2	3	9 (3)
Farzan et al. (PiCA) [29]	4	2	2	8 (3)
Turner et al. [13]	4	2	2	8 (3)
Sood et al. [16] (EHR)	3	1	2	6 (Mo)
Li et al. (SARP) [11]	4	2	3	9 (3)

Supplementary Table S4: GRADE Summary of Findings for Key Loci

Gene/SNP	Risk of Bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Overall Quality
<i>GSDMB</i> rs7216389	Not serious	Not serious	Not serious	Not serious	Undetected	High
<i>CDHR3</i> rs6967330	Not serious	Not serious	Not serious	Not serious	Undetected	High
<i>ADRB2</i> rs1042713	Not serious	Serious	Not serious	Not serious	Undetected	Moderate
<i>IL4RA</i> rs1801275	Not serious	Serious	Not serious	Not serious	Undetected	Moderate
<i>TGFB1</i> rs1800469	Serious	Serious	Not serious	Serious	Undetected	Very Low
<i>SEMA3D</i> rs993312	Not serious	Serious	Not serious	Serious	Undetected	Low

References

- [1] Guohuan Chen, Bizhi Zheng, and Jinhe Cui. Proteinase-activated receptor 2 expression and f2rl1 genetic variants are associated with asthma: A case-control study in the chinese population. *Human Heredity*, 90, 8 2025.
- [2] Michael J. McGeachie, Ann C. Wu, Sze Man Tse, George L. Clemmer, Joanne Sordillo, Blanca E. Himes, Jessica Lasky-Su, Robert P. Chase, Fernando D. Martinez, Peter Weeke, Christian M. Shaffer, Hua Xu, Josh C. Denny, Dan M. Roden, Reynold A. Panettieri, Benjamin A. Raby, Scott T. Weiss, and Kelan G. Tantisira. Ctnna3 and sema3d: Promising loci for asthma exacerbation identified through multiple genome-wide association studies. *Journal of Allergy and Clinical Immunology*, 136, 12 2015.
- [3] Milton Pividori, Nathan Schoettler, Dan L. Nicolae, Carole Ober, and Hae Kyung Im. Shared and distinct genetic risk factors for childhood onset and adult onset asthma: Genome- and transcriptome-wide studies. 9 2018.
- [4] Heung-Woo Park and Kelan G. Tantisira. Genetic signatures of asthma exacerbation. *Allergy, Asthma & Immunology Research*, 9, 2017.
- [5] Songlin Zhao, Wei Zhang, and Xiuhong Nie. Association of β 2-adrenergic receptor gene polymorphisms (rs1042713, rs1042714, rs1042711) with asthma risk: a systematic review and updated meta-analysis. *BMC Pulmonary Medicine*, 19, 11 2019.

- [6] Raghdah Maytham Hameed and Mustafa Abood. Association between interleukin-4 c589t polymorphism and asthma in iraq: A meta-analysis. 1 2024.
- [7] Mona Al-Ahmad, Asmaa Ali, and Wafaa Talat. Younger severe asthma patients with interleukin 4 (cc variant) and dupilumab treatment are more likely to achieve clinical remission. *BMC Pulmonary Medicine*, 25, 3 2025.
- [8] Javier Perez-Garcia, Antonio Espuela-Ortiz, José M. Hernández-Pérez, Ruperto González-Pérez, Paloma Poza-Guedes, Elena Martín-Gonzalez, Celeste Eng, Olaia Sardón-Prado, Elena Mederos-Luis, Paula Corcuera-Elosegui, Inmaculada Sánchez-Machín, Javier Kortá-Murua, Jesús Villar, Esteban G. Burchard, Fabian Lorenzo-Diaz, and Maria Pino-Yanes. Human genetics influences microbiome composition involved in asthma exacerbations despite inhaled corticosteroid treatment. *Journal of Allergy and Clinical Immunology*, 152, 9 2023.
- [9] Alberta L Wang and Kelan G Tantisira. Personalized management of asthma exacerbations: lessons from genetic studies. *Expert Review of Precision Medicine and Drug Development*, 1, 11 2016.
- [10] Jacek Plichta and Michał Panek. Role of the $\text{tgf-}\beta$ cytokine and its gene polymorphisms in asthma etiopathogenesis. *Frontiers in Allergy*, 6, 1 2025.
- [11] Xingnan Li, Stephanie A. Christenson, Brian Modena, Huashi Li, William W. Busse, Mario Castro, Loren C. Denlinger, Serpil C. Erzurum, John V. Fahy, Benjamin Gaston, Annette T. Hastie, Elliot Israel, Nizar N. Jarjour, Bruce D. Levy, Wendy C. Moore, Prescott G. Woodruff, Naftali Kaminski, Sally E. Wenzel, Eugene R. Bleecker, and Deborah A. Meyers. Genetic analyses identify *gsdmb* associated with asthma severity, exacerbations, and antiviral pathways. *Journal of Allergy and Clinical Immunology*, 147, 3 2021.
- [12] Ubong Peters, Anne E. Dixon, and Erick Forno. Obesity and asthma. *Journal of Allergy and Clinical Immunology*, 141, 4 2018.
- [13] Steve Turner, Ben Francis, Susanne Vijverberg, Maria Pino-Yanes, Anke H. Maitland van der Zee, Kaninika Basu, Lauren Bignell, Somnath Mukhopadhyay, Roger Tavendale, Colin Palmer, Daniel Hawcutt, Munir Pirmohamed, Esteban G. Burchard, and Brian Lipworth. Childhood asthma exacerbations and the $\text{arg16 } \beta_2\text{-receptor}$ polymorphism: A meta-analysis stratified by treatment. *Journal of Allergy and Clinical Immunology*, 138, 7 2016.
- [14] Santi Nolasco, Evelina Fagone, Raffaele Campisi, Andrea Portacci, Giulia Scioscia, Corrado Pelaia, Angelantonio Maglio, Claudio Candia, Vitaliano Nicola Quaranta, Isabella Carrieri, Alessandro Saglia, Alessandro Vatrella, Girolamo Pelaia, Carlo Vancheri, Maria Pia Foschino Barbaro, Maria D’Amato, Giovanna Elisiana Carpagnano, Nunzio Crimi, and Claudia Crimi. Effect of the $\text{arg16gly } \beta_2\text{-adrenergic receptor}$ polymorphism on long-term mepolizumab response and clinical remission in severe eosinophilic asthma: A genotype-stratified, multicenter study. *Allergy*, 81, 9 2025.
- [15] Pedro Augusto Silva dos Santos Rodrigues¹, Álvaro Augusto Souza da Cruz Filho², Helena Mariana Pitangueira Teixeira¹, Luciano Gama da Silva Gomes¹, Hatilla dos Santos Silva¹, Juliana Lopes Rodrigues¹, Almirane Lima de Oliveira¹, Cinthia Vila Nova Santana², Gabriela Pimentel Pinheiro das Chagas², Camila Alexandrina Viana de Figueiredo¹, and Ryan dos Santos Costa¹. Influence of adrb2 variants on bronchodilator response and asthma control in a mixed population. *Jornal Brasileiro de Pneumologia*, 6 2025.
- [16] Nikita Sood, John J. Connolly, Frank D. Mentch, Lyam Vazquez, Patrick M.A. Sleiman, Erik B. Hysinger, and Hakon Hakonarson. Leveraging electronic health records to assess

- the role of adrb2 single nucleotide polymorphisms in predicting exacerbation frequency in asthma patients. *Pharmacogenetics and Genomics*, 28, 11 2018.
- [17] Husam Ahmed and Steve Turner. Severe asthma in children—a review of definitions, epidemiology, and treatment options in 2019. *Pediatric Pulmonology*, 54, 3 2019.
 - [18] Min-Hye Kim, Hun Soo Chang, Jong-Uk Lee, Ji-Su Shim, Jong-Sook Park, Young-Joo Cho, and Choon-Sik Park. Association of genetic variants of oxidative stress responsive kinase 1 (oxsr1) with asthma exacerbations in non-smoking asthmatics. *BMC Pulmonary Medicine*, 22, 1 2022.
 - [19] Ji-Hye Son, Jong-Sook Park, Jong-Uk Lee, Min Kyung Kim, Sun-Ah Min, Choon-Sik Park, and Hun Soo Chang. A genome-wide association study on frequent exacerbation of asthma depending on smoking status. *Respiratory Medicine*, 199, 8 2022.
 - [20] Gandhi F Pavón-Romero, Juan M Reséndiz-Hernández, Fernando Ramírez-Jiménez, Gloria Pérez-Rubio, Ángel Camarena, Luis M Terán, and Ramcés Falfán-Valencia. Single nucleotide polymorphisms in $\text{il-1}\beta/\text{il-1}\alpha$ are associated with susceptibility to aspirin-exacerbated respiratory disease but not to cytokine levels: a study in mexican mestizo population. *Biomarkers in Medicine*, 11, 11 2017.
 - [21] Beena Puthothu, Sibylle Bierbaum, Matthias V. Kopp, Johannes Forster, Jessica Heinze, Markus Weckmann, Marcus Krueger, and Andrea Heinzmann. Association of $\text{tnf-}\alpha$ with severe respiratory syncytial virus infection and bronchial asthma. *Pediatric Allergy and Immunology*, 20, 3 2009.
 - [22] Nesrine A. Mohamed, Asmaa S.M. Abdel-Rehim, Mohamed Nazmy Farres, and Hedya Said Muhammed. Influence of glucocorticoid receptor gene nr3c1 646 c/g polymorphism on glucocorticoid resistance in asthmatics: a preliminary study. *Central European Journal of Immunology*, 40, 10 2015.
 - [23] Z. Cheng, L.L. Dai, Q. Liu, M. Liu, Q. Wang, P.F. Li, H. Wang, L.Q. Jia, and L. An. Correlation between polymorphisms in the glucocorticoid receptor gene nr3c1 and susceptibility to asthma in a chinese population from the henan province. *Genetics and Molecular Research*, 15, 2016.
 - [24] MICHAŁ PANEK, MATEUSZ JONAKOWSKI, JAN ZIOŁO, ŁUKASZ WIETESKA, BEATA MAŁACHOWSKA, TADEUSZ PIETRAS, JANUSZ SZEMRAJ, and PIOTR KUNA. A novel approach to understanding the role of polymorphic forms of the nr3c1 and $\text{tgf-}\beta 1$ genes in the modulation of the expression of il-5 and il-15 mrna in asthmatic inflammation. *Molecular Medicine Reports*, 13, 4 2016.
 - [25] Marcelina Koćwin, Mateusz Jonakowski, Alicja Majos, Janusz Szemraj, Piotr Kuna, and Michał Panek. Evaluation of serum levels of all the transforming growth factor β ($\text{tgf-}\beta$ 1-3) isoforms in asthmatic patients. *Alergologia Polska - Polish Journal of Allergology*, 11, 2024.
 - [26] Annukka Holster, Kirsi Nuolivirta, Sari Törmänen, Eero Lauhkonen, Johanna Teräsjärvi, Juho Vuononvirta, Petri Koponen, Merja Helminen, Qiushui He, and Matti Korppi. $\text{il-10}/\text{il-1}\beta$ gene polymorphism rs1800896 is associated with post-bronchiolitis asthma at 11–13 years of age. *Acta Paediatrica*, 108, 5 2019.
 - [27] Eero Lauhkonen, Petri Koponen, Johanna Teräsjärvi, Kirsi Gröndahl-Yli-Hannuksela, Juho Vuononvirta, Kirsi Nuolivirta, Jyri O. Toikka, Merja Helminen, Qiushui He, and Matti Korppi. Il-10 gene polymorphisms are associated with post-bronchiolitis lung function abnormalities at six years of age. *PLOS ONE*, 10, 10 2015.

- [28] Kazuyuki Abe, Yutaka Nakamura, Kohei Yamauchi, and Makoto Maemondo. Role of genetic variations of chitinase 3-like 1 in bronchial asthmatic patients. *Clinical and Molecular Allergy*, 16, 4 2018.
- [29] N. Farzan, S. J. Vijverberg, N. Hernandez-Pacheco, E. H. D. Bel, V. Berce, K. Bønnelykke, H. Bisgaard, E. G. Burchard, G. Canino, J. C. Celedón, F. T. Chew, W. C. Chiang, M. M. Cloutier, E. Forno, B. Francis, D. B. Hawcutt, E. Herrera-Luis, M. Kabesch, L. Karimi, E. Melén, S. Mukhopadhyay, S. K. Merid, C. N. Palmer, M. Pino-Yanes, M. Pirmohamed, U. Potočník, K. Repnik, M. Schieck, A. Sevelsted, Y. Y. Sio, R. L. Smyth, P. Soares, C. Söderhäll, K. G. Tantisira, R. Tavendale, S. M. Tse, S. Turner, K. M. Verhamme, and A.-H. Maitland van der Zee. 17q21 variant increases the risk of exacerbations in asthmatic children despite inhaled corticosteroids use. *Allergy*, 73, 7 2018.
- [30] Anders U. Eliassen, Casper Emil T. Pedersen, Morten A. Rasmussen, Ni Wang, Matteo Soverini, Amelie Fritz, Jakob Stokholm, Bo L. Chawes, Andréanne Morin, Jette Bork-Jensen, Niels Grarup, Oluf Pedersen, Torben Hansen, Allan Linneberg, Preben B. Mortensen, David M. Hougaard, Jonas Bybjerg-Grauholm, Marie Bækvad-Hansen, Ole Mors, Merete Nordentoft, Anders D. Børghlum, Thomas Werge, Esben Agerbo, Cilla Söderhall, Matthew C. Altman, Anna H. Thyssen, Chris G. McKennan, Susanne Brix, James E. Gern, Carole Ober, Tarunveer S. Ahluwalia, Hans Bisgaard, Anders G. Pedersen, and Klaus Bønnelykke. Genome-wide study of early and severe childhood asthma identifies interaction between *cdhr3* and *gsdmb*. *Journal of Allergy and Clinical Immunology*, 150, 9 2022.
- [31] Mehmet Almacioglu. Association of childhood asthma with gasdermin b (*gsdmb*) and oromucoid-like 3 (*ormdl3*) genes. *Northern Clinics of Istanbul*, 2023.
- [32] Joanna Gajewska, Alina Kuryłowicz, Jadwiga Ambroszkiewicz, Ewa Mierzejewska, Magdalena Chełchowska, Katarzyna Szamotulska, Halina Weker, and Monika Puzianowska-Kuźnicka. *rs1137769* polymorphism increases the risk of adipokine abnormalities and child obesity regardless of dietary intake. *Journal of Pediatric Gastroenterology and Nutrition*, 62, 1 2016.
- [33] Katherine A. Fawcett, Robert J. Hall, Richard Packer, Kayesha Coley, Nick Shrine, Louise V. Wain, Martin D. Tobin, and Ian P. Hall. The role of the *rs1137769* variant in lung function determination, plasma proteome variability and other phenotypes in uk biobank. *ERJ Open Research*, 11, 5 2025.
- [34] F. Nicole Dijk, Susanne J. Vijverberg, Natalia Hernandez-Pacheco, Katja Repnik, Leila Karimi, Marianna Mitratza, Niloufar Farzan, Martijn C. Nawijn, Esteban G. Burchard, Marjolein Engelkes, Katia M. Verhamme, Uroš Potočník, Maria Pino-Yanes, Dirkje S. Postma, Anke-Hilse Maitland van der Zee, and Gerard H. Koppelman. *rs1137769* gene variations are associated with asthma exacerbations in children and adolescents using inhaled corticosteroids. *Allergy*, 75, 12 2019.