Competitive Persuasion with a Noisy Judge

Igor Karpov*

November 3, 2026

Click here for the most recent version

Abstract

Can a firm choosing between two agents gain from judging them with error? I show that she can when the agents design their own evidence. Noiseless judging makes even a narrow lead decisive, so agents conceal bad news not to lose; noise makes a narrow lead worth little, so revealing pays more. For a broad class of noise and binary types, the equilibrium is unique: each agent sometimes admits being bad and otherwise pools, and if agents are likely good, both reveal fully, delivering the first best, unlike under noiseless judging. Moderate noise is best: it curbs concealment without reversing the ranking of the signals sent, often beating a noiseless rule. With a continuum of types, the case for noise survives: under regularity conditions, the firm prefers to judge with error unless strong types are at least as common as weak ones, and when weak types are more common, her best rule tosses a coin only among weak signals.

Keywords: Bayesian persuasion, multiple senders, competition in persuasion, full disclosure, noisy evaluation

JEL Classification: C72, D82, D83

*University of Chicago. I am very grateful to Doron Ravid, Ben Brooks and Emir Kamenica for their guidance and support. I also thank Alex Frankel, Joe Root, Phil Reny and the student workshop audience at the University of Chicago for useful comments and discussion. All errors are my own.

1. Introduction

A capital committee must fund one of two projects this quarter. Each sponsor decides how rigorous a pilot to run before the meeting, knowing that the committee's verdict will turn partly on things that have nothing to do with the projects: who presents second, how long the meeting has run. The committee would like to see past all of that. This paper asks whether it should want to.

Such imperfect verdicts, biased or merely arbitrary, are well recognized. In Scharfstein and Stein (2000), divisional rent-seeking leads to "socialist" internal capital markets, in which weaker divisions are subsidized by stronger ones. In Prendergast and Topel (1996), evaluators who lack objective performance measures act on personal preferences, favoring some subordinates beyond their true performance. And Pier et al. (2018) find essentially no agreement among experienced reviewers rating the same NIH grant applications, all of which were eventually funded: ratings of the same application were no more similar than ratings of different ones. The usual response is to engineer such imperfections away, through more reviewers, more structured evaluations, and more objective criteria. A few research funders have nonetheless moved the other way: they now fund by lot among proposals that clear a quality bar, citing reviewer bias, reviewer burden, and the hope of bolder proposals (Fang and Casadevall, 2016; Heyard et al., 2022; Röbbecke and Simon, 2023). Only the hope of bolder proposals concerns how applicants respond, and none concerns what the competitors choose to reveal about themselves, the channel I study.

The question is whether a decision-maker should eliminate all noise from her evaluation when the competitors know that it is noisy and adapt their reports to it. It arises whenever three features come together. First, a single prize is at stake, with no price attached: the decision-maker cannot pay more for the better competitor, so all she controls is whom she selects. Second, the competitors decide what and how much

to reveal about themselves after learning how the decision-maker will choose between them. Third, the decision-maker can make the verdict more or less accurate, so the competitor with the stronger evidence need not always win. I call the decision-maker *the firm*, a noisy judge when her verdict carries error, and the competitors *agents*. In the baseline, each agent is either a good or a bad type; Section 5 allows a continuum of types. Each agent simultaneously commits, at no cost, to a signal, that is, to a distribution of his posterior expected type; the firm observes the resulting posterior with a random error and selects the agent with the higher score. For many standard noise distributions and equal priors, this is also what she would do on seeing only the scores, so committing to the rule costs her nothing there.

My key observation is that noise of a suitable shape weakens each agent's incentive to conceal. An agent's chance of winning depends only on how far he looks ahead of his rival, in the same way for both agents; I call this dependence the firm's selection rule, and every such rule is generated by some noise. A noiseless firm always selects the agent who looks better, so her rule is a step: a narrow lead is as good as a wide one, and an agent revealed as bad loses unless his rival is too. Each agent therefore spreads his posterior to land just above his rival's, blending his bad type with his good one: Boleslavsky and Cotton (2015, Lemma 3) show that both agents reveal only partially, randomizing over a continuum of posteriors, and the firm obtains less than 80% of her first best whenever agents are good with probability below one third.

With noise, an agent's chance of winning instead rises smoothly with his lead rather than jumping from zero to one at a tie. A narrow lead is worth little, conceding bad news no longer means certain defeat, and the shape of the noise decides whether an agent gains more from gambling on being revealed as good or from pooling his evidence, and so how much information reaches the firm. The price is accuracy: where the chance of winning is near one half, selection is close to a coin flip, so the firm more

often passes over the agent who looks better.

An example. Suppose that each agent is equally likely to be good and bad. In the unique equilibrium of the noiseless rule, each agent's posterior is uniformly distributed on $[0, 1]$, so the firm sometimes selects a bad agent over a good one and obtains about 89% of her first best. Now suppose the firm scores each agent by his signal plus an independent draw from $U[0, 1]$. Against a rival who reveals his type, an agent's chance of winning is linear in his signal, so revealing is a best reply; against any other signal it is not, and, since the agents' game is constant-sum, full revelation by both agents is the unique equilibrium. As a good type always outscores a bad one, noise only decides between agents of the same type, so the firm attains her first best.

Main results. The example relies on uniform noise and an even prior; the results below show how far this extends. The first set of results, in Section 3, characterizes the equilibrium when the difference between the two evaluation errors has a concave density with a single peak, which I call a *well S-shaped* selection rule; normal and logistic noise qualify if its variance is large enough. Under such a rule, whatever his rival reveals, an agent's chance of winning first accelerates, then slows as his own signal rises, so below some signal he gains from gambling and above it from playing safe. Each agent uses a *bad news* signal: he sometimes reveals the bad type and otherwise pools, with full revelation and pooling as edge cases.

The equilibrium is unique whenever the priors are equal, and for every pair of priors unless the density is linear next to its peak. In the symmetric case, a single statistic governs it, the elasticity of the normalized selection rule at the pooled signal: an agent's bad signal is observed with probability equal to its inverse, and both agents reveal fully once the prior is high enough. With unequal priors (Subsection 3.2.3), each agent reveals more when his rival is better, and when one agent is much better, the favorite reveals nothing while the underdog reveals fully. Under every well S-shaped

rule, the firm gains from symmetrically better agents, even if they reveal less.

The second set of results, in Section 4, concerns the design of the selection rule with equal priors. Within the well S-shaped class, a more elastic rule suppresses information, and for normal and logistic noise more noise means more elastic. The firm's best rule is therefore at the edge of the class: it is generated by i.i.d. uniform shocks, induces the most information, and selects most accurately. It beats the noiseless benchmark once agents are good with probability above roughly 21%, and delivers the first best once they are at least as likely good as bad.

This does not mean that any noise is worth keeping. Scaling the uniform shocks, the firm does best at the least noise that keeps signals binary, which lies at or beyond the class's edge and depends on the agents' prior: more noise only blurs selection, and less noise pushes the equilibrium toward the noiseless one.

Section 4 then asks which rules of the signal gap attain the first best, which with binary types requires full revelation and a rule that always selects a revealed good agent over a revealed bad one. If agents may be both likely good and likely bad, with priors on either side of one half, only the linear rule works robustly, and it leaves agents indifferent among all signals, so robustness costs uniqueness. If the firm knows on which side of one half the prior lies, a rich family of rules makes full revelation the unique equilibrium, although for likely bad agents no i.i.d. shocks generate one.

These limits are specific to rules of the signal gap. A designer who could measure signals perfectly and commit to any rule avoids them: a pass/fail rule attains the first best for every pair of priors in the unique equilibrium, at the price of a strong commitment, as it tosses a coin between an agent who is almost surely good and one who is almost surely bad. With binary types, the linear rule of the signal gap does equally well in its firm-preferred equilibrium.

The third set of results, in Section 5, relaxes the binary prior, the most dispersed

prior with a given mean. Under a well S-shaped rule, each agent still reveals his type below a threshold and pools all types above it, for every prior, and he never reveals more than the lower half of his types; among well S-shaped rules, the firm still does best under the rule of Section 4, which makes him reveal exactly that half. The comparison with the noiseless rule now turns on the shape of the prior. Under a convex prior, the noiseless rule attains the first best, which essentially no other rule can. Under regularity conditions, every other prior admits a rule that beats it, so even a designer who measures signals perfectly and can commit to any rule should blur them. Noise helps by letting pooled agents spread their signals; when a pool follows revealed types, a little noise among the strongest revealed types frees the room for this.

With binary types, a deterministic pass/fail rule attains the first best. With a continuum of types, for concave priors and their close relatives, such as many single-peaked priors, the best rule makes the agents' scores uniform, as in the example, and generically strictly beats both the noiseless rule and every fully revealing rule. For concave priors, it is a random pass mark, which compares the agents exactly at the top and tosses a coin only among weak signals.

Appendix D collects extensions: under reverse S-shaped rules, agents reveal good news; beyond the well S-shaped class, the equilibrium stays unique under a weaker condition on the noise; and with a single agent, noise works the same way.

1.1 Literature review

This paper contributes to the Bayesian persuasion literature (Kamenica and Gentzkow, 2011; Kamenica, 2019), in particular to its strand on competing senders. Most results of Gentzkow and Kamenica (2017) require a Blackwell-connected information environment, which fails when each sender can only reveal information about himself, as here, and their comparison with collusion is vacuous, since the agents' game is constant-sum.

Each agent's best reply solves a single-sender problem with an S-shaped value function, so it is an upper censorship, as characterized by Kolotilin et al. (2022, Theorem 1) for smooth value functions; restated in quantiles, the result extends to every well S-shaped rule and an arbitrary prior (Lemma 13). With a general prior, I verify best replies through a convex function lying above the value function (Dworczak and Martini, 2019, Theorem 1), and restrict selection rules to posterior means, as is standard when the receiver's payoff is linear in the state (Gentzkow and Kamenica, 2016; Kolotilin, 2018). To a single agent, the rule acts like a receiver with a privately known threshold (Kolotilin et al., 2017), here the rival's signal plus noise, and for a single sender, making a bell-shaped threshold density more diffuse leads to more information (Curello and Sinander, 2025, Theorem 2 and Section 5.2). Here, the threshold's distribution is an equilibrium object, set by the rival's signal.

The closest paper is Boleslavsky and Cotton (2015), part of the literature on grading policies (Ostrovsky and Schwarz, 2010; Chan et al., 2007; Popov and Bernhardt, 2013; Bar et al., 2012): two schools design grades, and an evaluator hires the graduate with the higher posterior. I add evaluation noise to this game, and its shape decides what agents reveal. Under the noiseless rule, the equilibrium is unique, posteriors are uniform on an interval, possibly with atoms at 0 and 1, and schools never reveal fully. Informativeness increases with the number of senders, both in this setting, where it becomes full in the limit (Au and Kawai, 2020, Theorem 2), and when senders also compete in prices (Hwang et al., 2026); the latter also characterize the noiseless equilibrium with a continuous prior; with costly inspection, the same convexity condition governs full disclosure (Ding et al., 2023). Banerjee et al. (2026) bound the first-best loss of noiseless competition; a noisy judge can recover much of it, sometimes all. In search markets, whether competition raises informativeness depends on what sellers observe and how buyers search (Board and Lu, 2018; Au and Whitmeyer, 2023; He and Li, 2023).

An imperfect evaluation can also benefit the evaluator by strengthening investment incentives, as strategic grading does in Boleslavsky and Cotton (2015) and a noisy signal of universities' relative quality does in Boleslavsky and Li (2026). Here the gain comes through information instead: a suitably noisy evaluation makes signals more informative. In this, noise plays a role familiar from other settings: noise in prices preserves the incentive to acquire information that fully revealing prices would destroy (Grossman and Stiglitz, 1980), and noise in a cheap-talk channel makes the receiver less responsive to each message, which curbs the sender's exaggeration and improves communication (Blume et al., 2007). Closer in spirit, an organization that can fund only one of two proposals induces more informative evidence than an unconstrained one and is better off unless both are ex-ante strong (Boleslavsky and Cotton, 2018), and moderate attention costs can make a rationally inattentive consumer select the better of two disclosed products more often than when attention is free (Jain and Whitmeyer, 2026). In both, the gain comes from a constraint on the receiver; I instead treat the evaluation error as a primitive of the selection rule, whose shape determines the equilibrium and how much information the firm extracts.

My normative results also relate to the idea that observing an agent more closely can hurt the principal once the agent responds to it (Prat, 2005), and that a receiver can extract more information by committing to use it less: random costs of approval make a single sender indifferent among signals and elicit full revelation, at the price of burned utility (Tsakas et al., 2021), and when agents manipulate their data at a cost, so do a less responsive allocation rule, a test that passes some noncompliant agents, and a score that underweights some features (Frankel and Kartik, 2022; Perez-Richet and Skreta, 2022; Ball, 2025). Extra noise in an already noisy channel can benefit the sender and sometimes both players (Tsakas and Tsakas, 2021), and coarse grades can inform the public better by inducing participation (Harbaugh and Rasmusen, 2018). Here signals

are costless and flexible, and noise works only through competition, by changing the curvature of each agent's value function; it costs the firm nothing when it decides only between agents of the same type. Closest to the firm's commitment, Gao and Luo (2026) let a designer commit to a mechanism mapping each sender's experiment and posterior into a random action: robust mechanisms depend only on posteriors and induce full revelation, and with competing senders the efficient allocation is robustly implementable. Bergemann et al. (2023) show that a receiver uncertain about the sender's preferences optimally commits to a quota. Here the firm commits only to a comparison of the signals, which, for rules of the signal gap, makes robustness costly (Subsection 4.3) and, with a single agent and an absolute standard, caps her payoff (Appendix D.3). A single receiver who could commit to any decision rule would need no noise; with binary types, neither would a firm comparing two agents, as a pass/fail rule shows (Proposition 8), but with a continuum of types even her best rule blurs the signals (Subsection 5.2). In the game of Boleslavsky and Cotton (2018), Muratov (2025) lets the receiver choose how to break ties: asymmetric tie-breaking helps her and in some cases attains the first best; here the rule responds to every signal gap, and with a continuum of types even the best rule randomizes away from ties. When firms are restricted to threshold tests, randomized tests chosen by the principal beat those the firms choose (Banerjee et al., 2021); here signals are unrestricted, and the randomness lies in the comparison. Randomized choice also serves other purposes: breaking ties in school choice (Abdulkadiroğlu and Sönmez, 2003; Erdil and Ergin, 2008), saving on costly signaling (Hoppe et al., 2009), economizing on verification (Border and Sobel, 1987), deterring gaming (Ederer et al., 2018), and keeping improper reasons out of selection (Stone, 2011; Fang and Casadevall, 2016).

The agents' game is also a contest in which each agent chooses a distribution with a fixed mean, and noise gives it a difference-form success function (Lazear and Rosen,

1981; Hirshleifer, 1989; Che and Gale, 2000), the counterpart of the ratio form of Tullock (1980), with more noise playing the role of lower decisiveness. When contestants choose the riskiness of their performance, limiting competition helps a principal select the better one (Hvide and Kristiansen, 2003; Fang and Noe, 2022), and a noisy selection rule is a new instrument with the same effect. In effort contests, a noisy contest can also be optimal (Wang, 2010; Morgan et al., 2022; Letina et al., 2023), and the shape of the noise matters (Drugov and Ryvkin, 2020); see Fu and Wu (2019) for a survey.

2. Model

Agents. There are two agents and a firm (she). Each agent's type θ_i is drawn independently from $\{0,1\}$, with *prior* $p_i := \mathbb{P}(\theta_i = 1)$, written p if $p_1 = p_2$.

Information and Timing. Agents play a simultaneous-move game in which, before his type is realized, each agent i commits to a signal structure; once θ_i is realized, the firm observes a draw from it. As the prior is binary, a posterior is pinned down by its mean, so I identify each realization with its posterior mean $s_i := \mathbb{E}[\theta_i|s_i]$, and agent i chooses a distribution G_i of posterior means (a *signal*). Feasible signals for a prior p_i form $\mathcal{G}(p_i) := \{G_i : \text{supp}(G_i) \subseteq [0,1], \int_0^1 s_i \, dG_i(s_i) = p_i\}$ (Gentzkow and Kamenica, 2016, Remark 1 and Proposition 1). Section 5 allows general priors.

Selection. Agents choose signal structures knowing the firm's *selection rule*: a function $v : [-1,1] \to [0,1]$ such that $v(x) - \frac{1}{2}$ is odd, increasing[1] and continuous. Given signals (s_1, s_2), agent i is selected with probability $v(s_i - s_j)$. As $v(x) - \frac{1}{2}$ is odd, $v(s_1 - s_2) + v(s_2 - s_1) = 1$, so the firm always selects exactly one agent and treats the agents anonymously. The rule is exogenous until Section 4, where the firm chooses it.

[1]Throughout, increasing and decreasing are meant weakly; strict monotonicity is stated explicitly.

Noisy evaluation. Such selection rules arise naturally when the firm judges the agents with noise. Suppose that, instead of observing s_i without noise, the firm observes agent i's *score* $s_i + \varepsilon_i$, where ε_i is a random shock, independent of s_i, s_j, reflecting the idiosyncrasies of whoever is judging – a particular committee member's mood, the order in which proposals were read, an unmodeled personal rapport.[2] Taking the scores at face value, the firm selects the agent with the higher score. Suppose that $\xi := \varepsilon_j - \varepsilon_i$ has a continuous distribution, symmetric around 0; symmetry holds, for instance, whenever the shocks are exchangeable. Then she selects agent i with probability

$$\mathbb{P}(s_i + \varepsilon_i > s_j + \varepsilon_j) = \mathbb{P}(\xi < s_i - s_j) = v(s_i - s_j),$$

where $v(x)$, the CDF of ξ restricted to $[-1, 1]$, is a selection rule. Conversely, every selection rule arises in this way, although i.i.d. shocks generate only some (Corollary 2). I usually give $v(x)$ explicitly, and otherwise say that it *is generated by* the shocks.

Example 1. If $v(x)$ is generated by ε_i i.i.d. Gumbel or normal, then $v(x)$ is the CDF of a logistic or normal distribution, respectively.

Payoffs. Each agent receives 1 if selected and 0 otherwise. If the firm selects agent i, she receives θ_i, so her expected payoff is the selected agent's posterior s_i. The shocks that may generate the selection rule do not enter her payoff.

Commitment. The firm commits to her selection rule. The commitment need not be deliberate: some noise is unavoidable and undetectable, so the firm cannot filter it out, and everyone understands how it shapes the verdict. Nor, for the leading noise distributions, is it a restriction: if the shocks are i.i.d. with a log-concave density,

[2]The noise is additive on the scale of the firm's payoff; noise on another scale, such as log-odds, would give rules that depend on more than the signal gap.

such as normal, Gumbel or uniform, and both agents use the same signal, as in the equilibrium for equal priors, then by the monotone likelihood ratio property the firm's expected payoff $\mathbb{E}[s_i|s_i + \varepsilon_i]$ is non-decreasing in the score, so selecting the higher score is optimal even without commitment. For unequal priors, and for rules that i.i.d. shocks cannot generate, commitment can bind.

Equilibrium. As agent i commits before learning θ_i, he maximizes his ex-ante probability of being selected. An equilibrium of the game is a pair (G_1^*, G_2^*) such that

$$G_i^* \in \arg\max_{G_i \in \mathcal{G}(p_i)} \int_0^1 \int_0^1 v(s_i - s_j) dG_j^*(s_j) dG_i(s_i).$$

For any priors and rule, an equilibrium exists by Sion's minimax theorem: the game is constant-sum, as $v(x) - \frac{1}{2}$ is odd, payoffs are linear and continuous in G_i, and $\mathcal{G}(p_i)$ is convex and weakly compact, as a closed subset of the weakly compact set of distributions on $[0, 1]$. Equilibria are pairs of max-minimizers and form a product set.

Interpretation. The model is deliberately abstract and says nothing about a price: agent i's type θ_i is simply the value the firm realizes from picking i, with no wage or other transfer attached to that choice. This fits any setting in which a decision-maker allocates one indivisible prize between two competing options and cannot pay more to secure a better one. My running example is **internal capital budgeting**: a firm's capital committee (*the firm*) must fund exactly one of two project proposals this cycle, and each project's sponsor (*agent i*) controls how much hard evidence – the scope of a pilot, the rigor of a feasibility study, which numbers are led with – to place in front of the committee. The same primitives fit a **grant panel** choosing between two proposals, and **grading**, as in Boleslavsky and Cotton (2015): a university chooses a

grading policy for a graduate who competes for a single place and is evaluated by an admissions office. In this reading, the grading policy is the signal, the prior p_i is the strength of the school's cohort, and the noise is the office's judgment of what a transcript leaves out, such as interviews and essays; the school sets its policy before it learns how good its graduates are, which is the commitment the model assumes. I use *the firm* and *agent* as generic labels and switch between these readings.

Under any of these readings, the signal $G_i(s_i)$ is the design of the hard information about the project, applicant or student – the scope and pre-registration of a pilot study, the preliminary results in a grant proposal, a school's grading policy – summarized by the distribution of the posterior s_i that the agent is good. Equivalently, as in Boleslavsky and Cotton (2015), each type generates a more primitive random report – a grade, a pilot outcome, a preliminary result – whose distribution satisfies the monotone likelihood ratio property, so that, by Bayes' rule, the posterior increases in the report. I read θ_i as fit with what the firm needs, which an agent can hardly assess; what he controls, before any result exists, is how much evidence there will be. If he knew more than the firm, this would be a signaling problem, which I do not study.

The shocks are imperfections of the kind documented in the introduction: they move the firm's decision but are unrelated to the substance of the signals and to her payoff. An agent may also have outcome-relevant characteristics unrelated to his signal; I set these aside, since agents care only about the selection rule, and comparing the firm's utility across rules is cleaner when the shock is pure evaluation noise.

3. Equilibrium under Well S-Shaped Selection Rules

In this section, I study the agents' game under the additional assumption that the selection rule $v(x)$ has a concave density with a unique mode. Subsection 3.1 shows that this restricts every best reply to a bad news signal. Subsection 3.2 characterizes

the equilibrium: for equal priors, it is unique and governed by a single statistic of the rule, and for unequal priors, it takes one of four forms, in which the favorite plays safe and the underdog gambles. Subsection 3.3 shows that a more elastic rule suppresses information, and Subsection 3.4 that the firm gains from better agents.

3.1 Best replies

Definition 1. A function $v : [-1, 1] \to [0, 1]$ is called *well S-shaped* if $v(x) - \frac{1}{2}$ is odd and increasing, and $v'(x)$ is concave on $[-1, 1]$ and uniquely maximized at $x = 0$.

Since $v(x) - \frac{1}{2}$ is odd, $v'(x)$ is even, so a concave $v'(x)$ is always maximized at $x = 0$. Requiring this maximum to be unique only rules out densities that are flat around 0, including the linear selection rule. Without this requirement, the equilibrium may fail to be unique: if $v(x)$ is linear on some $[-b, b]$, then for priors (p, p) with $p \leq \frac{b}{2}$, both $(\{p\}; \{p\})$ and $(\{0, 2p\}; \{0, 2p\})$ are equilibria. If $v(x)$ is generated by ε_i, ε_j, well S-shapedness means that ξ has a concave density on $[-1, 1]$ with a unique mode, which holds for the distributions of Example 1 if their variance is large enough (Example 2).

Example 2. For $v(x)$ generated by i.i.d. Gumbel or normal shocks, there is a threshold $\overline{V}$ such that $v(x)$ is well S-shaped if $\text{Var}[\varepsilon_i] \geq \overline{V}$. So is the rule generated by i.i.d. $\varepsilon_i \sim U[0, u]$ with $u \geq 1$, although its triangular density is linear on $[0, 1]$.

For equal priors, the results below need only well S-shapedness, so they cover rules whose density is linear next to its mode, such as the uniform rules of Example 2; for $u = 1$, this is the firm's best rule (Theorem 2). For unequal or non-binary priors, uniqueness also requires that $v'(x)$ is *not linear near* 0, i.e., not linear on any $[0, \varepsilon]$, which makes every best reply unique (Lemma 1).

Let $\overline{v}(s_i, G_j) := \int_0^1 v(s_i - s_j) dG_j(s_j)$ be the *value function* of agent i, written $\overline{v}(s_i)$ from here on for simplicity when G_j is fixed. By choosing a signal G_i, agent i solves a

Bayesian persuasion problem as in Kamenica and Gentzkow (2011):

$$\max_{G_i \in \mathcal{G}(p_i)} \int_0^1 \overline{v}(s_i, G_j) dG_i(s_i) \tag{BP}$$

A signal G_i that solves (BP) is *an optimal signal* for $(\overline{v}(s_i), p_i)$, and *the optimal signal* if no other signal does. To find one, I use the *concavification* $\overline{v}^{cv}(s_i)$ of $\overline{v}(s_i)$, its smallest concave majorant. A signal G_i is optimal for $\overline{v}(s_i)$ if and only if it is optimal for $\overline{v}^{cv}(s_i)$ and supported on the *concavification support* $\text{supp}(\overline{v}^{cv}) := \{s_i : \overline{v}(s_i) = \overline{v}^{cv}(s_i)\}$. Optimality for the concave $\overline{v}^{cv}(s_i)$ is easy to check (Kamenica and Gentzkow, 2011, Corollary 2): if $\overline{v}^{cv}(s_i)$ is linear on an interval around p_i, every feasible signal supported on that interval is optimal; otherwise, only pooling at p_i is.

A well S-shaped $v(x)$ helps because if $v'(x)$ is concave on $[-1, 1]$, then $\overline{v}'(s_i)$ is concave on $[0, 1]$ for every G_j, which gives $\overline{v}(s_i)$ the following shape: $\overline{v}(s)$ is *S-shaped* on $[a, b]$ if it is convex on $[a, c]$ and concave on $[c, b]$ for some $c \in [a, b]$. As $v(x)$ is continuously differentiable, so is $\overline{v}(s_i)$, and then $\overline{v}(s_i)$ is S-shaped if and only if $\overline{v}'(s_i)$ is quasi-concave. Since $\overline{v}'(s_i)$ is in fact concave, $\overline{v}(s_i)$ is S-shaped for every well S-shaped $v(x)$ and every G_j. As quasi-concave functions are not closed under addition, there is no simple extension of the well S-shaped class for which $\overline{v}(s_i)$ is always S-shaped. Characterizing the equilibrium, however, needs this shape only against the rival's equilibrium signals. Using this, Proposition 21 in Appendix D.2 goes beyond the class, giving a weaker sufficient condition for a unique bad news equilibrium.


So $\overline{v}(s_i)$ has at most two regions of different convexity, and its concavification consists of at most two parts, which restricts the optimal signals (Lemma 1(ii)). Throughout, a signal with at most two points is identified with its support.

Lemma 1. *If $v(x)$ is well S-shaped, then*

(i) $\overline{v}(s_i, G_j)$ is S-shaped on $[0, 1]$ for all $p_j \in (0, 1)$ and $G_j \in \mathcal{G}(p_j)$.

(ii) An optimal signal for $(\overline{v}(s_i), p_i)$ is $\{0, a\}$ if $p_i < a$ and $\{p_i\}$ otherwise, where a is the largest $s_i \in [0,1]$ at which the secant of $\overline{v}$ from 0 to s_i is steepest, the secant at $s_i = 0$ being the tangent. If $v'(x)$ is not linear near 0, it is the optimal signal.

Figure 1: Illustration of Lemma 1 (ii)



(a) $\text{supp}(G_i) = \{0, a\}$ or $\{p_i\}$ (b) $\text{supp}(G_i) = \{0, 1\}$

This is not the image of $\overline{v}(s_i)$, but of $\overline{v}(s_i) - bs_i$ for some $b > 0$ for clarity. Since $cav(f(x) + bx) = cav(f(x)) + bx$ for all x, b, this does not influence the optimal signal.

Part (i) was shown above. For (ii), the concavification of an S-shaped $\overline{v}(s_i)$ coincides with the secant from $\overline{v}(0)$ to $\overline{v}(a)$ on $[0, a]$ and with $\overline{v}(s_i)$ on $[a, 1]$. Hence $\{0, a\}$ is an optimal signal if $a > p_i$, and $\{p_i\}$ if $a \leq p_i$, and if $v'(x)$ is not linear near 0, these are *the* optimal signals. Figure 1 illustrates both cases.

Lemma 1 thus restricts an agent's optimal signal to a very limited class, the *bad news* signals: the agent reveals the low type with some probability and otherwise sends a pooled signal, with full revelation ($\{0, 1\}$) and pooling ($\{p_i\}$) as the edge cases. Binary support itself is not special: two-point signals are the pure strategies of an agent, every other signal is their mixture, and against any continuous value function some signal with at most two support points is a best reply. What a well S-shaped $v(x)$ adds is that, unless pooling is optimal, such a best reply has 0 in its support, and that it is

unique if $v'(x)$ is not linear near 0. Moreover, the point a depends only on the rival's signal, and the agent's own prior only determines whether he pools at p_i or plays $\{0, a\}$. The concavity of the density is what makes agents concede bad news: under the mirror-image, reverse S-shaped rules of Appendix D.1, they reveal good news instead.

By the Bayesian constraint, positive scaling of $v(x)$ and adding a linear function to it do not change the concavification support, and hence do not change equilibria for any (p_1, p_2). I represent each such equivalence class of selection rules by the normalization

$$\tilde{v}(x) := \frac{v(x) - \frac{1}{2} - x \cdot v'(0)}{|v(1) - \frac{1}{2} - v'(0)|}.$$

It satisfies $\tilde{v}(0) = \tilde{v}'(0) = 0$ and $\tilde{v}(1) = -1$, which simplifies the exposition, and it is well defined for every well S-shaped $v(x)$: as $v'(x)$ is uniquely maximized at $x = 0$, $v(1) - \frac{1}{2} = \int_0^1 v'(x)dx < v'(0)$.

The key characteristic of $\tilde{v}(x)$ is its elasticity $\mathcal{E}_{\tilde{v}}(x) = \frac{x\tilde{v}'(x)}{\tilde{v}(x)}$. Since $\tilde{v}(0) = 0$, $\mathcal{E}_{\tilde{v}}(a)$ is the ratio of the slopes of the tangent to $\tilde{v}(x)$ at $x = a$ and of the secant from 0 to a, so it measures how curved $\tilde{v}(x)$ is on $[0, a]$. This curvature shapes $\overline{v}(s_i)$, and through it the pooled signal a of Lemma 1: the elasticity pins down when agents reveal fully (Proposition 1) and where they pool otherwise (Theorem 1).

3.2 Equilibrium

3.2.1 Full revelation

A natural first question is whether the firm can ever get the first best information, full revelation by both agents, in equilibrium. Under a well S-shaped $v(x)$, the answer is simple: both priors must be high enough.

Definition 2. For a given well S-shaped $v(x)$, let $\overline{p} := 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)} = 1 - \frac{1}{|\tilde{v}'(1)|}$.

Proposition 1. *Let $v(x)$ be well S-shaped. Then both agents playing full revelation is an equilibrium if and only if $p_1, p_2 \geq \overline{p}$, and then it is the unique equilibrium unless $v'(x)$ is linear near 0 and $\max(p_1, p_2) > \overline{p} = \min(p_1, p_2)$.*

Competing with a sufficiently good rival induces an agent to reveal fully, and the proof of the condition $p_1, p_2 \geq \overline{p}$ is built on exactly that. If agent j plays $\{0, 1\}$, the normalized value function of agent i is

$$\overline{v}(s_i) = (1 - p_j)\tilde{v}(s_i) + p_j\tilde{v}(s_i - 1) = (1 - p_j)\tilde{v}(s_i) - p_j\tilde{v}(1 - s_i).$$

By Lemma 1, it is S-shaped, so $\{0, 1\}$ is optimal if and only if the secant from 0 to 1 is weakly flatter than the tangent at $x = 1$ (see Figure 1b):

$$\overline{v}(1) - \overline{v}(0) \leq \overline{v}'(1) \Leftrightarrow \tilde{v}(1) \leq (1 - p_j)\tilde{v}'(1) \Leftrightarrow p_j \geq 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)}.$$

For uniqueness, as the game is constant-sum, all equilibria form a product set, so any signal played in equilibrium is a best reply to full revelation. An S-shaped $\overline{v}(s_i)$ can touch its secant from 0 to 1 at some $s \in (0, 1)$ only by coinciding with it on $[s, 1]$, which non-linearity of $v'(x)$ near 0, or a tangent at 1 strictly steeper than the secant ($p_j > \overline{p}$), rules out, and then full revelation is the only best reply. For $p_j = \overline{p}$, the best reply need not be unique, but uniqueness follows from Theorem 1.

Remark 1. Uniqueness can fail when both conditions of the exception hold. For $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$, $v'(x)$ is linear near 0 and $\overline{p} = \frac{1}{2}$; for $(p_1, p_2) = (0.75, \frac{1}{2})$, agent 1's value function against $\{0, 1\}$ is linear, and $\{0, 1\}$ remains a best reply to any $G_1 \in \mathcal{G}(0.75)$, since $\mathbb{E}_{G_1}[s_1^2] \geq \frac{1}{2}$. So $(G_1; \{0, 1\})$ is an equilibrium for every such G_1.

Proposition 1 contrasts with the noiseless benchmark[3] of Boleslavsky and Cotton

[3]The noiseless rule $v(x) = \frac{1}{2} + \frac{1}{2} \cdot \text{sign}(x)$ is not a selection rule, as it is discontinuous at 0; Appendix B.12 gives its equilibria.

(2015, Proposition 1), in which schools never use a fully informative grading policy. This is not a feature of well S-shaped rules alone: Section 4 shows that full revelation can be reached by selection rules far beyond this class.

How likely to be good must the agents be for full revelation? For every well S-shaped rule, the threshold $\overline{p}$ is at least $\frac{1}{2}$, because the density of the shock difference is concave with its peak at 0, and it equals $\frac{1}{2}$ only if $\tilde{v}(x) = -x^2$ on $[0,1]$ (Lemma 3). So, under a well S-shaped rule, full revelation requires both agents to be at least as likely good as bad, and, outside this knife-edge case, strictly more likely.

3.2.2 Equal priors

For equal priors, Lemma 1 and Proposition 1 characterize the equilibrium on the well S-shaped class. Unless it is full revelation, it is a bad news equilibrium in which the bad signal $s_i = 0$ has probability $1/\mathcal{E}_{\tilde{v}}(a)$, the inverse elasticity at the pooled signal a.

Theorem 1. *Let $v(x)$ be well S-shaped and the priors (p, p). If $p \geq \overline{p}$, the unique equilibrium is $(\{0,1\};\{0,1\})$. If $p < \overline{p}$, the unique equilibrium is $(\{0,a\};\{0,a\})$, where a solves $p = a(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)})$, and $\mathbb{P}(s_i = 0) = \frac{1}{\mathcal{E}_{\tilde{v}}(a)}$.*

The intuition is simple. Against $\{0,a\}$, an agent's normalized value function is

$$\overline{v}(s_i) = \mathbb{P}(s_i = 0)\tilde{v}(s_i) + \mathbb{P}(s_i = a)\tilde{v}(s_i - a) = \frac{a-p}{a}\tilde{v}(s_i) + \frac{p}{a}\tilde{v}(s_i - a),$$

where the second term $\tilde{v}(s_i - a) = -\tilde{v}(a - s_i)$ reflects $\tilde{v}(s_i)$ on $[0,a]$. For $\{0,a\}$ to be a best reply, the secant from $\overline{v}(0)$ to $\overline{v}(a)$ must be tangent at $s_i = a$. Since $\tilde{v}'(0) = 0$, the second term has zero derivative at a, so $\overline{v}'(a) = \frac{a-p}{a}\tilde{v}'(a)$, while the secant has the slope $\frac{\tilde{v}(a)}{a}$, the same for both terms. Hence, in equilibrium,

$$\frac{a-p}{a}\tilde{v}'(a) = \frac{\tilde{v}(a)}{a} \Leftrightarrow \frac{a-p}{a} = \frac{1}{\mathcal{E}_{\tilde{v}}(a)} \Leftrightarrow p = a\left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\right).$$

Such a is unique, continuous and strictly increasing in p: the pooled signal rises with the prior. Indeed, $p = a - \frac{\tilde{v}(a)}{\tilde{v}'(a)}$ is the point where the tangent to $\tilde{v}(x)$ at $x = a$, $y = \tilde{v}(a) + (x - a)\tilde{v}'(a)$, crosses the x-axis. Since $\tilde{v}(x)$ is strictly concave and decreasing on $[0, 1]$ with $\tilde{v}(0) = 0$, this point is continuous and strictly increasing in a, and so is its inverse $a(p)$. As $\mathbb{P}(s_i = 0) = \frac{a-p}{a}$, this also gives the inverse elasticity result.

Uniqueness needs one more step, since $\{0, a\}$ need not be the unique best reply to itself: if $\tilde{v}(x)$ is quadratic on $[0, 2p]$, every signal supported there is a best reply to $\{0, 2p\}$. But then $\{0, 2p\}$ is a best reply to none of them except itself, so, as equilibria form a product set, the equilibrium is still unique. Pooling can be approached in the limit: for $\tilde{v}_n(x) = -x^n$ on $[0, 1]$, the theorem gives $a = \frac{np}{n-1} \to p$, and more generally $a \to p$ whenever the elasticity of $\tilde{v}(x)$ diverges uniformly on $(0, 1)$ (Lemma 6).

3.2.3 Unequal priors

For unequal priors, let $p_2 \leq p_1$, so that agent 1 is the favorite and agent 2 the underdog, and let $v'(x)$ be not linear near 0. By Lemma 1, every best reply is unique and takes one of three forms: pooling $\{p_i\}$, full revelation $\{0, 1\}$, or bad news $\{0, c\}$ with $p_i < c < 1$. In general, an agent's best reply depends on his rival's whole signal, but in the following example it depends only on the rival's prior, which makes the equilibrium explicit.

Example 3. Let $\tilde{v}(x) = -x^3$. Agent i's best reply is $\{0, \min(\frac{3p_j}{2}, 1)\}$, or $\{p_i\}$ if $p_i \geq \frac{3p_j}{2}$. So for $p_2 \leq p_1$, the unique equilibrium is $(\{0, 1\}; \{0, 1\})$ if $p_2 \geq \frac{2}{3}$, and otherwise

(i) $(\{p_1\}; \{0, 1\})$ if $p_1 \geq \max(\frac{3p_2}{2}, \frac{2}{3})$;

(ii) $(\{p_1\}; \{0, \frac{3p_1}{2}\})$ if $\frac{3p_2}{2} \leq p_1 < \frac{2}{3}$;

(iii) $(\{0, \frac{3p_2}{2}\}; \{0, 1\})$ if $\frac{2}{3} \leq p_1 < \frac{3p_2}{2}$;

(iv) $(\{0, \frac{3p_2}{2}\}; \{0, \frac{3p_1}{2}\})$ if $p_1 < \min(\frac{3p_2}{2}, \frac{2}{3})$.

The example shows what asymmetry does. Each agent would ideally pool at one and a half times his rival's prior. When the priors are close and low enough, both concede bad news, and each pooled signal is set by the rival: the underdog, facing the better rival, pools higher and so reveals more. As the favorite's lead grows, the signal at which he would pool falls below his own prior, so once $p_1 \geq \frac{3p_2}{2}$ he pools at his prior. The underdog, facing a rival who reveals nothing, gambles harder and reveals fully once $p_1 \geq \frac{2}{3}$. So an agent nearly sure to win plays safe, and one likely to lose gambles; the reverse S-shaped rules of Appendix D.1 reverse the roles.

Proposition 2. *Let $v(x)$ be well S-shaped with $v'(x)$ not linear near 0, and let $p_1, p_2 \in (0,1)$ with $p_2 \leq p_1$. Then the equilibrium is unique: it is $(\{0,1\};\{0,1\})$ if $p_2 \geq \overline{p}$, and otherwise it has one of the following four forms:*

(i) $(\{p_1\};\{0,1\})$*: the favorite pools, the underdog reveals fully;*

(ii) $(\{p_1\};\{0,b\})$ *with $p_2 < b < 1$: the favorite pools, the underdog concedes bad news;*


(iii) $(\{0,a\};\{0,1\})$ *with $p_1 < a$: the favorite concedes bad news, the underdog reveals fully;*

(iv) $(\{0,a\};\{0,b\})$ *with $p_1 < a \leq b < 1$, and $a < b$ if $p_1 > p_2$: both concede bad news.*

In Example 3, regions (i)–(iv) are exactly the cases listed there (Figure 2). For a general rule, the two thresholds depend on the rival's whole signal rather than on his prior, so the regions have curved boundaries, given in Appendix A.7. Every pair of forms except pooling by both agents is played for some priors; against pooling, an agent's value function is convex below the rival's prior and concave above it, so pooling is a best reply only for the agent with the strictly higher prior (Lemma 5 in Appendix A.5), and the underdog never pools at his prior. Without non-linearity of $v'(x)$ near 0, neither uniqueness nor these forms are guaranteed (Remark 2).

In regions (ii) and (iii) of Proposition 2, an agent's bad news signal depends only on his rival's prior. More generally, the rival forces an agent to reveal.

Figure 2: Equilibria for unequal priors in Example 3, $\tilde{v}(x) = -x^3$ on $[0, 1]$



Colors are for the corresponding equilibrium type: black – $(\{0,1\};\{0,1\})$*; red* – $(\{p_1\};\{0,1\})$*; green* – $(\{0,a\};\{0,b\})$*; blue* – $(\{p_1\};\{0,b\})$*; yellow* – $(\{0,a\};\{0,1\})$.

Proposition 3. *Let $v(x)$ be well S-shaped with $v'(x)$ not linear near 0. Then agent i's equilibrium signal is non-decreasing in p_j, strictly so if it is $\{0,a\}$ with $p_i < a < 1$.*

An agent's own prior usually has the opposite effect: in Example 3, his high signal does not depend on p_i, so he reveals less as p_i increases. This effect need not be monotone, though (Example 6 in Appendix A.7).

3.3 Noise and informativeness

In the symmetric equilibrium of Theorem 1, a higher pooled signal a means, for a fixed p, a more informative signal: the agent reveals his bad type more often, and revealing it less often is a garbling (Blackwell, 1953). So the selection rule affects information through a. If $\tilde{v}(x)$ has a constant elasticity, then, for a fixed p, a higher elasticity lowers $\mathbb{P}(s_i = 0)$ and hence a, making the signal less informative. The next corollary extends this argument to any two selection rules.

Corollary 1. *Let $v_e(x)$ and $v_n(x)$ be well S-shaped, let the priors be (p, p) with $p \in (0, \min(\overline{p}_e, \overline{p}_n))$, and let $(\{0, a_k\}; \{0, a_k\})$ be the equilibrium for v_k, $k \in \{e, n\}$. If $\tilde{v}_e(x)$ is more elastic than $\tilde{v}_n(x)$, i.e. $\mathcal{E}_{\tilde{v}_e}(x) > \mathcal{E}_{\tilde{v}_n}(x)$ for all $x \in (0, 1)$, then $a_e < a_n$.*

So a more elastic selection rule suppresses information, and within the well S-shaped class, the most informative signal is induced by the rules with $\mathcal{E}_{\tilde{v}}(x) = 2$, i.e. $\tilde{v}(x) = -x^2$ on $[0, 1]$. For the natural examples, this reads intuitively in terms of noise.

Example 4. For $v(x)$ generated by i.i.d. Gumbel or normal shocks, $\mathcal{E}_{\tilde{v}}(x)$ increases in $\mathrm{Var}[\varepsilon_i]$ on $[\overline{V}, +\infty)$ for every $x \in (0, 1)$, so noisier shocks yield less information.

3.4 Better agents and the firm

It is natural to expect the firm to benefit from ex-ante better agents. But better agents need not reveal more: informativeness can fall in the prior (Appendix A.10), and this fall could outweigh the direct gain from a higher prior.

For a well S-shaped $v(x)$ and priors (p, p), let (G, G) be the unique equilibrium and $u_F(p) := \mathbb{E}_G[s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)]$ the firm's expected utility.

Proposition 4. *Let $v(x)$ be well S-shaped. Then $u_F(p)$ is strictly increasing in p.*

So, despite possibly decreasing informativeness, a firm that uses a well S-shaped selection rule is never harmed by ex-ante better agents. The result is not immediate: the equilibrium signals for different priors need not be ranked by first-order stochastic dominance or in the Blackwell order, and under a noisy selection rule the firm's payoff need not be monotone or convex in the signals. However, the change from p to $p' > p$ splits into two steps within the bad news class. First, keeping the prior p, the pooled signal rises from $a(p)$ to $a(p')$: this spread reveals the bad type more often and widens the gap between the signals, so the firm selects the worse agent less often. Second, keeping the pooled signal $a(p')$, the prior rises to p', which moves mass from 0 to $a(p')$,

and on $\{0, a\}$ the firm's payoff is monotone. Both steps benefit the firm. This logic rests on the shape of $v(x)$: under a well S-shaped rule, the pooled signal moves away from the revealed one as p increases. Under the reverse S-shaped selection rules of Appendix D.1, it moves toward it, and there exist $p_1 < p_2$ such that $u_F(p_1) > u_F(p_2)$ in the unique equilibrium. Better agents need not make her mistakes rarer: the probability of selecting the agent with the lower posterior falls in p if $\mathcal{E}_{\tilde{v}}(x)$ is non-decreasing, but can rise otherwise, as Appendix A.15 shows by example.

For unequal priors, this monotonicity extends to every region of Proposition 2 except region (iv), where both agents concede bad news: there, once the underdog stops revealing fully, his pooled signal can fall as his prior rises, and the firm's utility with it (Proposition 13 and Example 7 in Appendix A.7). For the agents, a better own prior always helps and a better rival always hurts (Lemma 7 in Appendix A.7).

4. Designing the Selection Rule

This section turns to the normative side: the firm's choice of the selection rule. Subsection 4.1 finds the best rule within the well S-shaped class, and Subsection 4.2 the optimal level of noise beyond it. Both depend on the prior, so Subsection 4.3 asks which rules implement full revelation robustly, and Subsection 4.4 compares rules of the signal gap with a benchmark designer who may use any rule.

4.1 The best well S-shaped rule

I now depart from an exogenous $v(x)$: for a fixed p, the firm chooses the selection rule to maximize $u_F(p)$, for instance, by choosing between committee structures, evaluation protocols or review panels. By changing the noise that shapes her selection rule, she changes the equilibrium signals and her utility.

Two questions arise. First, what is the best equilibrium utility within the well

S-shaped class for each p, and which rules attain it? Second, how does it compare with the noiseless rule? The answer to the latter is not obvious. In the example of the introduction, the noiseless rule leaves each agent's posterior uniformly distributed on $[0, 1]$, which is less informative than the binary signal under a well S-shaped rule; there noise costs no accuracy, but in general it trades selection accuracy for more information.

The natural examples show an important pattern: for $v(x)$ generated by i.i.d. Gumbel or normal shocks, $u_F(p)$ decreases in $\mathrm{Var}[\varepsilon_i]$ on $[\overline{V}, +\infty)$ for every $p \in (0, 1)$, so within each class the best rule lies at the "edge" of the well S-shaped class: the lowest level of noise that keeps $v(x)$ well S-shaped. The next theorem gives a tight upper bound on the firm's utility over the whole class and the rules that attain it.

Theorem 2. *Let $v(x)$ be well S-shaped. Then $u_F(p) \le u_F^{ws}(p) = p + \min(2p, 1) \cdot p(1-p)$, and the rule $v^*(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$ attains this bound for every $p \in (0, 1)$. Moreover, $v^*(x)$ gives the firm strictly more than the noiseless rule if and only if $p > \frac{3-\sqrt{3}}{6} \approx 0.21$.*

Since the first best utility is $u_F^*(p) := \mathbb{E}[\max(\theta_1, \theta_2)] = 2p - p^2 = p + p(1-p)$, $v^*(x)$ captures the share $\min(2p, 1)$ of the first best gain over random selection. No other well S-shaped rule attains the bound for $p < \frac{1}{2}$; for $p \ge \frac{1}{2}$, a rule attains it if and only if $p \ge \overline{p}$, so that agents reveal fully, and $v(1) = 1$ (Appendix B.1).

The rule $v^*(x)$ has a triangular density and is generated by two independent $\varepsilon_i \sim U[0, 1]$. Its optimality has a simple source. In the equilibrium $(\{0, a\}; \{0, a\})$, the firm's utility is $p + \mathbb{P}(s_1 \neq s_2) \cdot a \cdot \left(v(a) - \frac{1}{2}\right)$: she obtains p from a random pick and gains only when the signals differ, by their gap times her excess probability of selecting the higher one. This gain rises with the pooled signal a and, at a given a, with the accuracy $v(a)$, and $v^*(x)$ is best on both counts. It induces the highest possible pooled signal, $a = \min(2p, 1)$, and since a concave density on $[0, 1]$ declines no faster, in proportion, than the triangular density $1 - x$, and its mass there is at most $\frac{1}{2}$, $v^*(x)$ is also the most accurate well S-shaped rule at every gap: $v(x) \le v^*(x)$ on $[0, 1]$. So $v^*(x)$ lies at the




Figure 3: Losses from the best well S-shaped selection rules

far edge of the class, as in the natural examples, and the firm trades sharper selection against more information only beyond it.

Figure 3 shows the share of the first best utility $u_F^*(p)$ attained by $v^*(x)$ and by the noiseless rule. The extra information $v^*(x)$ induces outweighs the cost of sometimes selecting a bad agent over a good one, most of all at high priors, where agents reveal fully and $v^*(x)$ attains the first best.

4.2 The optimal level of noise

Theorem 2 leaves open what happens beyond the edge of the well S-shaped class, where the noise is too small for the density of the shock difference to be concave. The selection rules generated by i.i.d. $\varepsilon_i \sim U[0, u]$, which have a triangular density on $[-u, u]$, cover both sides of this edge. For $u \geq 1$, $v(x)$ is well S-shaped with $\tilde{v}(x) = -x^2$ on $[0, 1]$, and $u = 1$ gives $v^*(x)$; for $u < 1$, $v'(x)$ is not concave on $[0, 1]$, and as $u \to 0$, $v(x)$ converges pointwise to the noiseless rule. For every u and p, a symmetric equilibrium can be constructed explicitly, and it has binary support if and only if $u \geq 2\min(p, 1 - p)$; for smaller u, each agent randomizes over more than two posteriors. The construction,

in Appendix B.3, rests on u-partitions: signals on an interval $[a, b]$ such that, at every point s of $[a, b]$, the rival's signal is as likely to lie within u above s as within u below s. Under the triangular rule, the curvature of an agent's value function at s is proportional to the difference of these two probabilities, so against a u-partition it is linear on $[a, b]$. The next proposition concerns only informativeness and the firm's utility.

Proposition 5. *Let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$, $u \in (0, 1]$, and let $u_F(p, u)$ be the firm's utility in the equilibrium (G_u, G_u) constructed in Appendix B.3.*

(i) If $p \leq \frac{1}{2}$, G_u is strictly increasing in u in the Blackwell order on $(0, 2p]$, and $G_u = \{0, 2p\}$ for $u \in [2p, 1]$. If $p > \frac{1}{2}$, $G_u = \{0, 1\}$ for $u \in [2 - 2p, 1]$.

(ii) If $p \leq \frac{1}{2}$, $u_F(p, u)$ is uniquely maximized at $u = 2p$ and strictly decreasing on $[2p, 1]$. If $p > \frac{1}{2}$, $u_F(p, u)$ equals the first best $u_F^*(p) = 2p - p^2$ if and only if $u \in [2 - 2p, 1]$.

(iii) As $u \to 0$, G_u converges weakly to the equilibrium signal for the noiseless rule.

The proposition considers only $u \leq 1$: for $u \geq 1$, $v(x)$ is well S-shaped with $\tilde{v}(x) = -x^2$ on $[0, 1]$, so the equilibrium signal does not change with u, while $v(x) = 1 - \frac{(u-x)^2}{2u^2}$ decreases in u on $(0, 1]$, and so does the firm's utility.

Part (i) shows that for $p \leq \frac{1}{2}$, less noise only suppresses information. For $p > \frac{1}{2}$ and $u < 2 - 2p$, this ranking breaks down. For instance, for $p = \frac{2}{3}$, G_u puts $(\frac{1}{5}, \frac{1}{5}, \frac{3}{5})$ on $\{0, \frac{1}{3}, 1\}$ for $u = \frac{1}{3}$ and $(\frac{4}{21}, \frac{4}{21}, \frac{13}{21})$ on $\{0, \frac{1}{4}, 1\}$ for $u = \frac{1}{2}$. The latter reveals the good type more often but the bad type less often, and since a mean-preserving spread on $[0, 1]$ cannot lower the probabilities of 0 and 1, neither is more informative. The same holds for any two $u \in [1 - p, 2 - 2p)$ if $p \geq \frac{5}{9}$ (see the proof of Proposition 5). Still, on this range, the probability of revealing the good type and the variance increase in u.

Part (ii) shows that the firm's utility is maximized at the least noise at which agents play a binary signal, $\{0, 2p\}$ or $\{0, 1\}$; less noise breaks the binary structure and strictly lowers it. So the optimal noise is intermediate, trading off information from the agents

against accuracy of selection. At least within this family, the firm never benefits from non-binary signals, so she may not need to go far beyond the well S-shaped class.

Part (ii) also compares the best rule in this family with the noiseless rule at every prior. For $p \leq \frac{1}{2}$, the noise $u = 2p$ gives $v(2p) = 1$, so the firm selects the higher of two different signals for sure and noise changes only what agents reveal. Her utility is then $\frac{3p}{2}$, against $\frac{4p}{3}$ under the noiseless rule and $u_F^{ws}(p)$ under the best well S-shaped rule; as $p \to 0$, this is three quarters of the first best, against two thirds. For $p > \frac{1}{2}$, every $u \in [2 - 2p, 1]$ attains the first best. So the right amount of noise beats the noiseless rule at every prior, whereas $v^*(x)$ does so only for $p > 0.21$.

Part (iii) shows that there is no discontinuity at zero noise: as $u \to 0$, $v(x)$ converges pointwise to $\frac{1}{2} + \frac{1}{2} \cdot \text{sign}(x)$, and the equilibrium to the noiseless one of Boleslavsky and Cotton (2015). Simulations show the same for logistic and normal noise.

Two caveats, however, point to robust design. First, the optimal noise is tied to the prior: for $p \leq \frac{1}{2}$, more noise than $u = 2p$ blurs selection and less breaks the binary signals, so a firm that misjudges p loses either way. Second, for $u < 1$ I do not prove that the equilibrium is unique, so this comparison rests on the equilibrium of Proposition 14, to which the equilibria of the discretized game converge. A firm that does not know the priors exactly thus needs a robust design, the subject of Subsection 4.3.

4.3 Attaining the first best

Subsections 4.1 and 4.2 found the best rule within the well S-shaped class and the optimal level of uniform noise, and both depend on the prior. This subsection lets the firm choose any rule of the signal gap and asks which rules attain the first best, first for a known prior and then robustly, for every prior.

I say that $v(x)$ is *revelation-accurate* if $v(1) = 1$, i.e. whenever agent i is revealed to be good and agent j bad ($s_i = 1$, $s_j = 0$), the firm selects agent i with probability 1. This is

a simple necessary condition for optimality: if $v(1) < 1$, an affine modification of $v(x)$ leads to the same equilibria but gives the firm a strictly higher utility.

I also say that $v(x)$ *implements a fully revealing equilibrium (uniquely) for* (p_1, p_2) if, for these priors, full revelation by both agents is an (the unique) equilibrium.

The first result shows that, for symmetric agents, a fully revealing equilibrium and revelation accuracy are not only sufficient, but also necessary for the first best.

Lemma 2. *For every selection rule $v(x)$ and priors (p, p), the firm's utility in every equilibrium is at most $u_F^*(p)$, and some equilibrium attains it if and only if $v(x)$ implements a fully revealing equilibrium for (p, p) and is revelation-accurate.*

The proof is simple. For priors (p, p),

$$u_F^*(p) = \max_{G_1, G_2 \in \mathcal{G}(p)} \mathbb{E}_{G_1, G_2}[\max(s_1, s_2)].$$

As $\max(s_1, s_2)$ is convex and piecewise linear, the maximum is attained if and only if at least one agent plays full revelation. As all equilibria form a symmetric product set, if full revelation and some other signal form an equilibrium, so do two full revelations. Since $s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) \leq \max(s_1, s_2)$, the firm falls short of $u_F^*(p)$ in every equilibrium if $v(x)$ does not implement a fully revealing equilibrium, and in a fully revealing equilibrium, the firm gets the first best if and only if $v(x)$ is revelation-accurate. The result generally fails for $p_1 \neq p_2$: the set of equilibria is then not symmetric and may contain only equilibria in which exactly one agent reveals fully.

Against a fully revealing rival, an agent with signal x wins with probability $\frac{1}{2} + (1-p)\hat{v}(x) - p\hat{v}(1-x)$, where $\hat{v}(x) := v(x) - \frac{1}{2}$, so $v(x)$ implements a fully revealing equilibrium for (p, p) if and only if this function lies below its chord on $[0, 1]$:

$$(1-p)\hat{v}(x) - p\hat{v}(1-x) \leq (x-p)\hat{v}(1) \quad \forall x \in (0, 1).$$

Except for $p = \frac{1}{2}$, the rules satisfying this system have no simple structure, and the assumption that the firm knows the priors exactly is restrictive. Instead, suppose that the firm only knows that the priors lie in some exogenous set S.

Although a fully revealing equilibrium is not necessary for the first best under unequal priors, this approach covers that case too: a rule implements a fully revealing equilibrium for all pairs of priors in S if and only if it does so for all equal priors in S (Lemma 10 in Appendix B). Robust implementation strongly restricts $v(x)$.

Proposition 6. *Let a selection rule $v(x)$ implement a fully revealing equilibrium for (p_l, p_l) and (p_h, p_h), where $p_l < \frac{1}{2} < p_h$. Then*

(i) $v(x)$ implements a fully revealing equilibrium for all $(p_1, p_2) \in (0,1)^2$;

(ii) $v(x) = \frac{1}{2} + ax$, where $a \in [0, \frac{1}{2}]$.

To see why, note that the system of inequalities above holds for p_l and p_h. Adding it at x and at $1-x$ with weights $1-p$ and p gives $(1-2p)\big(\hat{v}(x) - x\hat{v}(1)\big) \leq 0$ for all $x \in (0,1)$. So p_l rules out any upward deviation of $\hat{v}(x)$ from the linear function $x\hat{v}(1)$, and p_h rules out any downward one. Hence, $v(x) = \frac{1}{2} + x(v(1) - \frac{1}{2})$ is linear, so it implements full revelation for all equal priors, and by Lemma 10 for all pairs of priors, and any $a = v(1) - \frac{1}{2} \in [0, \frac{1}{2}]$ is valid.

In particular, as the first best requires revelation accuracy (Lemma 2), a firm that attains her first best at one prior below $\frac{1}{2}$ and at one above must use $v(x) = \frac{1}{2} + \frac{1}{2}x$. But this rule has a crucial issue. Since $v(x)$, and hence $\overline{v}(s_i)$, is linear, every pair of signals is an equilibrium, and the firm need not get the first best: robustness comes at the cost of complete indifference. This indifference is fragile: any positive signal cost makes pooling the unique equilibrium (Proposition 16 in Appendix B.11).

The issue disappears under slightly less robustness. Suppose that the firm knows whether both agents are more or less likely to be good than bad. The rules implementing

a fully revealing equilibrium in either case are described by the deviation of $\hat{v}(x)$ from its secant from 0 to 1, i.e., by the function g below.

Definition 3. Let $v(x)$ be a selection rule, $a := v(1) - \frac{1}{2}$, and $g(x) := v(x) - \frac{1}{2} - ax$. Then $v(x)$ is *balanced* if $g(x) = g(1-x)$ on $(0,1)$; *positively* [*negatively*] *balanced* if also $g(x) \geq 0$ $[\leq 0]$ there; and *strictly* so if $g(x) > 0$ $[< 0]$.

Like $\tilde{v}(x)$, $g(x)$ removes a linear part of $v(x)$, but it subtracts the secant from 0 to 1 rather than the tangent at 0. So equilibria depend on $v(x)$ only through $g(x)$, and revelation accuracy only through a; a well S-shaped $v(x)$ is positively balanced if and only if $\tilde{v}(x) = -x^2$ on $[0,1]$. Balance alone is exactly the condition for a fully revealing equilibrium at $p = \frac{1}{2}$, as shown after Proposition 7; the sign of $g(x)$ then determines the half-interval of priors on which full revelation extends. Every selection rule has $a \in [0, \frac{1}{2}]$ and $g(0) = g(1) = 0$, and a balanced $v(x)$ is increasing if and only if $|g(x) - g(y)| \leq a|x - y|$ for all $x, y \in [0,1]$.

Proposition 7. *Let $v(x)$ be a selection rule.*

(i) $v(x)$ implements a fully revealing equilibrium for all $(p_1, p_2) \in [\frac{1}{2}, 1)^2$ [for all $(p_1, p_2) \in (0, \frac{1}{2}]^2$] if and only if it is positively [negatively] balanced.

(ii) If $v(x)$ is strictly positively [negatively] balanced, it implements a fully revealing equilibrium uniquely $\forall (p_1, p_2) \in (\frac{1}{2}, 1)^2$ $\left[\forall (p_1, p_2) \in (0, \frac{1}{2})^2\right]$.

Part (i) gives necessary and sufficient conditions for implementing full revelation for all priors on a half-interval. They impose a structure: $v(x)$ can deviate from the universal implementation $v(x) = \frac{1}{2} + ax$ only by a function that is symmetric around $\frac{1}{2}$ on $[0,1]$, non-negative for high priors and non-positive for low ones.

Indeed, the conditions hold for all $p \in (0, \frac{1}{2}]$ if and only if they hold for $p = 0$ and $p = \frac{1}{2}$, and the latter gives the symmetry: against a fully revealing rival with $p = \frac{1}{2}$,

agent i's value function $\frac{1}{2}\hat{v}(s_i) - \frac{1}{2}\hat{v}(1-s_i)$ is odd around $s_i = \frac{1}{2}$, so $\{0,1\}$ is a best reply only if it is linear, equal to $(s_i - \frac{1}{2})\hat{v}(1)$, i.e., only if $v(x)$ is balanced.

Part (ii) gives a sufficient condition for full revelation to be the unique equilibrium. As all equilibria form a product set, it suffices that $\{0,1\}$ is each agent's unique best reply to itself. This is also why $p = \frac{1}{2}$ is excluded: there, any $v(x)$ that implements full revelation makes $\overline{v}(s_i)$ linear, so the best reply is not unique. The condition is not necessary, but since it is only a slight strengthening of (i), the room for other rules with a unique fully revealing equilibrium is small, and checking uniqueness for them requires showing that no signal supported on $\{s_i : g(s_i) = 0\}$ is a max-minimizer. A strictly balanced rule whose $g(x)$ has a non-zero slope at 0 also survives slightly costly signals: for priors not too close to $\frac{1}{2}$, full revelation remains the unique equilibrium (Proposition 16), whereas under the linear rule any cost leads to pooling.

A firm facing a high prior needs a positively balanced rule, and some subclasses of these rules can be generated by i.i.d. shocks. A firm facing a low prior, however, needs a negatively balanced rule, and additive evaluation noise cannot deliver one: by the next result, which extends this consequence of Proposition 1 beyond the well S-shaped class, i.i.d. shocks rule out full revelation as soon as one prior is below $\frac{1}{2}$.

Corollary 2. *If $v'(0) > v'(1)$ and $\min(p_1, p_2) < \frac{1}{2}$, then $(\{0,1\};\{0,1\})$ is not an equilibrium. In particular, this holds for every $v(x)$ generated by i.i.d. shocks with a square-integrable density.*

4.4 A benchmark: general selection rules

Both limitations, indifference under full robustness and the failure of i.i.d. noise for low priors (Corollary 2), are specific to rules of the signal gap. To see how much they cost, consider a designer who may use any rule of the following kind.

Definition 4. A *general selection rule* is a measurable $H : [0,1]^2 \to [0,1]$ with $H(x,y) + H(y,x) = 1$ for all $x, y \in [0,1]$, so that $H(x,x) = \frac{1}{2}$: agent i is selected with probability

$H(s_i, s_j)$. Rules of the signal gap are the case $H(x, y) = v(x - y)$.

Neither continuity nor monotonicity in the own signal is required, and neither restriction would cost the firm anything. Every rule that attains a bound in this section or the next is nondecreasing in the own signal, and the bounds hold among all rules: if a value function decreases somewhere on the support, pooling the lowest signals makes it nondecreasing and weakly raises the firm's payoff. The optimal rules of Section 5 are discontinuous at ties, but smoothing them near ties gives continuous rules whose equilibrium payoffs converge to the same bounds (see Appendix C.8).

Many such rules have no natural interpretation, so I treat them as a benchmark. It is demanding in two ways. The designer commits to any rule, however unnatural ex post, and she measures each agent's signal, the posterior mean of his type, perfectly: the rule is applied to s_i itself rather than to a noisy reading of it. The preceding subsections showed that eliminating all noise from the measurement can hurt the firm. The benchmark asks the converse question: how much could a firm gain if it could measure perfectly and commit to any use of the measurement? The answer never requires an unnatural rule, because the optimum can be sought within a simple class. With binary types, a deterministic pass/fail rule attains the first best (Proposition 8). With a general prior, value functions can be taken nondecreasing without loss, every feasible such function is produced by a lottery over coarse gradings, and for concave priors the optimal one is produced by a single random pass mark (Subsection 5.2).

Proposition 8. *Let types be binary and H be nondecreasing in the own signal.*

(i) H attains the first best $2p - p^2$ in some equilibrium for all priors (p, p) with $p \in (0, 1)$ if and only if $H(1, 0) = 1$, $H(x, 0) \leq \frac{1+x}{2}$ and $H(x, 1) \leq \frac{x}{2}$ for all $x \in (0, 1)$. If the last two inequalities are strict, then for all priors $(p_1, p_2) \in (0, 1)^2$ the firm attains the first best in the unique equilibrium, which is fully revealing.

(ii) In particular, the pass/fail rule*, which selects an agent whose signal is* 1 *over one whose signal is below* 1 *and tosses a coin otherwise, attains the first best in the unique equilibrium for all* $(p_1, p_2) \in (0,1)^2$.

Against a fully revealing rival with prior p, a signal x wins with probability $(1-p)H(x,0)+p\,H(x,1)$, and full revelation is a best reply if and only if no intermediate signal wins more often than the chord $\frac{1-p+x}{2}$ through the winning probabilities at 0 and 1. This condition is linear in p, and its limits as $p \to 0$ and $p \to 1$ are the two inequalities in (i). For a rule of the signal gap, they read $v(x) \leq \frac{1+x}{2}$ and $v(1-x) \geq 1-\frac{x}{2}$, which together force the linear rule, as in Proposition 6. A general rule can satisfy both strictly: the rule $\frac{1}{2}+\frac{1}{2}(s_i^2 - s_j^2)$ does, and so does the pass/fail rule.

The pass/fail rule requires a strong commitment. Ex post, it ignores what the firm has learned: between an agent with signal 0.95 and one with signal 0.05, it tosses a coin, which is as noisy as selection can be. Ex ante, this is exactly what makes revelation pay: a partial signal, however close to 1, buys nothing over revealing a bad type. So, with binary types, the firm has two ways to curb concealment: she can commit to disregard every signal short of conclusive evidence, or she can prefer, ex ante, some deliberate noise in her observations, which is what a noisy rule of the signal gap amounts to. An imperfect measure can thus be just as good as a perfect one: the linear rule of the signal gap attains the benchmark's first best too. With binary types, the benchmark is thus trivial; Section 5 relaxes the binary prior and shows that with a concave prior the benchmark's own optimum deliberately blurs the measured signals.

5. Non-Binary Priors

With a general prior F, a signal G is feasible if and only if it is a mean-preserving contraction of F (Gentzkow and Kamenica, 2016, Remark 1 and Proposition 1), i.e. $\mathbb{E}_G[\phi] \leq \mathbb{E}_F[\phi]$ for every convex ϕ; the binary prior, the largest distribution on $[0,1]$

with mean p in the convex order, gives agents the richest set of feasible signals. I write $\underline{\theta} = \min \text{supp}(F)$ and $\overline{\theta} = \max \text{supp}(F)$, and count the noiseless rule $v(x) = \frac{1}{2} + \frac{1}{2} \cdot \text{sign}(x)$ among the selection rules of the signal gap.

With a general prior, two posteriors with the same mean can differ, for instance in how conclusive they are. Throughout, selection rules depend on the signals only through the posterior means, the standard restriction when the receiver's payoff is linear in the state (Gentzkow and Kamenica, 2016; Kolotilin, 2018; Dworczak and Martini, 2019; Kolotilin et al., 2022), as the firm's is here. For the benchmark designer of Subsection 4.4 the restriction has bite: a rule that prefers degenerate posteriors to non-degenerate ones makes full revelation weakly dominant and attains the first best for every prior. Subsection 5.1 keeps rules of the signal gap, and Subsection 5.2 returns to the benchmark designer of Subsection 4.4, who may choose any general rule.

5.1 Rules of the signal gap

Under a well S-shaped rule, an agent's value function is S-shaped against every signal of his rival, so, for a prior with a positive density, an upper censorship is a best reply (for smooth value functions, Kolotilin et al., 2022, Theorem 1): it reveals the types below a cutoff and pools all types above it into one signal. I extend this result to arbitrary priors, with atoms and gaps, by defining censorship on quantiles: the *quantile censorship* G_q reveals the lowest share q of types and pools the rest at their mean, splitting an atom at the cutoff if needed. Against every rival signal, some quantile censorship is a best reply (Lemma 13), so the equilibrium keeps the structure of Section 3: low types are revealed, and high types are pooled. Theorem 2 also extends to every prior, and accuracy and revelation do not conflict: the rule that ranks most accurately at every signal gap (Lemma 9) also reveals the most.

Proposition 9. *Let $v(x)$ be well S-shaped and F any prior. Then (G_q, G_q) is an equilibrium*

for some $q \in (0, \frac{1}{2}]$*, and the unique equilibrium if* $v'(x)$ *is not linear near* 0*. In every symmetric equilibrium in quantile censorships,* $q \leq \frac{1}{2}$*, and the firm's utility is at most its value in the median-censorship equilibrium of* $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$*.*

For any prior, the equilibrium is a quantile censorship that either solves a first-order condition, (QC′) in Appendix C, or reveals fully, which requires an atom at $\overline{\theta}$ of mass at least $1 - q \geq \frac{1}{2}$ (Lemmas 12 and 14). For the binary prior, the agents reveal fully exactly when the atom at 1 has mass $p \geq \overline{p}$, where $\overline{p} \geq \frac{1}{2}$ (Theorem 1). For $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$, the best well S-shaped rule of Theorem 2, the first-order condition collapses: $q = \frac{1}{2}$ for every prior, so each agent reveals the types below the median and pools those above it, splitting an atom at the median so that exactly half the mass is revealed (Appendix C.3).

5.2 General selection rules with a general prior

From here on, the firm chooses any general selection rule H of Definition 4, as in Proposition 8; rules of the signal gap are a special case, and *rule* means a general selection rule throughout. Against a rival who plays G, the agent's value function is $V_G(x) := \int H(x, y)\, dG(y)$, his probability of being selected with signal x.[4]

In a symmetric profile (G, G), the firm's payoff is $u_F = p + 2\,\mathrm{Cov}_G\big(s, V_G(s)\big)$, so she cares only about how steeply an agent's winning probability rises with his signal. Let V_F be the value function against a fully revealing rival, whose signal is his type. A function V equals V_F on $\mathrm{supp}(F)$ for some rule if and only if $V(\theta) \leq_{cx} U[0, 1]$ for $\theta \sim F$ (Appendix C.8): noise only makes selection probabilities less dispersed than exact ranking, and the noiseless rule, with $V_F = F$ for atomless F, is the boundary case.

[4]Without continuity an equilibrium need not exist; when equilibria exist, they are interchangeable and give each agent $\frac{1}{2}$, and I evaluate a rule at its firm-preferred equilibrium.

5.2.1 The first best

Proposition 10. *Let* $\text{supp}(F) = [\underline{\theta}, \overline{\theta}]$. *Some general selection rule nondecreasing in the own signal attains the first best in some equilibrium if and only if* F *is convex on* $[\underline{\theta}, \overline{\theta})$ *and has no atom below* $\overline{\theta}$. *Then the noiseless rule attains it, with full revelation as the unique equilibrium.*

A noisy rule of the signal gap selects the higher of two close signals with probability near one half, so for convex priors the comparison of Subsection 4.1 reverses. More generally, every equilibrium that attains the first best is fully revealing, so a rule attains it only if it coincides with the noiseless rule for $F \otimes F$-almost every pair of signals (Appendix C.8.3). For priors with a positive density, Hwang et al. (2026, Lemma 1, Theorem 1 and Corollary 1) show that full revelation is an equilibrium of the noiseless rule if and only if F is convex, and that it is then the unique symmetric equilibrium.

5.2.2 The best fully revealing rule

Define the affine value function

$$L(x) := \frac{1}{2} + \frac{x - p}{2(\overline{\theta} - p)}, \qquad \text{so } L(p) = \tfrac{1}{2},\ L(\overline{\theta}) = 1,$$

and let $\bar{U} := U[2p - \overline{\theta}, \overline{\theta}]$, the uniform distribution with the prior's mean and top.

Proposition 11. *Let* F *be any prior. For every general selection rule with upper semicontinuous* V_F *under which full revelation is an equilibrium,*

$$u_F \leq p + \frac{\text{Var}_F(\theta)}{\overline{\theta} - p},$$

with equality if and only if $V_F = L$ F*-almost surely, which is feasible if and only if* $F \leq_{cx} \bar{U}$.

So the best fully revealing rule is *the narrowest uniformizing noise*: when agents reveal fully, it makes their scores uniform on the narrowest interval that still contains the

prior's top and spreads the prior. This is the logic of the example in the introduction, where uniform shocks make the scores of fully revealing agents uniform. Every concave prior satisfies $F \leq_{cx} \bar{U}$; it fails for non-uniform convex priors, for which the noiseless rule attains the first best anyway.[5]

When $F \leq_{cx} \bar{U}$ fails, the bound is not attained. For a single-peaked prior, $F \leq_{cx} \bar{U}$ if and only if $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$; otherwise, the best fully revealing rule ranks weak types exactly and applies the narrowest uniformizing noise only above a cutoff (Appendix C.8.7).

Insisting on full revelation can be costly. For Beta$(2, 2)$, $p = \frac{1}{2}$, the bound is attained, and the best fully revealing rule gives 0.600, below the 0.626 of the noiseless rule.

The rule attaining the bound can be chosen nondecreasing in the own signal. Every feasible nondecreasing value function, L included, is produced by a lottery over coarse gradings: the firm draws a partition of the signals into intervals, selects the agent in the higher interval, and tosses a coin within an interval (Appendix C.8). So the benchmark designer loses nothing by restricting herself to such simple rules. General rules also avoid the indifference of Proposition 6.

The rule H^* that draws a bar uniformly between the lower signal and 1, selects the higher signal if it clears the bar, and tosses a coin otherwise makes full revelation the unique equilibrium for every prior. Among the rules under which full revelation is an equilibrium for every prior, it selects the higher signal with the highest probability at every pair of signals, so it is the best such rule for every prior (Appendix C.8.8).

5.2.3 The second best

Under the best fully revealing rule, scores are uniform on $[2p - \overline{\theta}, \overline{\theta}]$. Replacing the top $\overline{\theta}$ by any $\bar{s} \geq 2p - \underline{\theta}$ gives a family of rules. Let $\bar{U}_{\bar{s}} := U[2p - \bar{s}, \bar{s}]$ and $L_{\bar{s}}(x) := \frac{1}{2} + \frac{x-p}{2(\bar{s}-p)}$, its CDF on its support. For $\bar{s} < \overline{\theta}$, the prior cannot be spread into $\bar{U}_{\bar{s}}$, so the agents must

[5] A concave prior has a nonincreasing density, possibly unbounded at $\underline{\theta}$, and possibly an atom at $\underline{\theta}$.

pool some types. Let $G^*_{\bar{s}}$ be the most dispersed distribution that is a mean-preserving contraction of both F and $\bar{U}_{\bar{s}}$ (Appendix C.8). The *uniformizing rule of width* $\bar{s}$ adds noise that spreads $G^*_{\bar{s}}$ into $\bar{U}_{\bar{s}}$, so a signal x beats a rival drawn from $G^*_{\bar{s}}$ with probability $L_{\bar{s}}(x)$.

Proposition 12. *For every $\bar{s} \geq 2p - \underline{\theta}$, the uniformizing rule of width $\bar{s}$ has an equilibrium $(G^*_{\bar{s}}, G^*_{\bar{s}})$ with $u_F(\bar{s}) = p + \mathrm{Var}(G^*_{\bar{s}})/(\bar{s} - p)$. If $\bar{s} \in (2p - \underline{\theta}, \overline{\theta})$ and the set $\{x \in [\underline{\theta}, \bar{s}] : F(x) > L_{\bar{s}}(x)\}$ is an interval, as it is for concave F, then $G^*_{\bar{s}}$ reveals the types below some t_1, puts no mass on some (t_1, t_2), and has the density of $\bar{U}_{\bar{s}}$ on $[t_2, \bar{s}]$.*

Corollary 3. *Let F be concave on its support $[\underline{\theta}, \overline{\theta}]$. Then (G, G) with $G = U[\underline{\theta}, 2p - \underline{\theta}]$ is the unique equilibrium of the noiseless rule, and the firm's utility is $u_F = \frac{4p - \underline{\theta}}{3}$.*

For concave priors, the family runs between two familiar ends. At width $2p - \underline{\theta}$, the uniformizing rule adds no noise and is the noiseless rule, whose equilibrium Corollary 3 gives: a suitable uniform distribution is always feasible when the density is decreasing. For priors with a density, this symmetric equilibrium is also in Hwang et al. (2026, Corollary 1). At width $\overline{\theta}$, the equilibrium is fully revealing and gives the best fully revealing rule of Proposition 11. In between, agents reveal their low types and spread their high types uniformly, as all types do under noiseless competition.

This equilibrium has the shape of the censorship equilibrium of Proposition 9, except that a uniformizing rule replaces the concave part of the S by a straight line, so the pooled types are spread uniformly rather than sent to one signal.

Under the noiseless rule, the firm's payoff depends on the prior only through its mean and its lower support point, so it does not grow with dispersion, while the first best does. So either end of the family can win, as the following example shows.

Example 5. Let $p = \frac{1}{3}$ and $\mathrm{supp}(F) = [0, 1]$, and compare $F(x) = 2x - x^2$ with its mean-preserving spread $F(x) = \frac{1+2x}{3}$, an atom of $\frac{1}{3}$ at 0 plus a uniform density. Their

variances are $\frac{1}{18}$ and $\frac{1}{9}$. The noiseless rule gives $\frac{4}{9} \approx 0.444$ under both; the best fully revealing rule gives 0.417 and 0.500; and $v^*(x)$ of Theorem 2 gives 0.399 and 0.457.

Neither end of the family is optimal, and for concave priors, and more generally for the priors of the next result, the family beats every rule.

Theorem 3. *Let the prior F satisfy*

(i) $U[\underline{\theta}, 2p - \underline{\theta}] \leq_{cx} F$;

(ii) $F \leq_{cx} \bar{U}$;

(iii) for every $\bar{s} \in (2p - \underline{\theta}, \overline{\theta})$, the set $\{x \in [\underline{\theta}, \bar{s}] : F(x) > L_{\bar{s}}(x)\}$ is an interval.

In every symmetric equilibrium of every rule,

$$u_F \leq \max_{\bar{s} \in [2p - \underline{\theta}, \overline{\theta}]} u_F(\bar{s}).$$

If, moreover, $I_{U[\underline{\theta}, 2p-\underline{\theta}]} < I_F < I_{\bar{U}}$ on $(\underline{\theta}, \overline{\theta})$, the maximum is attained only at widths in $(2p - \underline{\theta}, \overline{\theta})$, so the best uniformizing rule strictly beats both ends of the family, the noiseless rule and the best fully revealing rule (Appendix C.8.10). Every concave prior satisfies (i)–(iii), and every non-uniform one satisfies these strict inequalities. Condition (iii) holds whenever F is single-peaked or U-shaped, and for a single-peaked prior, (ii) reduces to $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$ (Lemma 25).[6]

For the two priors of Example 5, the second best is 0.456 and 0.509, attained at widths $\bar{s} \approx 0.73$ and 0.94. At both ends the logic is first order against second order. Near full revelation, narrowing the width steepens winning probabilities, a first-order

[6]Condition (ii) cannot be dropped. Beta$(\frac{1}{2}, \frac{1}{2})$ satisfies (i) and (iii), and every uniformizing rule gives it at most 0.6976. Revealing the types below 0.371 and above 0.807 and spreading $(0.371, 0.807)$ uniformly over $[0.404, 0.788]$ gives 0.6987 in a symmetric equilibrium of a suitable rule, and no rule gives more than 0.7015 in a symmetric equilibrium (the first best is 0.7026).

gain, while agents pool only a sliver of top types; near the noiseless rule, widening it lets agents spread posteriors further, while only a sliver of the lowest types is revealed.

For concave priors, the best uniformizing rule is a *random pass mark*: the firm draws a bar from a fixed distribution that never exceeds t_2, selects the agent with the higher signal if it clears the bar, and tosses a coin otherwise. Agents who pool into the top range always clear it and are compared exactly; only revealed low types face the coin. For non-uniform concave F its equilibrium is unique, so every equilibrium gives the second best (Appendix C.8). For the other priors of Theorem 3, a pass mark may fail, but a lottery over coarse gradings always implements it (Appendix C.8.9). With binary types, a deterministic coarse grade attains the first best; with a continuum of types, coarse grading discards real distinctions.

5.2.4 When noise helps

Theorem 3 requires the noiseless equilibrium signal to be uniform.[7] In general, for a prior with a continuous positive density, the noiseless rule has at most one symmetric equilibrium (G_0, G_0): G_0 is convex with support $[\underline{\theta}, c]$, reveals the types in the contact set $E_0 := \{x \in [\underline{\theta}, c] : I_{G_0}(x) = I_F(x)\}$, and is uniform on each component of $(\underline{\theta}, c) \setminus E_0$, a *pool* (Lemma 29; for regular priors, Hwang et al., 2026, Theorem 1).

Definition 5. Let F be atomless, with $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$ and a density f that is continuous and positive on $(\underline{\theta}, \overline{\theta})$. Its noiseless equilibrium is *well behaved* if

(H1) the noiseless rule has a symmetric equilibrium (G_0, G_0); let $c := \max \operatorname{supp}(G_0)$ and $E_0 := \{x \in [\underline{\theta}, c] : I_{G_0}(x) = I_F(x)\}$;

(H2) $[\underline{\theta}, c]$ is the union of finitely many intervals, each of which is either a *revealed stretch*, a nondegenerate interval contained in E_0 on which F is twice continuously

[7] For Beta$(2, 5)$, $U[\underline{\theta}, 2p - \underline{\theta}] \leq_{cx} F$ fails, as the noiseless rule reveals the types below 0.110. The uniformizing rule of width $2p - \underline{\theta}$ still beats the noiseless rule, 0.3677 against 0.3672, but other rules beat the whole uniformizing family numerically for this prior.

differentiable and strictly convex, or a *block*, an interval on which G_0 is affine and which meets E_0 in finitely many points;

(H3) if $x_0 \in E_0$ is an endpoint of a component of $(\underline{\theta}, c) \setminus E_0$ on which G_0 is affine with slope γ, then $f \geq \gamma$ a.e. on a one-sided neighbourhood of x_0 in that component.

(H1) holds for regular priors (Hwang et al., 2026, Theorem 1), for convex and concave priors, and whenever $U[\underline{\theta}, 2p - \underline{\theta}] \leq_{cx} F$. (H2) and (H3) rule out only pathologies: infinitely many alternations of revealed and pooled types, and densities that oscillate infinitely often around the slope of a pool.

Theorem 4. *Let F have a well-behaved noiseless equilibrium (G_0, G_0). The noiseless rule is optimal among general selection rules, i.e., no rule gives the firm a higher utility in any equilibrium, if and only if F is convex on its support. If F is not convex, some general selection rule nondecreasing in the own signal has a symmetric equilibrium in which the firm's utility strictly exceeds its utility in (G_0, G_0).*

The improving rule perturbs the noiseless equilibrium on one run of adjacent pools. A pool at the bottom is widened by a little noise; a pool after revealed types spreads when noise among the strongest revealed types frees winning probability; where two pools meet, value moves from the upper pool to the lower one. The gain is first order in the amount of noise, while the cost, from keeping the spread signals contractions of F, is of higher order (Appendix C.8.14). In the Beta and power priors I computed, the gain is about 10^{-2} when the noiseless rule pools from the bottom, e.g., 0.041 for Beta$(\frac{1}{2}, 1)$, and mostly between 10^{-5} and $4 \cdot 10^{-3}$ when it reveals a bottom interval.

6. Conclusion

This paper asks whether a firm that chooses between two agents who design their own evidence should eliminate the noise in her evaluation. Under well S-shaped noise,

agents concede bad news and pool the rest, and the shape of the noise, summarized by a single elasticity, decides how much they reveal. The right amount of noise beats the noiseless rule and, with binary types, attains the first best at priors of at least one half. With a continuum of types, the case for noise survives even for a designer who measures signals perfectly and can commit to any rule: under mild regularity conditions, her best rule blurs the signals unless the prior is convex, and for concave priors it is a random pass mark that randomizes only among the weakest agents.

Appendix D shows how these conclusions extend. Under the mirror-image, reverse S-shaped rules, agents reveal good news instead, the favorite and the underdog switch roles, and better agents can hurt the firm. Beyond the well S-shaped class, the equilibrium stays unique with binary signals under a weaker condition on the noise, while further away agents randomize over several posteriors. With a single agent, noise works through the same channel, devaluing a narrow lead over an outside option, but there a firm that could commit to any rule would not need it.

References

Abdulkadiroğlu, A. and T. Sönmez (2003). School choice: a mechanism design approach. *American Economic Review 93*(3), 729–747.

Au, P. H. and K. Kawai (2020). Competitive information disclosure by multiple senders. *Games and Economic Behavior 119*, 56–78.

Au, P. H. and M. Whitmeyer (2023). Attraction versus persuasion: Information provision in search markets. *Journal of Political Economy 131*(1), 202–245.

Ball, I. (2025). Scoring strategic agents. *American Economic Journal: Microeconomics 17*(1), 97–129.

Banerjee, S., D. Kempe, and R. Kleinberg (2021). Threshold tests as quality signals:

Optimal strategies, equilibria, and price of anarchy. In *Web and Internet Economics (WINE 2021)*. arXiv:2110.10881.

Banerjee, S., K. Munagala, Y. Shen, and K. Wang (2026). The price of competitive information disclosure. Working paper, arXiv:2504.10459.

Bar, T., V. Kadiyali, and A. Zussman (2012). Putting grades in context. *Journal of Labor Economics 30*(2), 445–478.

Bergemann, D., T. Gan, and Y. Li (2023). Managing persuasion robustly: The optimality of quota rules. Working paper, arXiv:2310.10024.

Blackwell, D. (1953). Equivalent comparisons of experiments. *The Annals of Mathematical Statistics 24*(2), 265–272.

Blume, A., O. J. Board, and K. Kawamura (2007). Noisy talk. *Theoretical Economics 2*(4), 395–440.

Board, S. and J. Lu (2018). Competitive information disclosure in search markets. *Journal of Political Economy 126*(5), 1965–2010.

Boleslavsky, R. and C. Cotton (2015). Grading standards and education quality. *American Economic Journal: Microeconomics 7*(2), 248–279.

Boleslavsky, R. and C. Cotton (2018). Limited capacity in project selection: Competition through evidence production. *Economic Theory 65*(2), 385–421.

Boleslavsky, R. and C. Li (2026). Evaluating university graduates: Prestige versus merit. *Journal of Economics & Management Strategy*. Forthcoming.

Border, K. C. and J. Sobel (1987). Samurai accountant: a theory of auditing and plunder. *Review of Economic Studies 54*(4), 525–540.

Chan, W., H. Li, and W. Suen (2007). A signaling theory of grade inflation. *International Economic Review 48*(3), 1065–1090.

Che, Y.-K. and I. Gale (2000). Difference-form contests and the robustness of all-pay auctions. *Games and Economic Behavior 30*(1), 22–43.

Curello, G. and L. Sinander (2025). The comparative statics of persuasion. Working paper, arXiv:2204.07474.

Ding, B., Y. Feng, C.-J. Ho, W. Tang, and H. Xu (2023). Competitive information design for Pandora's box. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*. SIAM.

Drugov, M. and D. Ryvkin (2020). How noise affects effort in tournaments. *Journal of Economic Theory 188*, 105065.

Dworczak, P. and G. Martini (2019). The simple economics of optimal persuasion. *Journal of Political Economy 127*(5), 1993–2048.

Ederer, F., R. Holden, and M. Meyer (2018). Gaming and strategic opacity in incentive provision. *RAND Journal of Economics 49*(4), 819–854.

Erdil, A. and H. Ergin (2008). What's the matter with tie-breaking? Improving efficiency in school choice. *American Economic Review 98*(3), 669–689.

Fang, D. and T. Noe (2022). Less competition, more meritocracy? *Journal of Labor Economics 40*(3), 669–701.

Fang, F. C. and A. Casadevall (2016). Research funding: the case for a modified lottery. *mBio 7*(2), e00422–16.

Frankel, A. and N. Kartik (2022). Improving information from manipulable data. *Journal of the European Economic Association 20*(1), 79–115.

Fu, Q. and Z. Wu (2019). Contests: Theory and topics. In *Oxford Research Encyclopedia of Economics and Finance*. Oxford University Press.

Gao, E. and D. Luo (2026). (Robust) information acquisition design. Working paper, arXiv:2312.02465.

Gentzkow, M. and E. Kamenica (2016). A Rothschild-Stiglitz approach to Bayesian persuasion. *American Economic Review 106*(5), 597–601.

Gentzkow, M. and E. Kamenica (2017). Competition in persuasion. *The Review of Economic Studies 84*(1), 300–322.

Grossman, S. J. and J. E. Stiglitz (1980). On the impossibility of informationally efficient markets. *American Economic Review 70*(3), 393–408.

Harbaugh, R. and E. Rasmusen (2018). Coarse grades: Informing the public by withholding information. *American Economic Journal: Microeconomics 10*(1), 210–235.

He, W. and J. Li (2023). Competitive information disclosure in random search markets. *Games and Economic Behavior 140*, 132–153.

Heyard, R., M. Ott, G. Salanti, and M. Egger (2022). Rethinking the funding line at the Swiss National Science Foundation: Bayesian ranking and lottery. *Statistics and Public Policy 9*(1), 110–121.

Hirshleifer, J. (1989). Conflict and rent-seeking success functions: Ratio vs. difference models of relative success. *Public Choice 63*(2), 101–112.

Hoppe, H. C., B. Moldovanu, and A. Sela (2009). The theory of assortative matching based on costly signals. *Review of Economic Studies 76*(1), 253–281.

Hvide, H. K. and E. G. Kristiansen (2003). Risk taking in selection contests. *Games and Economic Behavior 42*(1), 172–179.

Hwang, I., K. Kim, and R. Boleslavsky (2026). Competitive advertising and pricing. *The Review of Economic Studies*. Forthcoming.

Jain, V. and M. Whitmeyer (2026). Competitive disclosure of information to a rationally inattentive agent. *Journal of Economic Behavior & Organization 242*, 107410.

Kamenica, E. (2019). Bayesian persuasion and information design. *Annual Review of Economics 11*, 249–272.

Kamenica, E. and M. Gentzkow (2011). Bayesian persuasion. *American Economic Review 101*(6), 2590–2615.

Karlin, S. (1959). *Mathematical Methods and Theory in Games, Programming, and Economics, Vol. II: The Theory of Infinite Games*. Reading, MA: Addison-Wesley.

Kleiner, A., B. Moldovanu, and P. Strack (2021). Extreme points and majorization: Economic applications. *Econometrica 89*(4), 1557–1593.

Kolotilin, A. (2018). Optimal information disclosure: A linear programming approach. *Theoretical Economics 13*(2), 607–635.

Kolotilin, A., T. Mylovanov, and A. Zapechelnyuk (2022). Censorship as optimal persuasion. *Theoretical Economics 17*(2), 561–585.

Kolotilin, A., T. Mylovanov, A. Zapechelnyuk, and M. Li (2017). Persuasion of a privately informed receiver. *Econometrica 85*(6), 1949–1964.

Lazear, E. P. and S. Rosen (1981). Rank-order tournaments as optimum labor contracts. *Journal of Political Economy 89*(5), 841–864.

Letina, I., S. Liu, and N. Netzer (2023). Optimal contest design: Tuning the heat. *Journal of Economic Theory 213*, 105616.

Morgan, J., J. Tumlinson, and F. Várdy (2022). The limits of meritocracy. *Journal of Economic Theory 201*, 105414.

Muratov, O. (2025). Mapping an information design game into an all-pay auction. *Review of Economic Design 29*(3), 513–543.

Niculescu, C. P. and L.-E. Persson (2003). Old and new on the Hermite–Hadamard inequality. *Real Analysis Exchange 29*(2), 663–686.

Ostrovsky, M. and M. Schwarz (2010). Information disclosure and unraveling in matching markets. *American Economic Journal: Microeconomics 2*(2), 34–63.

Perez-Richet, E. and V. Skreta (2022). Test design under falsification. *Econometrica 90*(3), 1109–1142.

Pier, E. L., M. Brauer, A. Filut, A. Kaatz, J. Raclaw, M. J. Nathan, C. E. Ford, and M. Carnes (2018). Low agreement among reviewers evaluating the same NIH grant applications. *Proceedings of the National Academy of Sciences 115*(12), 2952–2957.

Polyanin, A. D. and A. V. Manzhirov (2008). *Handbook of Integral Equations* (Second ed.). Boca Raton, FL: Chapman & Hall/CRC.

Popov, S. V. and D. Bernhardt (2013). University competition, grading standards, and grade inflation. *Economic Inquiry 51*(3), 1764–1778.

Prat, A. (2005). The wrong kind of transparency. *American Economic Review 95*(3), 862–877.

Prendergast, C. and R. H. Topel (1996). Favoritism in organizations. *Journal of Political Economy 104*(5), 958–978.

Röbbecke, M. and D. Simon (2023). Riskante Forschung und teilrandomisierte Begutachtungsverfahren: Neue Wege der Förderlinie "Experiment!" der VolkswagenStiftung. *Beiträge zur Hochschulforschung 45*(2), 8–30.

Sakhnovich, L. A. (2015). *Integral Equations with Difference Kernels on Finite Intervals* (Second ed.), Volume 84 of *Operator Theory: Advances and Applications*. Cham: Birkhäuser.

Scharfstein, D. S. and J. C. Stein (2000). The dark side of internal capital markets: Divisional rent-seeking and inefficient investment. *The Journal of Finance 55*(6), 2537–2564.

Stone, P. (2011). *The Luck of the Draw: The Role of Lotteries in Decision Making*. New York: Oxford University Press.

Tsakas, E. and N. Tsakas (2021). Noisy persuasion. *Games and Economic Behavior 130*, 44–61.

Tsakas, E., N. Tsakas, and D. Xefteris (2021). Resisting persuasion. *Economic Theory 72*(3), 723–742.

Tullock, G. (1980). Efficient rent seeking. In J. M. Buchanan, R. D. Tollison, and G. Tullock (Eds.), *Toward a Theory of the Rent-Seeking Society*, pp. 97–112. College Station: Texas A&M University Press.

Wang, Z. (2010). The optimal accuracy level in asymmetric contests. *The B.E. Journal of Theoretical Economics 10*(1), Article 13.

Overview of the Appendices

Appendix A proves the results of Section 3 and collects the details on unequal priors: the boundaries of the regions in Proposition 2, further examples, and how the priors

affect the payoffs of the firm and of the agents. Appendix B proves the results of Section 4, constructs the equilibrium for the triangular selection rules of Subsection 4.2, extends the robust design results to costly signals, and gives the equilibria of the noiseless rule. Appendix C proves the results of Subsection 4.4 and Section 5, starting with censorship under an arbitrary prior. Appendix D collects extensions: reverse S-shaped rules, the boundary of the well S-shaped class, and a single agent.

A. Proofs, Technical Details and Additions for Section 3

A.1 Properties of well S-shaped selection rules

Throughout the proofs, I use the following properties of a well S-shaped $v(x)$.

1) *Regularity.* Since $v'(x)$ is concave, it is continuous on $(-1,1)$ and has one-sided derivatives there; I set $v'(\pm 1)$ equal to its one-sided limits. I write $v''(x)$ and $\tilde{v}''(x)$ for either one-sided derivative, and every statement involving them holds for both.

2) *Shape.* Since $v(x)$ is increasing, $v'(1) \geq 0$, and since $v'(x)$ is even, concave and uniquely maximized at 0, $v'(x) \geq (1-|x|)v'(0) + |x|v'(1) > v'(1) \geq 0$ on $(-1,1)$, so $v(x)$ is strictly increasing. For $0 \leq x < y \leq 1$, the same argument gives $v'(x) \geq \frac{y-x}{y}v'(0) + \frac{x}{y}v'(y) > v'(y)$, so $v'(x)$ is strictly decreasing on $[0,1]$. Since $\tilde{v}'(x) = \frac{v'(x)-v'(0)}{|v(1)-\frac{1}{2}-v'(0)|}$, it is even, concave, and strictly decreasing on $[0,1]$ with $\tilde{v}'(0) = 0$. So $\tilde{v}(x)$ is negative, strictly decreasing and strictly concave on $(0,1]$. By the concavity of $\tilde{v}'(x)$, $\tilde{v}''(x) \leq \frac{\tilde{v}'(x)}{x} < 0$ on $(0,1)$, and $-\tilde{v}''(x)$ is non-decreasing on $(0,1)$.

3) *Elasticity.* The following lemma bounds the elasticity of $\tilde{v}(x)$ from below.

Lemma 3. *Let $v(x)$ be well S-shaped. Then $\mathcal{E}_{\tilde{v}}(x) \geq 2$ for all $x \in (0,1]$, with equality at x only if $\tilde{v}'(t)$ is linear on $[0,x]$. Hence $\frac{\tilde{v}(x)}{x^2}$ is non-increasing on $(0,1]$, so $\tilde{v}(x) \geq -x^2$ on $[0,1]$; if $v'(x)$ is not linear near 0, the ratio is strictly decreasing.*

Proof. Let $x \in (0,1]$ and $\phi := -\tilde{v}'$, which is convex on $[0,x]$ with $\phi(0) = 0$ by 2), so

$\phi(t) \leq \frac{t}{x}\phi(x)$ for $t \in [0, x]$. Hence $-\tilde{v}(x) = \int_0^x \phi(t)dt \leq \frac{x\phi(x)}{2}$, i.e., $\mathcal{E}_{\tilde{v}}(x) \geq 2$, with equality only if ϕ is linear on $[0, x]$. Finally, $\left(\frac{\tilde{v}(x)}{x^2}\right)' = \frac{\tilde{v}(x)(\mathcal{E}_{\tilde{v}}(x)-2)}{x^3} \leq 0$, with strict inequality if $v'(x)$ is not linear near 0, and $\tilde{v}(1) = -1$. □

4) *Non-linearity near* 0. If $v'(x)$ is not linear near 0, then it is not affine on any interval whose closure contains 0: on an interval with 0 in its interior, an affine $v'(x)$ would be constant, contradicting the unique maximum at 0, and an affine piece ending at 0 gives, by evenness, one starting at 0. Hence, for every $G \in \mathcal{G}(p)$, $\overline{v}'_G(s) := \int_0^1 v'(s-t)dG(t)$ is not affine on any interval $[\alpha, \beta]$ with $G([\alpha, \beta]) > 0$. Indeed, $\overline{v}'_G$ is a mixture of the concave functions $v'(s-t)$, so it is affine on $[\alpha, \beta]$ only if $v'(x)$ is affine on $[\alpha - t, \beta - t]$ for G-almost every t, and the closure of this interval contains 0 whenever $t \in [\alpha, \beta]$.

A.2 Proof of Lemma 1

(i) As shown in the text, $\overline{v}'(s_i) = \int_0^1 v'(s_i - s_j)dG_j(s_j)$ is concave on $[0, 1]$. For a maximizer c of $\overline{v}'$, $\overline{v}$ is convex on $[0, c]$ and concave on $[c, 1]$.

(ii) Let $slp(s) := \frac{\overline{v}(s)-\overline{v}(0)}{s}$ for $s \in (0, 1]$ and $slp(0) := \overline{v}'(0)$, so that a is the largest maximizer of slp on $[0, 1]$, which exists as slp is continuous. For $s \in (0, 1]$, differentiating gives $slp'(s) = \frac{N(s)}{s^2}$, where

$$N(s) := s\overline{v}'(s) - \overline{v}(s) + \overline{v}(0) = \int_0^s \left(\overline{v}'(s) - \overline{v}'(t)\right) dt.$$

By (i), $N(s) \geq 0$ on $[0, c]$, and $N(s)$ is non-increasing on $[c, 1]$, since for $c \leq s_1 < s_2 \leq 1$,

$$N(s_2) - N(s_1) = s_1 \left(\overline{v}'(s_2) - \overline{v}'(s_1)\right) + \int_{s_1}^{s_2} \left(\overline{v}'(s_2) - \overline{v}'(t)\right) dt \leq 0.$$

Hence slp is non-decreasing on $[0, c]$ and changes its monotonicity on $[c, 1]$ at most once, from non-decreasing to non-increasing. So $a \geq c$, and $\overline{v}$ is concave on $[a, 1]$. Let $l(s) := \overline{v}(0) + s \cdot slp(a)$. Then $l \geq \overline{v}$ on $[0, a]$ with equality at 0 and a, and if $a < 1$,

$N(a) = 0$, i.e., l is tangent to $\overline{v}$ at a. Therefore $\overline{v}^{cv}$ equals l on $[0, a]$ and $\overline{v}$ on $[a, 1]$. By Kamenica and Gentzkow (2011, Corollary 2), a signal is optimal if and only if it is supported where $\overline{v}$ touches a line supporting $\overline{v}^{cv}$ at p_i. So $\{0, a\}$ is optimal if $p_i < a$, and $\{p_i\}$ is optimal if $p_i \geq a$, where $\overline{v}^{cv}(p_i) = \overline{v}(p_i)$.

If $v'(x)$ is not linear near 0, the concave $\overline{v}'$ is strictly increasing on $[0, z_1]$, constant on $[z_1, z_2]$ and strictly decreasing on $[z_2, 1]$ for some $z_1 \leq z_2$, and $z_1 > 0$ unless $z_2 = 0$: if $\overline{v}'$ were constant on $[0, z_2]$ with $z_2 > 0$, then $G_j([0, z_2]) = 0$ by property 4) in Appendix A.1, so $\overline{v}''(s) = \int_{z_2}^{1} v''(s - s_j) dG_j(s_j) > 0$ on $(0, z_2)$, as $v'' > 0$ on $[-1, 0)$ by property 2), a contradiction. Hence $N > 0$ on $(0, z_2]$ and N is strictly decreasing on $[z_2, 1]$, so $\overline{v} < l$ on $(0, a)$, and $\overline{v}$ is strictly concave on $[a, 1]$. Hence the supporting line at p_i touches $\overline{v}$ only on $\{0, a\}$ if $p_i \leq a$ and only at p_i if $p_i > a$, so the optimum is unique.

A.3 Proof of Proposition 1

Let $\overline{v}(s_i)$ denote the value function of agent i for $\tilde{v}(x)$ rather than $v(x)$, which does not change best replies. If agent j plays $\{0, 1\}$, then $\overline{v}(s_i) = (1 - p_j)\tilde{v}(s_i) - p_j\tilde{v}(1 - s_i)$, which is S-shaped by Lemma 1, with $\overline{v}(1) - \overline{v}(0) = \tilde{v}(1) = -1$ and $\overline{v}'(1) = (1 - p_j)\tilde{v}'(1)$. By Lemma 1, $\{0, 1\}$ is a best reply if and only if $a = 1$, i.e., if and only if $\overline{v}$ lies below its secant $L(s) := \overline{v}(0) - s$ on $[0, 1]$. This implies $\overline{v}'(1) \geq -1$. Conversely, if $a < 1$, then $\overline{v}$ is concave on $[a, 1]$ by the proof of Lemma 1, so $\overline{v}(a) \leq \overline{v}(1) - (1 - a)\overline{v}'(1)$, while $slp(a) > slp(1)$ gives $\overline{v}(a) > \overline{v}(0) - a$; together, $\overline{v}'(1) < -1$. Hence

$$\{0, 1\} \text{ is a best reply } \Leftrightarrow\ (1 - p_j)\tilde{v}'(1) \geq -1 \ \Leftrightarrow\ p_j \geq 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)} = \overline{p},$$

and $(\{0, 1\}; \{0, 1\})$ is an equilibrium if and only if $p_1, p_2 \geq \overline{p}$.

For uniqueness, since all equilibria form a product set, it suffices that $\{0, 1\}$ is the unique best reply to $\{0, 1\}$ for both agents. Another best reply exists only if $D := \overline{v} - L \leq 0$ vanishes at some $s_0 \in (0, 1)$. Since D is convex on $[0, c]$ and concave on

$[c, 1]$ for some c, it then vanishes on $[s_0, 1]$: a convex D with an interior maximum at $s_0 < c$ vanishes on $[0, c]$, and a concave D that vanishes at both ends of $[\max(s_0, c), 1]$ is non-negative there. Hence $\overline{v}'(1) = -1$, i.e., $p_j = \overline{p}$, and $\overline{v}$ is affine on $[s_0, 1]$, which property 4) in Appendix A.1 rules out if $v'(x)$ is not linear near 0, as $\{0, 1\}$ puts mass at 1. This proves uniqueness if $v'(x)$ is not linear near 0 or $p_1, p_2 > \overline{p}$. For $p = \overline{p}$, the uniqueness follows from the proof of Theorem 1, which covers $p = \overline{p}$.

A.4 Proof of Theorem 1

If $p \geq \overline{p}$, full revelation is an equilibrium by Proposition 1, and it is unique for $p > \overline{p}$ by Proposition 1 and for $p = \overline{p}$ by the last paragraph of this proof. So let $p < \overline{p}$, and suppose agent j plays a signal $\{0, a\}$. Then $\overline{v}(s_i) = \frac{a-p}{a}\tilde{v}(s_i) + \frac{p}{a}\tilde{v}(s_i - a)$, and $\{0, a\}$ is a best reply only if the secant of $\overline{v}$ from 0 to a, of slope $\frac{\tilde{v}(a)}{a}$, is tangent at a, where $\overline{v}'(a) = \frac{a-p}{a}\tilde{v}'(a)$; that is, $(a - p)\tilde{v}'(a) = \tilde{v}(a)$, or, as $\tilde{v}' < 0$ on $(0, 1]$, $p = a - \frac{\tilde{v}(a)}{\tilde{v}'(a)}$. The RHS is the zero of the tangent $T_a(x) := \tilde{v}(a) + (x - a)\tilde{v}'(a)$, which is strictly decreasing in x, and it lies in $[0, a)$. For $a_1 < a_2$ and all $x \leq a_1$,

$$T_{a_2}(x) - T_{a_1}(x) = \int_{a_1}^{a_2} \left(\tilde{v}'(t) - \tilde{v}'(a_2)\right) dt + (a_1 - x)\left(\tilde{v}'(a_1) - \tilde{v}'(a_2)\right) > 0,$$

since $\tilde{v}'(x)$ is strictly decreasing on $[0, 1]$. So T_{a_2} is positive at the zero of T_{a_1}, and its own zero lies strictly to the right. Hence the RHS is strictly increasing in a, and the solution is unique and monotone in p, if it exists.

For existence, the RHS is continuous in a, equals $\overline{p}$ at $a = 1$, and equals $a\left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\right) \in [\frac{a}{2}, a)$ by Lemma 3, so it tends to 0 as $a \to 0_+$. So for every $p \in (0, \overline{p})$ the equation of the statement has a unique solution $a(p)$, which is continuous and strictly increasing, as the inverse of a continuous and strictly increasing function.

Best reply and uniqueness. The proof of Proposition 15 in Appendix B.11 never uses $\lambda > 0$, and neither does that of Lemma 11. For $\lambda = 0$, its normalized value function

f is $\overline{v}$, $\overline{p}_0 = \overline{p}$, condition (ii) of Proposition 15 fails as $\tilde{v}(p) < 0$, and its equation for a is the one above. So parts (i) and (iii) of that proof, with its Steps 1 and 2, show that $(\{0,1\};\{0,1\})$ is the unique equilibrium for $p = \overline{p}$ and $(\{0,a\};\{0,a\})$ the unique equilibrium for $p < \overline{p}$, including the case in which $\tilde{v}$ is quadratic on $[0,a]$.

Let me also introduce a bound on a that is used in later proofs.

Lemma 4. *Let $v(x)$ be well S-shaped and $p \in (0,\overline{p})$. In the symmetric equilibrium $(\{0,a\};\{0,a\})$, $p < a \leq 2p$, and $a < 2p$ if $v'(x)$ is not linear near 0.*

Proof. Since $\frac{p}{a} = 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}$, and $\mathcal{E}_{\tilde{v}}(a) \geq 2$ by Lemma 3, with strict inequality if $v'(x)$ is not linear near 0, $p < a \leq 2p$, and $a < 2p$ in the latter case. □

A.5 Pooling by both agents

Lemma 5. *Let $v(x)$ be well S-shaped. Then $(\{p_1\};\{p_2\})$ is never an equilibrium.*

Proof. Against $\{p_j\}$, $\overline{v}(s) = v(s - p_j)$, so $\overline{v}'(s) = v'(s - p_j)$ is strictly increasing on $[0,p_j]$, since $v'(x)$ is even and strictly decreasing on $[0,1]$ by property 2) in Appendix A.1. As in the proof of Lemma 1, let $slp(s) := \frac{\overline{v}(s)-\overline{v}(0)}{s}$, so that $slp'(s) = \frac{N(s)}{s^2}$ with $N(s) = \int_0^s (\overline{v}'(s) - \overline{v}'(t))dt$. Then $N > 0$ on $(0,p_j]$, and, by continuity, slp is strictly increasing on $(0,s]$ for some $s \in (p_j,1]$. If $p_i \leq p_j$, the signal $\{0,s\}$ with mean p_i yields $\overline{v}(0) + p_i\, slp(s) > \overline{v}(0) + p_i\, slp(p_i) = \overline{v}(p_i)$, so $\{p_i\}$ is not a best reply. Hence $\{p_i\}$ is a best reply only if $p_i > p_j$, which cannot hold for both agents.

A.6 Proof of Lemma 6

Lemma 6. *Let $p \in (0,1)$, let $v_n(x)$ be well S-shaped for all $n \in \mathbb{N}$, and let $(\{0,a_n\};\{0,a_n\})$ be the equilibrium for v_n and priors (p,p). If $\mathcal{E}_n(x) \to \infty$ uniformly on $(0,1)$, then $a_n \to p$.*

Since $\mathcal{E}_n(x) \to \infty$ uniformly on $(0,1)$ and $\mathcal{E}_n(x)$ is continuous on $(0,1]$, $\varepsilon_n := \sup_{x\in(0,1]} \frac{1}{\mathcal{E}_n(x)} \to 0$. Hence $p < \overline{p}_n = 1 - \frac{1}{\mathcal{E}_n(1)}$ for large n, and then, by Theorem 1,

$a_n = \frac{p}{1-1/\mathcal{E}_n(a_n)} \in \left[p, \frac{p}{1-\varepsilon_n}\right]$, so $a_n \to p$.

A.7 Unequal priors: regions and further results

The regions of Proposition 2 are bounded by best replies to the three forms of signals, listed below with their values in Example 3.

	definition	*Example 3*
$\overline{p}$	$1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)}$: full revelation for $p_1, p_2 \geq \overline{p}$	$\frac{2}{3}$
$b(p_1)$	agent 2's best reply to $\{p_1\}$ is $\{0, b(p_1)\}$ if $p_2 < b(p_1)$, else $\{p_2\}$	$\min(\frac{3p_1}{2}, 1)$
p^*	smallest p_1 with $b(p_1) = 1$	$\frac{2}{3}$
$a(p_2)$	agent 1's best reply to $\{0,1\}$ is $\{0, a(p_2)\}$ if $p_1 < a(p_2)$, else $\{p_1\}$	$\frac{3p_2}{2}$
$\check{p}_2(p_1)$	largest p_2 for which $\{p_1\}$ is agent 1's best reply to $\{0, b(p_1)\}$	$\frac{2p_1}{3}$
$\hat{p}_1(p_2)$	smallest p_1 for which $\{0,1\}$ is agent 2's best reply to $\{0, a(p_2)\}$	$\frac{2}{3}$

With $p_2 \leq p_1$ and $p_2 < \overline{p}$, the equilibrium is

(i) $(\{p_1\}; \{0,1\})$ if $p_1 \geq \max\{p^*, a(p_2)\}$;

(ii) $(\{p_1\}; \{0, b(p_1)\})$ if $p_1 < p^*$ and $p_2 \leq \check{p}_2(p_1)$;

(iii) $(\{0, a(p_2)\}; \{0,1\})$ if $\hat{p}_1(p_2) \leq p_1 < a(p_2)$;

(iv) $(\{0, a\}; \{0, b\})$ with $p_1 < a \leq b < 1$ otherwise.


Figure 4 depicts these regions for three other rules. In Example 3, $\overline{p} = p^*$ (Figure 2): it is the knife-edge case between $\overline{p} < p^*$ and $\overline{p} > p^*$ in panels (a) and (b). The boundary $a(\cdot)$ is monotone, but $\hat{p}_1(\cdot)$ and $\check{p}_2(\cdot)$ need not be: in panel (c), neither is monotone.

Region (i) has a simple explanation. Against full revelation, agent 1's value function $(1 - p_2)v(s_1) + p_2 v(s_1 - 1)$ is close to $v(s_1)$ for low p_2, so it is concave except near 0, and pooling at p_1 is optimal. Against pooling, agent 2's value function $v(s_2 - p_1)$ is convex on $[0, p_1]$ and concave on $[p_1, 1]$, and for high p_1, its concavification is supported on $\{0,1\}$ despite the concave part. For a fixed $v(x)$, not every pair of forms is played: full

revelation is a best reply to $\{0, c\}$ only if $c \geq p^*$, so $(\{0, c\}; \{0, 1\})$ is an equilibrium only if $c \geq p^*$.

Remark 2. Let $v(x) = v^*(x)$, for which $v'(x)$ is linear on $[0, 1]$, and $(p_1, p_2) = (0.4, 0.3)$. Then agent 2 plays $\{0, 0.6\}$ in every equilibrium, and agent 1's value function against it is linear on $[0, 0.6]$. So agent 1's equilibrium signals are exactly those with support in $[0, 0.6]$ and $\mathbb{E}[s_1^2] = 0.18$, which keep $\{0, 0.6\}$ a best reply of agent 2, such as $\{0, 0.45\}$, $\{0.3, 0.6\}$ and their mixtures.

Figure 4: Equilibria for unequal priors, with $\tilde{v}(x)$ given on $[0, 1]$



(a) $\tilde{v}(x) = -x^{2.5}$:
$\overline{p} = 0.6 < p^* \approx 0.68$

(b) $\tilde{v}(x) = -x^4$:
$\overline{p} = 0.75 > p^* \approx 0.64$

(c) $\tilde{v}(x) = -\frac{x^3 + 60 \max(x - 0.3, 0)^2}{30.4}$:
$p^* \approx 0.641 < \overline{p} \approx 0.651$

Colors are for the corresponding equilibrium type: black – $(\{0, 1\}; \{0, 1\})$; *red* – $(\{p_1\}; \{0, 1\})$; *green* – $(\{0, a\}; \{0, b\})$; *blue* – $(\{p_1\}; \{0, b\})$; *yellow* – $(\{0, a\}; \{0, 1\})$.

Example 6. Let $v(x)$ be well S-shaped with $\tilde{v}(x) = -x^{2.5}$ on $[0, 1]$, and let $p_j = 0.6$. Then agent i's equilibrium signal is non-monotone in p_i.

For the firm, the monotonicity of Proposition 4 extends to unequal priors everywhere except in region (iv) of Proposition 2.

Proposition 13. *Let $v(x)$ be well S-shaped with $v'(x)$ not linear near 0, and let $p_2 \leq p_1$. In the interior of the full revelation region and of each of the regions (i)–(iii) of Proposition 2, both partial derivatives of u_F are positive.*

Example 7. Let $v(x) = \frac{1}{2} + (20x - x^{20})/38$ on $[0,1]$, so that $\tilde{v}(x) = -x^{20}$ and $v(1) = 1$, and let $p_1 = 0.924$. Then u_F strictly decreases in p_2 on $[0.839, 0.873]$, where the equilibrium is in region (iv): once agent 2 stops revealing fully, his high signal b decreases in p_2.

For the agents, monotonicity in the priors is trivial under equal priors, as each is selected with probability $\frac{1}{2}$. Under unequal priors, since the game is constant-sum, each agent's equilibrium payoff $u_i(p_1, p_2) = \mathbb{E}_{G_1,G_2}[v(s_i - s_j)]$ is well defined for any selection rule, whether or not the equilibrium is unique.

Lemma 7. *For any selection rule, $u_i(p_1, p_2)$ is non-decreasing in p_i and non-increasing in p_j. If $v(x)$ is strictly increasing, in particular if it is well S-shaped, $u_i(p_1, p_2)$ is strictly increasing in p_i and strictly decreasing in p_j.*

A.8 Proof of Proposition 2

If $p_2 \geq \overline{p}$, the claim follows from Proposition 1, so let $p_2 < \overline{p}$.

Preliminaries. For a signal of agent j let $N(s) := s\overline{v}'(s) - \overline{v}(s) + \overline{v}(0)$, so that $slp'(s) = N(s)/s^2$. By the proof of Lemma 1 (ii), $N > 0$ on $(0, a)$ and $N < 0$ on $(a, 1]$, with a from Lemma 1. Hence agent i plays $\{p_i\}$ if and only if $N(p_i) \leq 0$, and $\{0, 1\}$ if and only if $N(1) \geq 0$. For $y, c \in [0,1]$, let

$$A(y) := \tilde{v}(y) - y\tilde{v}'(y) = -\int_0^y x\tilde{v}''(x)dx,$$
$$B(y;c) := y\tilde{v}'(c-y) + \tilde{v}(c-y) - \tilde{v}(c) = -\int_0^y x\tilde{v}''(c-x)dx.$$

Since $-\tilde{v}'' > 0$ on $(0,1]$, $A(y) > 0$, and $B(y;c) > 0$ for $0 < y \leq c$. If agent j plays $\{0, c\}$, so that $\mathbb{P}(s_j = c) = q := \frac{p_j}{c}$, then $\overline{v}(s) = (1-q)\tilde{v}(s) + q\tilde{v}(s-c)$ and

$$N(s) = -(1-q)A(s) + qB(s;c); \tag{N}$$

pooling is the case $c = p_j$, $q = 1$.

Step 1: $b(\cdot)$ and p^*. Against $\{p_1\}$, $N(s) = B(s; p_1)$. Hence agent 2 never plays $\{p_2\}$, as $p_2 \leq p_1$, and plays $\{0, b(p_1)\}$, where $b(p_1)$ solves $B(b; p_1) = 0$, i.e. $b\tilde{v}'(b - p_1) = \tilde{v}(b - p_1) + \tilde{v}(p_1)$, or $b(p_1) = 1$ if $B(1; p_1) \geq 0$. With $b = p_1 + \tilde{b}$ this equation reads

$$\varphi(\tilde{b}) := \tilde{v}(\tilde{b}) - (p_1 + \tilde{b})\tilde{v}'(\tilde{b}) + \tilde{v}(p_1) = 0,$$

where $\varphi'(\tilde{b}) = -(p_1 + \tilde{b})\tilde{v}''(\tilde{b}) > 0$, $\varphi(0) = \tilde{v}(p_1) < 0 < 2A(p_1) = \varphi(p_1)$ and $\partial_{p_1}\varphi = \tilde{v}'(p_1) - \tilde{v}'(\tilde{b}) < 0$ for $\tilde{b} < p_1$. So the root is unique, $p_1 < b(p_1) < 2p_1$, and b is continuous, strictly increasing while $b < 1$. Let p^* be the smallest p_1 with $b(p_1) = 1$, solving $\tilde{v}(1 - p^*) + \tilde{v}(p^*) = \tilde{v}'(1 - p^*)$; it exists and $p^* \in (\frac{1}{2}, 1)$, since $b(\frac{1}{2}) < 1$ and $B(1; 1) = 1 > 0$. Thus against $\{x\}$ the other agent plays $\{0, 1\}$ if and only if $B(1; x) \geq 0$, i.e. $x \geq p^*$, as b is non-decreasing.

Step 2: $a(\cdot)$. Against $\{0, 1\}$, by (N),

$$N(s) = \big(A(s) + B(s; 1)\big)\big(p_2 - \Phi(s)\big), \qquad \Phi := \frac{A}{A + B(\cdot\,; 1)}.$$

Φ is strictly increasing: writing $w = -\tilde{v}''$,

$$\partial_y B(y; 1)\, A(y) - B(y; 1)\, A'(y) = y \int_0^y x\big[w(1 - y)w(x) - w(1 - x)w(y)\big]dx < 0,$$

since $w(x) \leq w(y)$ and $w(1 - y) \leq w(1 - x)$ for $x < y$, and $w(x) < w(y)$ for x close to 0, as $v'(x)$ is not linear near 0. Moreover, $\Phi(1) = \frac{|\tilde{v}'(1)| - 1}{|\tilde{v}'(1)|} = \overline{p}$. Hence agent 1 plays $\{p_1\}$ if $p_1 \geq a(p_2)$ and $\{0, a(p_2)\}$ otherwise, where $a(p_2) = 0$ if $p_2 \leq \Phi(0^+)$, $a(p_2) = \Phi^{-1}(p_2)$ if $\Phi(0^+) < p_2 < \overline{p}$, and $a(p_2) = 1$ if $p_2 \geq \overline{p}$; a is continuous and non-decreasing, since Φ is continuous and strictly increasing.

Step 3: $\check{p}_2(\cdot)$ and $\hat{p}_1(\cdot)$. By (N), against $\{0, b(p_1)\}$, agent 1 plays $\{p_1\}$ if and only if

$$p_2 \le \check{p}_2(p_1) := \frac{b(p_1)A(p_1)}{A(p_1) + B\big(p_1; b(p_1)\big)},$$

and against $\{0, a\}$ with $a > p_1$, agent 2 plays $\{0, 1\}$ if and only if $-(a - p_1)A(1) + p_1 B(1; a) \ge 0$. By Step 1, $B(1; a) \ge 0$ if and only if $a \ge p^*$, so for $a < p^*$ this fails, and for $a \ge p^*$ it reads $p_1 \ge \frac{aA(1)}{A(1)+B(1;a)}$. For $a = a(p_2) \ge p^*$ denote this bound by $\hat{p}_1(p_2)$.

Step 4: cases (i)–(iii). By Steps 1 and 2, $(\{p_1\}; \{0, 1\})$ is an equilibrium if and only if $p_1 \ge p^*$ and $p_1 \ge a(p_2)$; by Steps 1 and 3, $(\{p_1\}; \{0, b\})$ with $b < 1$ is an equilibrium if and only if $b = b(p_1)$, $p_1 < p^*$ and $p_2 \le \check{p}_2(p_1)$; by Steps 2 and 3, $(\{0, a\}; \{0, 1\})$ is an equilibrium if and only if $a = a(p_2) \ge p^*$ and $\hat{p}_1(p_2) \le p_1 < a(p_2)$. An equilibrium exists by the minimax theorem, and as the game is constant-sum, all equilibria form a product set. By Lemma 1 every best reply is unique, hence so is the equilibrium.

Step 5: case (iv). For equal priors the game is symmetric, so the unique equilibrium is symmetric and none of the forms in (i)–(iii) is an equilibrium. By Step 4 this gives: $a(p) > p$ if $p \ge p^*$; $\check{p}_2(p) < p$ if $p < p^*$; and $\hat{p}_1(p) > p$ if $a(p) \ge p^*$ and $p < \overline{p}$, since then $p < a(p)$. Now let $p_2 \le p_1$ and suppose that none of (i)–(iii) holds. By Lemma 1 each agent plays $\{p_i\}$, $\{0, 1\}$ or $\{0, c\}$ with $p_i < c < 1$, and by Lemma 5 not both pool at their priors. Apart from (i)–(iv), this leaves the following forms, and each requires, by Steps 1–3 with the roles of the agents exchanged (the functions b, a, $\check{p}_2$ and $\hat{p}_1$ stay the same, only their arguments change), a condition incompatible with $p_2 \le p_1$:

- $(\{0, b(p_2)\}; \{p_2\})$ requires $p_2 < p^*$ and $p_1 \le \check{p}_2(p_2) < p_2$;
- $(\{0, 1\}; \{p_2\})$ requires $p_2 \ge p^*$ and $p_2 \ge a(p_1) \ge a(p_2) > p_2$;
- $(\{0, 1\}; \{0, a(p_1)\})$ with $a(p_1) < 1$ requires $a(p_1) \ge p^*$, $p_1 < \overline{p}$ and $p_2 \ge \hat{p}_1(p_1) > p_1$;
- $(\{0, 1\}; \{0, 1\})$ requires $p_2 \ge \overline{p}$ by Proposition 1.

Hence the equilibrium is $(\{0, a\}; \{0, b\})$ with $p_1 < a < 1$ and $p_2 < b < 1$.

Step 6: $a \leq b$ and continuity. The correspondence $p \mapsto \mathcal{G}(p)$ is continuous: upper hemicontinuous since the mean is weakly continuous, and lower hemicontinuous since any $G \in \mathcal{G}(p)$ is approximated by mixtures of G with δ_0 or δ_1. As payoffs are continuous, the equilibrium correspondence has a closed graph, and being single-valued with values in a compact set, it is continuous; so the equilibrium of Proposition 2 is continuous in (p_1, p_2). If $a = b = c$ in (iv), then, since $B(c; c) = -\tilde{v}(c)$, the indifference conditions $N(c) = 0$ of the two agents, with $q = \frac{p_2}{c}$ and $q = \frac{p_1}{c}$, read $\frac{c}{p_2} - 1 = -\frac{\tilde{v}(c)}{A(c)} = \frac{c}{p_1} - 1$, so $p_1 = p_2$; conversely $a = b$ on the diagonal by symmetry. On the convex set $U = \{0 < p_2 < p_1 < 1,\ p_2 < \overline{p}\}$, let R be the set where (iv) holds. R is open in U, as the strategies in (iv) are not weak limits of $\{0, 1\}$ or of point masses, and $a - b \neq 0$ is continuous on R. Since U is connected and (i) holds for p_1 close to 1, each connected component of R has a boundary point in $U \setminus R$. There one of (i)–(iii) holds, and continuity gives $(a, b) \to (p_1, 1)$, $(p_1, b(p_1))$ or $(a(p_2), 1)$ respectively, with $a(p_2) < 1$ as $p_2 < \overline{p}$. Each limit has $a < b$, hence $a < b$ on R.

A.9 Proof of Proposition 3

Describe a signal of agent i by its high signal $c_i \in [p_i, 1]$, where $c_i = p_i$ stands for $\{p_i\}$. For a fixed p_i, $\frac{\mathbb{P}(s_i=0)}{1-p_i} = \frac{c_i - p_i}{c_i(1-p_i)}$ is increasing in c_i, so it suffices to show that c_i is non-decreasing in p_j. By Lemma 1, the best reply to c_j is $c_i = \max\{r(c_j, \frac{p_j}{c_j}), p_i\}$, where $r(c, q)$ is the point at which N from (N) changes sign; it is continuous in (c, q). With $w = -\tilde{v}''$, which is odd and non-decreasing,

$$\partial_q N(s) = A(s) + B(s; c) = \int_0^s x\big[w(x) + w(c - x)\big]dx > 0,$$

since $w(x) + w(c - x)$ is at least $w(x) > 0$ if $x \leq c$ and equals $w(x) - w(x - c) \geq 0$ if $x > c$. Hence $r(c, q)$ is non-decreasing in q, and strictly increasing where $r(c, q) \in (0, 1)$, and

$r(c,1) = b(c)$ from Step 1 of the proof of Proposition 2.

Let $\rho(a) := r(a, \frac{p_1}{a})$, so that agent 2's best reply to a is $\max\{\rho(a), p_2\}$, and let $\Gamma(a) := \max\{r(c, \frac{p_2}{c}), p_1\}$ with $c = \max\{\rho(a), p_2\}$, for $a \in [p_1, 1]$. Γ is continuous and its fixed points are exactly the equilibria, so by Step 4 of the proof of Proposition 2 it has a unique fixed point a^*; since $\Gamma(p_1) \geq p_1$, $\Gamma(a) > a$ on $[p_1, a^*)$. As p_2 increases to p_2', $\Gamma(a)$ does not decrease for any a: if $\rho(a) \geq p_2'$, only the weight $\frac{p_2}{\rho(a)}$ increases; if $\rho(a) \leq p_2$, agent 2 pools and $b(p_2) \leq b(p_2')$; and if $p_2 < \rho(a) < p_2'$, then $r(\rho, \frac{p_2}{\rho}) \leq r(\rho, 1) = b(\rho) \leq b(p_2')$. Hence the fixed point does not decrease, and if $p_1 < a^* < 1$, $\Gamma(a^*)$ strictly increases, and so does the fixed point. As the argument does not use $p_2 \leq p_1$, it applies to both agents.

A.10 Informativeness and the prior

This subsection studies how informativeness in the symmetric equilibrium of Theorem 1 responds to the prior. For a fixed p, a higher a means a more informative bad news signal. In equilibrium, a rises with p, but this does not imply a more informative signal: if a rises only a little relative to p, the signal may become less informative. To compare them, I view signals as experiments $\pi_i : \{0,1\} \to [0,1]$ in the Blackwell order (Blackwell, 1953).

A bad news experiment sends $s_i = 0$ or $s_i = a$ if $\theta_i = 0$, and always $s_i = a$ if $\theta_i = 1$, so it is characterized by the conditional probability of revealing the bad type, $\frac{\mathbb{P}(s_i=0)}{\mathbb{P}(\theta_i=0)}$. A higher probability gives a more informative experiment, since a lower one is obtained by garbling. Hence, I say that agent i *reveals more for* p_{i1} *than for* p_{i2} if $\frac{\mathbb{P}(s_{i1}=0)}{1-p_{i1}} > \frac{\mathbb{P}(s_{i2}=0)}{1-p_{i2}}$ for the corresponding equilibrium strategies G_{i1}, G_{i2}. By Theorem 1, this measure depends on a single variable:

$$\mathbb{P}(s_i = 0) = \frac{1}{\mathcal{E}_{\tilde{v}}(a)} \text{ and } p = a\left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\right) \text{ hence } \frac{\mathbb{P}(s_i = 0)}{1-p} = \frac{1}{1 + (\mathcal{E}_{\tilde{v}}(a) - 1)(1-a)}.$$

This gives the following lemma.

Lemma 8. *Let $v(x)$ be well S-shaped, $p < \overline{p}$, and $\{0, a\}$ the equilibrium signal. Informativeness strictly increases in the prior at p if and only if $(\mathcal{E}_{\tilde{v}}(x) - 1)(1 - x)$ strictly decreases at $x = a$.*

Corollary 4. *Let $v(x)$ be well S-shaped with $\mathcal{E}_{\tilde{v}}(x)$ non-increasing. Then in equilibrium agents reveal more for p_2 than for p_1 whenever $0 < p_1 < p_2 \leq \overline{p}$.*

Some algebra shows that Corollary 4 applies to the natural examples.

Example 8. For $v(x)$ generated by i.i.d. Gumbel or normal shocks with $\text{Var}[\varepsilon_i] \geq \overline{V}$, $\mathcal{E}_{\tilde{v}}(x)$ is non-increasing, so in equilibrium agents reveal more as p increases.

However, the elasticity can vary flexibly, and better agents may then reveal less.

Example 9. There exists a well S-shaped $v(x)$ with $\tilde{v}(x) = -\frac{x^2}{16(1-x)^2}$ on $[0, \frac{1}{2}]$ such that in equilibrium agents reveal less for p_2 than for p_1 if $0 < p_1 < p_2 \leq 0.375$.

A.11 Proof of Lemma 8

By Theorem 1, $\frac{\mathbb{P}(s_i=0)}{1-p} = \frac{1}{1+(\mathcal{E}_{\tilde{v}}(a)-1)(1-a)}$, where, by its proof, $a(p)$ is continuous and strictly increasing, so it maps $(p, p+\varepsilon)$ and $(p-\varepsilon, p)$ onto a right and a left neighbourhood of $a(p)$. Therefore agents reveal more for all $p_h \in (p, p+\varepsilon)$ and less for all $p_l \in (p-\varepsilon, p)$ if and only if $(\mathcal{E}_{\tilde{v}}(x) - 1)(1 - x)$ is strictly decreasing at $x = a$.

A.12 Proof of Proposition 4

By Theorem 1 and Proposition 1, the equilibrium is $(\{0, a\}; \{0, a\})$ with $a = a(p) > p$, where $a(p)$ is non-decreasing on $(0, 1)$, as $a(p) = 1$ for $p \geq \overline{p}$. When both agents play $\{0, a\}$ with prior $p < a$, the firm gets

$$U(p, a) := \frac{p^2}{a} + \frac{2p(a-p)}{a} v(a) = p + p\Big(1 - \frac{p}{a}\Big)\big(2v(a) - 1\big).$$

For a fixed p, U is non-decreasing in a, as $1 - \frac{p}{a}$ and $2v(a) - 1$ are non-negative and non-decreasing. For a fixed a, $\partial_p U = 1 + \big(1 - \frac{2p}{a}\big)\big(2v(a) - 1\big) > 2\big(1 - v(a)\big) \geq 0$, as

$1 - \frac{2p}{a} > -1$ and $0 < 2v(a) - 1 \leq 1$. Hence, for $p_1 < p_2$,

$$u_F(p_1) = U\big(p_1, a(p_1)\big) \leq U\big(p_1, a(p_2)\big) < U\big(p_2, a(p_2)\big) = u_F(p_2).$$

A.13 Proof of Proposition 13

Since $v(-x) = 1 - v(x)$, $s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) = \frac{s_1 + s_2}{2} + \Psi(s_1 - s_2)$, where $\Psi(x) := x\hat{v}(x)$ is even, so the firm's expected utility is

$$u_F = \frac{p_1 + p_2}{2} + \mathbb{E}[\Psi(s_1 - s_2)].$$

For $x \in (0, 1]$, $0 < \Psi'(x) = \hat{v}(x) + xv'(x) \leq 2\hat{v}(x) \leq 2x$, since $v'(x)$ is decreasing on $[0, 1]$ and $v'(0) \leq 1$, as $\frac{1}{2} \geq \hat{v}(1) \geq \frac{v'(0)}{2}$ by the concavity of $v'(x)$. Also, $\Psi(x) \leq \frac{x}{2}$.

Full revelation. $u_F = \frac{p_1 + p_2}{2} + \Psi(1)(p_1 + p_2 - 2p_1 p_2)$, so $\partial_{p_1} u_F = \frac{1}{2} + \Psi(1)(1 - 2p_2) > \frac{1}{2} - \Psi(1) \geq 0$, and similarly for p_2.

Region (i). $u_F = \frac{p_1 + p_2}{2} + (1 - p_2)\Psi(p_1) + p_2\Psi(1 - p_1)$, so $\partial_{p_2} u_F = \frac{1}{2} - \Psi(p_1) + \Psi(1 - p_1) > 0$ and, as $p_2 \leq p_1$ and $\Psi'(x) \leq 2x$,

$$\partial_{p_1} u_F = \frac{1}{2} + (1 - p_2)\Psi'(p_1) - p_2\Psi'(1 - p_1) > \frac{1}{2} - 2p_1(1 - p_1) \geq 0.$$

Region (ii). Let $b = b(p_1)$ and $y := b - p_1$. By Step 1 of the proof of Proposition 2, $0 < y < p_1$ and $y' = -\frac{\partial_{p_1}\varphi}{\varphi'} > 0$. Then

$$u_F = \frac{p_1 + p_2}{2} + \Psi(p_1) - \frac{p_2}{b}\big(\Psi(p_1) - \Psi(y)\big),$$

where $0 < \Psi(p_1) - \Psi(y) \leq \frac{p_1}{2} < \frac{b}{2}$. Hence $\partial_{p_2} u_F > 0$ and

$$\partial_{p_1} u_F = \frac{1}{2} + \Big(1 - \frac{p_2}{b}\Big)\Psi'(p_1) + \frac{p_2}{b}\Psi'(y)y' + \frac{p_2(1 + y')}{b^2}\big(\Psi(p_1) - \Psi(y)\big) > 0.$$

Region (iii). Let $a = a(p_2)$, which is non-decreasing by Step 2 of the proof of Proposition 2, with $p_2 \leq p_1 < a$ and $a \geq p^* > \frac{1}{2}$. Then

$$u_F = \frac{p_1 + p_2}{2} + p_2\Psi(1) + \frac{p_1}{a}K, \quad \text{where} \quad K := (1-p_2)\Psi(a) + p_2\big(\Psi(1-a) - \Psi(1)\big),$$

so $\partial_{p_1} u_F = \frac{1}{2} + \frac{K}{a} \geq \frac{1}{2} - \frac{p_2\Psi(1)}{a} > 0$ and, with $K_a := \partial_a K$,

$$\partial_{p_2} u_F = \frac{1}{2} + \Psi(1) + \frac{p_1}{a}\big(\Psi(1-a) - \Psi(1) - \Psi(a)\big) + \frac{p_1 a'}{a^2}(aK_a - K).$$

The first three terms sum to at least $\frac{1}{2} - \Psi(a) > 0$, and

$$aK_a - K = (1-p_2)a^2 v'(a) + p_2\big(\hat{v}(1) - \hat{v}(1-a) - a(1-a)v'(1-a)\big) \geq 0,$$

since, by the concavity of $v'(x)$ and $a \geq \frac{1}{2}$,

$$\hat{v}(1) - \hat{v}(1-a) = \int_{1-a}^{1} v'(x)dx \geq \frac{a}{2}v'(1-a) \geq a(1-a)v'(1-a).$$

Example 7. Both signals solve the indifference conditions $N = 0$ from (N), which I solved numerically. For instance, for $p_2 = 0.84, 0.86, 0.873$, $(a, b) \approx (0.9375, 0.9996)$, $(0.9406, 0.9925)$ and $(0.9435, 0.9876)$, and $u_F \approx 0.96145$, 0.96092 and 0.96076.

A.14 Proof of Lemma 7

Write $u_i(p_i, p_j)$ for agent i's equilibrium payoff when his prior is p_i and agent j's prior is p_j. For $G_j \in \mathcal{G}(p_j)$, let $\overline{v}_{G_j}(x)$ be agent i's value function against G_j and $\overline{v}^{cv}_{G_j}$ its concavification on $[0,1]$. By Kamenica and Gentzkow (2011, Corollary 2), $\overline{v}^{cv}_{G_j}(p_i) = \max_{G' \in \mathcal{G}(p_i)} \mathbb{E}_{G'}[\overline{v}_{G_j}]$ is agent i's best payoff against G_j, where the maximum is attained as $\overline{v}_{G_j}$ is continuous. The game is constant-sum, so agent j's equilibrium

signal G_j^* minimizes i's payoff:

$$u_i(p_i, p_j) = \min_{G_j \in \mathcal{G}(p_j)} \overline{v}^{cv}_{G_j}(p_i) = \overline{v}^{cv}_{G_j^*}(p_i).$$

Step 1: each $\overline{v}^{cv}_{G_j}$ is non-decreasing on $[0,1]$, and strictly increasing if $v(x)$ is. As $v(x)$ is increasing, so is $\overline{v}_{G_j}$, so the constant $\overline{v}_{G_j}(1)$ is a concave majorant of $\overline{v}_{G_j}$, and $\overline{v}^{cv}_{G_j} \leq \overline{v}_{G_j}(1) = \overline{v}^{cv}_{G_j}(1)$, as $\{1\}$ is the only signal with mean 1. If $v(x)$ is strictly increasing, so is $\overline{v}_{G_j}$, and every $G' \in \mathcal{G}(x)$ with $x < 1$ puts positive mass on $[0,1)$, so $\overline{v}^{cv}_{G_j}(x) < \overline{v}^{cv}_{G_j}(1)$. For $0 \leq x < y \leq 1$, write $y = \lambda x + (1-\lambda)$ with $\lambda = \frac{1-y}{1-x} \in [0,1)$; concavity gives $\overline{v}^{cv}_{G_j}(y) \geq \lambda\, \overline{v}^{cv}_{G_j}(x) + (1-\lambda)\, \overline{v}^{cv}_{G_j}(1) \geq \overline{v}^{cv}_{G_j}(x)$, strictly in the second case.

Step 2: monotonicity of u_i. Let $p_i < p_i'$, and let G_j^* be agent j's equilibrium signal for priors (p_i', p_j). By Step 1,

$$u_i(p_i, p_j) \leq \overline{v}^{cv}_{G_j^*}(p_i) \leq (<)\, \overline{v}^{cv}_{G_j^*}(p_i') = u_i(p_i', p_j).$$

Finally, since $v(x) + v(-x) = 1$, $u_i(p_i, p_j) = 1 - u_j(p_j, p_i)$, so, by the above applied to agent j, u_i is non-increasing (strictly decreasing) in p_j.

A.15 Selection reversals

This subsection studies how often noise makes the firm select the agent with the lower posterior in the symmetric equilibrium. For $p < \overline{p}$, posteriors differ with probability $\frac{2}{\mathcal{E}_{\tilde{v}}(a)}\left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\right)$, and the firm then selects the agent with $s_i = 0$ with probability $1 - v(a)$. If $\mathcal{E}_{\tilde{v}}(x)$ is non-decreasing, the first factor does not increase in p, as $\frac{1}{\mathcal{E}_{\tilde{v}}(a)} \leq \frac{1}{2}$, and the second decreases, so such reversals become rarer. Under the condition of Corollary 4, which makes informativeness increase in p, the two factors move in opposite directions, and either can dominate. For the two classes from Example 1, which satisfy it by Example 8, a direct computation shows that reversals still become rarer, but for

$v(x) = \frac{1}{2} + 0.6x - 0.1e^{5-\frac{5}{x}}$ on $[0,1]$, the probability of a reversal increases in p for $p < 0.36$.

B. Proofs, Technical Details and Additions for Section 4

B.1 Proof of Theorem 2

Lemma 9. *Let $v(x)$ be well S-shaped and $\hat{v}(x) := v(x) - \frac{1}{2}$. Then $\hat{v}(x) \leq x(2-x)\hat{v}(1) \leq x - \frac{x^2}{2}$ for all $x \in [0,1]$, and if $\hat{v}(x) = x - \frac{x^2}{2}$ for some $x \in (0,1)$, then $v = v^*$.*

Proof. For $0 \leq t < s < 1$, s is a convex combination of t and 1 with weight $\frac{1-s}{1-t}$ on t, so the concavity of $v'(x)$ and $v'(1) \geq 0$ give $v'(s) \geq \frac{1-s}{1-t}v'(t)$. Hence $r(t) := \frac{v'(t)}{1-t}$ is non-decreasing on $[0,1)$: the density $v'(t)$ has a monotone likelihood ratio relative to the triangular density $1-t$. Let $T(x) := \int_0^x (1-t)dt = x - \frac{x^2}{2}$. For $x \in (0,1)$,

$$\frac{\hat{v}(x)}{T(x)} = \frac{\int_0^x r(t)(1-t)dt}{\int_0^x (1-t)dt} \leq r(x) \leq \frac{\int_x^1 r(t)(1-t)dt}{\int_x^1 (1-t)dt} = \frac{\hat{v}(1) - \hat{v}(x)}{T(1) - T(x)},$$

so $\hat{v}(x) \leq \frac{T(x)}{T(1)}\hat{v}(1) = x(2-x)\hat{v}(1) \leq x - \frac{x^2}{2}$, as $\hat{v}(1) \leq \frac{1}{2}$. If $\hat{v}(x) = x - \frac{x^2}{2}$ for some $x \in (0,1)$, then $\hat{v}(1) = \frac{1}{2}$ and both inequalities in the display are equalities, so r, which is continuous on $[0,1)$ by property 1) in Appendix A.1, is constant there and equal to $2\hat{v}(1) = 1$. Hence $v'(t) = 1 - t$ on $[0,1]$, and $v = v^*$, as $v'(x)$ is even. □

By Theorem 1 and Proposition 1, the equilibrium is $(\{0,a\};\{0,a\})$, where $a = 1$ if and only if $p \geq \overline{p}$, and $p < a \leq a^* := \min(2p,1)$ by Lemma 4 and $\overline{p} \geq \frac{1}{2}$ (Lemma 3). As in the proof of Proposition 4, $u_F(p) = p + 2p\left(1 - \frac{p}{a}\right)\hat{v}(a)$, so, by Lemma 9,

$$u_F(p) \leq p + 2p\left(1 - \frac{p}{a}\right)\left(a - \frac{a^2}{2}\right) \leq p + 2p\left(1 - \frac{p}{a^*}\right)\left(a^* - \frac{a^{*2}}{2}\right) = u_F^{ws}(p),$$

where the second inequality holds as both factors are positive and strictly increasing

in a on $(p, 1]$. This proves the bound, and equality holds if and only if $a = a^*$ and $\hat{v}(a) = a - \frac{a^2}{2}$.

If $p \geq \frac{1}{2}$, then $a^* = 1$, so equality means $a = 1$, i.e., $p \geq \overline{p}$, and $v(1) = 1$. For v^*, $\tilde{v}(x) = -x^2$ on $[0, 1]$, so $\overline{p} = \frac{1}{2}$ and $v^*(1) = 1$, and v^* attains the bound.

If $p < \frac{1}{2}$, then $a^* = 2p \in (0, 1)$, so equality requires $\hat{v}(2p) = 2p - 2p^2$, hence $v = v^*$ by Lemma 9. Conversely, for v^*, $\tilde{v}(x) = -x^2$ on $[0, 1]$, so $a = 2p$ by Theorem 1, and both inequalities hold with equality.

It remains to compare v^* with the noiseless rule. By the Fact in Appendix B.12, the utility of the firm under the noiseless rule is $\frac{4p}{3}$ for $p \leq \frac{1}{2}$ and $1 - \frac{(1-p)^2(4p-1)}{3p^2}$ for $p > \frac{1}{2}$. For $p \leq \frac{1}{2}$,

$$p + 2p^2(1-p) - \frac{4p}{3} = -\frac{p}{3}\left(6p^2 - 6p + 1\right) > 0 \Leftrightarrow p \in \left(\frac{3-\sqrt{3}}{6}, \frac{3+\sqrt{3}}{6}\right).$$

For $p > \frac{1}{2}$,

$$2p - p^2 - 1 + \frac{(1-p)^2(4p-1)}{3p^2} = \frac{(1-p)^2(4p-1-3p^2)}{3p^2} = \frac{(1-p)^3(3p-1)}{3p^2} > 0.$$

B.2 Proof of Proposition 5

Throughout, G_u is the equilibrium signal constructed in Proposition 14, and proper u-partitions are as defined in Appendix B.3.

(i) Let $p \leq \frac{1}{2}$, $u < 2p$, $k = \lceil \frac{2p}{u} \rceil \geq 2$ and $r = 2p - (k-1)u \in (0, u]$. The proper u-partition of $[0, 2p]$ has atoms iu with masses $\frac{k-i}{k(k+1)}$ and atoms $iu + r = 2p - (k-1-i)u$ with masses $\frac{i+1}{k(k+1)}$, $i = 0, \ldots, k-1$. For fixed k, as u increases, iu moves at rate i and $iu + r$ at rate $-(k-1-i)$, so $I_{G_u}(t) := \int_0^t G_u(x)dx$ satisfies $\partial_u I_{G_u}(t) = -M(t)$, where $M(t)$ sums rate times mass over the atoms below t. The pair $\{iu, iu + r\}$ contributes $\frac{2i+1-k}{k(k+1)}$ and the atom iu alone $\frac{i(k-i)}{k(k+1)}$, hence $M(t) = 0$ for $t \in (ju, ju + r]$ and

$M(t) = \frac{(j+1)(j+1-k)}{k(k+1)} < 0$ for $t \in (ju + r, (j+1)u]$, $j \leq k-2$. So $I_{G_u}(t)$ is non-decreasing in u, and strictly increasing for t in a set of positive measure, i.e. G_u is a strict mean-preserving spread of $G_{u'}$ for $u' < u$ with the same k. At $u = \frac{2p}{k}$, $r = u$ and G_u is the uniform distribution on $\{0, \frac{2p}{k}, \ldots, 2p\}$, which is also its limit as $u \uparrow \frac{2p}{k}$, where $k+1$ is used and $r \to 0$; so G_u is continuous in u and the ranking extends to $(0, 2p]$. By Proposition 14, $G_u = \{0, \min(2p, 1)\}$ if $u \geq 2p$ and $p \leq \frac{1}{2}$, or $u \geq 2 - 2p$ and $p > \frac{1}{2}$.

For the example after Proposition 5, let $p \geq \frac{5}{9}$ and $u \in [1-p, 2-2p)$. Then $k = 1$ in case (iii) of the full statement in the proof of Proposition 14, and $a = 2u - \frac{u^2}{1-p} \leq 1 - u$, as $3u - \frac{u^2}{1-p}$ is maximized at $u = \frac{3(1-p)}{2}$, where it equals $\frac{9(1-p)}{4} \leq 1$. The proper u-partition of $[0, a]$ is $\{0, a\}$ with equal probabilities, so G_u puts probabilities $\frac{1-p}{2-a}$, $\frac{1-p}{2-a}$ and $\frac{2p-a}{2-a}$ on $\{0, a, 1\}$. As $\frac{da}{du} = 2 - \frac{2u}{1-p} < 0$ for $u > 1 - p$, a higher u lowers the probability of 0 and raises that of 1, so no two of these signals are ranked in the Blackwell order.

(ii) Since $v(-x) = 1 - v(x)$, with $d = s_1 - s_2$,

$$u_F = \mathbb{E}[s_1 v(d) + s_2 v(-d)] = \mathbb{E}[\max(s_1, s_2)] - \mathbb{E}\big[|d|\big(1 - v(|d|)\big)\big],$$

and $v(x) = 1 - \frac{(u-x)^2}{2u^2}$ for $x \in [0, u]$. Let $p \leq \frac{1}{2}$. For $u \geq 2p$, $G_u = \{0, 2p\}$ and

$$u_F = \frac{p}{2} + p\, v(2p) = \frac{3p}{2} - \frac{p(u-2p)^2}{2u^2},$$

which equals $\frac{3p}{2}$ at $u = 2p$ and is strictly decreasing on $[2p, 1]$. For $u < 2p$, $u_F \leq \mathbb{E}[\max(s_1, s_2)] < \frac{3p}{2}$ by the chord argument of the proof of Lemma 2 on $[0, 2p]$ in place of $[0, 1]$, which gives $\mathbb{E}_{s_1}[\max(s_1, s_2)] \leq p + \frac{s_2}{2}$, strictly for $s_2 \in (0, 2p)$, as G_u puts mass on $(0, 2p)$.

Let $p > \frac{1}{2}$. For $u \in [2-2p, 1]$, $G_u = \{0, 1\}$ and $v(1) = 1$, so $u_F = u_F^*(p)$. For $u < 2-2p$, $v(x)$ does not implement a fully revealing equilibrium by the proof of Proposition 14 (ii), so $u_F < u_F^*(p)$ by Lemma 2.

(iii) The proper u-partition of $[0,a]$ with $k = \lceil \frac{a}{u} \rceil$ puts mass $\frac{1}{k}$ on each pair $\{iu, iu+r\} \subset [iu,(i+1)u]$, $i = 0,\dots,k-1$, so it converges weakly to $U[0,a]$ as $u \to 0$. If $p \le \frac{1}{2}$, G_u is the proper u-partition of $[0,2p]$ and converges to $U[0,2p]$. If $p > \frac{1}{2}$ and $u < \min(2p-1, 2-2p)$, then with k from Proposition 14 (iii), we have $a \in ((k-1)u, ku]$ and $ku < 2-2p \le (k+1)u$, so $a \in (2-2p-2u, 2-2p)$ and $a < 1-u$. Hence case (iii) applies, $a \to 2-2p$, the weight $\frac{2(1-p)}{2-a} \to \frac{1-p}{p}$, and G_u converges to the $(\frac{1-p}{p}, \frac{2p-1}{p})$ mix of $U[0,2-2p]$ and $\delta_{\{1\}}$ — exactly the distributions from Appendix B.12.

B.3 Triangular selection rules

This subsection constructs the symmetric equilibrium used in Subsection 4.2 for the selection rules generated by i.i.d. $\varepsilon_i \sim U[0,u]$ with $u < 1$. Since $v(x)$ is piecewise quadratic, $v'(x)$ is piecewise linear, so the functional equation from Appendix D.2 reduces to a difference equation, and one can find $G_i(s_i)$ s.t. $\overline{v}(s_j)$ is linear on $[a,b]$.

Definition 6. Say χ is a proper u-partition of the interval $[a,b]$ if its CDF is given by:

$$G(x) = \begin{cases} 0, x < a \\ \frac{i+1}{k+1}, x \in [a + i\cdot u, b - (k-1-i)\cdot u), i \in \{0,\dots,k-1\} \\ \frac{i+1}{k}, x \in [b-(k-1-i)\cdot u, a+(i+1)\cdot u), i \in \{0,\dots,k-2\} \\ 1, x \ge b \end{cases}, \text{with } k = \lceil \tfrac{b-a}{u} \rceil$$

Remark 3. This G solves the functional equation $G(x-u) + G(x+u) = 2G(x)$, i.e. $\mathbb{P}(x < \chi \le x+u) = \mathbb{P}(x-u < \chi \le x)$, for all $x \in [a,b)$, subject to $\text{supp}(\chi) \subseteq [a,b]$.

The proper u-partition of $[a,b]$ is the unique $G_i(s_i)$ supported in $[a,b]$ that makes $\overline{v}(s_j) = \int_0^1 v(s_j - s_i) dG_i(s_i)$ linear on $[a,b]$. Both agents playing it is often not an equilibrium, since $\overline{v}(s_j)$ beyond $[a,b]$ also matters, but it is the key ingredient: the next proposition constructs a symmetric equilibrium for all (u,p) from mixtures of proper u-partitions and, possibly, $\delta_{\{1\}}$.

Proposition 14. *Let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$. Then a symmetric equilibrium signal strategy is:*

(i) If $p \leq \frac{1}{2}$, the signal is a proper u-partition of $[0, 2p]$.

(ii) If $p > \frac{1}{2}$ and $u \geq 2 - 2p$, the signal is $\{0, 1\}$.

(iii) Otherwise, the signal mixes a proper u-partition of $[0, a]$ for some a with a mass point at 1 and (possibly) with a proper u-partition of $[0, 1 - u]$.

So binary support survives a slight violation of well S-shapedness for extreme priors, $p \leq \frac{1}{2}u$ or $p \geq 1 - \frac{1}{2}u$, but fails for all $p \in (\frac{1}{2}u, 1 - \frac{1}{2}u)$. The last case, whose exact equations are in the proof below, shows how complex equilibria can be for general $v(x)$: as the signal 1 must be treated separately, an equilibrium may require a three-way mixture of two proper u-partitions and a mass point.

B.4 Proof of Proposition 14

Full statement: let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$; then there exists a symmetric equilibrium in which each agent's signal G is:

(i) Proper u-partition of $[0, 2p]$ if $p \leq \frac{1}{2}$

(ii) $\{0, 1\}$, if $p > \frac{1}{2}, u \geq 2 - 2p$.

(iii) $(\frac{2(1-p)}{2-a}, \frac{2p-a}{2-a})$ mix of the proper u-partition of $[0, a]$ with $\delta_{\{1\}}$, if (i) $p > \frac{1}{2}, u < 2-2p$, and (ii) $a \leq 1 - u$, where $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}$, $k = \lceil \frac{2-2p}{u} \rceil - 1$.

(iv) If, in the solution of the equations in (iii), $a > 1 - u$, the true equilibrium G is constructed as follows: let $k = \lceil \frac{1}{u} \rceil$, and let $\tilde{G}$ be a $(\frac{k-1}{k}, \frac{1}{k})$ mix of the proper u-partition of $[0, 1 - u]$ and $\delta_{\{1\}}$. Then let G be the

$$\left(\frac{(2p - a)k}{(k-1)(1-u) + 2 - ka}, \frac{(k-1)(1-u) + 2 - 2pk}{(k-1)(1-u) + 2 - ka} \right)$$

mix of $\tilde{G}$ and the proper u-partition of $[0, a]$, where $a \in (1-u, (k-1)u)$ is the unique solution of $(1-a)\overline{v}(0) + a\overline{v}(1) = \overline{v}(a)$.

If $u \geq 1$, then $v(x)$ is quadratic and $\tilde{v}(x) = -x^2$ on $[0,1]$; hence I refer the reader to Proposition 1 and Theorem 1. Consider now $u < 1$. Then $v'(x) = \frac{\max(u-|x|,0)}{u^2}$, which is continuous, so $\overline{v}'$ is continuous; $v(x) = 0$ for $x < -u$, $v(x) = \frac{(u+x)^2}{2u^2}$ on $[-u, 0]$, and $v(x) = 1 - v(-x)$.

(i) Suppose that $p \leq \frac{1}{2}, u < 1$, and let $G(x)$ be the CDF of the proper u-partition χ on $[0, 2p]$ with $k = \lceil \frac{2p}{u} \rceil$. As $\mathbb{P}(\chi = iu) = \mathbb{P}(\chi = 2p - iu) = \frac{k-i}{k(k+1)}$ for $i < k$, and the terms $i = k$ below vanish,

$$\overline{v}'(x) = \int_0^{2p} v'(x-s)dG(s) = \underbrace{\sum_{i=0}^{k} \frac{\max(u-|x-iu|,0)}{u^2} \frac{k-i}{k(k+1)}}_{S_L} + \underbrace{\sum_{i=0}^{k} \frac{\max(u-|x-2p+iu|,0)}{u^2} \frac{k-i}{k(k+1)}}_{S_R}.$$

Consider S_L and say that $a_i = \max(u - |x - iu|, 0)$ is *active* if $a_i > 0$. Then, as long as $x \in (iu, (i+1)u)$, only a_i and a_{i+1} are active, so

$$S_L = \frac{(u-(x-iu))(k-i) + (u-((i+1)u-x))(k-i-1)}{u^2 k(k+1)} = \frac{ku-x}{u^2 k(k+1)}.$$

By the continuity of S_L at $x = iu$, $S_L = \frac{ku-x}{u^2k(k+1)}$ on $[0, ku]$, and hence on $[0, 2p]$, since $ku \geq 2p$. Analogously, $S_R = \frac{ku+x-2p}{u^2k(k+1)}$ on $[0, 2p]$, hence $\overline{v}'(x) = \frac{2ku-2p}{u^2k(k+1)}$ on $[0, 2p]$.

For $x > 2p$, $S_L = \frac{(ku-x)^+}{u^2k(k+1)}$ and $S_R = \frac{k(u-x+2p)^+}{u^2k(k+1)}$, both nonincreasing and convex. So $\overline{v}'(x)$ is constant on $[0, 2p]$ and nonincreasing and convex on $[2p, 1]$; hence $\overline{v}(x)$ is concave on $[0, 1]$ and linear on $[0, 2p]$, and a pair of proper u-partitions of $[0, 2p]$ is an equilibrium.

(ii) Suppose that $p > \frac{1}{2}$, $2 - 2p \le u < 1$; then $\overline{v}(x) = (1-p)v(x) - pv(1-x)$. Then

$$\overline{v}'(x) = \frac{(1-p)\max(u-x,0) + p\max(u+x-1,0)}{u^2},$$

where the numerator is a piecewise linear convex function, hence $\overline{v}(x)$ is concave and then convex, and we need to check that $\overline{v}'(0) \le \overline{v}(1) - \overline{v}(0)$.

$\overline{v}'(0) = \frac{1-p}{u}$, $\overline{v}(1) - \overline{v}(0) = v(1) - v(0) = \frac{1}{2}$, so $\{0,1\}$ is a best reply iff $2 - 2p \le u$.

(iii) Suppose $p > \frac{1}{2}, u < 2 - 2p$, and $a \le 1 - u$, where $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}$ for $k = \lceil \frac{2-2p}{u} \rceil - 1$. Then consider the $(\frac{2(1-p)}{2-a}, 1 - \frac{2(1-p)}{2-a})$ mix of the proper u-partition of $[0,a]$ with $\delta_{\{1\}}$; notice that this is a proper G, since the proper u-partition of $[0,a]$ has a mean $\frac{1}{2}a$, and $\frac{2(1-p)}{2-a} \cdot \frac{1}{2}a + 1 - \frac{2(1-p)}{2-a} = p$.

First, $\lceil \frac{2-2p}{u} \rceil - 1 = \lceil \frac{a}{u} \rceil$, so k is the same in the proper u-partition of $[0,a]$:

$$\left\lceil \frac{a}{u} \right\rceil = \left\lceil 2k - \frac{k(k+1)u}{2(1-p)} \right\rceil = k,$$

since $k < \frac{2(1-p)}{u} \le k+1$, hence $k \le \frac{k(k+1)u}{2(1-p)} < k+1$.

Since $\overline{v}(x)$ is linear in the strategy of the other agent,

$$\overline{v}'(x) = \frac{2(1-p)}{2-a} S(x) + \frac{2p-a}{2-a} \cdot \frac{\max(u+x-1,0)}{u^2},$$

where $S(x)$ is $\overline{v}'(x)$ from (i) against the proper u-partition χ of $[0,a]$. Hence $\overline{v}(x)$ is linear on $[0,a]$ with slope $\overline{v}'(0) = \frac{2(1-p)}{2-a} \cdot \frac{2ku-a}{u^2 k(k+1)}$. Since $v(1-s) = 1$ for $s \le a \le 1-u$,

$$\overline{v}(1) - \overline{v}(0) = \frac{2(1-p)}{2-a}\mathbb{E}[v(\chi)] + \frac{2p-a}{2(2-a)}, \qquad \mathbb{E}[v(\chi)] = 1 - \frac{1}{2(k+1)} - \frac{(ku-a)^2}{2u^2 k(k+1)},$$

and the equation $\overline{v}(1) - \overline{v}(0) = \overline{v}'(0)$ reduces to a linear equation in a with the root $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}$. Hence $\int_a^1 (\overline{v}'(x) - \overline{v}'(a))dx = 0$.

On $[a,1]$ both summands of $\overline{v}'(x)$ are convex: the first by (i), the second as a maximum of linear functions. Hence $\overline{v}'(x) - \overline{v}'(a)$ is convex on $[a,1]$, vanishes at a and integrates to zero, so $\exists c : \overline{v}'(x) \le \overline{v}'(a)\ \forall x \in [a,c]$ and $\overline{v}'(x) \ge \overline{v}'(a)\ \forall x \in [c,1]$. Therefore $\overline{v}(x) \le \overline{v}(a) + (x-a)\overline{v}'(a)$ on $[a,1]$, with equality at $x = 1$, so $[0,a] \cup \{1\}$ is the concavification support, and (G,G) is an equilibrium.

(iv) Notice that G is a feasible distribution, since the mean of $\tilde{G}$ is $\frac{k-1}{k}\frac{1-u}{2} + \frac{1}{k} = \frac{Q}{2k}$ with $Q := (k-1)(1-u)+2$, the mean of the proper u-partition of $[0,a]$ is $\frac{a}{2}$, and $\beta := \frac{(2p-a)k}{Q-ka}$ solves $\beta\frac{Q}{2k} + (1-\beta)\frac{a}{2} = p$.

Suppose that a from (iii) satisfies $a > 1-u$. I first show that if the other agent plays the $(\frac{k-1}{k}, \frac{1}{k})$ mix of the proper u-partition of $[0, 1-u]$ and $\delta_{\{1\}}$, where $k = \lceil \frac{1}{u} \rceil$, then $\overline{v}(x)$ is linear on $[0,(k-1)u]$. Again, consider $\overline{v}'(x)$, noting that $\lceil \frac{1-u}{u} \rceil = k-1$:

$$\overline{v}'(x) = \frac{k-1}{k}\Bigg[\underbrace{\sum_{i=0}^{k-1} \frac{\max(u-|x-iu|,0)}{u^2}\frac{k-1-i}{k(k-1)}}_{S_L} + \underbrace{\sum_{i=0}^{k-1} \frac{\max(u-|x-1+(i+1)u|,0)}{u^2}\frac{k-1-i}{k(k-1)}}_{S_R}\Bigg] + \frac{1}{k}\frac{\max(u-|x-1|,0)}{u^2}.$$

By the proof of (i), $\overline{v}'(x)$ is constant on $[0,1-u]$ (since the last term is zero on that interval) and equal to $\frac{k-1}{k}\frac{2(k-1)u-(1-u)}{u^2k(k-1)} = \frac{(2k-1)u-1}{u^2k^2}$. On $[1-u,(k-1)u]$, the only active terms are a_{k-2} in S_L, a_0 in S_R and the last term.

$$\begin{aligned}\overline{v}'(x) &= \frac{k-1}{k}\left[\frac{((k-1)u-x)}{u^2}\frac{1}{k(k-1)} + \frac{1-x}{u^2}\frac{k-1}{k(k-1)}\right] + \frac{1}{k}\frac{x+u-1}{u^2} \\ &= \frac{((k-1)u-x)+(k-1)(1-x)+k(x+u-1)}{k^2u^2} = \frac{(2k-1)u-1}{k^2u^2}.\end{aligned}$$

Finally, on $[(k-1)u, 1]$, only a_0 in S_R and the last term are active.

$$\overline{v}'(x) = \frac{k-1}{k}\frac{1-x}{u^2}\frac{k-1}{k(k-1)} + \frac{1}{k}\frac{x+u-1}{u^2} = \frac{(k-1)(1-x)+k(x+u-1)}{k^2u^2} = \frac{x+ku-1}{k^2u^2}.$$

So against $\tilde{G}$, $\overline{v}'(x)$ is constant on $[0,(k-1)u]$ and increasing linear on $[(k-1)u,1]$, while against the proper u-partition of $[0,a]$ it is constant on $[0,a]$ and convex on $[a,1]$ by (i). Hence, if the other agent plays G, $\overline{v}(x)$ is linear on $[0,a]$ and $\overline{v}'(x)$ is convex on $[a,1]$, and as in (iii), if a solves $(1-a)\overline{v}(0) + a\overline{v}(1) = \overline{v}(a)$, then $[0,a]\cup\{1\}$ is the concavification support, and (G,G) is an equilibrium.

It remains to show that such an $a \in (1-u,(k-1)u)$ exists and is unique, and that $\beta \in (0,1)$. Let a_3 and $k_3 = \lceil \frac{2-2p}{u} \rceil - 1$ be a and k from (iii). Since $a_3 \le k_3 u$ and $k_3 \le k-1$, $a_3 > 1-u > (k-2)u$ requires $k_3 = k-1$, and then $a_3 > 1-u$ is equivalent to $p < p_u := 1 - \frac{k(k-1)u^2}{2((2k-1)u-1)}$. Let χ be the proper u-partition of $[0,a]$ and, for a signal G' of the other agent, let $s_{G'}$ be the slope of $\overline{v}(x)$ on $[0,a]$ and $C_{G'} := \overline{v}(1) - \overline{v}(0)$, so that the condition on a is $s_G = C_G$. A direct computation gives

$$s_{\tilde{G}} - C_{\tilde{G}} = -\frac{((k-1)u-1)^2}{2k^2u^2}, \qquad \frac{d}{da}(s_\chi - C_\chi) = -\frac{1-a}{(k-1)u^2},$$

and $s_\chi = \frac{2(k-1)u-a}{u^2k(k-1)}$ by (i). Hence

$$\Lambda(a) := (Q-ka)(s_G - C_G) = -(2p-a)\frac{((k-1)u-1)^2}{2ku^2} + (Q-2kp)(s_\chi - C_\chi)$$

has the sign of $s_G - C_G$, as $Q - ka \ge (k+1)(1-(k-1)u) > 0$, and $\Lambda(a)$ is quadratic in a with the leading coefficient $\frac{Q-2kp}{2(k-1)u^2}$. Since $p < p_u$ and $\frac{Q}{2k} - p_u = \frac{(k-1)((k-1)u-1)^2}{2k((2k-1)u-1)} > 0$,

$2kp < Q$ and Λ is strictly convex. Moreover,

$$\Lambda(1-u) = \frac{(1+u)((2k-1)u-1)}{k(k-1)u^2}(p_u - p) > 0,$$
$$\Lambda((k-1)u) = -\frac{(k+1)(2p-1)((k-1)u-1)^2}{2ku^2} < 0,$$

so Λ has a unique root in $(1-u, (k-1)u)$. And $a < 1 < 2p$ and $2kp < Q$ give $\beta \in (0,1)$.

B.5 Proof of Lemma 2

Since $v(-x) = 1 - v(x)$, for $s_1 \geq s_2$,

$$\begin{aligned} s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) &= s_2 + (s_1 - s_2)v(s_1 - s_2) \leq s_2 + (s_1 - s_2)v(1) \\ &= v(1)\max(s_1, s_2) + (1 - v(1))\min(s_1, s_2) \leq \max(s_1, s_2), \end{aligned}$$

where the first inequality holds as $v(x)$ is increasing and $s_1 - s_2 \leq 1$, and the second as $v(1) \leq 1$, strictly if $s_1 \neq s_2$ and $v(1) < 1$. Hence, for any signals, the firm's utility is at most $\mathbb{E}[\max(s_1, s_2)]$. For $s_2 \in [0,1]$, $\max(s, s_2)$ lies below its secant $s_2 + s(1 - s_2)$ on $[0,1]$, strictly on $(0,1)$ if $s_2 \in (0,1)$. Hence $\mathbb{E}_{G_1}[\max(s_1, s_2)] \leq s_2 + p(1 - s_2)$, with equality for all s_2 if G_1 is full revelation, and for $s_2 \in (0,1)$ only then. Averaging over s_2, $\mathbb{E}[\max(s_1, s_2)] \leq 2p - p^2$, with equality if and only if at least one agent plays full revelation. Moreover, $2p - p^2 = \mathbb{E}[\max(\theta_1, \theta_2)] = u_F^*(p)$.

So the firm's utility in every equilibrium is at most $u_F^*(p)$. Let (G_1^*, G_2^*) be an equilibrium with $\mathbb{E}[s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)] = u_F^*(p)$. Then all inequalities above are equalities, so some agent plays full revelation, and since then $s_1 \neq s_2$ with positive probability, $v(1) = 1$. As all equilibria form a symmetric product set, $(\{0,1\}; \{0,1\})$ is then an equilibrium too. Conversely, if $(\{0,1\}; \{0,1\})$ is an equilibrium and $v(1) = 1$, then $s_1 - s_2 \in \{-1, 0, 1\}$ and both inequalities are equalities, so the firm attains $u_F^*(p)$.

B.6 Robustness across unequal priors

Lemma 10. *Let $S \subseteq (0,1)$. A selection rule $v(x)$ implements a fully revealing equilibrium for all $(p_1, p_2) \in S^2$ if and only if it does so for all (p, p) with $p \in S$.*

B.7 Proof of Lemma 10

Let agent j with prior p_j play full revelation. Then $\overline{v}(s_i) - \frac{1}{2} = (1-p_j)\hat{v}(s_i) - p_j\hat{v}(1-s_i)$ does not depend on p_i, and full revelation is a best reply of agent i for any $p_i \in (0,1)$ if and only if $\overline{v}$ lies below its secant from 0 to 1:

$$(1-p_j)\hat{v}(x) - p_j\hat{v}(1-x) \leq \hat{v}(1)(x-p_j) \quad \forall x \in (0,1).$$

Hence $(\{0,1\};\{0,1\})$ is an equilibrium for (p_1, p_2) if and only if this condition holds for $p_j \in \{p_1, p_2\}$, so it is an equilibrium for all $(p_1, p_2) \in S^2$ if and only if the condition holds for all $p_j \in S$, i.e., if and only if it is an equilibrium for all (p, p), $p \in S$.

B.8 Proof of Proposition 6

Let $a := \hat{v}(1)$; since $v(x) \leq 1$ is increasing, $a \in [0, \frac{1}{2}]$. By the proof of Lemma 10, $v(x)$ implements a fully revealing equilibrium for (p, p) if and only if $(1-p)\hat{v}(x) - p\hat{v}(1-x) \leq a(x-p)$ for all $x \in (0,1)$. Applying it also at $1-x$ and summing the two inequalities with weights $1-p$ and p gives $(1-2p)\big(\hat{v}(x) - ax\big) \leq 0$. Taking $p = p_l$ and $p = p_h$ yields $\hat{v}(x) = ax$, which proves (ii); then the condition holds with equality for all $p \in (0,1)$, and Lemma 10 extends full revelation to all pairs of priors, which proves (i).

B.9 Proof of Proposition 7

(i) With a and g as in Definition 3, $\hat{v}(x) = ax + g(x)$ on $[0,1]$ and $g(0) = g(1) = 0$, so the condition $(1-p)\hat{v}(x) - p\hat{v}(1-x) \leq a(x-p)$ for all $x \in (0,1)$ from the proof of

Lemma 10 reads $(1-p)g(x) - pg(1-x) \leq 0$ for all $x \in (0,1)$. It is linear in p, so it holds for all $p \in [\frac{1}{2}, 1)$ if and only if it holds at $p = \frac{1}{2}$ and $p = 1$. At $p = 1$ it reads $g \geq 0$ on $(0,1)$; at $p = \frac{1}{2}$ it reads $g(x) \leq g(1-x)$, which, applied at x and $1-x$, means $g(x) = g(1-x)$. The case $p \in (0, \frac{1}{2}]$ is the same with $p = 0$, where the condition reads $g \leq 0$. For a balanced $v(x)$ and $x < y$ in $[0,1]$, $\hat{v}(x) \leq \hat{v}(y)$ and $\hat{v}(1-y) \leq \hat{v}(1-x)$ read $g(x) - g(y) \leq a(y-x)$ and $g(y) - g(x) \leq a(y-x)$, so $\hat{v}$ is increasing if and only if $|g(x) - g(y)| \leq a|x-y|$.

(ii) As all equilibria form a product set, it suffices that full revelation is each agent's unique best reply to full revelation, i.e., that the condition above is strict on $(0,1)$. For a balanced $v(x)$ it reads $(1-2p)g(x) < 0$, which holds for all $p > \frac{1}{2}$ [$p < \frac{1}{2}$] if and only if $g > 0$ [$g < 0$] on $(0,1)$.

B.10 Proof of Corollary 2

Suppose that agent j plays $\{0,1\}$. Then $\overline{v}(s_i) = (1-p_j)v(s_i) + p_j v(s_i - 1)$, and $\{0,1\}$ is a best reply of agent i only if the secant from 0 to 1 lies weakly above $\overline{v}(s_i)$ on $[0,1]$, which requires $\overline{v}'(0) \leq \overline{v}(1) - \overline{v}(0) \leq \overline{v}'(1)$. Since $v'(x)$ is even, $\overline{v}'(0) = (1-p_j)v'(0) + p_j v'(1)$ and $\overline{v}'(1) = (1-p_j)v'(1) + p_j v'(0)$, so this requires $(2p_j - 1)\big(v'(0) - v'(1)\big) \geq 0$, i.e., $p_j \geq \frac{1}{2}$. Hence in a fully revealing equilibrium both priors are at least $\frac{1}{2}$.

If $v(x)$ is generated by i.i.d. shocks with a square-integrable density f, then $v'(x) = \int f(t)f(t+x)dt$ is continuous, and by the Cauchy–Schwarz inequality,

$$v'(1) = \int f(t)f(t+1)dt \leq \int f^2(t)dt = v'(0),$$

with equality only if $f(t+1) = f(t)$ almost everywhere, which is impossible for a probability density. Hence $v'(0) > v'(1)$.

B.11 Costly signals

Suppose that signals are costly: agent i pays $\kappa\, c(G_i)$, where $\kappa > 0$ and $c(G_i) = \int_0^1 \phi\, dG_i - \phi(p_i)$ for a strictly convex ϕ, continuous on $[0,1]$, so that his payoff is $\mathbb{E}[v(s_i - s_j)] - \kappa\, c(G_i)$. The firm does not bear the cost, and her utility is unchanged.

Lemma 11. *For any priors and selection rule, equilibria exist and form a product set.*

Proof. Since agent i's cost does not depend on G_j, both agents have the same best replies as in the zero-sum game in which agent 1 receives $W(G_1, G_2) := \mathbb{E}[v(s_1 - s_2)] - \kappa c(G_1) + \kappa c(G_2)$ and agent 2 receives $-W(G_1, G_2)$. W is affine in each G_i and continuous in the weak* topology, as v and ϕ are continuous, and each $\mathcal{G}(p_i)$ is convex and weak* compact. By Sion's minimax theorem, this game has a value and optimal strategies, and its equilibria are exactly the pairs of optimal strategies. □

Unlike the value of this zero-sum game, the agents' own payoffs need not be the same in all equilibria.

B.11.1 The variance cost

Let $\phi(x) = x^2$, so that $c(G_i) = \mathrm{Var}_{G_i}[s_i]$ is the expected reduction in the residual variance of θ_i. As the linear part of $v(x)$ adds only a linear function of s_i to agent i's value function, best replies depend on $v(x)$ only through $\tilde{v}(x)$ and

$$\lambda := \frac{\kappa}{|v(1) - \frac{1}{2} - v'(0)|},$$

and the normalized value function against G_j is $f(s_i) := \overline{v}(s_i) - \lambda s_i^2$.

Proposition 15. *Let $v(x)$ be well S-shaped, the priors (p, p), and $c(G_i) = Var_{G_i}[s_i]$.*

(i) If $p \geq \overline{p}_\lambda := 1 - \frac{1-\lambda}{\mathcal{E}_{\tilde{v}}(1)}$, the unique equilibrium is $(\{0,1\}; \{0,1\})$.

(ii) If $\lambda \geq \frac{|\tilde{v}(p)|}{p^2}$*, the unique equilibrium is* $(\{p\};\{p\})$*.*

(iii) Otherwise, the unique equilibrium is $(\{0,a\};\{0,a\})$*, where* $a \in (p,1)$ *solves* $p = a - \frac{|\tilde{v}(a)|-\lambda a^2}{|\tilde{v}'(a)|}$*, and* $\mathbb{P}(s_i = 0) = \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\left(1 - \frac{\lambda a^2}{|\tilde{v}(a)|}\right)$*.*

The conditions in (i) and (ii) are mutually exclusive.

For $\lambda = 0$, (ii) never applies, and Proposition 15 reduces to Theorem 1. Since λ is the cost relative to the curvature of $v(x)$, noise that flattens $v(x)$ makes the same cost weigh more. For instance, for $\varepsilon_i \sim U[0,u]$ with $u \geq 1$, $\tilde{v}(x) = -x|x|$ and $\lambda = 2\kappa u^2$: agents reveal fully if $p \geq \frac{1+\lambda}{2}$, pool if $\lambda \geq 1$, and otherwise play $\{0, \frac{2p}{1+\lambda}\}$ with $\mathbb{P}(s_i = 0) = \frac{1-\lambda}{2}$. Without the cost, informativeness does not depend on $u \geq 1$; with it, it decreases in u.

Proof. Exclusivity. By the concavity of $\tilde{v}(x)$ on $[0,1]$ (property 2 in Appendix A.1), $\tilde{v}(p) \leq \tilde{v}(1) + (p-1)\tilde{v}'(1) = -1 + (1-p)\mathcal{E}_{\tilde{v}}(1)$. So if (i) holds, $\lambda \leq 1 - (1-p)\mathcal{E}_{\tilde{v}}(1) \leq |\tilde{v}(p)| < \frac{|\tilde{v}(p)|}{p^2}$.

Uniqueness. By Lemma 11 and symmetry, all equilibria form a product set $E \times E$. So if $G^* \in E$ is the unique best reply to itself, then $E = \{G^*\}$, as every $G \in E$ is a best reply to G^*.

Best reply to $\{0,a\}$. Let agent j play $\{0,a\}$ with $a \in (p,1]$, and let $\gamma := \frac{a-p}{a} \in (0,1)$. Then

$$f(s) = \gamma\tilde{v}(s) + (1-\gamma)\tilde{v}(s-a) - \lambda s^2.$$

Let $l(s)$ be the line through $(0, f(0))$ and $(a, f(a))$, whose slope is $\frac{\tilde{v}(a)-\lambda a^2}{a}$, and $d(s) := l(s) - f(s)$ on $[0,a]$, so that $d(0) = d(a) = 0$. As $\tilde{v}'(x)$ is even,

$$d'(s) = \frac{\tilde{v}(a) - \lambda a^2}{a} - \gamma\tilde{v}'(s) - (1-\gamma)\tilde{v}'(a-s) + 2\lambda s,$$

which is convex on $[0,a]$, since $\tilde{v}'(x)$ is concave. If $d'(a) \leq 0$, then $d \geq 0$ on $[0,a]$: with $z := \inf\{s \in [0,a] : d'(s) \leq 0\}$, convexity gives $d' \leq 0$ on $[z,a]$, while $d' > 0$ on $[0,z)$. If

$a < 1$ and $d'(a) = 0$, then for $s \in (a, 1]$,

$$f'(s) = \gamma\tilde{v}'(s) + (1-\gamma)\tilde{v}'(s-a) - 2\lambda s < \gamma\tilde{v}'(a) + (1-\gamma)\tilde{v}'(0) - 2\lambda a = f'(a),$$

as $\tilde{v}'(x)$ is strictly decreasing on $[0, 1]$, so f lies strictly below its tangent l on $(a, 1]$. Hence, if $d'(a) \leq 0$ for $a = 1$, or $d'(a) = 0$ for $a < 1$, then $\{0, a\}$ is a best reply to itself. Since $f'(a) = \gamma\tilde{v}'(a) - 2\lambda a$,

$$d'(a) = 0 \;\Leftrightarrow\; \gamma\tilde{v}'(a) = \frac{\tilde{v}(a)}{a} + \lambda a \;\Leftrightarrow\; \gamma = \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\Big(1 - \frac{\lambda a^2}{|\tilde{v}(a)|}\Big) \;\Leftrightarrow\; p = a - \frac{|\tilde{v}(a)| - \lambda a^2}{|\tilde{v}'(a)|},$$

where the last step uses $\gamma a = a - p$.

Step 1: other best replies. Suppose that $d'(a) = 0$ and $d(c) = 0$ for some $c \in (0, a)$. Then c minimizes d on $[0, a]$, so $d'(c) = 0$; the convex d' vanishes at c and a, so $d' \leq 0$ on $[c, a]$, and $\int_c^a d' = 0$ gives $d' \equiv 0$ there; by convexity again $d' \geq 0$ on $[0, c]$, and $d(0) = d(c)$ gives $d' \equiv 0$ on $[0, a]$. So $\gamma\tilde{v}'(s) + (1-\gamma)\tilde{v}'(a-s)$ is affine on $[0, a]$; as a sum of two concave functions, it is affine only if both are, so $\tilde{v}'(s) = -Ks$ on $[0, a]$ for some $K > 0$, and $d' \equiv 0$ then requires $K(1-2\gamma) = 2\lambda$, i.e., $\frac{p}{a} = 1 - \gamma = \frac{1}{2} + \frac{\lambda}{K}$. Hence, either $d > 0$ on $(0, a)$, and $\{0, a\}$ is the unique best reply to itself, or $\tilde{v}(x) = -\frac{K}{2}x|x|$ on $[-a, a]$ and $p = a\big(\frac{1}{2} + \frac{\lambda}{K}\big)$, and every $G \in \mathcal{G}(p)$ supported on $[0, a]$ is a best reply to $\{0, a\}$.

Step 2: the quadratic case. Take such a $G \neq \{0, a\}$ and let $w(s) := \int_0^a \tilde{v}(s-t)dG(t) - \lambda s^2$ be the normalized value function against it. For $s \in [0, a]$, $w'(s) = -K\int_0^a |s - t|dG(t) - 2\lambda s$, hence $w'(a) = -K(a-p) - 2\lambda a$, while $w(0) = \frac{K}{2}\mathbb{E}_G[t^2]$ and $w(a) = -\frac{K}{2}\mathbb{E}_G[(a-t)^2] - \lambda a^2$. So the secant slope from $(0, w(0))$ to $(a, w(a))$ exceeds $w'(a)$ by

$$\frac{K}{a}\Big(\frac{a^2}{2} - \mathbb{E}_G[t^2]\Big) + \lambda a > \frac{K}{a}\Big(\frac{a^2}{2} - ap\Big) + \lambda a = 0,$$

since $t^2 \leq at$ on $[0, a]$, with equality only at $t \in \{0, a\}$, and $p = a\big(\frac{1}{2} + \frac{\lambda}{K}\big)$. Hence $w(s)$

lies strictly above the secant for $s < a$ close enough to a, so, for such $s > p$, $\{0, s\}$ yields a strictly higher payoff against G than $\{0, a\}$. Thus $\{0, a\}$ is not a best reply to G, $G \notin E$, and the equilibrium is unique.

(i) For $a = 1$, $\gamma = 1 - p$ and $d'(1) = -1 + \lambda + (1 - p)\mathcal{E}_{\tilde{v}}(1)$, so $d'(1) \leq 0$ if and only if $p \geq \overline{p}_\lambda$, and then $\{0, 1\}$ is a best reply to itself. If $p > \overline{p}_\lambda$, then $d > 0$ on $(0, 1)$: otherwise, $d(c) = 0$ for some $c \in (0, 1)$ gives $d'(c) = 0$, so $d' \leq 0$ on $[c, 1]$ by convexity, and $\int_c^1 d'(s)ds = 0$ gives $d' \equiv 0$ on $[c, 1]$, contradicting $d'(1) < 0$. So $\{0, 1\}$ is the unique best reply to itself, and the equilibrium is unique. If $p = \overline{p}_\lambda$, then $d'(1) = 0$, and Steps 1 and 2 with $a = 1$ give uniqueness.

(ii) Against $\{p\}$, $f(s) = \tilde{v}(s - p) - \lambda s^2$, and $f'(p) = -2\lambda p$. So $\{p\}$ is a best reply if and only if f lies below its tangent at p, i.e., if and only if $\tilde{v}(s - p) \leq \lambda(s - p)^2$ on $[0, 1]$. For $s > p$, this holds strictly, as $\tilde{v} < 0$ on $(0, 1]$. For $s < p$, with $t = p - s$, it reads $\frac{|\tilde{v}(t)|}{t^2} \leq \lambda$, and as this ratio is non-decreasing on $(0, 1]$ by Lemma 3, it holds for all $t \in (0, p]$ if and only if $\lambda \geq \frac{|\tilde{v}(p)|}{p^2}$. Then every best reply to $\{p\}$ is supported on $[0, p]$ and has mean p, so it is $\{p\}$, and the equilibrium is unique.

(iii) Let $p < \overline{p}_\lambda$ and $\lambda < r(p)$, where $r(t) := \frac{|\tilde{v}(t)|}{t^2}$ is continuous and non-decreasing on $(0, 1]$. Let $h(a) := a - \frac{|\tilde{v}(a)| - \lambda a^2}{|\tilde{v}'(a)|} = a\big(1 - \frac{1 - \lambda/r(a)}{\mathcal{E}_{\tilde{v}}(a)}\big)$, continuous on $(0, 1]$, with $h(1) = \overline{p}_\lambda > p$. Let $a_0 := \sup\{t \in (0, 1] : r(t) \leq \lambda\}$, and $a_0 := 0$ if this set is empty; by continuity and monotonicity of r, $a_0 < p$. If $a_0 > 0$, then $r(a_0) = \lambda$ and $h(a_0) = a_0 < p$. If $a_0 = 0$, then $\lambda < r(a)$ for all $a \in (0, 1]$, so $h(a) \in [\frac{a}{2}, a)$ by Lemma 3, and $h(a) \to 0$ as $a \to 0$. In both cases, by continuity, $h(a) = p$ for some $a \in (a_0, 1)$. As $r(a) > \lambda$ there, $p = h(a) < a$, so $\gamma \in (0, 1)$, and $d'(a) = 0$ by the equivalence above. Hence $\{0, a\}$ is a best reply to itself, with $\mathbb{P}(s_i = 0) = \gamma$ as stated, and Steps 1 and 2 give uniqueness. □

B.11.2 Robust implementation

For the rules of Section 4, let ϕ be continuously differentiable on $[0,1]$, including the endpoints, and let $\psi(s) := (1-s)\phi(0) + s\phi(1) - \phi(s)$, which is positive on $(0,1)$ by strict convexity, with $\lim_{s\to 0} \frac{\psi(s)}{s} = \phi(1) - \phi(0) - \phi'(0) > 0$ and $\lim_{s\to 1} \frac{\psi(s)}{1-s} = \phi'(1) - \phi(1) + \phi(0) > 0$.

Proposition 16. *Let ϕ be continuously differentiable on $[0,1]$.*

(i) If $v(x) = \frac{1}{2} + ax$, the unique equilibrium is $(\{p_1\};\{p_2\})$ for all (p_1, p_2).

(ii) Let $v(x)$ be strictly positively [negatively] balanced with $\liminf_{s\to 0_+} \frac{|g(s)|}{s} > 0$. Then $m := \inf_{s\in(0,1)} \frac{|g(s)|}{\psi(s)} > 0$, and full revelation is the unique equilibrium for all (p_1, p_2) with $p_1, p_2 > \frac{1}{2} + \frac{\kappa}{2m}$ $\left[p_1, p_2 < \frac{1}{2} - \frac{\kappa}{2m}\right]$.

(iii) Let $v(x)$ be positively or negatively balanced with $\liminf_{s\to 0_+} \frac{|g(s)|}{s} = 0$. Then $(\{0,1\};\{0,1\})$ is not an equilibrium for any (p_1, p_2).

For the variance cost, $\psi(s) = s(1-s)$; for $v^*(x)$, $g(s) = \frac{s(1-s)}{2}$, so $m = \frac{1}{2}$ and (ii) gives full revelation for $p > \frac{1}{2} + \kappa$, in line with Proposition 15(i), as $\lambda = 2\kappa$. If instead $\phi'(0) = -\infty$, as for entropy, then $\frac{\psi(s)}{s} \to \infty$ as $s \to 0$, while $\frac{|g(s)|}{s} \leq a$ for every balanced rule, so by (FR) below, no balanced rule implements full revelation for any $\kappa > 0$.

Proof. (i) Agent i's payoff is $\frac{1}{2} + a(p_i - p_j) - \kappa c(G_i)$, and by strict convexity, $c(G_i) > 0$ unless $G_i = \{p_i\}$.

(ii)–(iii) Let agent j play $\{0,1\}$. With $\hat{v}(x) = ax + g(x)$ on $[0,1]$ and $g(x) = g(1-x)$, agent i's value function is, up to a constant,

$$(1-p_j)\hat{v}(s) - p_j\hat{v}(1-s) = a(s-p_j) + (1-2p_j)g(s),$$

and he maximizes the expectation of this function minus $\kappa\phi(s)$. By Kamenica and Gentzkow (2011, Corollary 2), $\{0,1\}$ is a best reply if and only if this function lies below

its chord on $[0,1]$. As $g(0) = g(1) = 0$, this reads

$$(2p_j - 1)g(s) \geq \kappa\psi(s) \quad \forall s \in (0,1), \tag{FR}$$

which does not depend on p_i; if (FR) is strict on $(0,1)$, $\{0,1\}$ is the unique best reply.

(ii) Consider the positive case. The ratio $\frac{g}{\psi}$ is continuous and positive on $(0,1)$, and $\liminf_{s\to 0} \frac{g(s)}{\psi(s)} > 0$ by the limit of $\frac{\psi(s)}{s}$ above. As $g(s) = g(1-s)$, $\liminf_{s\to 1} \frac{g(s)}{1-s} > 0$ too, so $\liminf_{s\to 1} \frac{g(s)}{\psi(s)} > 0$, and hence $m > 0$. If $p_j > \frac{1}{2} + \frac{\kappa}{2m}$, then $(2p_j - 1)g(s) \geq (2p_j - 1)m\psi(s) > \kappa\psi(s)$ on $(0,1)$, so (FR) is strict. Hence, if both priors exceed $\frac{1}{2} + \frac{\kappa}{2m}$, full revelation is each agent's unique best reply to itself, and by Lemma 11, it is the unique equilibrium. The negative case is symmetric.

(iii) Consider the positive case. If $p_j \leq \frac{1}{2}$, the left side of (FR) is non-positive, and the right side is positive. If $p_j > \frac{1}{2}$, take $s_n \to 0$ with $\frac{g(s_n)}{s_n} \to 0$; then $(2p_j - 1)\frac{g(s_n)}{s_n} \to 0$, while $\kappa\frac{\psi(s_n)}{s_n} \to \kappa\big(\phi(1) - \phi(0) - \phi'(0)\big) > 0$, so (FR) fails. The negative case is symmetric. □

B.12 Equations for the noiseless rule

Fact (Boleslavsky and Cotton 2015, Lemma 3)**.** *The unique symmetric equilibrium for* $v(x) = \frac{1}{2} + \frac{1}{2} \cdot \operatorname{sign}(x)$ *has the form* (G, G)*, where for* $p \geq \frac{1}{2}$

$$G(x) = \begin{cases} \frac{x}{2p}, & x \leq 2 - 2p, \\ \frac{1}{p} - 1, & 2 - 2p \leq x < 1, \\ 1, & x = 1, \end{cases}$$

while for $p \leq \frac{1}{2}$

$$G(x) = \begin{cases} \frac{x}{2p}, & x \leq 2p, \\ 1, & x \in [2p, 1]. \end{cases}$$

Hence the firm's equilibrium utility is

$$u_F^0(p) = \mathbb{E}_G[\max(s_1, s_2)] = \begin{cases} \frac{4p}{3}, & 0 < p \le \frac{1}{2}, \\ 1 - \frac{(1-p)^2(4p-1)}{3p^2}, & \frac{1}{2} < p < 1. \end{cases}$$

C. Proofs, Technical Details and Additions for Subsection 4.4 and Section 5

C.1 Censorship under a general prior

For an arbitrary prior F, let $F^{-1}(u) := \inf\{x \in [0,1] : F(x) \ge u\}$ for $u \in (0,1]$, $F^{-1}(0) := 0$, and let $F^{-1}(u^+)$ be its right limit, with $F^{-1}(1^+) := F^{-1}(1)$. For $q \in [0,1)$ let $m(q) := \frac{1}{1-q}\int_q^1 F^{-1}(u)du$, and $m(1) := \overline{\theta}$. With $U \sim U[0,1]$, the quantile censorship G_q reveals the type $F^{-1}(U)$ if $U < q$ and sends $m(q)$ otherwise:

$$G_q(x) = \begin{cases} \min\{F(x), q\}, & x < m(q), \\ 1, & x \ge m(q). \end{cases}$$

The existence and uniqueness in Proposition 9 follow from the next lemma; the bound $q \le \frac{1}{2}$ and the utility bound are proved in Appendix C.5.

Lemma 12. *Let $v(x)$ be well S-shaped and F any prior.*

(i) There are $q^* \in (0,1]$ and $\theta^* \in [F^{-1}(q^*), F^{-1}(q^{*+})]$ such that (G_{q^*}, G_{q^*}) is an equilibrium and either $\theta^* < m^* := m(q^*)$ and

$$\int_0^{q^*} \big[\tilde{v}(m^* - F^{-1}(u)) - \tilde{v}(\theta^* - F^{-1}(u)) - (m^* - \theta^*)\tilde{v}'(m^* - F^{-1}(u))\big]du + \tilde{v}(m^* - \theta^*)(1 - q^*) = 0, \tag{QC'}$$

or $q^* = 1$, so that both agents reveal fully, and $\int_0^1 v'(x-t)\,dF(t)$ is nondecreasing on $[\underline{\theta}, \overline{\theta}]$.

(ii) Every q and $\theta \in [F^{-1}(q), F^{-1}(q^+)]$ with $\theta < m(q)$ that solve (QC′) *give an equilibrium (G_q, G_q).*

(iii) If $v'(x)$ is not linear near 0, the equilibrium is unique; all solutions in (ii) then have the same q and induce the same signal.

Lemma 13. *Let $v(x)$ be well S-shaped and F any prior on $[0,1]$. Against every rival signal K, some quantile censorship G_q, $q \in [0,1]$, is a best reply.*

Proof. Let $\underline{\theta} < \overline{\theta}$, $w := w_K$, let L_x be the tangent to w at x, and let $e(q,\theta) := L_{m(q)}(\theta) - w(\theta) = \int_\theta^{m(q)} (w'(z) - w'(m(q)))dz$, which is continuous on Γ.

(a) If $e(0,\underline{\theta}) \geq 0$, then pooling $G_0 = \{\mu\}$ is a best reply. Let $D := L_\mu - w$, so $D(\underline{\theta}) \geq 0 = D(\mu) = D'(\mu)$. If $w'(x) > w'(\mu)$ for some $x > \mu$, then $w' < w'(\mu)$ on $[0,\mu)$, so $D(\underline{\theta}) < D(\mu) = 0$, a contradiction. So $D \geq 0$ on $[\mu,1]$. On $[\underline{\theta},\mu]$, D first increases and then decreases, so $D \geq \min(D(\underline{\theta}), D(\mu)) \geq 0$. Hence $L_\mu \geq w$ on $[\underline{\theta},\overline{\theta}]$, and $\int w\,dG' \leq L_\mu(\mu) = w(\mu)$ for every mean-preserving contraction G' of F.

(b) If w' is non-decreasing on $[\underline{\theta},\overline{\theta}]$, then $F = G_1$ is a best reply by Step 2 of the proof of Lemma 12.

(c) Otherwise, $e(0,\underline{\theta}) < 0$, and as in Step 3 there are $x_1 < x_2 < \overline{\theta}$ with $w'(x_1) > w'(x_2)$. So w' is strictly decreasing on $[x_2,1]$, and $e(q,\theta) > 0$ for $(q,\theta) \in \Gamma$ with $x_2 < \theta < m(q)$, which holds near $(\overline{q},\overline{\theta})$. Since the path from $(0,\underline{\theta})$ to $(\overline{q},\overline{\theta})$ is continuous, e has a zero (q,θ) with $\theta < m(q)$, and G_q is a best reply by Step 1. □

C.2 Proof of Lemma 12

Preliminaries. By property 1) in Appendix A.1, v is C^1 on $[-1,1]$ with a concave derivative, but in general it is not C^2 (e.g. $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$). For a distribution K

on $[0,1]$, let $w_K(x) := \int_0^1 \tilde{v}(x-y)dK(y)$. Since $v(x) = \frac{1}{2} + v'(0)x + |v(1) - \frac{1}{2} - v'(0)|\tilde{v}(x)$ and every feasible signal has mean $\mathbb{E}[\theta] =: \mu$, w_K has the same best replies as $\overline{v}(\cdot, K)$. It is C^1 on $[0,1]$, and $w_K'(x) = \int_0^1 \tilde{v}'(x-y)dK(y)$ is concave.

Quantile facts: F^{-1} is non-decreasing and left-continuous, and $F^{-1}(u) \leq x \Leftrightarrow u \leq F(x)$. Let $\overline{q} := F(\overline{\theta}-)$, so that $F^{-1} < \overline{\theta}$ on $(0, \overline{q})$ and $F^{-1} = \overline{\theta}$ on $(\overline{q}, 1]$. As F^{-1} is integrable, m is continuous on $[0,1)$, and $F^{-1}(q^+) \leq m(q) \leq \overline{\theta}$ gives $m(q) \to \overline{\theta} = m(1)$ as $q \to 1$. For $q < \overline{q}$, F^{-1} is not a.e. constant on $(q,1)$, so $m(q) > F^{-1}(q^+)$; for $q \geq \overline{q}$, $m(q) = \overline{\theta}$ and $G_q = F$. G_q is the distribution of $\mathbb{E}[F^{-1}(U)|\, U\mathbf{1}\{U < q\}]$, so it is feasible.

The curve. Let $\Gamma := \{(q,\theta) : q \in [0,\overline{q}],\ F^{-1}(q) \leq \theta \leq F^{-1}(q^+)\}$, with $F^{-1}(\overline{q}^+) := \overline{\theta}$, i.e. the graph of F^{-1} on $[0,\overline{q}]$ with its jumps filled in. As $\phi(q) := q + F^{-1}(q)$ is strictly increasing, for each $r \in [0, \overline{q} + \overline{\theta}]$ there is a unique $q(r)$ with $\phi(q(r)) \leq r \leq \phi(q(r)^+)$, and $q(r)$ is 1-Lipschitz, since $q(r) > q(r')$ gives $r - r' \geq q(r) - q(r') + F^{-1}(q(r)) - F^{-1}(q(r')^+) \geq q(r) - q(r')$. So $r \mapsto (q(r), r - q(r))$ is a continuous path that traverses Γ from $(0,0)$ to $(\overline{q}, \overline{\theta})$. By the quantile facts, $\theta < m(q)$ on $\Gamma \setminus \{(\overline{q}, \overline{\theta})\}$.

For $(q,\theta) \in \Gamma$, let $w_q := w_{G_q}$, let L_q be the tangent to w_q at $m(q)$, and let $E(q,\theta) := L_q(\theta) - w_q(\theta)$. Since $\tilde{v}$ is odd with $\tilde{v}'(0) = 0$, $E(q,\theta)$ is the left-hand side of (QC'), and

$$E(q,\theta) = \int_\theta^{m(q)} \big(w_q'(z) - w_q'(m(q))\big)dz.$$

E is continuous on Γ, as m is continuous and $\tilde{v}, \tilde{v}'$ are uniformly continuous, and $E(\overline{q}, \overline{\theta}) = 0$.

Step 1 (verification). Let K be any distribution on $[0,1]$, $w := w_K$, $(q,\theta) \in \Gamma$ with $\theta < m := m(q)$, and L the tangent to w at m with $L(\theta) = w(\theta)$. Then G_q is a best reply to K. Indeed, let $D := L - w$, so that $D(\theta) = D(m) = D'(m) = 0$ and $D' = w'(m) - w'$. If $w'(x) > w'(m)$ for some $x > m$, then $w' < w'(m)$ on $[0,m)$ by concavity, so D is strictly increasing on $[0,m]$, contradicting $D(\theta) = D(m)$. Hence D is non-decreasing,

and non-negative, on $[m, 1]$. The set of $x \leq m$ with $w'(x) \geq w'(m)$ is an interval $[\alpha, m]$, so D is non-decreasing on $[\theta, \alpha]$ and non-increasing on $[\max(\alpha, \theta), m]$, and $D(\theta) = D(m) = 0$ gives $D \geq 0$ on $[\theta, m]$. If $\alpha > \theta$, then $w'(\theta) < w'(\alpha)$, so w' is non-decreasing on $[0, \theta]$ and $w'(\theta) < w'(m)$. If $\alpha \leq \theta$, then $D = 0$ on $[\theta, m]$, so w' is constant on $[\theta, m]$, hence maximal there and non-decreasing on $[0, \theta]$. In both cases

$$p(x) := \begin{cases} w(x), & x \leq \theta, \\ L(x), & x \geq \theta, \end{cases}$$

is convex, $p \geq w$, and $p = w$ on $[0, \theta] \cup \{m\}$. For $u < q$, $F^{-1}(u) \leq F^{-1}(q) \leq \theta$, and for $u > q$, $F^{-1}(u) \geq F^{-1}(q^+) \geq \theta$, where p is affine. Hence $\operatorname{supp}(G_q) \subseteq [0, \theta] \cup \{m\}$, and

$$\int p \, dF = \int_0^q p(F^{-1}(u))du + (1 - q)p(m) = \int p \, dG_q = \int w \, dG_q.$$

So every mean-preserving contraction G' of F satisfies $\int w \, dG' \leq \int p \, dG' \leq \int p \, dF = \int w \, dG_q$.

Step 2 (full revelation). If w'_K is non-decreasing on $[\underline{\theta}, \overline{\theta}]$, then F is a best reply to K: w_K is convex on $[\underline{\theta}, \overline{\theta}]$, which contains the support of every mean-preserving contraction G' of F, so $\int w_K \, dG' \leq \int w_K \, dF$.

Step 3 (existence). Let $\underline{\theta} < \overline{\theta}$. At the start of the path, $E(0, \theta) = \tilde{v}(\mu - \theta) < 0$ for $\theta \in [0, \underline{\theta}]$, as $\theta \leq \underline{\theta} < \mu$. Let $g := w'_F$, which is concave.

If g is non-decreasing on $[\underline{\theta}, \overline{\theta}]$, then $F = G_1$ is a best reply to F by Step 2, so (G_1, G_1) is an equilibrium, which is the second alternative in (i).

Otherwise, there are $\underline{\theta} \leq x_1 < x_2 < \overline{\theta}$ with $g(x_1) - g(x_2) =: 2\gamma > 0$ (if $x_2 = \overline{\theta}$, move x_2 slightly to the left, by continuity). Let $Y_q := F^{-1}(U)\mathbf{1}\{U < q\} + m(q)\mathbf{1}\{U \geq q\} \sim G_q$. As $q \to \overline{q}$, $Y_q \to Y_{\overline{q}}$ in probability, since they differ only on $\{q \leq U < \overline{q}\}$ and by $|m(q) - m(\overline{q})|$. Since $G_{\overline{q}} = F$ and $\tilde{v}'$ is uniformly continuous, $w'_q \to g$ uniformly.

Hence there is a neighborhood N of $(\overline{q}, \overline{\theta})$ in Γ on which $w'_q(x_1) > w'_q(x_2)$ and $\theta > x_2$. For $(q, \theta) \in N$, the concave w'_q is then strictly decreasing on $[x_2, 1] \supseteq [\theta, m(q)]$, so $E(q, \theta) > 0$ whenever $\theta < m(q)$. By the intermediate value theorem along the path, E has a zero $(q^*, \theta^*) \in \Gamma \setminus \{(\overline{q}, \overline{\theta})\}$, where $\theta^* < m(q^*)$, and $q^* > 0$ since $E < 0$ for $q = 0$. By Step 1 with $K = G_{q^*}$, (G_{q^*}, G_{q^*}) is an equilibrium. This proves (i), and Step 1 with $K = G_q$ proves (ii).

By Lemma 14, the second alternative requires an atom of mass at least $\frac{1}{2}$ at $\overline{\theta}$.

Step 4 (uniqueness). Now suppose that $v'(x)$ is not linear near 0. First, let (q, θ) be as in (ii), so $q < 1$, and let $w := w_q$, $m := m(q)$, with D, α, p as in Step 1 for $K = G_q$. By property 4) in Appendix A.1, w' is affine on no interval that G_q charges, and $G_q(\{m\}) \geq 1 - q > 0$. Hence $\alpha > \theta$, as otherwise w' is constant on $[\theta, m]$. Also $D > 0$ on $(\theta, 1] \setminus \{m\}$: a zero of D in $(m, 1]$, or in $[\alpha, m)$, makes w' constant on an interval with endpoint m, and a zero in (θ, α) would give $D = 0$ on $[\theta, x]$, i.e. $w' = w'(m)$ there, while $w' < w'(m)$ on $[0, \alpha)$. Finally, w' is strictly increasing on $[0, \theta]$: it is non-decreasing there and concave, so if it were constant on a subinterval of $[0, \theta]$, it would attain its maximum there, but $w'(\theta) < w'(m)$.

Let G' be a best reply to G_q, so that both inequalities in Step 1 are equalities. The first one gives $\operatorname{supp}(G') \subseteq [0, \theta] \cup \{m\}$. Writing $p(x) = p(0) + p'(0)x + \int_0^1 (x - y)^+ dp'(y)$ and $I_K(y) := \int_0^y K(x)dx$, the second one gives $\int_0^1 \big(I_F(y) - I_{G'}(y)\big)dp'(y) = 0$, where $I_{G'} \leq I_F$. As dp' charges every subinterval of $[0, \theta]$, $I_{G'} = I_F$ on $[0, \theta]$, so $G' = F$ on $[0, \theta)$. The rest of G', of mass $1 - F(\theta-)$, lies on $\{\theta, m\}$, and its first moment is $\int_{[\theta,1]} x\, dF$. As G_q also equals F on $[0, \theta)$ and puts $q - F(\theta-)$ on θ and $1 - q$ on m, these two moment conditions, with $\theta \neq m$, give $G' = G_q$. So G_q is the unique best reply to itself.

Second, if $g = w'_F$ is non-decreasing on $[\underline{\theta}, \overline{\theta}]$, it is strictly increasing there: a concave non-decreasing g that is constant on a subinterval is constant up to $\overline{\theta}$, i.e. affine on an interval that F charges, contradicting property 4). So w_F is strictly convex

on $[\underline{\theta}, \overline{\theta}]$, and the same argument gives $G' = F$ for every best reply G' to F.

So in both cases of (i), the equilibrium signal is the unique best reply to itself. As all equilibria form a product set, every equilibrium strategy is a best reply to it, so the equilibrium is unique, and every solution in (ii) induces the same signal. A signal G_q with $\theta < m(q)$ determines q, as its top atom is at $m(q)$ with mass $1-q$, and the revealed types lie below $m(q)$. So q is unique, and θ can vary only within $[F^{-1}(q), F^{-1}(q^+)]$, i.e. within a gap in $\operatorname{supp}(F)$.

C.3 Median censorship

This subsection shows that, for $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$ and every prior, $(G_{1/2}, G_{1/2})$ is an equilibrium, and $q = \frac{1}{2}$ is the only solution of (QC′) with $\theta < m(q)$. Here $\tilde{v}(x) = -x|x|$. For $(q, \theta) \in \Gamma$ with $\theta < m$ and $u < q$, $F^{-1}(u) \leq \theta < m$, so the integrand in (QC′) is

$$-(m - F^{-1}(u))^2 + (\theta - F^{-1}(u))^2 + 2(m-\theta)(m - F^{-1}(u)) = (m-\theta)^2,$$

and the left-hand side of (QC′) equals $(m-\theta)^2(2q-1)$. So $q = \frac{1}{2}$ is the only solution with $\theta < m(q)$. If $\overline{q} > \frac{1}{2}$, then $(\frac{1}{2}, F^{-1}(\frac{1}{2})) \in \Gamma$ with $F^{-1}(\frac{1}{2}) < m(\frac{1}{2})$ solves it, and $(G_{1/2}, G_{1/2})$ is an equilibrium by Lemma 12 (ii). If $\overline{q} \leq \frac{1}{2}$, then $G_{1/2} = F$, and $w'_F(x) = -2\int |x-t| dF(t)$ has right derivative $2(1-2F(x)) \geq 0$ on $[\underline{\theta}, \overline{\theta})$, as $F(x) \leq F(\overline{\theta}-) = \overline{q} \leq \frac{1}{2}$ there. So (F, F) is an equilibrium by Step 2 of the proof of Lemma 12. If $\underline{\theta} = \overline{\theta}$, the claim is trivial.

C.4 Full revelation and the top atom

Lemma 14. *Let $v(x)$ be well S-shaped, F a prior on $[0,1]$ with $\underline{\theta} < \overline{\theta}$, and $\alpha := F(\overline{\theta}) - F(\overline{\theta}-)$. If F is a best reply to F, then $\alpha \geq \frac{1}{2}$. More precisely:*

(i) If $\overline{\theta}$ is isolated in $\operatorname{supp}(F)$, let $t := \max\big(\operatorname{supp}(F) \cap [0, \overline{\theta})\big)$. Then $\alpha \geq 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(\overline{\theta}-t)}$, and if $\operatorname{supp}(F) = \{t, \overline{\theta}\}$, this condition is also sufficient.

(ii) Otherwise, F is a best reply to F if and only if $\alpha|\tilde{v}''_+(0)| \geq \int_{[0,\overline{\theta})} |\tilde{v}''_-(\overline{\theta}-y)|\,dF(y)$; this requires $\tilde{v}''_+(0) \neq 0$.

Proof. Let $w := w_F$, $g := w'$ and $L(y) := w(\overline{\theta}) - w'(\overline{\theta})(\overline{\theta}-y)$, the tangent to w at $\overline{\theta}$, with w_F as in the proof of Lemma 12. Throughout, I use that for $0 \leq b \leq a \leq 1$,

$$R(a,b) := \tilde{v}(a) - \tilde{v}(b) - \tilde{v}'(a)(a-b) \geq \tilde{v}(a-b) - (a-b)\tilde{v}'(a-b) \geq -\tilde{v}(a-b). \qquad \text{(R)}$$

Indeed, let $\delta := a-b$. Then $R(a,b) = \int_0^\delta [\tilde{v}'(b+t) - \tilde{v}'(a)]\,dt$. As $\tilde{v}'$ is concave on $[0,1]$, $\tilde{v}'(t+b) - \tilde{v}'(t)$ is non-increasing in t, so for $t \in [0,\delta]$, $\tilde{v}'(b+t) - \tilde{v}'(a) \geq \tilde{v}'(t) - \tilde{v}'(\delta)$, and hence $R(a,b) \geq \int_0^\delta [\tilde{v}'(t) - \tilde{v}'(\delta)]\,dt = \tilde{v}(\delta) - \delta\tilde{v}'(\delta)$, which is at least $-\tilde{v}(\delta)$ as $\mathcal{E}_{\tilde{v}}(\delta) \geq 2$ (Lemma 3).

(i) As $\overline{\theta} \in \operatorname{supp}(F)$ is isolated, $\alpha > 0$, $\delta := \overline{\theta} - t \in (0,1]$ and $F((t,\overline{\theta})) = 0$. First, $w \geq L$ on $\operatorname{supp}(F) \cap [0,\overline{\theta})$. Let $B \subseteq [0,\overline{\theta})$ be a Borel set with $\beta := F(B) > 0$ and $b := \mathbb{E}_F[\theta|\theta \in B]$. For $\varepsilon \in (0,\beta]$, the signal that pools the atom at $\overline{\theta}$ with a share $\frac{\varepsilon}{\beta}$ of the types in B at $m_\varepsilon := \frac{\alpha\overline{\theta}+\varepsilon b}{\alpha+\varepsilon}$ and reveals all other types is a mean-preserving contraction of F. Its gain over F is

$$(\alpha+\varepsilon)w(m_\varepsilon) - \alpha w(\overline{\theta}) - \frac{\varepsilon}{\beta}\int_B w\,dF = \frac{\varepsilon}{\beta}\int_B (L-w)\,dF + o(\varepsilon),$$

since $m_\varepsilon - \overline{\theta} = -\frac{\varepsilon(\overline{\theta}-b)}{\alpha+\varepsilon}$, w is differentiable at $\overline{\theta}$, and $L(b) = \frac{1}{\beta}\int_B L\,dF$ as L is affine. So $\int_B (w-L)\,dF \geq 0$ for every such B, i.e. $w \geq L$ F-a.e. on $[0,\overline{\theta})$, and on $\operatorname{supp}(F) \cap [0,\overline{\theta})$ by continuity. In particular, as $\tilde{v}$ is odd and $\tilde{v}(0) = \tilde{v}'(0) = 0$,

$$0 \leq w(t) - L(t) = -\alpha\tilde{v}(\delta) - \int_{[0,t]} R(\overline{\theta}-y, t-y)\,dF(y) \leq -\alpha\tilde{v}(\delta) - (1-\alpha)\big(\tilde{v}(\delta) - \delta\tilde{v}'(\delta)\big),$$

by (R) with $a = \overline{\theta}-y$, $b = t-y$. Dividing by $-\tilde{v}(\delta) > 0$ gives $\alpha \geq (1-\alpha)(\mathcal{E}_{\tilde{v}}(\delta) - 1)$, i.e. $\alpha \geq 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(\delta)}$, which is at least $\frac{1}{2}$ by Lemma 3. If $\operatorname{supp}(F) = \{t, \overline{\theta}\}$, every feasible signal

is supported on $[t, \overline{\theta}]$, and the condition is equivalent, as in the proof of Proposition 1, to the secant ℓ of w from t to $\overline{\theta}$ being weakly flatter than $w'(\overline{\theta})$. Then $D := \ell - w$ has $D(t) = D(\overline{\theta}) = 0$ and a convex derivative with $D'(\overline{\theta}) \leq 0$, so D' is positive and then non-positive on $[t, \overline{\theta}]$, $D \geq 0$ there, and $\int w\, dG' \leq \int \ell\, dG' = \int \ell\, dF = \int w\, dF$ for every feasible G'.

(ii) Now $\overline{\theta}$ is a limit point of $\mathrm{supp}(F) \cap [0, \overline{\theta})$, so $F([0, \overline{\theta})) = 1 - \alpha > 0$. Suppose that $g'_-(\overline{\theta}) < 0$. As the left derivative of the concave g is left-continuous, g is strictly decreasing on some $[\overline{\theta} - \eta, \overline{\theta}]$, so w is strictly concave there. This interval contains infinitely many points of $\mathrm{supp}(F)$, so pooling all types in it at their mean is feasible and, by the strict Jensen inequality, strictly better than F. Hence F is a best reply only if $g'_-(\overline{\theta}) \geq 0$. For $h \in (0, 1)$, $\frac{g(\overline{\theta}) - g(\overline{\theta} - h)}{h} = \int \frac{\tilde{v}'(\overline{\theta} - y) - \tilde{v}'(\overline{\theta} - y - h)}{h}\, dF(y)$, and as $\tilde{v}'$ is concave, each integrand is non-increasing as $h \downarrow 0$ and converges to $\tilde{v}''_-(\overline{\theta} - y)$, so

$$g'_-(\overline{\theta}) = \alpha |\tilde{v}''_+(0)| - \int_{[0, \overline{\theta})} |\tilde{v}''_-(\overline{\theta} - y)|\, dF(y),$$

by monotone convergence, using $\tilde{v}''_-(0) = |\tilde{v}''_+(0)|$ and $\tilde{v}''_-(z) < 0$ for $z \in (0, 1]$ (property 2) in Appendix A.1). So the condition is necessary. Conversely, if it holds, then $g'_-(\overline{\theta}) \geq 0$, so the concave g is non-decreasing on $[0, \overline{\theta}]$, and F is a best reply to F by Step 2 of the proof of Lemma 12. As $-\tilde{v}'$ is convex on $[0, 1]$ with $-\tilde{v}'(0) = 0$, $|\tilde{v}''_-(z)| \geq |\tilde{v}''_+(0)|$ and $|\tilde{v}''_-(z)| > 0$ for $z \in (0, 1]$, so the condition requires $\tilde{v}''_+(0) \neq 0$ and $\alpha \geq 1 - \alpha$. □

For $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$, $\tilde{v}(x) = -x|x|$, so both bounds read $\alpha \geq \frac{1}{2}$, and by Appendix C.3 full revelation is an equilibrium if and only if $\alpha \geq \frac{1}{2}$. For rules with $\tilde{v}''_+(0) = 0$, such as those generated by normal or Gumbel shocks, (ii) excludes full revelation whenever types accumulate at $\overline{\theta}$. For the binary prior, (i) is Proposition 1: $t = 0$, $\overline{\theta} = 1$ and $\alpha = p$.

C.5 Proof of the bounds in Proposition 9

The proof also shows that the utility bound holds with equality if and only if $v(x) = \frac{1}{2} + x - \frac{1}{2}x|x|$, F is degenerate, or $v(1) = 1$ and F puts mass $p \geq \overline{p}$ on 1 and $1 - p$ on 0.

Write $x_u := F^{-1}(u)$, $p := \mathbb{E}_F[\theta]$ and $\overline{q} := F(\overline{\theta}-)$, so that $G_q = F$ for every $q \geq \overline{q}$ (Appendix C.2). If $\underline{\theta} = \overline{\theta}$, every feasible signal is $\{\overline{\theta}\} = G_0$, and the firm's utility is $\overline{\theta}$ under every rule; so let $\underline{\theta} < \overline{\theta}$, hence $\overline{q} > 0$. Let (G, G) be a symmetric equilibrium with $G = G_{q_0}$, and let $q := \min(q_0, \overline{q})$, so that $G = G_q$, and $q < \overline{q}$ unless $G = F$.

Step 1: $q \leq \frac{1}{2}$. If $G = F$, then $q = \overline{q} = 1 - \alpha \leq \frac{1}{2}$ by Lemma 14. Let $q < \overline{q}$, so that $x_{q^+} < m := m(q) < \overline{\theta}$, and let $w := w_{G_q}$. I show that G_q solves (QC′). As G_q is a best reply to G_q and w is Lipschitz, as v' is bounded, there is a convex $\phi \geq w$ on $[\underline{\theta}, \overline{\theta}]$ with $\int \phi \, dF = \int w \, dG_q$ (Dworczak and Martini, 2019, Theorem 2). Then $\int w \, dG_q \leq \int \phi \, dG_q \leq \int \phi \, dF$ are equalities, so $\phi = w$ G_q-a.e., and Jensen's inequality $(1 - q)\phi(m) \leq \int_q^1 \phi(x_u) \, du$ is an equality. As the law of x_U given $U > q$ has x_{q^+} and $\overline{\theta}$ as the infimum and supremum of its support, ϕ coincides on $(x_{q^+}, \overline{\theta})$ with an affine L. Since $\phi \geq w$ with equality at the atom m of G_q, L is the tangent to w at m, and $\phi \geq L$ on $[\underline{\theta}, \overline{\theta}]$, as ϕ is convex. Let $\Delta(\theta) := L(\theta) - w(\theta)$, the left-hand side of (QC′) at (q, θ). On $(x_{q^+}, m]$, $\Delta = \phi - w \geq 0$, so $\Delta(x_{q^+}) \geq 0$. Moreover, $\phi(x_q) = w(x_q)$: if $x_q > \underline{\theta}$, then ϕ is continuous at $x_q < \overline{\theta}$, $\phi(x_u) = w(x_u)$ for almost every $u < q$, and $x_u \to x_q$ as $u \uparrow q$; if $x_q = \underline{\theta}$, then G_q has an atom of mass at least q at $\underline{\theta}$. If $q = 0$, then $x_{0^+} = \underline{\theta}$, $m = p$ and $w(\theta) = \tilde{v}(\theta - p)$, so $L \equiv 0$ and $\Delta(\underline{\theta}) = \tilde{v}(p - \underline{\theta}) < 0$, which contradicts $\Delta(\underline{\theta}) \geq 0$; so $q > 0$. So $\Delta(x_q) \leq \phi(x_q) - w(x_q) = 0$, and $\Delta(\theta) = 0$ for some $\theta \in [x_q, x_{q^+}]$, where $\theta < m$.

So (q, θ) solves (QC′). Let $d := m - \theta \in (0, 1]$. By (QC′), $E(q) := \int_0^q R(m - x_u, \theta - x_u) \, du + (1 - q)\tilde{v}(d) = 0$, with R as in (R), and $0 \leq \theta - x_u \leq m - x_u \leq 1$ for $u < q$. By (R), $R(m - x_u, \theta - x_u) \geq -\tilde{v}(d)$, and since $\tilde{v}(d) < 0$, this gives $E(q) \geq -(2q - 1)\tilde{v}(d) > 0$ for $q > \frac{1}{2}$, so every root satisfies $q \leq \frac{1}{2}$.

Step 2: accuracy. For any rule of the signal gap and any symmetric profile, as $v(x) - \frac{1}{2}$ is odd, $s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) = \frac{s_1+s_2}{2} + h_v(|s_1 - s_2|)$ with $h_v(\delta) := \delta\big(v(\delta) - \frac{1}{2}\big)$, so $u_F = p + \mathbb{E}[h_v(|s_1 - s_2|)]$. By Lemma 9, $h_v(\delta) \le h^*(\delta) := \delta^2 - \frac{\delta^3}{2}$ on $[0,1]$, with equality at some $\delta \in (0,1)$ only if $v = v^*$, and at $\delta = 1$ if and only if $v(1) = 1$. Hence $u_F \le U(q) := p + \mathbb{E}_{G_q \times G_q}[h^*(|s_1 - s_2|)]$, with equality if and only if $h_v(|s_1 - s_2|) = h^*(|s_1 - s_2|)$ almost surely.

Step 3: U is non-decreasing on $[0, \frac{1}{2}]$, and $U(q) < U(\frac{1}{2})$ for $q < \min(\frac{1}{2}, \overline{q})$. In this step, $\theta := x_q$, $m := m(q)$ and $d := m - \theta$, so $d > 0$ if and only if $q < \overline{q}$. $U(q) - p = \int_0^q \int_0^q h^*(|x_u - x_w|)\, du\, dw + 2(1-q) \int_0^q h^*(m - x_u)\, du$. As $(1-q)m(q) = \int_q^1 x_u\, du$, m and hence U are Lipschitz on $[0, \frac{1}{2}]$, and at every continuity point q of x_u, hence for almost every q, $m'(q) = \frac{d}{1-q}$, and $\frac{1}{2}U'(q) = -\int_0^q R^*(m - x_u, \theta - x_u)\, du + (1-q)h^*(d)$, $R^*(a,b) := h^*(a) - h^*(b) - h^{*\prime}(a)(a-b)$. As h^* is a cubic, $R^*(a, a-d) = -d^2 + \frac{3}{2}ad^2 - \frac{d^3}{2}$, so, with $\mu := \frac{1}{q}\int_0^q x_u\, du$, $\frac{1}{2}U'(q) = d^2[1 + \frac{d}{2}(2q-1) - \frac{3}{2}q(m-\mu)] \ge d^2[1 - \frac{1-2q}{2} - \frac{3}{2}q] = d^2\,\frac{1-q}{2}$, using $d \le 1$, $m - \mu \le 1$ and $q \le \frac{1}{2}$. So $U' \ge 0$ almost everywhere on $[0, \frac{1}{2}]$, and $U' > 0$ almost everywhere on $(0, \min(\frac{1}{2}, \overline{q}))$, which proves Step 3. For $q \in [\overline{q}, \frac{1}{2}]$, $G_q = G_{1/2}$ and $U(q) = U(\frac{1}{2})$.

Conclusion. Under $v^*(x)$, $h_{v^*} = h^*$, and $(G_{1/2}, G_{1/2})$ is an equilibrium (Appendix C.3). Conversely, it is its only symmetric equilibrium in quantile censorships: by Step 1, an equilibrium G_q with $q < \overline{q}$ solves (QC′) with $\theta < m(q)$, so $q = \frac{1}{2}$ by Appendix C.3, and $G = F$ requires $\overline{q} \le \frac{1}{2}$ by Lemma 14, so that $F = G_{1/2}$. Hence $U(\frac{1}{2})$ is the firm's utility in the median-censorship equilibrium of $v^*(x)$. By Steps 1–3, $u_F \le U(q) \le U(\frac{1}{2})$.

Equality requires $U(q) = U(\frac{1}{2})$, i.e. $G = G_{1/2}$ by Steps 1 and 3, and equality in Step 2. If $v \ne v^*$, the latter means that $|s_1 - s_2| \in \{0, 1\}$ almost surely under $G_{1/2} \times G_{1/2}$, and $v(1) = 1$ if the gap 1 has positive probability. So $\mathrm{supp}(G_{1/2})$ is a single point or $\{0, 1\}$. In the first case, the revealed types and the pooled mean $m(\frac{1}{2})$ coincide, so F is degenerate. In the second, $m(\frac{1}{2}) = 1$, as it is the largest point of the support, so $x_u = 1$

for $u > \frac{1}{2}$ and $x_u \in \{0,1\}$ for $u < \frac{1}{2}$: F puts mass $p \geq \frac{1}{2}$ on 1 and $1-p$ on 0, and $G = F$, which by Proposition 1 requires $p \geq \overline{p}$; moreover $v(1) = 1$. Conversely, equality holds in these cases: trivially if F is degenerate, and for the binary prior with $p \geq \overline{p}$ and $v(1) = 1$, both $v(x)$ and $v^*(x)$ induce full revelation, as $\overline{p} \geq \frac{1}{2}$ is the threshold of $v(x)$ and $\frac{1}{2}$ that of $v^*(x)$, and $u_F = p + 2p(1-p)\big(v(1) - \frac{1}{2}\big) = 2p - p^2$ under both.

C.6 A convexity lemma

The following lemma extends the converse of the Hermite–Hadamard inequality (see Niculescu and Persson (2003)), which is the case of a uniform K.

Lemma 15. *Let a CDF K and a function h be continuous on an interval J, with K strictly increasing. If $h\big(\mathbb{E}_K[\theta|\theta \in J']\big) \leq \mathbb{E}_K[h(\theta)|\theta \in J']$ for every interval $J' \subseteq J$, then h is convex.*

Proof. Let $x < z$ in J, let ℓ be the secant of h through x and z, and suppose that $g := h - \ell$ has a positive maximum on $[x,z]$, first attained at $c \in (x,z)$. For $\delta > 0$ small enough that $\mathbb{E}_K[\theta|\theta \in [c-\delta, z]] > c$, there is $\delta' \in (0, z-c)$ such that $J' := [c-\delta, c+\delta']$ satisfies $\mathbb{E}_K[\theta|\theta \in J'] = c$, as K is continuous and strictly increasing. Since $g < g(c)$ on $[c-\delta, c)$, which has positive K-mass, and ℓ is affine, $\mathbb{E}_K[h(\theta)|\theta \in J'] = \ell(c) + \mathbb{E}_K[g(\theta)|\theta \in J'] < h(c)$, a contradiction. Hence $h \leq \ell$ on $[x,z]$. □

Lemma 16. *Let F be a prior, $\underline{\theta} = \min \operatorname{supp}(F)$, $\overline{\theta} = \max \operatorname{supp}(F)$, and let V be bounded and upper semicontinuous on $[\underline{\theta}, \overline{\theta}]$. If $\int V\, dG \leq \int V\, dF$ for every $G \leq_{cx} F$, then there is a bounded convex $\phi \geq V$ on $[\underline{\theta}, \overline{\theta}]$ with $\phi = V$ on $\operatorname{supp}(F)$. In particular, if $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$, then V is convex on $[\underline{\theta}, \overline{\theta}]$.*

Proof. Let $S := \operatorname{supp}(F)$, let $x < y < z$ with $x, z \in S$, and let $y = \lambda x + (1-\lambda)z$. For $0 < \varepsilon < \min(y-x, z-y)$, let $A := [x-\varepsilon, x+\varepsilon]$ and $B := [z-\varepsilon, z+\varepsilon]$; then $F(A), F(B) > 0$, and the conditional means $m_A \in A$, $m_B \in B$ satisfy $m_A < y < m_B$. Choose $\alpha, \beta \in (0,1]$ with $\alpha \int_A \theta\, dF + \beta \int_B \theta\, dF = (\alpha F(A) + \beta F(B))\, y$, and let G be obtained from F by moving

the mass $\alpha F|_A + \beta F|_B$ to an atom at y. By Jensen's inequality $G \leq_{cx} F$, so with $\lambda_\varepsilon := \alpha F(A)/(\alpha F(A) + \beta F(B))$, $V(y) \leq \lambda_\varepsilon \sup_A V + (1 - \lambda_\varepsilon) \sup_B V$. As $\varepsilon \to 0$, $\lambda_\varepsilon \to \lambda$, and by upper semicontinuity $V(y) \leq \lambda V(x) + (1 - \lambda)V(z)$. Let $\tilde{V} := V$ on S and $\tilde{V} := +\infty$ elsewhere, and let $\phi(y) := \inf \sum_i \lambda_i \tilde{V}(x_i)$ over finite convex combinations with $\sum_i \lambda_i x_i = y$, the convex hull of $\tilde{V}$. For fixed points x_i, the minimum over the polytope $\{\lambda \geq 0 : \sum_i \lambda_i = 1,\ \sum_i \lambda_i x_i = y\}$ is attained at a vertex, which has at most two positive entries. So ϕ is convex, $\phi \geq V$ on $[\underline{\theta}, \overline{\theta}]$ by the chord inequality, $\phi \leq V$ on S, and $\inf V \leq \phi \leq \sup V$, as $\underline{\theta}, \overline{\theta} \in S$. If $S = [\underline{\theta}, \overline{\theta}]$, then $V = \phi$. □

C.7 Proof of Corollary 3

Let $c = 2p - \underline{\theta}$ and $G = U[\underline{\theta}, c]$. As F is concave, it lies above the uniform CDF on $[\underline{\theta}, \overline{\theta}]$, so $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$, i.e. $c \leq \overline{\theta}$. Moreover, $F - G$ is concave on $[\underline{\theta}, c]$ and non-negative at $\underline{\theta}$, while $F - G = F - 1 \leq 0$ on $[c, \overline{\theta}]$, so $F - G$ changes sign at most once, from positive to negative. Since F and G have the same mean, G is a mean-preserving contraction of F by the single-crossing condition, i.e. it is feasible. Against G, $\overline{v}(s) = G(s)$, which is concave on $[\underline{\theta}, \overline{\theta}]$, so by Jensen's inequality every feasible G' gives $\mathbb{E}_{G'}[G] \leq G(p) = \frac{1}{2} = \mathbb{E}_G[G]$, and (G, G) is an equilibrium. Finally,

$$u_F = \mathbb{E}[\max(s_1, s_2)] = \underline{\theta} + \frac{2}{3}(c - \underline{\theta}) = \frac{4p - \underline{\theta}}{3}.$$

For uniqueness, let (G, G') be an equilibrium, so that (G', G') and (G', G) are equilibria too. As in Appendix C.8.4, G' has no atom in $(\underline{\theta}, \overline{\theta}]$, and an atom $a > 0$ at $\underline{\theta}$ would not be a best reply to G' either: pooling it with a small mass ε of G' above $\underline{\theta}$ raises the payoff per unit of the atom from $\frac{a}{2}$ to at least a, at a total cost of at most $\varepsilon < \frac{a^2}{2}$. So G' is atomless. As G' is a best reply to G and $G(s) = \min\{\frac{s-\underline{\theta}}{c-\underline{\theta}}, 1\}$ is concave, $\mathbb{E}_{G'}[G] \leq G(p) = \frac{1}{2}$ with equality only if $\mathrm{supp}(G') \subseteq [\underline{\theta}, c]$. As G is a best reply to G' and every mean-preserving contraction of G is feasible, pooling the types of G in any interval $J \subseteq [\underline{\theta}, c]$ does not

pay, so G' is convex on $[\underline{\theta}, c]$ by Lemma 15 with $K = G$. Hence $G' \leq G$ on $[\underline{\theta}, c]$, as G is the secant of G' there, and $\int_{\underline{\theta}}^{c}(G - G')dx = \mathbb{E}_{G'}[s] - \mathbb{E}_G[s] = 0$, so $G' = G$.

C.8 Proofs of the benchmark results

Throughout, a rule is a general selection rule, and $V_G(x) = \int H(x, y)\, dG(y)$ is the agent's value function against a rival who plays G. For a CDF K, let $I_K(x) := \int_{-\infty}^{x} K(t)\, dt$; for distributions with the same mean, $G \leq_{cx} F$ if and only if $I_G \leq I_F$.

C.8.1 Two tools

Lemma 17. *In every symmetric profile (G, G), the firm's utility is $u_F = p + 2\,\mathrm{Cov}_G\big(s, V_G(s)\big)$.*

Proof. With s_1, s_2 i.i.d. G, $u_F = \mathbb{E}[s_2] + \mathbb{E}[(s_1 - s_2)H(s_1, s_2)]$. Since $\kappa := H - \frac{1}{2}$ is anti-symmetric, $\mathbb{E}[s_2\kappa(s_1, s_2)] = -\mathbb{E}[s_1\kappa(s_1, s_2)]$, so $\mathbb{E}[(s_1 - s_2)H(s_1, s_2)] = 2\mathbb{E}[s_1\kappa(s_1, s_2)] = 2\mathbb{E}\big[s_1\big(V_G(s_1) - \frac{1}{2}\big)\big]$, and $\mathbb{E}[V_G(s)] = \frac{1}{2}$. □

Lemma 18. *Let F be any prior and $V : [\underline{\theta}, \overline{\theta}] \to [0, 1]$. There is a rule with $V_F = V$ on* $\mathrm{supp}(F)$ *if and only if $V(\theta) \leq_{cx} U[0, 1]$ for $\theta \sim F$, i.e.,*

$$\mathbb{E}_F[V(\theta)] = \tfrac{1}{2} \qquad \textit{and} \qquad \mathbb{E}_F\big[(V(\theta) - t)^+\big] \leq \tfrac{(1-t)^2}{2} \ \textit{ for all } t \in [0, 1].$$

Then one such rule is $H(x, y) = \mathbb{P}(S_x > S_y) + \frac{1}{2}\mathbb{P}(S_x = S_y)$, where the agents' scores are independent, the score S_x of signal x has mean $V(x)$, and $S_\theta \sim U[0, 1]$ for $\theta \sim F$.

Proof. Necessity: $\mathbb{E}_F[V] = \frac{1}{2}$ by symmetry. For every set A, the winner's type lies in A with probability $2\int_A V\, dF$, at most the probability $1 - (1 - F(A))^2$ that some type lies in A. Applied to the upper level sets of V, this bounds the sum of the top m quantiles of $V(\theta)$ by $m - \frac{m^2}{2}$, the same sum for $U[0, 1]$, at every mass m of such a set, and hence at every $m \in [0, 1]$: between two such masses, $V(\theta)$ has an atom, so the left side is linear in m, while the right side is concave. With equal means, this is the stated condition.

Sufficiency: by Strassen's theorem, there is a kernel $\kappa(\cdot|z)$ with mean z that spreads the distribution of $V(\theta)$ into $U[0,1]$; let $S_x \sim \kappa(\cdot|V(x))$, and for x with $V(x)$ outside the support of the law of $V(\theta)$, let S_x be any law on $[0,1]$ with mean $V(x)$. Against a fully revealing rival, whose score is $U[0,1]$, $\mathbb{P}(S_x > S_y) = \mathbb{E}[S_x] = V(x)$, and the rule is symmetric and constant-sum by construction. □

The second lemma is the feasibility claim in Subsection 5.2: noise can only make winning probabilities less dispersed than exact ranking does, and the noiseless rule is the boundary case, with $V_F = F$ and $F(\theta) \sim U[0,1]$.

C.8.2 Proof of Proposition 8

I first show that, for priors (p,p), H attains the first best $2p - p^2$ in some equilibrium if and only if $H(1,0) = 1$ and $(1-p)H(x,0) + p\,H(x,1) \leq \frac{1-p+x}{2}$ for all $x \in (0,1)$. For any signals, $u_F = \mathbb{E}[s_2 + (s_1 - s_2)H(s_1,s_2)] \leq \mathbb{E}[\max(s_1,s_2)] \leq \mathbb{E}[s_1 + s_2 - s_1 s_2] = 2p - p^2$, where the second inequality holds as $\mathbb{E}[\max(\theta_1,\theta_2)|s_1,s_2] = s_1 + s_2 - s_1 s_2$, and it is an equality if and only if $\min(s_1,s_2)\big(1 - \max(s_1,s_2)\big) = 0$ almost surely. If both agents put mass on $(0,1)$, this fails with positive probability, so in an equilibrium that attains the first best, some agent, say agent 1, reveals fully. As the game is symmetric and constant-sum, its equilibrium strategies form the same set for both agents, so $(\{0,1\};\{0,1\})$ is an equilibrium too. The first inequality is an equality only if $H(1,s) = 1$ for almost every signal $s < 1$ of agent 2, and agent 2 sends some $s < 1$, as $p < 1$. As H is nondecreasing in the own signal and constant-sum, $H(1,0) = 1 - H(0,1) \geq 1 - H(s,1) = H(1,s) = 1$. Against a fully revealing rival, the agent's value function is $V(x) = (1-p)H(x,0) + p\,H(x,1)$, with $V(0) = \frac{1-p}{2}$ and $V(1) = 1 - \frac{p}{2}$ when $H(1,0) = 1$, so full revelation is a best reply if and only if V lies below its chord from 0 to 1, which is $\frac{1-p+x}{2}$. Conversely, if $H(1,0) = 1$ and this holds, $(\{0,1\};\{0,1\})$ is an equilibrium, and the firm selects a good agent over a bad one for

sure, which attains $2p - p^2$.

(i) This condition is linear in p, so it holds for all $p \in (0,1)$ if and only if it holds in the limits $p \to 0$ and $p \to 1$, which give $H(x,0) \leq \frac{1+x}{2}$ and $H(x,1) \leq \frac{x}{2}$. If both are strict on $(0,1)$, then against a fully revealing rival with any prior p_j, $V(x) < \frac{1-p_j+x}{2}$ on $(0,1)$ while V equals the chord at 0 and 1, so full revelation is the unique best reply for every own prior, as every other signal puts mass on $(0,1)$. So $(\{0,1\};\{0,1\})$ is an equilibrium for all $(p_1,p_2) \in (0,1)^2$; if (G_1,G_2) is another, then by interchangeability so is $(G_1,\{0,1\})$, so G_1 is a best reply to full revelation, and $G_1 = \{0,1\}$, and likewise $G_2 = \{0,1\}$.

(iii) The pass/fail rule is nondecreasing in the own signal, $H(1,0) = 1$, $H(x,0) = \frac{1}{2} < \frac{1+x}{2}$ and $H(x,1) = 0 < \frac{x}{2}$ for $x \in (0,1)$, so (i) applies.

C.8.3 The first best for a general support

Lemma 19. *Let F be any prior, $S := \operatorname{supp}(F) \subseteq [\underline{\theta}, \overline{\theta}]$, and $\bar{F}(x) := F(x^-) + \frac{1}{2}F(\{x\})$.*

(i) If $\bar{F}$ coincides on S with some function convex on $[\underline{\theta}, \overline{\theta}]$, the rule that rounds each signal down to the support, $r(x) := \max\left(S \cap [\underline{\theta}, x]\right)$, and selects the higher rounded signal, tossing a coin at ties, attains the first best, with full revelation as an equilibrium.

(ii) If some general selection rule nondecreasing in the own signal attains the first best in a fully revealing equilibrium, then $\bar{F}$ coincides on S with some function convex on $[\underline{\theta}, \overline{\theta}]$.

(iii) If $S = [\underline{\theta}, \overline{\theta}]$, every equilibrium of every general selection rule that attains the first best is fully revealing, and some general selection rule nondecreasing in the own signal attains the first best in some equilibrium if and only if F is convex on $[\underline{\theta}, \overline{\theta})$ and has no atom except possibly at $\overline{\theta}$; the noiseless rule then attains it, with full revelation as an equilibrium.

Proposition 10 is part (iii), which also shows that every equilibrium that attains

the first best is fully revealing, with the uniqueness claim proved in Appendix C.8.4, and parts (i) and (ii) extend it to supports that are not intervals. Every binary prior satisfies the condition of part (i), and the rounding rule is then the pass/fail rule of Proposition 8.

Proof. Here $\bar{F}(x)$ is the winning probability of type x against a fully revealing rival when the higher type is selected and ties are split, as symmetry forces. In terms of F, the condition in (i) and (ii) says that F is convex on every interval contained in S, every atom of F below $\overline{\theta}$ is an isolated point of S, and the chords of $\bar{F}$ across the gaps of S have nondecreasing slopes.

(i) *Sufficiency, for any support.* Let ψ be convex with $\psi = \bar{F}$ on S, and let ϕ equal $\bar{F}$ on S and be affine on each gap (a, b) of S. Then ϕ is convex: on each interval contained in S it equals ψ, and at the ends of a gap its slopes are ordered, as the chord of the convex ψ over $[a, b]$ has slope between $\psi'(a^+)$ and $\psi'(b^-)$, and consecutive chords of ψ have nondecreasing slopes. As $\bar{F}$ is nondecreasing on S, $\phi \geq \bar{F}(a)$ on a gap (a, b). Let $r(x) := \max\left(S \cap [\underline{\theta}, x]\right)$ and let the rule select the higher rounded signal, tossing a coin at ties; it is nondecreasing in the own signal. Against a fully revealing rival, $V_F = \bar{F}$ on S and $V_F(x) = \bar{F}(r(x)) \leq \phi(x)$ off S. So for every $G \leq_{cx} F$, $\int V_F \, dG \leq \int \phi \, dG \leq \int \phi \, dF = \int V_F \, dF$, and full revelation is an equilibrium. Distinct types are never rounded to the same point, so the higher type is always selected.

(ii) *Necessity, for any support and a rule nondecreasing in the own signal.* Under full revelation, the first best requires $H(x, y) = 1$ for $F \otimes F$-almost all $x > y$, and $H(x, x) = \frac{1}{2}$ by symmetry, so $V_F = \bar{F}$ F-almost everywhere and at every atom; as V_F is nondecreasing, $V_F(x) \in [F(x^-), F(x)]$ for every $x \in S$. Three deviations, each a mean-preserving contraction of F, rule out the three ways in which $\bar{F}|_S$ can fail to extend convexly. (a) An atom $a < \overline{\theta}$ of mass m that is a limit of S from the right: pooling a fraction α of the atom with all types in $(a, a + \delta)$, of mass $\mu_\delta > 0$, gives a

signal in $(a, a+\delta)$ with winning probability at least $F(a)$, so the atom gains at least $\alpha\frac{m}{2}$, while the other types lose at most $\mu_\delta(F(a+\delta)-F(a))$, which is smaller for small δ and $\alpha = \mu_\delta$. (b) An atom $a < \overline{\theta}$ that is a limit of S from the left only: then a is the left end of a gap (a, b), and pooling the types in $(a-\delta, a)$, of mass μ_δ, with a mass $w = O(\mu_\delta\delta)$ of the types at b into the signal a raises the former from at most $F(a^-)$ to $\bar{F}(a) = F(a^-) + \frac{m}{2}$ and costs the latter at most w. (c) Points $x < y < z$ of S with $\bar{F}(y)$ above the chord of $\bar{F}$ through x and z: pooling types near x and near z (at x and z themselves if they are atoms) into the signal y in the right proportions pays, as V_F is continuous at non-atoms of S by (a) and (b). If none of (a)–(c) applies, $\bar{F}|_S$ is convex on S and continuous at every limit point of S below $\overline{\theta}$, and its extension by chords is convex. This proves (ii).

(iii) *Interval support.* Let $S = [\underline{\theta}, \overline{\theta}]$ and let (G_1, G_2) attain the first best. The selected type must be the higher type almost surely, and selection depends only on the signals, so for almost all pairs of signals the conditional supports of θ_1 and θ_2 are ordered. If a signal of agent 1 pooled two types $a < b$ of S, the conditional supports of agent 2 would avoid (a, b), whose F-mass is positive. So both agents reveal fully, and (ii) applies: by (a), F has no atom below $\overline{\theta}$, and by (c), F is convex on $[\underline{\theta}, \overline{\theta})$. Conversely, then $\bar{F} = F$ on $[\underline{\theta}, \overline{\theta})$, and an atom at $\overline{\theta}$ only raises $\bar{F}(\overline{\theta})$ above $F(\overline{\theta}-)$, which keeps $\bar{F}$ convex on $[\underline{\theta}, \overline{\theta}]$; with no gaps, the rounding rule of (i) is the noiseless rule. □

C.8.4 Uniqueness under the noiseless rule

Let $\mathrm{supp}(F) = I := [\underline{\theta}, \overline{\theta}]$, and let F be convex on $[\underline{\theta}, \overline{\theta})$ with no atom except possibly one of mass $m \geq 0$ at $\overline{\theta}$. For a distribution K, let $\bar{K}(x) := K(x^-) + \frac{1}{2}K(\{x\})$ be its mid-CDF, the payoff of a signal x against a rival who plays K under the noiseless rule. A signal G is feasible if and only if $I_G \leq I_F$ on I with equality at $\overline{\theta}$; then $\int_x^{\overline{\theta}} G \geq \int_x^{\overline{\theta}} F$ for $x < \overline{\theta}$, so $G(\overline{\theta}-) \geq F(\overline{\theta}-)$ and $G(\{\overline{\theta}\}) \leq m$. Let $\phi = \phi_c + j\mathbb{1}\{\overline{\theta}\}$, with ϕ_c convex

and continuous on I and a jump $j \geq 0$ at $\overline{\theta}$. Integrating by parts twice,

$$\int \phi \, dF - \int \phi \, dG = \int \big(I_F(y) - I_G(y)\big) d\phi_c'(y) + j\big(m - G(\{\overline{\theta}\})\big),$$

and both terms are nonnegative.

Let $(F, \hat{G})$ be an equilibrium. As all equilibria form a symmetric product set, $(\hat{G}, \hat{G})$ and $(\hat{G}, F)$ are equilibria too, and each agent gets $\frac{1}{2}$ in all of them, so $\int \bar{\hat{G}} \, d\hat{G} = \int \bar{\hat{G}} \, dF = \int \bar{F} \, d\hat{G} = \int \bar{F} \, dF = \frac{1}{2}$.

First, $\hat{G}$ has no atom in $[\underline{\theta}, \overline{\theta})$. An atom a at $\underline{\theta}$ would give $I_{\hat{G}}(x) \geq a(x - \underline{\theta}) > I_F(x)$ for x close to $\underline{\theta}$, as F is continuous at $\underline{\theta}$ with $F(\underline{\theta}) = 0$. If $\hat{G}$ had an atom $a > 0$ at $x_0 \in (\underline{\theta}, \overline{\theta})$, then $I_{\hat{G}}(x_0) < I_F(x_0)$, since otherwise $I_F - I_{\hat{G}} \geq 0$ would have a minimum at x_0 with left derivative $F(x_0) - \hat{G}(x_0^-)$ above its right derivative $F(x_0) - \hat{G}(x_0)$. So replacing the atom by masses $\frac{2a}{3}$ at $x_0 + \varepsilon$ and $\frac{a}{3}$ at $x_0 - 2\varepsilon$ is feasible for small ε, as $I_F - I_{\hat{G}} > 0$ near x_0, and against $\hat{G}$ it raises the payoff by $a^2(\frac{2}{3} - \frac{1}{2}) + o(1) > 0$, contradicting that $\hat{G}$ is a best reply to $\hat{G}$. So $\bar{\hat{G}} = \hat{G}$ on $[\underline{\theta}, \overline{\theta})$, and $\bar{\hat{G}}$ jumps at $\overline{\theta}$ by $\frac{1}{2}\hat{G}(\{\overline{\theta}\})$, which is zero if $m = 0$.

Second, $\hat{G}$ is convex on $[\underline{\theta}, \overline{\theta})$. As F is a best reply to $\hat{G}$, pooling the types in an interval $J \subseteq [\underline{\theta}, \overline{\theta})$ into their mean m_J does not pay, i.e., $\hat{G}(m_J) \leq \mathbb{E}_F[\hat{G}(\theta) | \theta \in J]$; as F is convex with support I, it is continuous and strictly increasing on $[\underline{\theta}, \overline{\theta})$, and Lemma 15 applies with $K = F$ and $h = \hat{G}$ on every $[\underline{\theta}, x]$ with $x < \overline{\theta}$.

Hence $\bar{F}$ and $\bar{\hat{G}}$ both have the form of ϕ, with jumps $\frac{m}{2}$ and $\frac{1}{2}\hat{G}(\{\overline{\theta}\})$ at $\overline{\theta}$. The equalities above give $\int \bar{F} \, d(F - \hat{G}) = 0$ and $\int \bar{\hat{G}} \, d(F - \hat{G}) = 0$, so all four nonnegative terms of the identity vanish. If $m > 0$, the jump term for $\bar{F}$ gives $\hat{G}(\{\overline{\theta}\}) = m$. The other terms give $I_F = I_{\hat{G}}$ on the supports in $(\underline{\theta}, \overline{\theta})$ of dF' and $d\hat{G}'$, and there $F = \hat{G}$, since each such point is a minimum of $I_F - I_{\hat{G}} \geq 0$, which is differentiable on $(\underline{\theta}, \overline{\theta})$, as F and $\hat{G}$ are continuous there. On each open interval of $(\underline{\theta}, \overline{\theta})$ off these supports, F and $\hat{G}$ are affine, and they agree at its endpoints: at points of the supports, at $\underline{\theta}$, where $F = \hat{G} = 0$, and at $\overline{\theta}$, where $F(\overline{\theta}-) = 1 - m = \hat{G}(\overline{\theta}-)$. Hence $\hat{G} = F$, and (F, F) is the

unique equilibrium.

C.8.5 Rules of the signal gap

This proves the claim after Proposition 10: with $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$, a rule of the signal gap attains the first best only if it is the noiseless rule. Let $d = s_1 - s_2$. For any rule of the signal gap and feasible G_1, G_2, since $v(-x) = 1 - v(x)$,

$$u_F = \mathbb{E}[\max(s_1, s_2)] - \mathbb{E}\big[|d|\big(1 - v(|d|)\big)\big] \leq \mathbb{E}[\max(s_1, s_2)] \leq \mathbb{E}[\max(\theta_1, \theta_2)].$$

By Lemma 19(iii), the first best requires full revelation by both agents and an F that is convex on $[\underline{\theta}, \overline{\theta})$ with no atom below $\overline{\theta}$, so F has a density on $[\underline{\theta}, \overline{\theta})$; then also $|d|\big(1 - v(|d|)\big) = 0$ almost surely. As $0 < |d| < \varepsilon$ with positive probability for every $\varepsilon > 0$, $v(x) = 1$ for arbitrarily small $x > 0$, i.e., $v(x)$ is the noiseless rule, as it is increasing and $v(x) \leq 1$.

C.8.6 Proof of Proposition 11

For any prior, full revelation is an equilibrium if there is a convex $\phi \geq V_F$ on $[\underline{\theta}, \overline{\theta}]$ with $\phi = V_F$ on $\operatorname{supp}(F)$, by Jensen's inequality, and, when V_F is upper semicontinuous, only if (Lemma 16, which extends the necessity part of Dworczak and Martini, 2019, Theorem 2); it is the unique equilibrium if V_F is strictly convex.

As full revelation is a best reply and V_F is upper semicontinuous, Lemma 16 gives a bounded convex $\phi \geq V_F$ on $[\underline{\theta}, \overline{\theta}]$ with $\phi = V_F$ on $\operatorname{supp}(F)$. Let $g := \phi - L$. It is convex, $g(\overline{\theta}) \leq 0$ as $\phi(\overline{\theta}) = V_F(\overline{\theta}) \leq 1 = L(\overline{\theta})$, and $\int g \, dF = \frac{1}{2} - \frac{1}{2} = 0$. If $g(x_1) \leq 0$, then $g \leq \max(g(x_1), g(\overline{\theta})) \leq 0$ on $[x_1, \overline{\theta}]$, so $g > 0$ exactly on some $[\underline{\theta}, x^*)$ and $g \leq 0$ on $[x^*, \overline{\theta}]$. Hence $\int x \, g \, dF = \int (x - x^*) g \, dF \leq 0$, with equality only if $g = 0$ F-almost everywhere, and by Lemma 17, $u_F = p + 2\operatorname{Cov}_F(\theta, \phi(\theta)) \leq p + 2\operatorname{Cov}_F(\theta, L(\theta)) =$

$p + \frac{\mathrm{Var}_F(\theta)}{\overline{\theta}-p}$. By Lemma 18, L is implementable if and only if $L(\theta) \leq_{cx} U[0,1]$, i.e., $F \leq_{cx} L^{-1}(U[0,1]) = \bar{U}$. The implementing kernel of Lemma 18, written in signal units, is a mean-preserving spread of each signal whose scores under full revelation are distributed as $\bar{U}$; Lemma 20 gives a monotone implementation. The bound is also approached by rules with full revelation as the unique equilibrium. Under L, the agent is indifferent among all feasible signals; mixing the rule with H^* of Proposition 19, whose V_F is strictly convex below the top type (see Appendix C.8.8), makes full revelation the unique equilibrium at a payoff arbitrarily close to the bound.

Every concave prior satisfies $F \leq_{cx} \bar{U}$: it lies above the uniform CDF on $[\underline{\theta}, \overline{\theta}]$, so $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$, and $F - \bar{U}$ is concave on $[\underline{\theta}, \overline{\theta}]$, vanishes at $\overline{\theta}$ and is non-positive below $\underline{\theta}$, so it crosses zero once, from below. A convex prior other than the uniform has $p > \frac{\underline{\theta}+\overline{\theta}}{2}$, so $2p - \overline{\theta} > \underline{\theta}$, and F puts mass below the support of $\bar{U}$.

Lemma 20. *Let F be any prior; in (iii), let F be atomless.*

(i) Every nondecreasing V that satisfies the condition of Lemma 18 equals V_F on $\mathrm{supp}(F)$ *for a rule nondecreasing in the own signal. One such rule is a lottery over coarse gradings: the firm draws a partition of $[\underline{\theta}, \overline{\theta}]$ into intervals, selects the agent whose signal lies in the higher interval, and tosses a coin if both lie in the same one; a signal at an atom of F draws its quantile uniformly from the atom's quantile interval, so the partition may split an atom.*

(ii) In particular, whenever $F \leq_{cx} \bar{U}$, the bound of Proposition 11 is attained by such a rule.

(iii) L is generated by a single random pass mark C, drawn independently of the signals, under which the higher signal is selected if it is at least C and a coin is tossed otherwise, if and only if $\sqrt{I_F}$ is concave on $[\underline{\theta}, \overline{\theta}]$; the pass mark then has the CDF $\mathbb{P}(C \leq z) = \frac{I_F(z)}{(\overline{\theta}-p)\,F(z)^2}$. Every concave prior satisfies this condition.

Proof. (i) In quantiles $q = F(x)$, a nondecreasing V that satisfies Lemma 18 is a nondecreasing $v(q) = V(F^{-1}(q))$ majorized by the identity, the interim rule of the noiseless comparison. By Kleiner et al. (2021, Theorems 1 and 3), the extreme points of this set equal the identity outside a collection of disjoint intervals and are constant inside each, and each is implemented by selecting the higher quantile across intervals and randomizing uniformly within an interval; this rule is nondecreasing in the own signal, and so is every mixture of such rules. If F has an atom at x, v is constant on its quantile interval $[F(x^-), F(x)]$, and drawing the quantile of signal x uniformly from that interval, independently across agents, gives signal x the average of v over the interval, which is $V(x)$; the drawn quantile is stochastically increasing in the signal, so the rule stays nondecreasing. (ii) follows from (i), as L is increasing. (iii) Under a pass mark with CDF K, $H(x, y) = \frac{1}{2} + K(\max(x, y))\left(\mathbb{1}\{x > y\} - \frac{1}{2}\right)$, which is nondecreasing in x, and $V_F(x) = \frac{1}{2} + \frac{1}{2}F(x)K(x) - \frac{1}{2}\int_x^{\overline{\theta}} K\, dF$. So $V_F' = Kf + \frac{1}{2}FK'$, i.e., $(F^2K)' = 2F\,V_F'$, and $V_F = L$ if and only if $F^2K = I_F/(\overline{\theta} - p)$, using $I_F(\overline{\theta}) = \overline{\theta} - p$ for $K(\overline{\theta}) = 1$. This K is a CDF if and only if $I_F/F^2 = I_F/(I_F')^2$ is nondecreasing, i.e., $(I_F')^2 \geq 2I_F I_F''$, i.e., $\sqrt{I_F}$ is concave. If F is concave, it lies below its tangent at z, which vanishes at or below $\underline{\theta}$, so $I_F(z) \leq \frac{F(z)^2}{2f(z)}$, which is the same inequality. □

So for concave priors the best fully revealing rule is a lottery among agents who *fail* a random pass mark. A lottery among agents who *clear* a bar instead makes V_F flat at the top, so V_F is not convex and agents gain by pooling their best types.

C.8.7 The best fully revealing rule for single-peaked priors

Proposition 17. *Let F have a density that is positive on $(\underline{\theta}, \overline{\theta})$, nondecreasing below some $m \in [\underline{\theta}, \overline{\theta}]$ and nonincreasing above it. Among general selection rules nondecreasing in the own signal, with upper semicontinuous V_F, under which full revelation is an equilibrium, a rule maximizes the firm's utility if and only if, F-almost surely,*

(i) $V_F = L$, *if* $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$;

(ii) $V_F = F$, *if* $p > \frac{\underline{\theta}+\overline{\theta}}{2}$ *and* F *is convex;*

(iii) otherwise, $V_F(\theta) = F(\theta)$ *for* $\theta \leq c^*$ *and* $V_F(\theta) = F(c^*) + \big(1 - F(c^*)\big)\frac{\theta - c^*}{\overline{\theta} - c^*}$ *for* $\theta \geq c^*$, *where* c^* *is the smallest solution in* $(\underline{\theta}, m]$ *of* $\mathbb{E}_F[\theta \mid \theta \geq c] = \frac{c+\overline{\theta}}{2}$, *and* $c^* < m$.

Let F have a density f that is positive on $(\underline{\theta}, \overline{\theta})$. For a rule nondecreasing in the own signal, V_F is nondecreasing. As $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$, full revelation is an equilibrium under a rule with upper semicontinuous V_F if and only if V_F is convex on $[\underline{\theta}, \overline{\theta}]$, by Jensen's inequality and Lemma 16. By Lemmas 18 and 20(i), a nondecreasing V equals V_F for some rule nondecreasing in the own signal if and only if $V(\theta) \leq_{cx} U[0,1]$, and by Lemma 17, $u_F = 2\int \theta\, V_F(\theta)\, dF(\theta)$. As F is atomless, the value of V_F at $\overline{\theta}$ does not matter. So Proposition 17 concerns the problem (FR) below, which I first solve for every such F and then for single-peaked F. For a bounded measurable V on $[\underline{\theta}, \overline{\theta}]$ let

$$T_V(\theta) := \int_{[\theta,\overline{\theta}]} V\, dF, \qquad M(\theta) := \frac{1 - F(\theta)^2}{2} = T_F(\theta), \qquad \theta \in [\underline{\theta}, \overline{\theta}].$$

Let C be the set of convex nondecreasing $V : [\underline{\theta}, \overline{\theta}] \to [0,1]$; such a V is continuous on $[\underline{\theta}, \overline{\theta})$, and I always take $V(\overline{\theta}) := V(\overline{\theta}-)$, which changes nothing below. For $V \in C$ let $V'(\underline{\theta}^+)$ be its right derivative at $\underline{\theta}$ and ν_V its second-derivative measure on $(\underline{\theta}, \overline{\theta})$ (the Lebesgue–Stieltjes measure of the right derivative), so that

$$V(\theta) = V(\underline{\theta}) + V'(\underline{\theta}^+)(\theta - \underline{\theta}) + \int_{(\underline{\theta},\overline{\theta})} (\theta - a)^+\, d\nu_V(a), \qquad \theta \in [\underline{\theta}, \overline{\theta}]. \tag{1}$$

The feasible set of the fully revealing problem is

$$\mathcal{V} := \Big\{ V \in C : \int V\, dF = \tfrac{1}{2},\ \ T_V \leq M \text{ on } [\underline{\theta}, \overline{\theta}] \Big\},$$

and the problem is

$$\text{(FR)} \qquad \max_{V \in \mathcal{V}} \int \theta \, V(\theta) \, dF(\theta).$$

For $V \in \mathcal{V}$ let $B(V) := \{\theta \in (\underline{\theta}, \overline{\theta}) : T_V(\theta) = M(\theta)\}$ be the set where the implementability constraint binds. By Lemma 21(iii), $T_V \leq M$ is Border's condition $V(\theta) \leq_{cx} U[0,1]$. Integrating by parts,

$$\int \theta V \, dF = \frac{\theta}{2} + \int_{\underline{\theta}}^{\overline{\theta}} T_V(\theta) \, d\theta, \tag{2}$$

so the firm wants T_V as close to its cap $M = T_F$ as possible; without convexity the cap is attained by $V = F$, and convexity is what forces T_V below M.

Lemma 21. *(i) $\mathcal{V}$ is nonempty and convex, compact for pointwise convergence, and the maximum in (FR) is attained.*

(ii) If $V \in \mathcal{V}$ and $\theta \in B(V)$, then $V(\theta) = F(\theta)$, $\int_x^\theta (V - F) \, dF \leq 0$ for $x < \theta$ and $\int_\theta^x (V - F) \, dF \geq 0$ for $x > \theta$.

(iii) For nondecreasing $V : [\underline{\theta}, \overline{\theta}] \to [0,1]$ with $\int V \, dF = \frac{1}{2}$, $V(\theta) \leq_{cx} U[0,1]$ if and only if $T_V \leq M$ on $[\underline{\theta}, \overline{\theta}]$.

Proof. (i) $V \equiv \frac{1}{2}$ is feasible, as $T_{1/2} - M = -\frac{1}{2}F(1-F) \leq 0$. By Helly's theorem every sequence in $\mathcal{C}$ has a subsequence converging pointwise on $[\underline{\theta}, \overline{\theta})$ to some $V \in \mathcal{C}$; F is atomless, so $\int V_n \, dF \to \int V \, dF$ and $T_{V_n} \to T_V$ pointwise by dominated convergence, and so does the objective. Hence $\mathcal{V}$ is compact and the objective is continuous on it. (ii) For $x < \theta$, $T_V(x) - M(x) = T_V(\theta) - M(\theta) + \int_x^\theta (V - F) \, dF = \int_x^\theta (V - F) \, dF \leq 0$; dividing by $F(\theta) - F(x) > 0$ and letting $x \uparrow \theta$ gives $V(\theta) \leq F(\theta)$ by continuity. For $x > \theta$, $T_V(x) - M(x) = -\int_\theta^x (V - F) \, dF \leq 0$ gives $V(\theta) \geq F(\theta)$. (iii) For random variables with the same mean and values in $[0,1]$, $V(\theta) \leq_{cx} U[0,1]$ if and only if $\mathbb{E}_F[(V-t)^+] \leq \mathbb{E}[(U-t)^+] = \frac{(1-t)^2}{2}$ for all $t \in [0,1]$ (for $t \notin [0,1]$ both sides agree). As V is nondecreasing, its upper level sets are intervals $[\theta, \overline{\theta}]$, so

$\mathbb{E}_F[(V-t)^+] = \sup_\theta \int_{[\theta,\overline{\theta}]}(V-t)\,dF = \sup_\theta \big(T_V(\theta) - t(1-F(\theta))\big)$. So the condition is $T_V(\theta) \le \inf_{t\in[0,1]} \left[\frac{(1-t)^2}{2} + t(1-F(\theta))\right] = M(\theta)$ for all θ, the infimum being attained at $t = F(\theta)$. □

A dual characterization. A *multiplier* is a triple (K, λ, D): a nondecreasing $K : (\underline{\theta}, \overline{\theta}) \to \mathbb{R}$ with $\int |K|\,dF < \infty$, a constant $\lambda \in \mathbb{R}$, and a constant $D \ge 0$. Let μ be the Lebesgue–Stieltjes measure of K on $(\underline{\theta}, \overline{\theta})$ (a Radon measure, possibly infinite near the endpoints), and define

$$\psi(s) := s - K(s) - \lambda, \qquad \Psi(\theta) := \int_\theta^{\overline{\theta}} \psi\,dF, \qquad \Psi_1(a) := \int_a^{\overline{\theta}} \Psi(\theta)\,d\theta, \qquad E(a) := \Psi_1(a) - D(\overline{\theta} - a).$$

Then $\psi \in L^1(F)$, Ψ is absolutely continuous with $\Psi' = -\psi f$ a.e., $E \in C^1[\underline{\theta}, \overline{\theta}]$ with $E' = D - \Psi$, $E'' = \psi f$ a.e., and $E(\overline{\theta}) = 0$, $E'(\overline{\theta}) = D$. Let $\mathcal{K} := \{a \in [\underline{\theta}, \overline{\theta}] : E(a) = 0\}$, a closed set containing $\overline{\theta}$.

Lemma 22. *Let F have a density that is positive on $(\underline{\theta}, \overline{\theta})$, and let (K, λ, D) be a multiplier with $E \le 0$ on $[\underline{\theta}, \overline{\theta}]$ and $\Psi(\underline{\theta}) \le D$. Every $W \in \mathcal{V}$ satisfies*

$$\int \theta W\,dF \le D + \frac{\lambda}{2} + \int F\,K\,dF, \tag{3}$$

with equality if and only if

(i) μ is carried by $B(W)$, i.e. $\mu\big((\underline{\theta}, \overline{\theta}) \setminus B(W)\big) = 0$;

(ii) $\nu_W(\{E < 0\}) = 0$; $E(\underline{\theta}) = 0$ if $W'(\underline{\theta}^+) > 0$; $\Psi(\underline{\theta}) = D$ if $W(\underline{\theta}) > 0$; and $W(\overline{\theta}) = 1$ if $D > 0$.

In particular, a $W \in \mathcal{V}$ that satisfies (i) and (ii) solves (FR), and then every solution of (FR) satisfies (i) and (ii).

The proof rests on two identities.

Lemma 23. *Let (K, λ, D) be a multiplier and W nondecreasing with values in $[0,1]$ and $\int W\,dF = \frac{1}{2}$. Then*

$$\int \theta W\,dF = \int W\psi\,dF + \int_{(\underline{\theta},\overline{\theta})} \big(T_W - M\big)\,d\mu + \frac{\lambda}{2} + \int F\,K\,dF, \tag{4}$$

all integrals being absolutely convergent.

Proof. As $\theta = \psi(\theta) + K(\theta) + \lambda$ and $\int W\,dF = \int F\,dF = \frac{1}{2}$, $\int \theta W\,dF = \int W\psi\,dF + \int (W - F)K\,dF + \int FK\,dF + \frac{\lambda}{2}$. Fix $s_0 \in (\underline{\theta}, \overline{\theta})$. For $s > s_0$, $K(s) = K(s_0) + \mu((s_0, s])$, and for $s \le s_0$, $K(s) = K(s_0) - \mu((s, s_0])$. As $\int (W - F)\,dF = 0$, the constant $K(s_0)$ drops out, and

$$\int (W - F)K\,dF = \int_{s > s_0} (W - F)(s)\,\mu((s_0, s])\,dF(s) - \int_{s \le s_0} (W - F)(s)\,\mu((s, s_0])\,dF(s).$$

Both double integrals converge absolutely: $|W - F| \le 1$, and $\int_{s>s_0} \mu((s_0, s])\,dF(s) = \int_{(s_0,\overline{\theta})} (1 - F)\,d\mu \le \int_{s_0}^{\overline{\theta}} |K - K(s_0)|\,dF < \infty$, and likewise below s_0. By Fubini the first term equals $\int_{(s_0,\overline{\theta})} \big[\int_{[\theta,\overline{\theta}]} (W - F)\,dF\big]\,d\mu(\theta) = \int_{(s_0,\overline{\theta})} (T_W - M)\,d\mu$, and the second equals $-\int_{(\underline{\theta},s_0]} \big[\int_{[\underline{\theta},\theta)} (W - F)\,dF\big]\,d\mu(\theta) = \int_{(\underline{\theta},s_0]} (T_W - M)\,d\mu$, as $\int_{[\underline{\theta},\theta)} (W - F)\,dF = -(T_W - M)(\theta)$. □

Lemma 24. *Let (K, λ, D) be a multiplier with $E \le 0$ on $[\underline{\theta}, \overline{\theta}]$ and $\Psi(\underline{\theta}) \le D$. For every $W \in C$,*

$$\int W\psi\,dF = W(\underline{\theta})\,\Psi(\underline{\theta}) + W'(\underline{\theta}^+)\,\Psi_1(\underline{\theta}) + \int_{(\underline{\theta},\overline{\theta})} \Psi_1(a)\,d\nu_W(a) \;\le\; D\,W(\overline{\theta}) \;\le\; D, \tag{5}$$

with equality throughout if and only if $\nu_W(\{E < 0\}) = 0$, $E(\underline{\theta}) = 0$ if $W'(\underline{\theta}^+) > 0$, $\Psi(\underline{\theta}) = D$ if $W(\underline{\theta}) > 0$, and $W(\overline{\theta}) = 1$ if $D > 0$.

Proof. Integrating by parts, for $a \in [\underline{\theta}, \overline{\theta})$, $\int (\theta - a)^+\psi\,dF = \int_a^{\overline{\theta}} (\theta - a)(-\Psi'(\theta))\,d\theta = \Psi_1(a)$, and $\int \psi\,dF = \Psi(\underline{\theta})$. Insert (1) into $\int W\psi\,dF$; Fubini applies because $\int (\theta - a)^+ d\nu_W(a) = W(\theta) - W(\underline{\theta}) - W'(\underline{\theta}^+)(\theta - \underline{\theta}) \le 1$ and $\psi \in L^1(F)$. This gives the identity

in (5). By (i), $\Psi(\underline{\theta}) \leq D$ and $\Psi_1(a) = E(a) + D(\overline{\theta} - a) \leq D(\overline{\theta} - a)$ for all $a \in [\underline{\theta}, \overline{\theta})$; as $W(\underline{\theta})$, $W'(\underline{\theta}^+)$ and ν_W are nonnegative and $D \geq 0$,

$$\int W\psi \, dF \leq D\left[W(\underline{\theta}) + W'(\underline{\theta}^+)(\overline{\theta} - \underline{\theta}) + \int (\overline{\theta} - a)\, d\nu_W(a)\right] = D\, W(\overline{\theta}) \leq D,$$

by (1) at $\theta = \overline{\theta}$. Equality in the first inequality holds if and only if each weight is positive only where its coefficient is maximal: $W(\underline{\theta}) > 0 \Rightarrow \Psi(\underline{\theta}) = D$; $W'(\underline{\theta}^+) > 0 \Rightarrow \Psi_1(\underline{\theta}) = D(\overline{\theta} - \underline{\theta})$, i.e. $E(\underline{\theta}) = 0$; and $\nu_W(\{E < 0\}) = 0$. Equality in the second holds if and only if $D = 0$ or $W(\overline{\theta}) = 1$. □

Proof of Lemma 22. Let $W \in \mathcal{V}$. By Lemmas 23 and 24, $\int \theta W \, dF = \int W\psi \, dF + \int (T_W - M)\, d\mu + \frac{\lambda}{2} + \int FK \, dF$, where $\int W\psi \, dF \leq D$ and, as $T_W \leq M$ and $\mu \geq 0$, $\int (T_W - M)\, d\mu \leq 0$. So (3) holds, with equality if and only if $\int (T_W - M)\, d\mu = 0$, which is (i), and $\int W\psi \, dF = D$, which is (ii) by Lemma 24. A $W \in \mathcal{V}$ that satisfies (i) and (ii) attains the bound, so it solves (FR), and then so does every solution. □

Call F *single-peaked with mode* $m \in [\underline{\theta}, \overline{\theta}]$ if f is positive on $(\underline{\theta}, \overline{\theta})$, nondecreasing on $(\underline{\theta}, m]$ and nonincreasing on $[m, \overline{\theta})$; so F is convex on $[\underline{\theta}, m]$ and concave on $[m, \overline{\theta}]$. Concave priors ($m = \underline{\theta}$) and convex priors ($m = \overline{\theta}$) are included; L and $\bar{U}$ are as in Subsection 5.2.

Lemma 25. *Let F be single-peaked. Then $F \leq_{cx} \bar{U}$, i.e. $L \in \mathcal{V}$, if and only if $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$.*

Proof. If $F \leq_{cx} \bar{U}$, the support of F lies in that of $\bar{U}$, so $\underline{\theta} \geq 2p - \overline{\theta}$. Conversely let $p \leq \frac{\underline{\theta}+\overline{\theta}}{2}$ and $g := L - F$ on $[\underline{\theta}, \overline{\theta}]$. Then $g(\underline{\theta}) = L(\underline{\theta}) = \frac{1}{2} - \frac{p-\underline{\theta}}{2(\overline{\theta}-p)} \geq 0$, $g(\overline{\theta}) = 0$, g is concave on $[\underline{\theta}, m]$ and convex on $[m, \overline{\theta}]$, and $\int g \, dF = 0$. On $[\underline{\theta}, m]$ the superlevel set $\{g \geq 0\}$ is an interval containing $\underline{\theta}$; on $[m, \overline{\theta}]$ the sublevel set $\{g \leq 0\}$ is an interval containing $\overline{\theta}$. So g is first ≥ 0 and then ≤ 0 on $[\underline{\theta}, \overline{\theta}]$ (if $g(m) > 0$ the first interval is all of $[\underline{\theta}, m]$; if $g(m) \leq 0$ the second is all of $[m, \overline{\theta}]$). A function that changes sign

once, from + to −, and integrates to zero has nonnegative partial integrals from the left: $\int_{\underline{\theta}}^{\theta} g\,dF \geq 0$ for all θ, i.e. $T_L \leq M$. As L is affine and nondecreasing with $\int L\,dF = \frac{1}{2}$, $L \in \mathcal{V}$, and by Lemma 21(iii) $L(\theta) \leq_{cx} U[0,1]$, i.e. $F \leq_{cx} L^{-1}(U[0,1]) = \bar{U}$. □

For $c \in [\underline{\theta}, \overline{\theta})$ write $F_c := F(\cdot \mid \theta \geq c)$ for the prior truncated from below at c and $p_c := \mathbb{E}_F[\theta \mid \theta \geq c]$.

Proof of Proposition 17. Let F be single-peaked with mode m. By the reduction above, the proposition states that (FR) has a unique solution V^*, the function given in (i), (ii) or (iii).

(i) By Lemma 25, $L \in \mathcal{V}$; by Proposition 11, whose bound is attained only by $V_F = L$, $V^* = L$.

(ii) *Convex F.* If F is convex, then $F \in \mathcal{V}$ ($T_F = M$), and by (2) every $W \in \mathcal{V}$ has $\int \theta W\,dF \leq \frac{\theta}{2} + \int_{\underline{\theta}}^{\overline{\theta}} M\,d\theta = \int \theta F\,dF$, with equality only if the continuous functions T_W and M coincide, i.e. only if $W = F$ F-almost everywhere; so $V^* = F$.

(iii) From now on $p > \frac{\underline{\theta}+\overline{\theta}}{2}$ and F is not convex; then $m < \overline{\theta}$, as otherwise f is nondecreasing on $(\underline{\theta}, \overline{\theta})$.

Existence of c^*, and $c^* < m$. Let $\phi(c) := p_c - \frac{c+\overline{\theta}}{2}$ on $[\underline{\theta}, \overline{\theta})$, continuous, with $\phi(\underline{\theta}) = p - \frac{\underline{\theta}+\overline{\theta}}{2} > 0$. For $c \geq m$ the density of F_c is nonincreasing on $[c, \overline{\theta}]$, so $p_c \leq \frac{c+\overline{\theta}}{2}$ and $\phi(m) \leq 0$. Hence the zero set of ϕ in $(\underline{\theta}, m]$ is nonempty and closed in $[\underline{\theta}, m]$ and does not contain $\underline{\theta}$, so it has a smallest element $c^* \in (\underline{\theta}, m]$. If $\phi(m) < 0$, then $c^* < m$ by continuity. Suppose $\phi(m) = 0$ and let $\omega := \frac{m+\overline{\theta}}{2}$, $h := \frac{\overline{\theta}-m}{2}$. Then $0 = (1 - F(m))\phi(m) = \int_0^h t\,[f(\omega + t) - f(\omega - t)]\,dt$, and the integrand is ≤ 0 as f is nonincreasing on $[m, \overline{\theta})$; so $f(\omega + t) = f(\omega - t)$ for a.e. $t \in (0, h)$, and by monotonicity f is a.e. equal to a constant k on $(m, \overline{\theta})$. If $f(m^-) \leq k$, then f is nondecreasing on $(\underline{\theta}, \overline{\theta})$ and F is convex, which is excluded; so $f(m^-) > k$. Let $N(c) := (1 - F(c))\phi(c) = \int_c^{\overline{\theta}} \left(\theta - \frac{c+\overline{\theta}}{2}\right) dF(\theta)$. It is absolutely continuous with $N'(c) = \frac{\overline{\theta}-c}{2} f(c) - \frac{1-F(c)}{2}$ a.e., and as $c \uparrow m$, $f(c) \to f(m^-)$ and $1 - F(c) \to k(\overline{\theta} - m)$, so $N'(c) \to \frac{\overline{\theta}-m}{2}\big(f(m^-) - k\big) > 0$. Hence $N' > 0$ a.e. on some

$(m-\delta, m)$, and $N(c) = N(m) - \int_c^m N' < 0$ there; so $\phi < 0$ on $(m-\delta, m)$, ϕ has a zero in $(\underline{\theta}, m-\delta)$, and $c^* < m$.

Feasibility. Fix a zero $c \in (\underline{\theta}, m)$ of ϕ (for instance c^*), let $q := F(c)$ and let V be the function in (iii) with c in place of c^*. On $[c, \overline{\theta}]$, $V = q + (1-q)U_c$ with $U_c(\theta) := \frac{\theta - c}{\overline{\theta} - c}$ the CDF of $U[c, \overline{\theta}]$, and $F = q + (1-q)F_c$, so for $\theta \geq c$

$$T_V(\theta) - M(\theta) = \int_\theta^{\overline{\theta}} (V - F)\, dF = (1-q)^2 \int_\theta^{\overline{\theta}} (U_c - F_c)\, dF_c.$$

Now F_c is single-peaked on $[c, \overline{\theta}]$ (its density is $f/(1-q)$ and $c \leq m$) with mean $p_c = \frac{c + \overline{\theta}}{2}$, the midpoint of its support, and U_c is the function L of Lemma 25 for the prior F_c (indeed $\frac{1}{2} + \frac{\theta - p_c}{2(\overline{\theta} - p_c)} = \frac{\theta - c}{\overline{\theta} - c}$). By Lemma 25 and Lemma 21(iii) applied to F_c, $\int_\theta^{\overline{\theta}} (U_c - F_c)\, dF_c \leq 0$ for all $\theta \geq c$, with equality at $\theta = c$. So $T_V \leq M$ on $[c, \overline{\theta}]$ and $T_V(c) = M(c)$; for $\theta \leq c$, $T_V(\theta) - M(\theta) = T_V(c) - M(c) + \int_\theta^c (F - F)\, dF = 0$. In particular $\int V\, dF = T_V(\underline{\theta}) = \frac{1}{2}$, and $B(V) \supseteq [\underline{\theta}, c]$. V is nondecreasing with values in $[0,1]$, convex on $[\underline{\theta}, c]$ (as F is) and affine on $[c, \overline{\theta}]$; at c the slope rises: the chord slope $s_c := \frac{1-q}{\overline{\theta} - c}$ satisfies $s_c \geq f(c^+)$, since otherwise, F being convex on $[c, m]$ with right derivative $f(c^+)$ at c, $V(\theta) - F(\theta) \leq (s_c - f(c^+))(\theta - c) < 0$ for $\theta \in (c, m]$, and then $T_V(\theta) - M(\theta) = -\int_c^\theta (V - F)\, dF > 0$ for $\theta \in (c, m]$; and $f(c^+) \geq f(c^-)$ because $c < m$ and f is nondecreasing on $(\underline{\theta}, m]$. So $s_c \geq f(c^-)$, V is convex, and $V \in \mathcal{V}$.

A multiplier. Let

$$\kappa := \frac{\int_c^{\overline{\theta}} (\overline{\theta} - s)\, s\, dF(s)}{\int_c^{\overline{\theta}} (\overline{\theta} - s)\, dF(s)} \in [c, \overline{\theta}],$$

and let $K(s) := s - \underline{\theta}$ for $s \in (\underline{\theta}, c]$ and $K(s) := \kappa - \underline{\theta}$ for $s \in (c, \overline{\theta})$; $\lambda := \underline{\theta}$. So μ is Lebesgue measure on $(\underline{\theta}, c)$ plus an atom of mass $\kappa - c$ at c, carried by $[\underline{\theta}, c] \subseteq B(V)$: this is (i) of Lemma 22. Then $\psi = 0$ on $(\underline{\theta}, c)$ and $\psi(s) = s - \kappa$ on $(c, \overline{\theta})$. Set $D := \Psi(c) = \int_c^{\overline{\theta}} (s - \kappa)\, dF(s)$. Since $\int_c^{\overline{\theta}} (\overline{\theta} - s)(s - \kappa)\, dF = 0$ by the choice of κ, the function

$(s-\kappa)(\overline{\theta}-s)f(s)$ is negative on (c,κ) and positive on $(\kappa,\overline{\theta})$ and integrates to zero, while $1/(\overline{\theta}-s)$ is increasing; hence $D=\int_c^{\overline{\theta}}(s-\kappa)(\overline{\theta}-s)\,\frac{1}{\overline{\theta}-s}\,dF \geq \frac{1}{\overline{\theta}-\kappa}\int_c^{\overline{\theta}}(s-\kappa)(\overline{\theta}-s)\,dF=0$, with strict inequality because $f>0$. So $D>0$.

The hypothesis and condition (ii) of Lemma 22. On $[\underline{\theta},c]$, $\Psi\equiv\Psi(c)=D$, so $\Psi_1(a)=D(c-a)+\Psi_1(c)$ and $\Psi_1(c)=\int_c^{\overline{\theta}}(s-c)(s-\kappa)\,dF=\int_c^{\overline{\theta}}(\overline{\theta}-c)(s-\kappa)\,dF-\int_c^{\overline{\theta}}(\overline{\theta}-s)(s-\kappa)\,dF=D(\overline{\theta}-c)$; hence $E\equiv 0$ on $[\underline{\theta},c]$ and $\Psi(\underline{\theta})=D$. On $[c,\overline{\theta}]$, $E(c)=0$, $E'(c)=D-\Psi(c)=0$, $E(\overline{\theta})=0$, and $E''=(s-\kappa)f$ is ≤ 0 on (c,κ) and ≥ 0 on $(\kappa,\overline{\theta})$: E is concave on $[c,\kappa]$ with $E(c)=E'(c)=0$, so $E\leq 0$ there, and convex on $[\kappa,\overline{\theta}]$ with $E(\kappa)\leq 0=E(\overline{\theta})$, so $E\leq 0$ there. This is the hypothesis. For (ii): ν_V is carried by $[\underline{\theta},c]\subseteq\mathcal{K}$ (it is ν_F on $(\underline{\theta},c)$ plus the atom $s_c-f(c^-)$ at c); $E(\underline{\theta})=0$; $V(\underline{\theta})=0$; and $V(\overline{\theta})=1$. By Lemma 22, V solves (FR), and every solution W satisfies (i) and (ii) of Lemma 22 for this multiplier. By (i), $T_W=M$ almost everywhere on $(\underline{\theta},c)$, hence on $[\underline{\theta},c]$ by continuity, so $W=F$ F-almost everywhere on $[\underline{\theta},c]$, hence everywhere there, as $f>0$ and W and F are continuous on $[\underline{\theta},c]$. As $f>0$ and $\kappa\in(c,\overline{\theta})$, $E'(s)=\int_c^s(t-\kappa)f(t)\,dt<0$ on $(c,\kappa]$, so $E<0$ there, and $E<0$ on $(\kappa,\overline{\theta})$ by convexity; so by (ii), ν_W is carried by $(\underline{\theta},c]$, W is affine on $[c,\overline{\theta}]$, and $W(c)=F(c)$ and $W(\overline{\theta})=1$ (as $D>0$) give $W=V$. So V is the unique solution, the same for every zero $c\in(\underline{\theta},m)$ of ϕ, in particular for c^*.

□

Proposition 18. *Let F be single-peaked with mode m, $p>\frac{\underline{\theta}+\overline{\theta}}{2}$ and F not convex. Let c^* be the smallest solution in $(\underline{\theta},m]$ of $\mathbb{E}_F[\theta\mid\theta\geq c]=\frac{c+\overline{\theta}}{2}$, $q:=F(c^*)$, and V^* the solution of (FR).*

(i) Let $\kappa(\cdot\mid x)$, $x\in[c^*,\overline{\theta}]$, be a martingale kernel that spreads F_{c^*} into $U[c^*,\overline{\theta}]$: $\int S\,\kappa(dS\mid x)=x$ and $\int\kappa(\cdot\mid x)\,dF_{c^*}(x)=U[c^*,\overline{\theta}]$ (it exists by Strassen's theorem, as $F_{c^*}\leq_{cx}U[c^*,\overline{\theta}]$, and is supported in $[c^*,\overline{\theta}]$). The rule that selects the higher signal if $\min(x,y)<c^*$ and otherwise selects the agent with the higher independent score $S_x\sim\kappa(\cdot\mid x)$, $S_y\sim\kappa(\cdot\mid y)$ (coin at ties), has $V_F=V^*$, and full revelation is an equilibrium under it. A rule

nondecreasing in the own signal with the same V_F exists: a lottery over coarse gradings whose cells are the singletons below c^* and a random partition of $[c^*, \overline{\theta}]$ into intervals.

(ii) No bar rule implements V^*. (For convex F with $p > \frac{\underline{\theta}+\overline{\theta}}{2}$, $V^* = F$ is implemented by the noiseless rule, a degenerate pass mark.) (a) Lottery above a bar. *Under any rule that selects the higher signal unless both signals are at least B, and then tosses a coin, with B drawn independently and $\mathbb{P}(B < \overline{\theta}) > 0$, $V_F(x) \leq 1 - \frac{1}{2}\mathbb{P}(\theta \geq B,\ x \geq B)$, so $V_F(\overline{\theta}-) < 1 = V^*(\overline{\theta})$; this holds for every prior. (b)* Pass mark above c^*. *Under any rule that selects the higher signal unless both signals lie in $[c^*, B)$, and then tosses a coin, with B independent and $\mathbb{P}(B > c^*) > 0$, $\liminf_{x \downarrow c^*} V_F(x) - \lim_{x \uparrow c^*} V_F(x) \geq \frac{1}{2}\mathbb{P}\big(\theta \in [c^*, B)\big) > 0$, so V_F is not convex and full revelation is not an equilibrium. (c) Under a random pass mark (the higher signal is selected if it is at least an independent bar B, and a coin is tossed otherwise), $V_F = F$ on $(\underline{\theta}, c^*)$ only if $B \leq \underline{\theta}$ a.s.; then the rule is noiseless and $V_F = F \neq V^*$.*

Proof. (i) Against a fully revealing rival, a signal $x < c^*$ wins if and only if $y < x$, so $V_F(x) = F(x)$; a signal $x \geq c^*$ wins against every $y < c^*$ and, conditionally on $y \geq c^*$, with probability $\mathbb{P}(S_y < S_x) = \mathbb{E}[U_{c^*}(S_x)] = U_{c^*}(x)$, as $S_y \sim U[c^*, \overline{\theta}]$ unconditionally and U_{c^*} is affine on $[c^*, \overline{\theta}]$, which contains the support of S_x. So $V_F(x) = q + (1-q)\frac{x-c^*}{\overline{\theta}-c^*} = V^*(x)$. As V^* is convex on $[\underline{\theta}, \overline{\theta}]$, full revelation is a best reply to itself by Jensen's inequality. For the monotone implementation, apply Lemma 20(i) to the prior F_{c^*} on $[c^*, \overline{\theta}]$ and the nondecreasing feasible value function U_{c^*}: it gives a rule $\tilde{H}$ on $[c^*, \overline{\theta}]^2$, nondecreasing in the own signal, with $\int \tilde{H}(x, y)\, dF_{c^*}(y) = U_{c^*}(x)$. Let $H(x, y) := \mathbb{1}\{x > y\} + \frac{1}{2}\mathbb{1}\{x = y\}$ if $\min(x, y) < c^*$ and $H := \tilde{H}$ if both signals are at least c^*. For fixed $y < c^*$, $x \mapsto H(x, y)$ is nondecreasing; for fixed $y \geq c^*$ it is 0 below c^* and nondecreasing and nonnegative above, so H is nondecreasing in the own signal, and against a fully revealing rival $V_F(x) = F(x)$ for $x < c^*$ and $V_F(x) = q + (1-q)U_{c^*}(x)$ for $x \geq c^*$, i.e. $V_F = V^*$. (ii)(a) A signal x loses the coin with probability $\frac{1}{2}$ whenever

$y \geq B$ and $x \geq B$, so $V_F(x) \leq 1 - \frac{1}{2}\mathbb{P}(\theta \geq B, x \geq B) \to 1 - \frac{1}{2}\mathbb{P}(\theta \geq B) < 1$ as $x \uparrow \overline{\theta}$, since $\mathbb{P}(\theta \geq B) = \mathbb{E}[1 - F(B^-)] > 0$ when $\mathbb{P}(B < \overline{\theta}) > 0$. (b) For $x < c^*$, $V_F(x) = F(x) \to F(c^*)$; for $x > c^*$, x beats every $y < c^*$ and, when $B > x$, gets the coin against every $y \in [c^*, B)$, so $V_F(x) \geq F(c^*) + \frac{1}{2}\mathbb{P}(\theta \in [c^*, B),\ B > x)$, which tends to $F(c^*) + \frac{1}{2}\mathbb{P}(\theta \in [c^*, B))$ as $x \downarrow c^*$ by monotone convergence; this exceeds $F(c^*)$ as $f > 0$ and $\mathbb{P}(B > c^*) > 0$. A convex function is continuous on $(\underline{\theta}, \overline{\theta})$, so V_F is not convex; it is upper semicontinuous and $\text{supp}(F) = [\underline{\theta}, \overline{\theta}]$, so by Lemma 16 full revelation is not an equilibrium. (c) Let K be the CDF of B. As in the proof of Lemma 20(iii), $V_F(x) = \frac{1}{2} + \frac{1}{2}K(x)F(x) - \frac{1}{2}\int_x^{\overline{\theta}} K\, dF$, so $d(F^2K) = 2F\, dV_F$; if $V_F = F$ on $(\underline{\theta}, c^*)$, then $d(F^2K) = d(F^2)$ there, so $F^2K = F^2 + C$ on $(\underline{\theta}, c^*)$ for a constant C. As $x \downarrow \underline{\theta}$, $F(x) \to 0$ and $K \in [0, 1]$, so $C = 0$; hence $K \equiv 1$ on $(\underline{\theta}, c^*)$ and $B \leq \underline{\theta}$ a.s. Then the higher signal is never below B (up to a null event), the rule is noiseless, and $V_F = F$ on $[\underline{\theta}, \overline{\theta}]$, which differs from V^* because V^* is convex and F is not. □

C.8.8 Robust full revelation

Proposition 19. *If a general selection rule H has $H(\cdot, y)$ convex for every y, full revelation is an equilibrium for every prior on $[0, 1]$; conversely, if H is continuous and full revelation is an equilibrium for every prior, each $H(\cdot, y)$ is convex. Among such rules,*

$$H^*(x, y) = \frac{1}{2} + \frac{x - y}{2\big(1 - \min(x, y)\big)}$$

selects the higher signal with the highest probability at every pair of signals, so it maximizes the firm's utility under full revelation for every prior simultaneously. For every prior on $[0, 1]$, full revelation is the unique equilibrium under H^*.

Proof. Sufficiency: convexity of each $H(\cdot, y)$ makes V_F convex for every F, so full revelation is a best reply to itself by Jensen's inequality. Necessity: for atomless

full-support priors $F_n \to \delta_y$, full revelation is a best reply to itself, so pooling the types in any interval does not pay, and V_{F_n} is convex by Lemma 15; as H is continuous, V_{F_n} converges pointwise to $H(\cdot, y)$, which is therefore convex. Optimality: for $x > y$, $H(\cdot, y)$ is convex on $[y, 1]$ with $H(y, y) = \frac{1}{2}$ and $H(1, y) \leq 1$, so it lies below the chord, $H(x, y) \leq \frac{1}{2} + \frac{x-y}{2(1-y)} = H^*(x, y)$. H^* is admissible: for $x < y$, $H^*(x, y) = \frac{1-y}{2(1-x)}$, which is convex in x and has the same slope $\frac{1}{2(1-y)}$ at $x = y$ as the linear branch, and $H^*(x, y) + H^*(y, x) = 1$. Since $u_F = \mathbb{E}[m + (M - m)H(M, m)]$ with $M = \max(\theta_1, \theta_2)$ and $m = \min(\theta_1, \theta_2)$, pointwise dominance gives the ranking. Uniqueness: H^* is continuous except at $(1, 1)$, so V_F is continuous on $[\underline{\theta}, 1)$, and $V_F''(x) = \int_{(x,1]} \frac{1-y}{(1-x)^3} dF(y) > 0$ on $[\underline{\theta}, t)$, where $t := \sup(\mathrm{supp}(F) \cap [0, 1))$; on $[t, 1)$, which contains no type, V_F is affine, and if F has an atom at 1, V_F jumps up there, as $H^*(x, 1) = 0$ for $x < 1$ and $H^*(1, 1) = \frac{1}{2}$. A mean-preserving contraction $G \neq F$ pools some types $a < b$; if $a < t$, strict convexity on $[a, \min(b, t))$ makes the pooling strictly costly, and otherwise $a = t$ and $b = 1$, and the jump at 1 does. So $\int V_F \, dG < \int V_F \, dF$, F is the unique best reply to F, and by interchangeability every equilibrium (G_1, G_2) has (G_1, F) as an equilibrium, so $G_1 = F$, and likewise $G_2 = F$. □

C.8.9 Proof of Proposition 12

The rule. The greatest common mean-preserving contraction $G^*_{\bar{s}}$ of F and $\bar{U}_{\bar{s}}$ is defined by $I_{G^*_{\bar{s}}}$ being the convex minorant of $\min(I_F, I_{\bar{U}_{\bar{s}}})$. For $\bar{s} \geq 2p - \underline{\theta}$, the uniformizing rule of width $\bar{s}$ scores an agent with signal x by an independent draw S_x with mean x: a draw from a martingale kernel that spreads $G^*_{\bar{s}}$ into $\bar{U}_{\bar{s}}$ if $x \in \mathrm{supp}(G^*_{\bar{s}})$, and $S_x = x$ otherwise; the higher score is selected. For concave F, Lemma 27 gives a monotone implementation. For every prior, Lemma 20(i), applied with $G^*_{\bar{s}}$ in place of F to the nondecreasing $L_{\bar{s}}$, which satisfies the condition of Lemma 18 as $G^*_{\bar{s}} \leq_{cx} \bar{U}_{\bar{s}}$, gives a lottery over coarse gradings, nondecreasing in the own signal, under which the value

function against $G^*_{\bar{s}}$ equals $L_{\bar{s}}$ on $\text{supp}(G^*_{\bar{s}})$. Off the support, a signal is graded with the highest support point below it, so its value is at most the certificate $L_{\bar{s}}$, because the certificate is nondecreasing, and the bound by $\int L_{\bar{s}}\, dG' = \frac{1}{2}$ below applies to this rule too.

Equilibrium and payoff. Against $G^*_{\bar{s}}$, the rival's score is distributed as $\bar{U}_{\bar{s}}$, whose CDF equals $L_{\bar{s}}$ on $[2p - \bar{s}, \bar{s}]$. For $x \le \bar{s}$, S_x lies in this interval, as $\underline{\theta} \ge 2p - \bar{s}$, so the agent wins with probability $\mathbb{E}[L_{\bar{s}}(S_x)] = L_{\bar{s}}(x)$; for $x > \bar{s}$, he wins with probability $1 < L_{\bar{s}}(x)$. Hence every feasible G' yields at most $\int L_{\bar{s}}\, dG' = \frac{1}{2}$, which $G^*_{\bar{s}}$ attains, and $u_F(\bar{s})$ follows from Lemma 17.

Shape. Let $\bar{s} \in (2p - \underline{\theta}, \overline{\theta})$ and let $\{x \in [\underline{\theta}, \bar{s}] : F(x) > L_{\bar{s}}(x)\}$ be an interval. For concave F it is, as $F - L_{\bar{s}}$ is then concave on $[\underline{\theta}, \bar{s}]$. Let $D := I_F - I_{\bar{U}_{\bar{s}}}$. Then $D' = F - \bar{U}_{\bar{s}}$ is non-positive below $\underline{\theta}$ and above $\bar{s}$, and on $[\underline{\theta}, \bar{s}]$, where $\bar{U}_{\bar{s}} = L_{\bar{s}}$ as $2p - \bar{s} < \underline{\theta}$, it is positive exactly on an interval. As D vanishes at $\pm\infty$, $D(\underline{\theta}) = -I_{\bar{U}_{\bar{s}}}(\underline{\theta}) < 0$ and $D > 0$ on $(\bar{s}, \overline{\theta})$, D changes sign once, from negative to positive, at some x^*. So $\min(I_F, I_{\bar{U}_{\bar{s}}})$ is convex on each side of x^*, and every interval on which its convex minorant lies strictly below it contains x^*, as on such an interval the minorant is affine and touches the minimum at both ends. Hence the minorant bridges the crossing by a single common supporting line, touching I_F at t_1 (which may be the kink of I_F at $\underline{\theta}$ if F has an atom there) and $I_{\bar{U}_{\bar{s}}}$ at t_2. For concave F, at the ends, $\bar{U}_{2p-\underline{\theta}} \le_{cx} F$ (proof of Corollary 3) and $F \le_{cx} \bar{U}_{\overline{\theta}}$ (Appendix C.8.6), so $G^*_{2p-\underline{\theta}} = \bar{U}_{2p-\underline{\theta}}$ and $G^*_{\overline{\theta}} = F$.

C.8.10 Proof of Theorem 3

The bound holds equilibrium by equilibrium:

Lemma 26. *Let $U[\underline{\theta}, 2p - \underline{\theta}] \le_{cx} F \le_{cx} \bar{U}$, let the set $\{x \in [\underline{\theta}, \bar{s}'] : F(x) > L_{\bar{s}'}(x)\}$ be an interval for every $\bar{s}' \in (2p - \underline{\theta}, \overline{\theta})$, and let (G, G) be a symmetric equilibrium of a general selection rule in which V_G is nondecreasing on* $\text{supp}(G)$ *and* $\text{Cov}_G(s, V_G(s)) > 0$. *Let*

$\beta := \frac{\mathrm{Cov}_G(s,V_G(s))}{\mathrm{Var}_G(V_G(s))}$ *and* $\bar{s} := p + \frac{\beta}{2}$. *Then* $u_F \le p + \frac{\mathrm{Var}(G^*_{\bar{s}})}{\bar{s}-p}$, *which equals* $u_F(\bar{s})$ *if* $\bar{s} \in [2p - \underline{\theta}, \overline{\theta}]$, *and is at most* $u_F(2p - \underline{\theta})$ *if* $\bar{s} < 2p - \underline{\theta}$ *and at most* $u_F(\overline{\theta})$ *if* $\bar{s} > \overline{\theta}$.

Steps 1, 2 and 4–6 below prove Lemma 26; Step 3 reduces the bound of Theorem 3 for every symmetric equilibrium to it, and Step 7 proves that the maximum is attained only at interior widths under the strict inequalities stated after the theorem.

Let (G, G) be a symmetric equilibrium, $s \sim G$ and $V := V_G$. By Lemma 17, $u_F = p + 2\,\mathrm{Cov}(s, V(s))$. The maximum in the statement exists, as $\mathrm{Var}(G^*_{\bar{s}})$ is continuous in $\bar{s}$.

Step 1 (convexity on the support). By Lebesgue's differentiation theorem for G, the set $A \subseteq \mathrm{supp}(G)$ of points at which V is approximately continuous with respect to G has $G(A) = 1$. For $s_1 < s_2 < s_3$ in A, run the chord argument of the proof of Lemma 16 with G in place of F, with the two neighbourhoods replaced by sets of positive G-mass within δ of s_1 and s_3 on which V is within ε of $V(s_1)$ and $V(s_3)$, and with the target replaced by a point x within δ of s_2 with $|V(x) - V(s_2)| \le \varepsilon$; as every contraction of G is feasible, letting $\varepsilon, \delta \to 0$ gives $V(s_2) \le \frac{s_3 - s_2}{s_3 - s_1} V(s_1) + \frac{s_2 - s_1}{s_3 - s_1} V(s_3)$. So V is the restriction to A of a convex function φ on the convex hull of A, and $V(s) = \varphi(s)$ almost surely.

Step 2 (implementability). As in the proof of Lemma 18, with G in place of F, $V(s) \le_{cx} U[0,1]$.

Step 3 (monotone pooling). Let Q be the quantile function of G and $r(t) := \varphi(Q(t))$, which is quasi-convex on $[0,1]$, as φ is convex and Q nondecreasing. Let q^* be the largest minimizer of the running average $c(q) := \frac{1}{q}\int_0^q r$ on $[0,1]$, with $c(0) := r(0^+)$, and $c^* := c(q^*)$. Minimality of c^* and quasi-convexity of r, whose sublevel sets are intervals, give three facts: $r \ge c^*$ almost everywhere on $(q^*, 1]$, as otherwise c would fall below c^* to the right of q^*; r is nondecreasing there; and on $[0, q^*]$, $r - c^*$ is positive below some j and nonpositive above it, as otherwise c would fall below c^* just left of q^*. Now pool the lowest q^* quantiles into one signal, their mean μ, with winning probability c^*.

The new signal distribution is a contraction of G, so it is still $\leq_{cx} F$; the new winning probability is a conditional expectation of $V(s)$, so it is still $\leq_{cx} U[0,1]$ by Step 2; and $\mathbb{E}[(s-p)V(s)]$ rises by $-\int_0^{q^*}(Q(t)-\mu)(r(t)-c^*)\,dt = -\int_0^{q^*}(Q(t)-Q(j))(r(t)-c^*)\,dt \geq 0$, so u_F weakly rises. The new value function is the restriction of $\max(\varphi, c^*)$ to $[\mu,\infty)$, which is convex and nondecreasing there, as $\varphi(\mu) \leq c^*$ by Jensen's inequality. As Steps 4–6 use only these properties, assume from now on that φ is nondecreasing and that $\mathrm{Cov}(s, V(s)) > 0$, as otherwise $u_F \leq p$, below every $u_F(\bar{s})$. This also proves the claim in Subsection 4.4 that monotonicity in the own signal is without loss for the firm.

Step 4 (the width). Let $\beta := \frac{\mathrm{Cov}(s,V(s))}{\mathrm{Var}(V(s))}$, $\bar{s} := p + \frac{\beta}{2}$, $\psi := p + \beta(\varphi - \frac{1}{2})$, which is convex and nondecreasing, and $Y := \psi(s)$. By Step 2, $Y \leq_{cx} U[p - \frac{\beta}{2}, p + \frac{\beta}{2}] = \bar{U}_{\bar{s}}$, so $\mathbb{E}Y = p$; by the choice of β, $\mathrm{Cov}(Y, s-Y) = \beta\,\mathrm{Cov}(V,s) - \beta^2\mathrm{Var}(V) = 0$; and $\mathbb{E}[(Y-p)(s-p)] = 2(\bar{s}-p)\,\mathrm{Cov}(s,V(s))$. So it suffices to show that $\mathbb{E}[(Y-p)(s-p)] \leq \mathrm{Var}(G^*_{\bar{s}})$.

Step 5 (a dual bound). Suppose that w_1, w_2 are convex, $\int w_1\,dF + \int w_2\,d\bar{U}_{\bar{s}} = \mathrm{Var}(G^*_{\bar{s}})$, a is concave, and

$$R(s,y) := w_1(s) + w_2(y) - (y-p)(s-p) - a(y)(s-y) \geq 0 \qquad \text{for all } s, y.$$

As $G \leq_{cx} F$ and $Y \leq_{cx} \bar{U}_{\bar{s}}$,

$$\mathbb{E}[(Y-p)(s-p)] = \mathbb{E}w_1(s) + \mathbb{E}w_2(Y) - \mathbb{E}[a(Y)(s-Y)] - \mathbb{E}R(s,Y) \leq \mathrm{Var}(G^*_{\bar{s}}) - \mathbb{E}[a(Y)(s-Y)].$$

Such functions exist. For $\bar{s} \in (2p - \underline{\theta}, \overline{\theta})$, let t_1, t_2 be as in Proposition 12, which applies

by the hypothesis on $\{F > L_{\bar{s}}\}$, let C be the chord of $(x-p)^2$ over $[t_1, t_2]$, and let

$$w_1(x) := \begin{cases} (x-p)^2, & x \le t_1, \\ C(x), & x \ge t_1, \end{cases} \qquad w_2(x) := \begin{cases} 0, & x \le t_2, \\ (x-p)^2 - C(x), & x \ge t_2, \end{cases}$$

$$a(y) := \min\{y - p,\ t_1 - p,\ t_1 + t_2 - p - y\}.$$

Here w_1 is affine where $I_{G^*} < I_F$ (above t_1), so $\int w_1\, dF = \int w_1\, dG^*$; w_2 is affine where $I_{G^*} < I_{\bar{U}_{\bar{s}}}$ (below t_2), so $\int w_2\, d\bar{U}_{\bar{s}} = \int w_2\, dG^*$; and $w_1 + w_2 = (x-p)^2$ on $\operatorname{supp}(G^*) \subseteq [\underline{\theta}, t_1] \cup [t_2, \bar{s}]$, so the two integrals sum to $\operatorname{Var}(G^*_{\bar{s}})$. For $y \le t_1$, $R(s,y)$ is the gap between w_1 and its tangent at y; for $y \ge t_2$, $R(s,y) = w_1(s) - C(s) \ge 0$; and for $t_1 < y < t_2$, $R(s,y) = (s - t_1)(t_2 - y)$ if $s \ge t_1$ and $R(s,y) = (s - t_1)(s - y)$ if $s \le t_1$, both nonnegative. For $\bar{s} \le 2p - \underline{\theta}$, take $w_1 = 0$, $w_2 = (x-p)^2$ and $a(y) = p - y$, so that $R = 0$ and the value is $\operatorname{Var}(\bar{U}_{\bar{s}}) = \operatorname{Var}(G^*_{\bar{s}})$, as $\bar{U}_{\bar{s}} \le_{cx} \bar{U}_{2p-\underline{\theta}} \le_{cx} F$; for $\bar{s} \ge \overline{\theta}$, take $w_1 = (x-p)^2$, $w_2 = 0$ and $a(y) = y - p$, so that $R = (s-y)^2$ and the value is $\operatorname{Var}(F) = \operatorname{Var}(G^*_{\bar{s}})$, as $F \le_{cx} \bar{U}_{\overline{\theta}} \le_{cx} \bar{U}_{\bar{s}}$. In these two cases, $p + \operatorname{Var}(G^*_{\bar{s}})/(\bar{s} - p)$ equals $p + \frac{\bar{s}-p}{3}$, which is increasing, and $p + \frac{\operatorname{Var}(F)}{\bar{s}-p}$, which is decreasing, so a width outside $[2p - \underline{\theta}, \overline{\theta}]$ is weakly beaten by the nearest end of the family.

Step 6 (the sign). Let $\nu(B) := \mathbb{E}[(s - Y)\mathbb{1}\{Y \in B\}]$, so that $\nu(\mathbb{R}) = 0$ and $\int y\, d\nu = \operatorname{Cov}(Y, s - Y) = 0$. As $x - \psi(x)$ is concave, the signals with $s > \psi(s)$ lie in an interval J that contains every other signal between two of them. If J contains no signal, $\nu = 0$. Otherwise, let (y_a, y_b) be the interior of the convex hull of the values $\psi(s)$ of signals $s \in J$. If $\psi(s) \in (y_a, y_b)$, then $\psi(s') < \psi(s) < \psi(s'')$ for some signals $s', s'' \in J$, so $s' < s < s''$ as ψ is nondecreasing, and $s \in J$. Hence $\nu \ge 0$ on (y_a, y_b) and $\nu \le 0$ outside $[y_a, y_b]$. With ℓ the chord of a over $[y_a, y_b]$ (a supporting line if $y_a = y_b$), $a - \ell$ is nonnegative on $[y_a, y_b]$ and nonpositive outside it, so $\mathbb{E}[a(Y)(s - Y)] = \int (a - \ell)\, d\nu \ge 0$. With Step 5, $\mathbb{E}[(Y - p)(s - p)] \le \operatorname{Var}(G^*_{\bar{s}})$, which proves the bound.

Step 7 (interior widths). Let $I_{\bar{U}_{2p-\underline{\theta}}} < I_F < I_{\bar{U}_{\overline{\theta}}}$ on $(\underline{\theta}, \overline{\theta})$. Every concave F that is not uniform satisfies this: then $p < \frac{\underline{\theta}+\overline{\theta}}{2}$, $F \leq_{cx} \bar{U}_{\overline{\theta}}$ and $\bar{U}_{2p-\underline{\theta}} \leq_{cx} F$, and both inequalities are strict on $(\underline{\theta}, \overline{\theta})$, since in both cases the CDFs cross once. The strict inequalities give $2p - \underline{\theta} < \overline{\theta}$, as otherwise $\bar{U}_{2p-\underline{\theta}} = \bar{U}_{\overline{\theta}}$, so $p < \frac{\underline{\theta}+\overline{\theta}}{2}$ and $I_{\bar{U}_{\overline{\theta}}}(\underline{\theta}) > 0 = I_F(\underline{\theta})$. Two facts are used. For distributions with mean p, $\mathrm{Var}(K_1) - \mathrm{Var}(K_2) = 2\int (I_{K_1} - I_{K_2})\, dx$. For a uniform distribution of half-width r around p, $\frac{\partial I}{\partial r} = \frac{(x-p+r)(r-x+p)}{4r^2}$ on $[p-r, p+r]$ and 0 elsewhere.

Near the full revelation end, let $\eta := \overline{\theta} - \bar{s}$ and $h := \max_x (I_F - I_{\bar{U}_{\bar{s}}})^+$. Then $\max(I_F - h,\ x - p,\ 0)$ is a convex minorant of $\min(I_F, I_{\bar{U}_{\bar{s}}})$, so $\mathrm{Var}(F) - \mathrm{Var}(G^*_{\bar{s}}) \leq 2(\overline{\theta} - \underline{\theta})h$. Write $I_F - I_{\bar{U}_{\bar{s}}} = -(I_{\bar{U}_{\overline{\theta}}} - I_F) + (I_{\bar{U}_{\overline{\theta}}} - I_{\bar{U}_{\bar{s}}})$. The derivative formula, with $r - x + p \leq \overline{\theta} - x$ and $x - p + r \leq 2r$, bounds the second term by $\frac{\eta(\overline{\theta}-x)^+}{2(\bar{s}-p)}$. Fix $\delta > 0$: on $[\underline{\theta}, \overline{\theta} - \delta]$ the first term is bounded away from zero, so there is no violation for small η; on $[\overline{\theta} - \delta, \overline{\theta}]$ the violation is at most $\frac{\eta\delta}{2(\bar{s}-p)}$. Hence $h = o(\eta)$, and

$$u_F(\bar{s}) - u_F(\overline{\theta}) \geq \frac{\mathrm{Var}(F)\,\eta}{(\bar{s}-p)(\overline{\theta}-p)} - \frac{2(\overline{\theta} - \underline{\theta})\, o(\eta)}{\bar{s}-p} > 0 \quad \text{for small } \eta.$$

Near the noiseless end, let $\bar{s}_0 := 2p - \underline{\theta}$, $\eta := \bar{s} - \bar{s}_0$, and $h' := \max_x (I_{\bar{U}_{\bar{s}}} - I_F)^+$. Symmetrically, $\max(I_{\bar{U}_{\bar{s}}} - h',\ x - p,\ 0)$ is a convex minorant of the minimum, so $\mathrm{Var}(\bar{U}_{\bar{s}}) - \mathrm{Var}(G^*_{\bar{s}}) \leq 4(\bar{s} - p)h'$. Write $I_{\bar{U}_{\bar{s}}} - I_F = (I_{\bar{U}_{\bar{s}}} - I_{\bar{U}_{\bar{s}_0}}) - (I_F - I_{\bar{U}_{\bar{s}_0}})$. The derivative formula, with $r - x + p \leq 2r$ and $x - p + r \leq x - \underline{\theta} + \eta$, bounds the first term by $\frac{\eta(x-\underline{\theta}+\eta)^+}{2(p-\underline{\theta})}$; below $\underline{\theta}$, the violation is at most $I_{\bar{U}_{\bar{s}}}(\underline{\theta}) = O(\eta^2)$; and near $\overline{\theta}$ the first term vanishes for $x \geq \bar{s}$, while on $[\underline{\theta} + \delta, \bar{s}]$, a compact subset of $(\underline{\theta}, \overline{\theta})$ as $\bar{s}_0 < \overline{\theta}$, the second is bounded away from zero and the first is at most $\frac{\eta(\overline{\theta}-\underline{\theta}+\eta)}{2(p-\underline{\theta})} \to 0$. So $h' = o(\eta)$. As $\mathrm{Var}(\bar{U}_{\bar{s}}) = \frac{(\bar{s}-p)^2}{3}$ and $G^*_{\bar{s}_0} = \bar{U}_{\bar{s}_0}$,

$$u_F(\bar{s}) - u_F(\bar{s}_0) \geq \frac{\bar{s}-p}{3} - \frac{\bar{s}_0 - p}{3} - 4h' = \frac{\eta}{3} - o(\eta) > 0 \quad \text{for small } \eta.$$

So neither end maximizes $u_F(\bar{s})$, and the maximum is attained only at interior widths.

C.8.11 The random pass mark

For concave priors, a random pass mark implements the whole uniformizing family, and its equilibrium is unique.

Lemma 27. *Let F be concave, $\bar{s} \in [2p - \underline{\theta}, \overline{\theta}]$, and $G^* := G^*_{\bar{s}}$. Draw a pass mark C independently of the signals with*

$$\mathbb{P}(C \le z) = K_{\bar{s}}(z) := \min\left\{1, \frac{I_{G^*}(z)}{(\bar{s} - p)\, G^*(z)^2}\right\},$$

and select the higher signal if it is at least C, tossing a coin otherwise. Then $K_{\bar{s}}$ is a CDF with $K_{\bar{s}} = 1$ on $[t_2, \infty)$, and (G^*, G^*) is an equilibrium in which the agent's value function is $L_{\bar{s}}$ on $[\underline{\theta}, \bar{s}]$, so the rule attains $u_F(\bar{s})$. If F is not uniform and $\bar{s} < \overline{\theta}$, it is the unique equilibrium.

Proof. Equilibrium. Against a rival G that is atomless on $(\underline{\theta}, \infty)$, $(G^2 K)' = 2G\, V'_G$ on $(\underline{\theta}, \infty)$, as in the proof of Lemma 20(iii). With $G = G^*$ and $K = K_{\bar{s}}$, $(G^{*2}K)' = 2dG^*$ with $d = \frac{1}{2(\bar{s}-p)}$, i.e., $V'_{G^*} = d$, and $V_{G^*}(\bar{s}) = K(\bar{s}) = 1$, so $V_{G^*} = L_{\bar{s}}$ on $(\underline{\theta}, \bar{s}]$. If G^* has an atom at $\underline{\theta}$, signals there tie with each other, and $V_{G^*}(\underline{\theta}) = V_{G^*}(\underline{\theta}^+)$, as the gap $\frac{1}{2}K(\underline{\theta}^+)G^*(\{\underline{\theta}\})$ vanishes: $I_{G^*}(x) \le x - \underline{\theta}$, while $G^*(x)$ stays above the mass of the atom, so $K(\underline{\theta}^+) = 0$. So $V_{G^*} = L_{\bar{s}}$ on $[\underline{\theta}, \bar{s}]$, while $V_{G^*} = 1 < L_{\bar{s}}$ above $\bar{s}$; hence every feasible signal yields at most $\int L_{\bar{s}}\, dG' = \frac{1}{2}$. On $[t_2, \bar{s}]$, $I_{G^*} = \frac{d}{2}(z - 2p + \bar{s})^2$ and $G^* = d(z - 2p + \bar{s})$, so $K = 1$. K is nondecreasing if and only if $\sqrt{I_{G^*}}$ is concave, as in Lemma 20(iii), which holds: $\sqrt{I_F}$ is concave on $[\underline{\theta}, t_1]$, as the argument of Lemma 20(iii) uses only concavity of F, I_{G^*} is affine on $[t_1, t_2]$, $\sqrt{I_{G^*}}$ is affine on $[t_2, \bar{s}]$, and I_{G^*} is continuously differentiable on $(\underline{\theta}, \infty)$.

Uniqueness. Write $K := K_{\bar{s}}$ and $m_R := G^*(t_1)$, which is $F(t_1)$ unless G^* keeps only part of an atom at $t_1 = \underline{\theta}$. Against a rival G, a signal x wins with probability $V_G(x) =$

$\frac{1}{2} + \frac{1}{2}K(x)G(x^-) - \frac{1}{2}\int_{(x,\infty)} K\,dG$. If (G, G') is an equilibrium, then by interchangeability so is (G, G^*), so G is a best reply to G^* and G^* is a best reply to G. The first gives $\mathrm{supp}(G) \subseteq [\underline{\theta}, \bar{s}]$. The second is used through deviations from G^* that remain feasible: any mean-preserving contraction of G^*, and any small mean-preserving perturbation of G^* compactly supported in $(t_1, \overline{\theta})$, where $I_{G^*} < I_F$.

Step 1. Two-sided perturbations inside $(t_2, \bar{s})$ make V_G affine there. As $K = 1$ there, V_G is the mid-CDF of G, so G has a constant density d' on $(t_2, \bar{s})$ and no atoms there. Spreading mass from near $\bar{s}$ to just above it, where $V_G = 1$, is unprofitable only if G has no atom at $\bar{s}$. So $V_G = \ell$ on $(t_2, \bar{s})$ with $\ell(x) := 1 - d'(\bar{s} - x)$.

Step 2. G has no atom in $(\underline{\theta}, t_1)$: at an atom a, V_G jumps up by $K(a)\,G(\{a\})$, so pooling a small interval of types around a with mean just above a would pay. Hence V_G is continuous on $(\underline{\theta}, t_1)$. Merging a small mass of G^* at a signal $u < t_2$ with pooled mass at $x \in (t_2, \bar{s})$ into a point $y \in (t_2, x)$ is a contraction of G^* and changes the payoff by a positive multiple of $\ell(u) - V_G(u)$. Hence $V_G \geq \ell$ G^*-almost everywhere below t_2.

Step 3. If $d' > d$, then $G < G^*$ on $(t_2, \bar{s})$, and $G \leq 1 - d'(\bar{s} - t_2) < m_R = G^*$ on (t_1, t_2), while $\int_{\underline{\theta}}^{t_1} (G - G^*)\,dx = I_G(t_1) - I_F(t_1) \leq 0$. So $\int (G - G^*)\,dx < 0$, contradicting equal means. Hence $d' \leq d$.

Step 4. The equilibrium value gives $\frac{1}{2} = \int V_G\,dG^* \geq \int \ell\,dG^* = \ell(p) = 1 - \frac{d'}{2\bar{d}}$, by Steps 1 and 2. Hence $d' = d$, $\ell = L_{\bar{s}}$, $G = G^*$ on $(t_2, \bar{s}]$, and, as the inequality is now an equality, $V_G = L_{\bar{s}}$ G^*-almost everywhere below t_2.

Step 5. K is nondecreasing and positive on $(\underline{\theta}, \infty)$. If $\bar{s} = 2p - \underline{\theta}$, then $G^* = \bar{U}_{\bar{s}}$ and Step 4 already gives $G = G^*$. If $t_1 = \underline{\theta}$ otherwise, then G^* has an atom at $\underline{\theta}$, and $V_G(\underline{\theta}) = L_{\bar{s}}(\underline{\theta}) = V_{G^*}(\underline{\theta})$ with $G = G^*$ above t_2 gives $\int_{(\underline{\theta}, t_2]} K\,dG = 0$, so G has no mass on $(\underline{\theta}, t_2]$ and $G = G^*$. Otherwise, letting $x \uparrow t_1$ in $V_G(x) = L_{\bar{s}}(x) = V_{G^*}(x)$, with $G = G^*$ above t_2 and G^* putting no mass on $[t_1, t_2]$, gives $K(t_1)\big(G(t_1^-) - m_R\big) = \int_{[t_1, t_2]} K\,dG \geq K(t_1)\,G([t_1, t_2]) = K(t_1)\big(m_R - G(t_1^-)\big)$, so $G(t_1^-) \geq m_R$; as $G(t_1^-) \leq G(t_2) = m_R$, G has no

mass on $[t_1, t_2]$. On $[\underline{\theta}, t_1)$, with $E := G - F$, the same comparison and an integration by parts give $2K(x)E(x) = -\int_x^{t_1} E\, dK =: -\Phi(x)$. For every $x_0 \in (\underline{\theta}, t_1)$, $K \geq K(x_0) > 0$ on $[x_0, t_1]$, so $|\Phi(x)| \leq \int_x^{t_1} \frac{|\Phi|}{2K(x_0)}\, dK$ there, and $\Phi \equiv 0$ on $[x_0, t_1]$ by Grönwall's inequality. So $E \equiv 0$ on $(\underline{\theta}, t_1)$, and also at $\underline{\theta}$ by right-continuity. Hence $G = G^*$, and every equilibrium is (G^*, G^*). □

C.8.12 Continuous rules

Lemma 28. *Let F be concave and not uniform, let $\bar{s}^*$ be the best width, and let K be the pass-mark CDF of Lemma 27 at $\bar{s}^*$, restricted to signals in $[\underline{\theta}, \overline{\theta}]$. For $\varepsilon > 0$, let*

$$H_1^\varepsilon(x, y) := \tfrac{1}{2} + K(\max(x, y))\, \rho\Big(\frac{x - y}{\varepsilon}\Big), \qquad \rho(t) := \max\{-\tfrac{1}{2}, \min\{\tfrac{t}{2}, \tfrac{1}{2}\}\}.$$

H^ε is continuous and equals the pass-mark rule when $|x - y| \geq \varepsilon$. It has an equilibrium, and as $\varepsilon \to 0$, every equilibrium strategy of H^ε converges weakly to $G^*_{\bar{s}^*}$, and the firm's payoff in every equilibrium converges to $u_F(\bar{s}^*)$. Likewise, if F is convex and atomless with $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$ *and $K \equiv 1$, every equilibrium strategy converges weakly to F, and the firm's payoff converges to the first best.*

Proof. Write H for the pass-mark rule and $G^* := G^*_{\bar{s}^*}$. K, with $K(\underline{\theta}) := K(\underline{\theta}^+)$, is continuous on $[\underline{\theta}, \overline{\theta}]$, so H^ε is continuous, and an equilibrium exists by Sion's theorem; each equilibrium strategy guarantees $\frac{1}{2}$. H is continuous off the diagonal, $H^\varepsilon \to H$ uniformly off every neighborhood of the diagonal, and at a diagonal point (y, y) all limits of H^ε lie between $\frac{1}{2} - \frac{1}{2}K(y)$ and $\frac{1}{2} + \frac{1}{2}K(y)$. Hence, if $\mu_\varepsilon \to \mu$ weakly and μ puts no mass on the diagonal, $\int H^\varepsilon\, d\mu_\varepsilon \to \int H\, d\mu$; the same holds if μ puts mass on the diagonal only at $(\underline{\theta}, \underline{\theta})$ and $K(\underline{\theta}^+) = 0$.

Let G_ε be equilibrium strategies; the feasible set is weakly compact, so take a convergent subsequence, $G_\varepsilon \to G$. (i) G is a best reply to G^* under H: $\iint H^\varepsilon\, dG_\varepsilon\, dG^* \geq$

$\frac{1}{2}$, and $G \otimes G^*$ puts mass on the diagonal only where G^* has an atom, which is at most at $\underline{\theta}$, where $K(\underline{\theta}^+) = 0$; so $\int V_{G^*}\, dG \geq \frac{1}{2}$. (ii) G^* is a best reply to G under H against every deviation G' that has no atom in common with G except possibly at $\underline{\theta}$: $\iint H^\varepsilon\, dG'\, dG_\varepsilon \leq \frac{1}{2}$, and $G' \otimes G$ puts no mass on the diagonal, so $\int V_G\, dG' \leq \frac{1}{2}$, while $\int V_G\, dG^* = \frac{1}{2}$. The uniqueness part of the proof of Lemma 27 uses only (i) and (ii): every deviation in its Steps 1 and 2 can avoid the countably many atoms of G other than $\underline{\theta}$. Hence $G = G^*$, and the whole family converges to G^*. The firm's payoff is $\iint c_\varepsilon\, dG_\varepsilon\, dG'_\varepsilon$ with $c_\varepsilon(x, y) := y + (x - y)H_1^\varepsilon(x, y)$, which converges uniformly, as $|c_\varepsilon - c| \leq \varepsilon$, to $c(x, y) := y + (x - y)H(x, y)$, a continuous function; so the payoff converges to $u_F(\bar{s}^*)$.

For convex F, (i) and (ii) hold with F in place of G^*, as full revelation is an equilibrium of the noiseless rule and F is atomless; $V_G(x) = G(x^-) + \frac{1}{2}G(\{x\})$ is the mid-CDF. G has no atom: it has none at $\underline{\theta}$ or $\overline{\theta}$, as $G \leq_{cx} F$ and F is atomless, and at an interior atom x_0, V_G jumps up, and pooling the types near x_0 into a point just above x_0 that is not an atom of G would pay. So $V_G = G$ is continuous, and every $G' \leq_{cx} F$ is admissible in (ii). Pooling the types in an interval at their mean creates a single atom, which G does not have, and F is continuous and strictly increasing on $[\underline{\theta}, \overline{\theta}] = \mathrm{supp}(F)$. So G is convex on $[\underline{\theta}, \overline{\theta}]$ by Lemma 15 with $K = F$ and $h = G$. By (i), (ii) and constant-sum, $\int G\, dF = \int F\, dG = \frac{1}{2} = \int F\, dF = \int G\, dG$, so $\int F\, d(F - G) = \int G\, d(F - G) = 0$, and the last paragraph of Appendix C.8.4, with $m = 0$ and $\hat{G} = G$, gives $G = F$, and the payoff converges to $\mathbb{E}\max(\theta_1, \theta_2)$. □

C.8.13 The noiseless equilibrium for a general prior

For regular priors, the following structure is due to Hwang et al. (2026, Theorem 1). I prove it under the weaker conditions used in Subsection 5.2.4, as the proof of Theorem 4 uses its details.

Lemma 29. *Let F be atomless, with* $\operatorname{supp}(F) = [\underline{\theta}, \overline{\theta}]$ *and a density f that is continuous and positive on $(\underline{\theta}, \overline{\theta})$. A symmetric profile (G, G) is an equilibrium of the noiseless rule if and only if*

(a) G is atomless and $\operatorname{supp}(G) = [\underline{\theta}, c]$ *for some $c \in (\underline{\theta}, \overline{\theta}]$;*

(b) G is convex on $[\underline{\theta}, c]$;

(c) $G \leq_{cx} F$;

(d) G is affine on every component of $(\underline{\theta}, c) \setminus E$, where $E := \{x \in [\underline{\theta}, c] : I_G(x) = I_F(x)\}$.

In that case, $G = F$ on E; $c \notin E$ if $c < \overline{\theta}$; the function ψ that equals G on $[\underline{\theta}, c]$ and $1 + G'(c^-)(x - c)$ on $[c, \overline{\theta}]$ is convex, $\psi \geq G$, and $d\psi'$ is carried by E; the firm's utility is $\mathbb{E}_G[\max(s_1, s_2)]$; and (G, G) is the unique symmetric equilibrium. If (a, b) is a component of $(\underline{\theta}, c) \setminus E$ and a is a limit of points of E from the left, the slope of G on (a, b) is $f(a)$.

Proof. Against a rival who plays K, the noiseless rule gives a signal x the payoff $\bar{K}(x) = K(x^-) + \frac{1}{2}K(\{x\})$, and G is a best reply to itself if and only if it maximizes $\int \bar{G}\, dG'$ over feasible G'.

No atoms. Let (G, G) be an equilibrium. An atom at $\underline{\theta}$ or at $\overline{\theta}$ is infeasible, as in Appendix C.8.4: at $\underline{\theta}$ because $I_F(x) = o(x - \underline{\theta})$, and at $\overline{\theta}$ because $I_F - I_G \geq 0$ vanishes there, so its left derivative $F(\overline{\theta}-) - G(\overline{\theta}-)$ is not positive. An interior atom is ruled out by the deviation of Appendix C.8.4. So $\bar{G} = G$ is continuous.

The certificate. For a continuous convex ψ on $[\underline{\theta}, \overline{\theta}]$ with $\psi'(\underline{\theta}^+)$ finite and distributions K_1, K_2 on $[\underline{\theta}, \overline{\theta}]$ with equal means, writing $\psi(x) = \psi(\underline{\theta}) + \psi'(\underline{\theta}^+)(x - \underline{\theta}) + \int (x - a)^+\, d\psi'(a)$ (at $x = \overline{\theta}$ by monotone convergence) and using $\int (x-a)^+\, dK(x) = p - a + I_K(a)$ gives

$$\int \psi\, dK_1 - \int \psi\, dK_2 = \int \left(I_{K_1} - I_{K_2}\right) d\psi', \tag{6}$$

which extends by linearity to differences of such functions. Hence, if G is continuous and such a $\psi \geq G$ has $\psi = G$ on $\operatorname{supp}(G)$ and $d\psi'$ carried by $\{I_F = I_G\}$, then $\int G\,dG' \leq \int \psi\,dG' \leq \int \psi\,dF = \int \psi\,dG = \int G\,dG$ for every feasible G', so G is a best reply to itself.

Necessity. Let $\ell := \min \operatorname{supp}(G) < c := \max \operatorname{supp}(G)$. Every $G' \leq_{cx} G$ is feasible, so Lemma 16, applied to the prior G and $V = G$, gives a convex $\phi \geq G$ on $[\ell, c]$ with $\phi = G$ on $\operatorname{supp}(G)$. If $a < b$ are in $\operatorname{supp}(G)$ with $G(a) = G(b)$, then ϕ equals $G(a)$ at a and b and is at least $G \geq G(a)$ between them, so it is constant on $[a, b]$ and nonincreasing on $[\ell, a]$. Then $0 = G(\ell) = \phi(\ell) \geq \phi(a) = G(b)$, which contradicts $\ell \in \operatorname{supp}(G)$. So $\operatorname{supp}(G) = [\ell, c]$, and $G = \phi$ is convex on $[\ell, c]$; extended by 0, it is convex on $[\underline{\theta}, c]$. Let $D := I_F - I_G \geq 0$, and let $[x_1, x_3] \subseteq [\underline{\theta}, c]$ with $\kappa := \min_{[x_1,x_3]} D > 0$. Then G is affine on $[x_1, x_3]$. Otherwise, being convex, G lies strictly below its secant s through x_1 and x_3 on (x_1, x_3) and is not constant there. So some $[y_1, y_2] \subset (x_1, x_3)$ has G-mass $m > 0$, and $\delta := \min_{[y_1,y_2]}(s - G) > 0$. Splitting a share ε of that mass, each point onto $\{x_1, x_3\}$ with its own mean, gives G_ε with $0 \leq I_{G_\varepsilon} - I_G \leq \varepsilon m(x_3 - x_1)$, and the difference vanishes off $[x_1, x_3]$. So $G_\varepsilon \leq_{cx} F$ for small ε, while $\int G\,dG_\varepsilon - \int G\,dG = \varepsilon \int_{[y_1,y_2]}(s - G)\,dG \geq \varepsilon m\delta > 0$, a contradiction. If $\ell > \underline{\theta}$, then $D = I_F > 0$ on $(\underline{\theta}, \ell]$, so $D > 0$ on some $[x_1, \ell + \eta]$ with $\underline{\theta} < x_1 < \ell < \ell + \eta < c$. But G vanishes on $[x_1, \ell]$ and is positive at $\ell + \eta$, so it is not affine there; hence $\ell = \underline{\theta}$, which proves (a)–(c). Each compact subinterval of a component of $(\underline{\theta}, c) \setminus E$ has $\min D > 0$, which proves (d). If $c < \overline{\theta}$, then $D(x) = \int_x^{\overline{\theta}} (1 - F)\,dt > 0$ on $[c, \overline{\theta})$, so $c \notin E$, and by (d) G is affine on a left neighbourhood of c, so $G'(c^-) < \infty$. At every $x \in E \cap (\underline{\theta}, \overline{\theta})$, D is differentiable and minimal, so $F(x) = G(x)$.

Sufficiency and payoff. Given (a)–(d), the function ψ of the statement is convex, $\psi \geq G$, $\psi = G$ on $[\underline{\theta}, c]$, and $d\psi'$ is carried by E, so (G, G) is an equilibrium. The firm selects the higher signal, so $u_F = \mathbb{E}_G[\max(s_1, s_2)]$.

Uniqueness. Let G and $\hat{G}$ satisfy (a)–(d), with certificates $\psi, \hat{\psi}$, tops $c, \hat{c}$ and contact sets $E, \hat{E}$. By interchangeability $(G, \hat{G})$ is an equilibrium, so $\frac{1}{2} = \int G\, d\hat{G} \leq \int \psi\, d\hat{G} \leq \int \psi\, dF = \frac{1}{2}$. Hence $\psi = G$ on $[\underline{\theta}, \hat{c}]$, and, by (6), $d\psi'$ is carried by $\hat{E}$. If $\hat{c} > c$, then $\psi = 1$ on $[c, \hat{c}]$, so $G'(c^-) = 0$ and $G = 0$ on $[\underline{\theta}, c)$, which is absurd. So $\hat{c} \leq c$, symmetrically $c \leq \hat{c}$, and dG', $d\hat{G}'$ are carried by $E \cap \hat{E}$, where $G = F = \hat{G}$. Off this set, G and $\hat{G}$ are affine on each interval and agree at its endpoints, which lie in $E \cap \hat{E}$ or are $\underline{\theta}$ or c, so $G = \hat{G}$.

Tangency. Let γ be the slope of G on (a, b). As $I_G \leq I_F$ with equality at a and $G(a) = F(a)$, $\gamma \frac{(x-a)^2}{2} \leq \int_a^x (F - F(a))\, dt$ for $x \in (a, b)$, so $\gamma \leq f(a)$. If $x_n \uparrow a$ with $x_n \in E$, the left derivative of G at a is $\lim \frac{F(a)-F(x_n)}{a-x_n} = f(a)$, and convexity gives $\gamma \geq f(a)$. □

C.8.14 Proof of Theorem 4

Lemma 30. *Let F be a prior, $G \leq_{cx} F$ with $\underline{\theta} \in \operatorname{supp}(G)$, and $\psi : [\underline{\theta}, \overline{\theta}] \to \mathbb{R}$ convex, continuous and nondecreasing, with $\psi(\underline{\theta}) \geq 0$, $\psi \leq 1$ on $\operatorname{supp}(G)$, $\int \psi\, dG = \frac{1}{2}$, and $d\psi'$ carried by $\{I_G = I_F\}$. Suppose that, for every t,*

$$\int_{[t,\overline{\theta}]} \psi\, dG \leq \frac{1 - G(t^-)^2}{2} \qquad \text{and} \qquad \int_{(t,\overline{\theta}]} \psi\, dG \leq \frac{1 - G(t)^2}{2}.$$

Then some general selection rule nondecreasing in the own signal has the equilibrium (G, G), the agent's value function equals ψ on $\operatorname{supp}(G)$, and the firm's utility is $2\int s\, \psi(s)\, dG(s)$.

Proof. Let $Y := \psi(s)$ for $s \sim G$; it has mean $\frac{1}{2}$ and values in $[0, 1]$. As ψ is nondecreasing, its upper level sets are intervals $[t, \overline{\theta}]$ or $(t, \overline{\theta}]$, so the two inequalities are the bound of the necessity part of the proof of Lemma 18 at every upper level set, and the interpolation argument there gives $Y \leq_{cx} U[0, 1]$; then Lemmas 18 and 20(i), applied with G in place of F, give a rule nondecreasing in the own signal, a lottery over coarse gradings, with $V_G = \psi$ on $\operatorname{supp}(G)$. A signal x off the support has the same quantile as

the highest support point $r(x)$ below it, which exists as $\underline{\theta} \in \operatorname{supp}(G)$, so it is graded like $r(x)$, and $V_G(x) = \psi(r(x)) \le \psi(x)$, as ψ is nondecreasing. For every $G' \le_{cx} F$, by (6), $\int V_G \, dG' \le \int \psi \, dG' \le \int \psi \, dF = \int \psi \, dG = \frac{1}{2} = \int V_G \, dG$. So (G, G) is an equilibrium, and Lemma 17 gives $u_F = p + 2\operatorname{Cov}_G(s, \psi(s)) = 2\int s\,\psi \, dG$. □

From now on, let F have a well-behaved noiseless equilibrium (G_0, G_0) with firm's utility u_0, and let c, E_0 and the certificate ψ_0 be as in Lemma 29; let g_0 be the right derivative of G_0 and $\Delta := I_F - I_{G_0} \ge 0$. A *pool* is a component of $(\underline{\theta}, c) \setminus E_0$. By (H2), merging adjacent blocks with equal slopes, $[\underline{\theta}, c]$ is a finite union of maximal revealed stretches and maximal blocks. Two adjacent maximal blocks meet at a point of E_0 where g_0 jumps up, a *kink*, as G_0 is convex and dg_0 is carried by E_0. A *chain* is a maximal interval $[u, v]$ that is a union of adjacent maximal blocks $B_i = [b_{i-1}, b_i]$, $i = 1, \dots, K$, with $b_0 = u$, $b_K = v$, slopes $\gamma_1 < \cdots < \gamma_K$ and widths $W_i := b_i - b_{i-1}$. Then $u = \underline{\theta}$ or u is the upper end of a revealed stretch, and $v = c$ or v is the lower end of one.

Proposition 20. *Let F have a well-behaved noiseless equilibrium (G_0, G_0), and let F not be convex. Let u_0 be the firm's utility in (G_0, G_0), $c := \max \operatorname{supp}(G_0)$, $\psi_0 := G_0$ on $[\underline{\theta}, c]$ and $\psi_0(x) := 1 + G_0'(c^-)(x - c)$ on $[c, \overline{\theta}]$, and let P and S be the unions of the components of $(\underline{\theta}, c) \setminus \{I_{G_0} = I_F\}$ and of the revealed stretches. For every $\theta \in [0, \frac{1}{2})$, there are a continuous, piecewise twice continuously differentiable h on $[\underline{\theta}, \overline{\theta}]$ and $\varepsilon_0 > 0$ such that, with $H(t) := \int_{[\underline{\theta}, t]} h \, dG_0$,*

$$\Phi_\theta := (2 - 4\theta) \int_P H(t)\, dt - 2 \int_S H(t)\, dt > 0,$$

and, for every $\varepsilon \in (0, \varepsilon_0)$, some general selection rule nondecreasing in the own signal has a symmetric equilibrium $(G_\varepsilon, G_\varepsilon)$ in which the agent's value function equals $\psi_0 + \varepsilon h$ on

$\operatorname{supp}(G_\varepsilon)$ *and the firm's utility* $u(\varepsilon)$ *satisfies*

$$\varepsilon\,\Phi_\theta - o(\varepsilon) \le u(\varepsilon) - u_0 \le \varepsilon\,\Phi_\theta.$$

The equilibrium signal is built from the following tangency lemma, applied at every contact point of a chain.

Lemma 31. *Let* $r > 0$*, and let* A *and* Γ *be continuously differentiable and convex on* $[a, a+r]$*, with* A' *strictly increasing,* $\Gamma'' = \gamma > 0$*,* $A(a) = \Gamma(a)$*,* $A'(a) = \Gamma'(a)$*,* $A > \Gamma$ *on* $(a, a+r]$ *and* $A' - \Gamma'$ *nondecreasing. Let* $T(x) = T_0 + T_1(x-a) + \frac{\tau}{2}(x-a)^2$ *with* $\tau \le 0$*,* $T \ge 0$ *on* $[a, a+r]$*, and either* $T_0 > 0$ *or* $T_0 = 0 < T_1$*, and let* $\Lambda := \Gamma + \varepsilon T$*. For all sufficiently small* $\varepsilon > 0$*, there are* $a < x_1 < x_2 < a + r$ *such that*

(i) the tangent ℓ *of* A *at* x_1 *is the tangent of* Λ *at* x_2*;*

(ii) $\Lambda \le A$ *on* $[x_2, a+r]$*;*

(iii) $\Gamma \le \ell \le \Lambda$ *on* $[x_1, x_2]$ *and* $A \le \Lambda$ *on* $[a, x_1]$*;*

(iv) $x_2 \to a$ *as* $\varepsilon \to 0$*.*

Proof. Let $\varepsilon < \gamma/(2|\tau|)$, so that $\Lambda'' \ge \frac{\gamma}{2}$, and let $\rho := \varepsilon T - (A - \Gamma)$. As $\rho' = \varepsilon T' - (A' - \Gamma')$ is nonincreasing, ρ is concave; $\rho(a) = \varepsilon T_0 \ge 0$, $\rho > 0$ just right of a (if $T_0 = 0$, as $\rho'(a) = \varepsilon T_1 > 0$), and $\rho(a+r) < 0$ for small ε. So ρ has a unique zero $x^* \in (a, a+r)$, $\rho > 0$ on (a, x^*), $\rho < 0$ on $(x^*, a+r]$ and $\rho'(x^*) < 0$; and $x^* \to a$ as $\varepsilon \to 0$, as $A - \Gamma$ is bounded away from zero on $[a+\eta, a+r]$ for every $\eta > 0$. By $\rho'(x^*) < 0$, $A'(x^*) > \Lambda'(x^*) \ge \Lambda'(a)$, and $A'(x^*) > A'(a)$. For $\sigma \in [\sigma_0, A'(x^*)]$ with $\sigma_0 := \max\{A'(a), \Lambda'(a)\}$, let $x_1(\sigma) \in [a, x^*]$ and, for small ε, $x_2(\sigma) \in [a, a+r]$ solve $A'(x_1) = \sigma$ and $\Lambda'(x_2) = \sigma$, and let $\tau(\sigma)$ be the intercept of the tangent of A at x_1 minus that of the tangent of Λ at x_2; it is continuous. At σ_0, $\tau < 0$: if $\sigma_0 = \Lambda'(a)$, then $x_2 = a$, and if moreover $x_1 > a$, the tangent of A at x_1

lies strictly below $A(a) = \Gamma(a) \leq \Lambda(a)$ at a, while if $x_1 = a$, then $T_0 > 0$, as $T_0 = 0$ gives $\Lambda'(a) > A'(a)$, and $\tau = -\varepsilon T_0$; if $\sigma_0 = A'(a) > \Lambda'(a)$, then $x_1 = a$, $x_2 > a$, the tangent of A at a is that of Γ, and $\tau \leq \Gamma(x_2) - \Lambda(x_2) = -\varepsilon T(x_2) < 0$, as $T > 0$ on $(a, a + r)$ near a. At $A'(x^*)$, $x_1 = x^*$ and $\tau \geq A(x^*) - \Lambda(x^*) = -\rho(x^*) = 0$, as the tangent of the convex Λ lies below it. Take a zero of τ: the two tangents coincide, ℓ. As $\ell \leq A$, $\rho(x_2) = \ell(x_2) - A(x_2) \leq 0$, and $x_2 \neq a$ as $\tau(\sigma_0) < 0$, so $x_2 \geq x^* \geq x_1$, and $x_1 = x_2$ would give $A'(x^*) = \Lambda'(x^*)$; so $x_1 < x_2$, which proves (i). (ii) holds as $\rho \leq 0$ on $[x^*, a + r]$. In (iii), $\ell \leq \Lambda$ by convexity, $\ell \geq \Gamma$ at x_1 and x_2 and hence between them, as Γ is convex, and $\rho \geq 0$ on $[a, x_1]$. For (iv), $\frac{\gamma}{2}(x_2 - a) \leq \Lambda'(x_2) - \Lambda'(a) \leq A'(x^*) - A'(a) + \varepsilon|T_1| \to 0$. □

By the reflection $x \mapsto 2b - x$, which preserves all hypotheses, the same holds at an upper endpoint b, on $[b - r, b]$, with either $T(b) > 0$ or $T(b) = 0 > T'(b)$, and $b - r < x_2 < x_1 < b$.

Proof of Proposition 20. As F is not convex, G_0 has a block: otherwise $[\underline{\theta}, c] \subseteq E_0$, so $G_0 = F$ on $[\underline{\theta}, c]$, and $c = \overline{\theta}$, as $c \notin E_0$ if $c < \overline{\theta}$; then $F = G_0$ would be convex. So a chain $[u, v]$ exists; fix one and $\theta \in [0, \frac{1}{2})$.

Step 1 (the direction on the chain). On each block B_i of the chain, let H be the following polynomial of degree at most two and $h := H'/\gamma_i$:

(a) if $K = 1$, $u = \underline{\theta}$ and $v = c$: $H(t) = 4(t - \underline{\theta})(c - t)/W_1^2$;

(b) if $K = 1$, $u > \underline{\theta}$ and $v = c$: $H(t) = 1 - (t - u)/W_1$;

(c) otherwise: $H(t) = 1 - \big((b_1 - t)/W_1\big)^2$ on B_1 if $u = \underline{\theta}$, $H(t) = 1 - \big((t - b_{K-1})/W_K\big)^2$ on B_K if $v = c$, and $H \equiv 1$ on every other block.

If $v = c$, extend h affinely to $[b_{K-1}, \overline{\theta}]$. Then h is affine and nonincreasing on each block, $H \geq 0$ on $[u, v]$, and $H > 0$ on the interior of each block. At each kink, $H = 1$ and $h = 0$ on both sides; $h(\underline{\theta}) > 0$ if $u = \underline{\theta}$, and $h(c) < 0$ if $v = c$. If $v < c$, then $h(v) = 0 \geq h'(v^-)$. If $u > \underline{\theta}$, then $h'(u^+) = 0$, and $h(u) = 0$ except in case (b), where $h(u) = -1/(\gamma_1 W_1)$.

Step 2 (sinks). Let $\delta > 0$. If $u > \underline{\theta}$, let $[s, u]$ be the revealed stretch below u. As f' is continuous and nonnegative on $[s, u]$ and vanishes on no interval, as F is strictly convex, there is a window $[w, w'] \subseteq (u - \delta, u)$ on which $f' \geq m$ for some $m > 0$. Outside case (b), let $h := \lambda(t - w)^2(w' - t)^2$ on $[w, w']$ and $h := 0$ elsewhere on $[s, u]$, with $\lambda > 0$ such that $\int_w^{w'} h \, dF = 1$; then H rises from 0 to 1 on $[w, w']$ and equals 1 on $[w', u]$. In case (b), with $h_u := -1/(\gamma_1 W_1)$, take the window $[w', w]$ with $w - w' \leq u - w$, let $h := 0$ on $[s, w']$, $h(t) := h_u + k(t - u)$ on $[w, u]$, and let h on $[w', w]$ be the cubic Hermite interpolant from value 0 and slope 0 at w' to value $h_u + k(w - u)$ and slope k at w. Then $\int_{w'}^{u} h \, dF$ is affine and strictly decreasing in k, with slope at most $2 \sup f \, (u - w')^2$ in absolute value, and nonpositive at $k = 0$. So k with $\int_{w'}^{u} h \, dF = 1$ is negative and $h(w) = h_u + |k|(u - w) \geq h_u + \frac{1}{4 \sup f \, (u - w')} > 0$ for small δ. Then $h \geq 0$ on $[w', w]$, as both Hermite terms are nonnegative, and h changes sign once on $[w, u]$, so H rises from 0 and then falls to $H(u) = 1$, and $h'(u^-) = k < 0 = h'(u^+)$. If $v < c$, let $[v, e]$ be the revealed stretch above v, take a window $[w, w'] \subseteq (v, v + \delta)$ with $f' \geq m$, and let $h := -\mu(t - w)^2(w' - t)^2$ there and $h := 0$ elsewhere on $[v, e]$, with $\int_w^{w'} h \, dF = -1$; then $h(v) = h'(v^+) = 0$ and H falls from 1 to 0. Let $h := 0$ everywhere else. Then $H \geq 0$, H vanishes outside the chain and the at most two revealed stretches next to it, $H(c) = 0$, and $\int_S H \leq 2\delta(1 + |h_u| \sup f \, \delta)$. As $\int_P H > 0$ does not depend on δ, $\Phi_\theta > 0$ for small δ; fix such a δ.

Step 3 (the certificate). Let $\psi_\varepsilon := \psi_0 + \varepsilon h$. On revealed stretches, $\psi_0'' = f' \geq 0$, and h is continuously differentiable except at the ends of the stretches, with $h'' \neq 0$ only on the windows, where $\psi_\varepsilon'' \geq m - \varepsilon \sup |h''| \geq 0$ for small ε; in case (b), h is affine on $[w, u]$, so $\psi_\varepsilon'' = f' \geq 0$ there even where $f'(u) = 0$. On blocks and above c, ψ_ε is affine, also across contact points inside a block, where h has no kink. At a kink b_i, ψ_0' jumps up by $\gamma_{i+1} - \gamma_i > 0$, which dominates the jump of $\varepsilon h'$ for small ε; at u and v, $h'(u^-) \leq h'(u^+)$ and $h'(v^-) \leq h'(v^+)$, and ψ_0 is convex. So ψ_ε is convex and continuous.

It is nondecreasing for small ε, as $h' \neq 0$ only where ψ_0' is bounded away from zero, and $\psi_\varepsilon(\underline{\theta}) = \varepsilon h(\underline{\theta}) \geq 0$. As $\psi_\varepsilon(c) = 1 + \varepsilon h(c) \leq 1$, $\psi_\varepsilon \leq 1$ on $[\underline{\theta}, c]$. Finally, $d\psi_\varepsilon'$ is carried by E_0, on which $d\psi_0'$ is carried, together with the sink windows and the kinks of h, all of which lie in $S \cup E_0$.

Step 4 (the signal). On each pool Q of the chain, with slope γ, let $T := (1-\theta)H/\gamma$, a polynomial of degree at most two with $T' = (1-\theta)h$ nonincreasing, $T \geq 0$, and $T > 0$ inside Q; let $\Lambda_Q := I_{G_0} + \varepsilon T$. At each endpoint $x_0 \in E_0$ of Q, apply Lemma 31, or its reflection, with $A := I_F$ and $\Gamma := I_{G_0}$, on a one-sided neighbourhood of x_0 in Q of length r on which $\Delta > 0$ by (H2) and $f \geq \gamma$ almost everywhere by (H3), so that $A' - \Gamma' = F - G_0$ is nondecreasing; $A' = F$ is strictly increasing as $f > 0$. Its conditions on T hold: $T(x_0) > 0$ at every contact point in $(\underline{\theta}, \overline{\theta})$, as $H(x_0) > 0$ there; $T(\underline{\theta}) = 0 < T'(\underline{\theta})$ if $x_0 = \underline{\theta}$; and $T(\overline{\theta}) = 0 > T'(\overline{\theta})$ if $x_0 = \overline{\theta} = c$. The *revealed part* of the resulting layer is the interval between x_0 and x_1, and its *gap* is (x_1, x_2), or (x_2, x_1). Define K by

$$K := \begin{cases} I_F & \text{on revealed parts,} \\ \ell & \text{on gaps,} \\ \Lambda_Q & \text{on the rest of each pool } Q \text{ of the chain,} \\ I_{G_0} & \text{elsewhere.} \end{cases}$$

For small ε, $\Lambda_Q \leq I_F$ on the rest of Q: near the contact points by Lemma 31(ii), and elsewhere, also near c when $c < \overline{\theta}$, as Δ is bounded away from zero there while $\varepsilon T \leq \varepsilon \sup T$. The pieces of K are convex and meet with equal values and slopes, except at c when $v = c < \overline{\theta}$, where $K'(c^-) = 1 + \varepsilon(1-\theta)h(c) < 1 = K'(c^+)$. So K is convex, $K \leq I_F$, $K(\overline{\theta}) = I_F(\overline{\theta})$, and $G_\varepsilon := K'$ is a CDF with $G_\varepsilon \leq_{cx} F$, $\operatorname{supp}(G_\varepsilon) \subseteq [\underline{\theta}, c]$ and $\underline{\theta} \in \operatorname{supp}(G_\varepsilon)$. It is continuous except for an atom of mass $\alpha := -\varepsilon(1-\theta)h(c)$ at c if $v = c < \overline{\theta}$; let $\alpha := 0$ otherwise. Let $J := (K - I_{G_0})/\varepsilon$ and $D := J' = (G_\varepsilon - G_0)/\varepsilon$, which

is continuous on $[\underline{\theta}, c)$. Then $J = 0$ outside the pools of the chain and on E_0; on these pools, $0 \le J \le T$ by Lemma 31(iii), and $J = T$ outside the layers, whose total length $\lambda(\varepsilon)$ tends to zero by Lemma 31(iv). As $K = I_F$ on $S \cup E_0$, $d\psi'_\varepsilon$ is carried by $\{K = I_F\}$.

Step 5 (totals). As $J = 0$ on E_0 and $dh' = 0$ on the pools, (6) gives $\int \psi_0 \, dG_\varepsilon = \frac{1}{2} + \varepsilon \int J \, d\psi'_0 = \frac{1}{2}$ and $\int h \, dG_\varepsilon = \int h \, dG_0 + \varepsilon \int J \, dh' = H(c) = 0$. So $\int \psi_\varepsilon \, dG_\varepsilon = \frac{1}{2}$.

Step 6 (the rival's scores). Let $B(t) := \int_{[t,\overline{\theta}]} \psi_\varepsilon \, dG_\varepsilon - \frac{1-G_\varepsilon(t^-)^2}{2}$. For $t > c$, $B(t) = 0$; $B(c) = \alpha\big(\varepsilon h(c) + \frac{\alpha}{2}\big) = \frac{1+\theta}{2}\alpha\varepsilon h(c) \le 0$, and with the atom excluded the value is 0. For $t < c$, as G_ε is continuous on $[t, c)$ and $\psi_\varepsilon - G_\varepsilon = \varepsilon(h - D)$ there,

$$B(t) = B(c) + \int_{[t,c)} (\psi_\varepsilon - G_\varepsilon) \, dG_\varepsilon = B(c) + \varepsilon \int_t^c (h - D) \, dG_0 + \varepsilon^2 \int_{[t,c)} (h - D) \, dD.$$

Integrating by parts, with $H(c) = 0$, $J(c) = 0$, $D(c^-) = -\alpha/\varepsilon$, and $\int_t^c J \, dg_0 = \int_t^c J \, dh' = 0$, gives

$$B(t) = -\varepsilon\big(H - g_0 J\big)(t) + \varepsilon^2\Big[D\Big(\frac{D}{2} - h\Big) + h'J\Big](t),$$

as the terms at c^- cancel $B(c)$. (i) On the rest of a pool, $J = T$, $g_0 = \gamma$ and $D = (1 - \theta)h$, so $B = -\varepsilon\theta H - \varepsilon^2 \frac{1-\theta^2}{2} h^2 + \varepsilon^2 h'T \le 0$. (ii) On a gap, G_ε is constant, so B is constant and equals its value at the end x_2, which is ≤ 0 by (i). (iii) On a maximal interval R within the closure of the pools of the chain on which $G_\varepsilon = F$, which contains at most one contact point, $B' = -f(\psi_\varepsilon - F) = f\varphi$ with $\varphi := F - G_0 - \varepsilon h$, which is continuous and nondecreasing, as $\varphi' = f - \gamma - \varepsilon h' \ge 0$ almost everywhere on each side of the contact point by (H3) and $h' \le 0$. So B is nonincreasing and then nondecreasing on R, and is at most its value at an end of R: a gap end, a point of a revealed stretch, $\underline{\theta}$, where $B = \int \psi_\varepsilon \, dG_\varepsilon - \frac{1}{2} = 0$, or $\overline{\theta}$, where $B = 0$. (iv) Elsewhere, $J = D = 0$, so $B = -\varepsilon H \le 0$. Hence both inequalities of Lemma 30 hold, as they differ only at the atom. With Steps 3–5, Lemma 30 gives a rule nondecreasing in the own signal with the equilibrium $(G_\varepsilon, G_\varepsilon)$ and value function ψ_ε on $[\underline{\theta}, c]$.

Step 7 (payoff). By Lemma 30, $u(\varepsilon) - u_0 = 2\int s\psi_\varepsilon \, dG_\varepsilon - 2\int s\psi_0 \, dG_0$. By (6), with $d(s\psi_0)' = 2\psi_0' \, ds + s \, d\psi_0'$, $d(sh)' = 2h' \, ds + s \, dh'$, $J = 0$ on E_0 and above c, and $\int s \, h \, dG_0 = -\int H \, ds$,

$$u(\varepsilon) - u_0 = \varepsilon\Big[4\int g_0 J \, ds - 2\int H \, ds\Big] + 4\varepsilon^2 \int Jh' \, ds.$$

As $J \le T = (1-\theta)H/\gamma$ on the pools of the chain, $J = 0$ off them, $H = 0$ outside $P \cup S$, and $Jh' \le 0$, $u(\varepsilon) - u_0 \le \varepsilon\Phi_\theta$. As $J = T$ off the layers, $u(\varepsilon) - u_0 \ge \varepsilon\Phi_\theta - 4\varepsilon\gamma_K \sup T \, \lambda(\varepsilon) - 4\varepsilon^2 \sup T \sup|h'|(c - \underline{\theta}) = \varepsilon\Phi_\theta - o(\varepsilon)$. □

Proof of Theorem 4. By Lemma 29 and interchangeability, (G_0, G_0) is the only equilibrium of the noiseless rule. If F is convex, $G_0 = F$, and the noiseless rule attains the first best by Proposition 10; no rule exceeds it, as $u_F \le \mathbb{E}[\max(s_1, s_2)] \le \mathbb{E}[\max(\theta_1, \theta_2)]$ in every equilibrium. This direction holds for every prior, by Proposition 10. If F is not convex, Proposition 20, with any $\theta \in [0, \frac{1}{2})$, gives $u(\varepsilon) > u_0$ for small $\varepsilon > 0$. □

The conclusion of Theorem 4 holds whenever (H1) holds and (H2) and (H3) hold on one chain and the revealed stretches next to it, as the proof uses them only there.

D. Extensions

This appendix relaxes the remaining assumptions. Appendices D.1 and D.2 leave the well S-shaped class: the first studies its mirror image, and the second describes equilibria around its boundary. Appendix D.3 removes the rival.

D.1 Reverse S-shaped selection rules

The well S-shaped class has a mirror image, in which the density of the selection rule is convex rather than concave. A concave density puts an agent's marginal gain from a higher signal near a tie, whereas a convex density puts it at large margins.

Definition 7. A function $v : [-1, 1] \to [0, 1]$ is called *well reverse S-shaped* if $v(x) - \frac{1}{2}$ is odd and increasing, and $v'(x)$ is convex on $[-1, 1]$ and uniquely minimized at $x = 0$.

The following lemma transfers the equilibrium analysis to this class.

Lemma 32. *(i) For every well reverse S-shaped $v_2(x)$, there is a well S-shaped $v_1(x)$ with $\tilde{v}_2(x) = -\tilde{v}_1(x)$.*

(ii) Let $\tilde{v}_2(x) = -\tilde{v}_1(x)$. If $(G_1(s_1), G_2(s_2))$ is an equilibrium for $v_1(x)$ and priors (p_1, p_2), then $(\check{G}_1, \check{G}_2)$ is an equilibrium for $v_2(x)$ and priors $(1 - p_1, 1 - p_2)$, where $\check{G}_i$ is the distribution of $1 - s_i$ for $s_i \sim G_i$.

Hence, for equal priors, the unique equilibrium of a well reverse S-shaped $v(x)$ is $(\{b, 1\}; \{b, 1\})$, where b increases in p analogously to Theorem 1: each agent reveals good news rather than conceding bad news, and full revelation is reached for $p \leq \frac{1}{\mathcal{E}_{\tilde{v}}(1)}$, mirroring Proposition 1. In particular, such rules can deliver the first best for $p \leq \frac{1}{2}$: on $[0, 1]$, $v(x) = \frac{1}{2} + (\frac{1}{2} - \eta)x + \eta x|x|$ with $\eta \in (0, \frac{1}{2}]$ equals $\frac{1}{2} + \frac{1}{2}x - \eta x(1 - x)$, so it is revelation-accurate and strictly negatively balanced (Proposition 7).

The guarantee of Proposition 4, however, fails for many such rules.

Example 10. For $v(x) = \frac{1}{2} + \frac{1}{2}x^3$, $u_F(p)$ is non-monotone in p.

For these rules, informativeness often decreases in p unless the elasticity increases sharply, mirroring Lemma 8 and Corollary 4, and since $v(x)$ is convex on $[0, 1]$, this effect can outweigh the increase in the prior. Equilibria for unequal priors are flipped too: under a well reverse S-shaped rule with $v'(x)$ not linear near 0, the agent with the high prior reveals fully and the other pools at his prior, the opposite of the pattern in Subsection 3.2.3.

D.2 Around the boundary of the well S-shaped class

This appendix describes what changes as the selection rule leaves the well S-shaped class: under a weaker condition, the equilibrium keeps the forms of Subsection 3.2.3, whereas further away equilibrium signals may have non-binary support.

D.2.1 Binary support equilibrium

Although $\overline{v}(s_i)$ need not stay S-shaped outside the class, the equilibrium forms of Lemma 1 and Proposition 2 survive outside it. Their proofs use the shape of $v(x)$ only through the shape of agent i's value function against agent j's equilibrium signals, $\{p_j\}$ and $\{0, c\}$. Against $\{0, c\}$, the sign of $\overline{v}''(s_i)$ on $(0, c)$ depends on $w(s_i)$ and $w(c - s_i)$, where $w(x) := -v''(x)$, so only pairs of points with sum at most 1 matter.

Proposition 21. *Let $v(x)$ be a selection rule with $v'(x)$ differentiable on $(0, 1]$, $w(x) > 0$ on $(0, 1]$, and $\frac{w(c-x)}{w(x)}$ strictly decreasing in x on $(0, c)$ for every $c \in (0, 1]$. Then for all priors, the equilibrium is unique, and each agent i plays $\{p_i\}$ or $\{0, a_i\}$ with $a_i \in (p_i, 1]$.*

These conditions hold for every well S-shaped $v(x)$ with strictly concave and differentiable $v'(x)$, but they do not require $v'(x)$ to be concave. Unlike well S-shapedness, however, they are not straightforward to verify, since the monotonicity of $\frac{w(c-x)}{w(x)}$ must hold for every $c \in (0, 1]$. For $\xi \sim N(0, \sigma^2)$, they hold if and only if $\sigma \geq \frac{1}{2}$, a quarter of the variance that well S-shapedness requires ($\sigma \geq 1$). This bound is nearly tight: in simulations, for $\sigma = 0.46$ and $(p_1, p_2) = (0.62, 0.48)$, agent 1 sends three signals in every equilibrium.

D.2.2 Non-binary support equilibrium

Outside the well S-shaped class, equilibrium signals need not have binary support, and their structure varies considerably with the selection rule.

Lemma 33. *Let $v(x)$ be real-analytic on $[-1,1]$ with $v''(x) > 0$ for all $x \in [-1,0)$. Then for all $p_i, p_j \in (0,1)$ and $G_j \in \mathcal{G}(p_j)$, every optimal signal for $(\overline{v}(s_i, G_j), p_i)$ has finite support.*

These conditions hold for the selection rules generated by i.i.d. Gumbel or normal shocks of any variance, so for these natural examples, equilibrium signals are discrete. By the identity theorem, an analytic $\overline{v}(s_i)$ coincides with a line either on a finite set or on all of $[0,1]$, and the latter is ruled out by the convexity of $v(x)$ on $[-1,0]$. Still, simulations[8] show that the number of support points grows as the noise decreases, mirroring the classical result for games on the unit square with bell-shaped kernels (see Karlin (1959)), in which optimal strategies are finitely supported and the number of support points grows without bound as the kernel sharpens.

In contrast, a selection rule that is not analytic can induce an equilibrium signal with a continuous part, already for exponential shocks.

Example 11. Let $v(x)$ be generated by i.i.d. $\varepsilon_i \sim Exp(\lambda)$, i.e. the Laplace CDF $v(x) = \frac{1}{2} + \frac{1}{2} \cdot \text{sign}(x) \left(1 - e^{-\lambda|x|}\right)$. Then for all $p \leq \frac{1}{2}$, both agents playing

$$G = \frac{1}{2(1+\lambda p)} \left(\delta_{\{0\}} + \delta_{\{2p\}}\right) + \frac{\lambda p}{1+\lambda p} U[0, 2p]$$

is a symmetric equilibrium.

In general, such equilibria are hard to find explicitly. If an equilibrium signal of agent j has an interval $[a,b]$ in its support, $\overline{v}(s_j)$ must be linear on $[a,b]$, which is an integral equation for G_i^*:

$$\overline{v}'(s_j) = \int_0^1 v'(s_j - s_i) dG_i^*(s_i) = const \quad \forall s_j \in [a,b].$$

Closed-form solutions are known only for special kernels,[9] such as those with a rational

[8] For $p = 0.3$ and $\xi \sim N(0, \sigma^2)$, the equilibrium G_i^* has 5 support points for $\sigma = 0.1$ and 9 for $\sigma = 0.05$.

[9] This is a Fredholm equation of the first kind with a difference kernel on a finite interval; see

Fourier transform, as in Example 11, or piecewise polynomial ones, which include the triangular selection rules of Appendix B.3.

D.3 A single agent

Without a rival, the firm compares the agent with an outside option of known value $c \in (p, 1)$: she hires him if his score exceeds c, i.e., with probability $a(s) := \mathbb{P}(s + \varepsilon \geq c)$, where ε has a continuous CDF F with $F(0) = \frac{1}{2}$. Her gain over not hiring is $W := \mathbb{E}_G[a(s)(s - c)]$, and her first best is $p(1 - c)$. The agent maximizes $\mathbb{E}_G[a(s)]$ over $G \in \mathcal{G}(p)$.

Proposition 22. *Let $c \in (p, 1)$.*

(i) Without noise, $W = 0$. With noise, $W \leq p \min(\frac{1}{2}, 1 - c)$.

(ii) If $\varepsilon \sim U[-u, u]$ with $u < c$ and $c + u \leq 1$, the agent's unique best reply is $\{0, c + u\}$, and $W = \frac{pu}{c+u}$. In particular, if $c > \frac{1}{2}$ and $u = 1 - c$, he reveals fully, and the firm attains her first best.

(iii) The firm can attain her first best only if $c \geq \frac{1}{2}$.

Without noise, the agent splits the prior into $\{0, c\}$, and the firm hires him exactly when she is indifferent, so she gains nothing from the evidence. Noise devalues a narrow lead over c, as it devalues a narrow lead over a rival in the main model, so the agent must prove more. The bound in (i) holds because the agent can always split into $\{0, c\}$, which wins with probability at least $\frac{p}{c}a(c) = \frac{p}{2c}$: at the outside option, the noise makes hiring a coin flip. The firm thus hires with probability at least $\frac{p}{2c}$, each hire costs her c, and she can hire at most the mass p of good types. By (ii), uniform noise of half-width $\min(c, 1 - c)$ attains this bound, strictly for $c > \frac{1}{2}$ and in the limit for $c \leq \frac{1}{2}$: it is the widest noise that never hires a revealed bad agent and never rejects a revealed

Sakhnovich (2015) for the general theory and Polyanin and Manzhirov (2008) for solvable kernels.

good one. As after Proposition 1, a tie is a coin flip, and this is why full revelation requires $c \geq \frac{1}{2}$.[10] A privately known threshold can likewise force a single sender to reveal fully (Kolotilin et al., 2017); here, the threshold is random to both sides, and the noise is unbiased.

More generally, if the density of ε is unimodal with its peak at 0 and is not constant on $[0, c]$, then $a(s)$ is convex on $[0, c]$ and concave on $[c, 1]$, and the agent splits the prior into $\{0, y\}$ with $y > c$. The firm then gains if and only if p exceeds

$$\hat{p} := \frac{y\, c\, (1 - F(c))}{a(y)(y - c) + c\, (1 - F(c))}$$

Here $1 - F(c)$ is the probability that the noise hires a revealed bad agent, so $\hat{p} = 0$ if $\varepsilon \leq c$, and for normal noise, $\hat{p}$ is close to 0 for small variances and rises with the variance. Two differences from the main model stand out. Without a rival, the noiseless firm learns nothing useful, whereas with two agents, she already learns much (Boleslavsky and Cotton, 2015). And a single agent's threshold is fixed, so a unimodal density suffices for bad news signal, whereas a rival's randomization makes the threshold random, and the well S-shaped class requires a concave density. The comparison also shows the role of commitment: a firm facing a single agent that could commit to any rule would hire only on conclusive evidence and attain her first best without noise. With binary types, a firm comparing two agents can do the same with a pass/fail rule (Proposition 8); with a continuum of types, she cannot, and her best rule blurs the signals (Theorem 3).

[10] The bound uses only $a(c) \geq \frac{1}{2}$, so it holds for any noise with median 0. With mean-zero noise instead, rare shocks far beyond the posterior scale can move the mean without affecting decisions, and the firm can approach her first best for every c.

D.4 Proofs

D.4.1 Proof of Lemma 32

(i) Consider $v_1(x) = \frac{1}{2} + \frac{1}{2}x - a\tilde{v}_2(x)$. The convexity of $\tilde{v}_2'(x)$ and the uniqueness of its minimum at 0 are simply inverted into the concavity of $v_1'(x)$ and the uniqueness of its maximum at 0, and $v_1(x) - \frac{1}{2}$ is trivially odd and increasing for sufficiently small a, since $\tilde{v}_2'(x)$ is bounded. So $v_1(x)$ is well S-shaped for such a, while

$$\tilde{v}_1(x) = \frac{\frac{1}{2}x - a\tilde{v}_2(x) - x(\frac{1}{2} - a\tilde{v}_2'(0))}{|\frac{1}{2} - a - (\frac{1}{2} - a\tilde{v}_2'(0))|} = -\tilde{v}_2(x).$$

(ii) Since a normalization changes payoffs by a positive factor and terms independent of the signals, $v_k(x)$ and $\tilde{v}_k(x)$ have the same equilibria, $k \in \{1, 2\}$. The map $G \mapsto \check{G}$, where $\check{G}$ is the distribution of $1 - s$ for $s \sim G$, is a bijection between $\mathcal{G}(p)$ and $\mathcal{G}(1-p)$. Since $\tilde{v}_1(x)$ is odd, $\tilde{v}_2\big((1 - s_i) - (1 - s_j)\big) = -\tilde{v}_1(s_j - s_i) = \tilde{v}_1(s_i - s_j)$, so each agent's payoff from $(\check{G}_1', \check{G}_2')$ under $\tilde{v}_2$ equals that from (G_1', G_2') under $\tilde{v}_1$ for all G_1', G_2'. Hence $(\check{G}_1, \check{G}_2)$ is an equilibrium for $\tilde{v}_2$ and priors $(1 - p_1, 1 - p_2)$.

D.4.2 Proof of Proposition 21

Let $G_j = \{0, c\}$ with $c \in (p_j, 1]$, so that $\mathbb{P}(s_j = c) = q := \frac{p_j}{c} \in (0, 1)$. Since $v''(x)$ is odd, $\overline{v}''(s) = (1-q)v''(s) + q\, v''(s-c) = -(1-q)w(s) - q\, w(s-c) < 0$ on $(c, 1]$, and on $(0, c)$,

$$\overline{v}''(s) = -(1-q)w(s) + q\, w(c-s) = w(s)\left(q\,\frac{w(c-s)}{w(s)} - (1-q)\right),$$

which, as the ratio is strictly decreasing, is first positive and then negative, with at most one zero. If $G_j = \{p_j\}$, then $\overline{v}''(s) = v''(s - p_j)$ is positive on $[0, p_j)$ and negative on $(p_j, 1]$. In both cases, there is $c^* \in [0, 1]$ such that $\overline{v}$ is strictly convex on $[0, c^*]$ and strictly concave on $[c^*, 1]$, so, as in the proof of Lemma 1, agent i's best reply is unique

and is either $\{p_i\}$ or $\{0, a_i\}$ with $a_i \in (p_i, 1]$.

Now restrict each agent i to these signals, indexed by $a_i \in [p_i, 1]$, where $a_i = p_i$ stands for $\{p_i\}$. Payoffs are continuous in (a_1, a_2), and, by the above, the best reply within this family is unique and is also a best reply among all signals. By the maximum theorem, each agent's best reply is continuous in the other agent's index, so Brouwer's theorem gives a pair of mutual best replies, which is an equilibrium. Since all equilibria form a product set and each agent's best reply to the other's equilibrium signal is unique, this equilibrium is unique.

D.4.3 Proof of Lemma 33

Since $v(x)$ is real-analytic on $[-1, 1]$, it extends analytically to $\{z \in \mathbb{C} : dist(z, [-1, 1]) < \delta\}$ for some $\delta > 0$, hence $\overline{v}(s_i) = \int_0^1 v(s_i - s_j) dG_j(s_j)$ extends analytically to $\{z \in \mathbb{C} : dist(z, [0, 1]) < \delta\}$. Let $l(s_i)$ be a line supporting $\overline{v}^{cv}(s_i)$ at p_i. Then $l(s_i) \geq \overline{v}(s_i)$ on $[0, 1]$, and every optimal signal is supported on $\{s_i \in [0, 1] : \overline{v}(s_i) = l(s_i)\}$. If this set were infinite, it would have an accumulation point in $[0, 1]$, and by the identity theorem $\overline{v}(s_i) = l(s_i)$ on $[0, 1]$, so $\overline{v}''(0) = 0$. However, $\overline{v}''(0) = \int_0^1 v''(-s_j) dG_j(s_j) > 0$, since $v''(-s_j) > 0$ for $s_j \in (0, 1]$ and G_j puts positive mass on $(0, 1]$ as $p_j > 0$.

D.4.4 Proof of Proposition 22

(i) Without noise, $a(s) = \mathbb{1}_{s \geq c}$ lies below $\frac{s}{c}$ on $[0, 1]$, touching it only at 0 and c, so the agent's unique best reply is $\{0, c\}$, and $W = \frac{p}{c}(c - c) = 0$. With noise, the split $\{0, c\}$ is feasible since $p < c$, and it gives the agent $(1 - \frac{p}{c})a(0) + \frac{p}{c}a(c) \geq \frac{p}{2c}$, as $a(c) = 1 - F(0) = \frac{1}{2}$ by continuity. Hence $\mathbb{E}_G[a(s)] \geq \frac{p}{2c}$ in equilibrium, and since $a \leq 1$ and $s \geq 0$,

$$W = \mathbb{E}_G[a(s)s] - c\,\mathbb{E}_G[a(s)] \leq p - c \cdot \frac{p}{2c} = \frac{p}{2}.$$

Also $W \leq \mathbb{E}_G[(s - c)^+] \leq p(1 - c)$, since $(s - c)^+ \leq s(1 - c)$ on $[0, 1]$.

(ii) Here $a(s) = 0$ on $[0, c-u]$, $a(s) = \frac{s-c+u}{2u}$ on $[c-u, c+u]$, and $a(s) = 1$ on $[c+u, 1]$. Let $l(s) := \frac{s}{c+u}$. Then $l > a$ on $(0, c-u]$, $l - a$ is affine on $[c-u, c+u]$, positive at $c-u$ and zero at $c+u$, and $l > a$ on $(c+u, 1]$. So l supports the concavification of a and touches a only at 0 and $c+u$, and by Kamenica and Gentzkow (2011, Corollary 2), the agent's unique best reply is $\{0, c+u\}$, with $\mathbb{P}(s = c+u) = \frac{p}{c+u}$. As $a(0) = 0$ and $a(c+u) = 1$, $W = \frac{p}{c+u}(c+u-c) = \frac{pu}{c+u}$. If $c > \frac{1}{2}$ and $u = 1-c$, then $u < c$, $c+u = 1$, and $W = p(1-c)$.

(iii) Since $a \leq 1$ and $(s-c)^+ \leq s(1-c)$ with equality only at $s \in \{0, 1\}$, $W = p(1-c)$ requires $G = \{0, 1\}$, $a(0) = 0$ and $a(1) = 1$. Full revelation is a best reply only if a lies below its chord from $(0, 0)$ to $(1, 1)$, i.e., $a(s) \leq s$ on $[0, 1]$, and at $s = c$ this gives $\frac{1}{2} \leq c$.