

Competitive Persuasion with a Noisy Judge

Igor Karpov*

October 8, 2026

Abstract

Can a firm choosing between two agents benefit from judging their evidence with error, even though on given evidence error can only hurt her? I show that she can when the agents design their evidence themselves: suitably shaped noise weakens each agent's incentive to conceal, because a narrow lead no longer guarantees selection. For a broad class of noise distributions and equal priors, the equilibrium is unique, with each agent conceding that he is bad with some probability and otherwise sending one pooled signal. Within this class, the firm's utility rises with the agents' ex-ante strength, though they may disclose less, and, with binary types, the best judge outperforms a perfectly accurate one unless agents are very likely bad and attains the first best once they are more likely good than bad. Among rules depending on the signal gap, the only one that can attain the first best for every prior leaves agents indifferent among all disclosures, whereas knowing only whether the pool is strong or weak allows a unique full disclosure equilibrium, though not under i.i.d. noise for weak pools.

Keywords: flexible information disclosure, competition, information design

JEL Classification: C72, D82, D83

*University of Chicago. I am very grateful to Doron Ravid, Ben Brooks and Emir Kamenica for their guidance and support. I also thank Alex Frankel, Joe Root, Phil Reny and the student workshop audience at the University of Chicago for useful comments and discussion. All errors are my own.

1. INTRODUCTION

A capital committee must fund one of two projects this quarter. Each sponsor decides how rigorous a pilot to run before the meeting, knowing that the committee's verdict will turn partly on things that have nothing to do with the projects: who presents second, how long the meeting has run. The committee would like to see past all of that. This paper asks whether it should want to.

Such imperfect verdicts, biased or merely arbitrary, are well recognized. In Scharfstein and Stein (2000), divisional rent-seeking leads to “socialist” internal capital markets, in which weaker divisions are subsidized by stronger ones. In Prendergast and Topel (1996), evaluators who lack objective performance measures act on personal preferences, favoring some subordinates beyond their true performance. And Pier et al. (2018) find essentially no agreement among experienced reviewers rating the same NIH grant applications, all of which were eventually funded: ratings of the same application were no more similar than ratings of different ones. The usual response is to engineer such imperfections away, through more reviewers, more structured evaluations, and more objective criteria. A few research funders have nonetheless moved the other way: they now fund by lot among proposals that clear a quality bar, citing reviewer bias, reviewer burden, and the hope of bolder proposals (Fang and Casadevall, 2016; Heyard et al., 2022; Röbbcke and Simon, 2023). Only the hope of bolder proposals concerns how applicants respond, and none concerns what the competitors choose to reveal about themselves, the channel I study.

The question is whether a decision-maker would ever want to make her evaluation noisier rather than sharper once the competitors adjust what they reveal to it. It arises whenever three features come together. First, a single prize is at stake, with no price attached: the decision-maker cannot pay more for the stronger competitor, and a close second earns nothing. Second, the competitors decide what and how much to reveal

about themselves after learning how the decision-maker will choose between them. Third, the decision-maker can make the verdict more or less accurate, so the competitor with the stronger evidence need not always win. I call the decision-maker *the firm* and the competitors *agents*. Each agent is either a good or a bad type and simultaneously commits, at no cost, to a disclosure policy, that is, to a distribution of the posterior probability that he is good; the firm observes the resulting posterior with a random error and selects the agent with the higher score. For normal, Gumbel or uniform noise and equal priors, this is also what she would do on seeing only the scores, so committing to the rule costs her nothing there (Section 2).

My key observation is that noise of a suitable shape weakens each agent's incentive to conceal. An agent's chance of winning depends on how far he looks ahead of his rival, in the same way for both agents; I call this dependence the firm's selection rule, and choosing the noise and choosing the selection rule are one question, since every selection rule is generated by some noise (Section 2). A perfectly accurate firm always selects the agent who looks better, so her rule is a step: a narrow lead is as good as a wide one, and an agent revealed as bad loses unless his rival is too. Each agent therefore spreads his posterior to land just above his rival's, blending his bad type with his good one: Boleslavsky and Cotton (2015) show that both agents disclose only partially, randomizing over a continuum of posteriors, and the firm obtains less than 80% of her first best whenever agents are good with probability below one third.

With noise, the rule instead rises gradually with the lead. A narrow lead is worth little, conceding bad news no longer means certain defeat, and the shape of the noise decides whether an agent gains more from gambling on being revealed as good or from pooling his evidence, and so how much information reaches the firm. The price is accuracy: a rule near one half is close to a coin flip, so the firm passes over the agent who looks better more often, and noise trades selection accuracy for information.

An example. Suppose that each agent is good with probability 0.6. Under the noiseless rule, each agent's posterior equals 1 with probability $1/3$ and is otherwise uniformly distributed on $[0, 0.8]$, so the firm sometimes selects a bad agent over a good one and obtains about 94% of her first best. Now let the firm score each agent by his signal plus an independent draw from $U[0, 1]$. Against a rival who reveals his type and is more likely good than bad, an agent's chance of winning rises faster the stronger his own signal already is, so what he gains when revealed as good outweighs what he loses when revealed as bad: revealing his type is his unique best reply, and, since the agents' game is constant-sum, full disclosure by both agents is the unique equilibrium. Since a revealed good type always outscores a revealed bad one, the noise only decides between agents of the same type, and the firm attains her first best. The noise is large enough to make concealment unprofitable, but never large enough to overturn the difference between a good and a bad agent.

Main results. The first set of results, in Section 3, characterizes the equilibrium when the difference between the two evaluation errors has a concave density with a single peak over possible signal gaps, which I call a *well S-shaped* selection rule; normal and logistic noise qualify if its variance is large enough. Under such a rule, whatever his rival discloses, an agent's chance of winning accelerates, then slows as his own signal rises, so below some signal he gains from gambling and above it from playing safe; concavity is what preserves this shape when the rival randomizes. Each agent thus uses a *bad news* disclosure: he reveals the bad type with some probability and otherwise sends one pooled signal, with full and no disclosure as edge cases.

The equilibrium is unique whenever the priors are equal, and for every pair of priors unless the density is linear next to its peak. In the symmetric case, a single statistic governs it, the elasticity of the normalized selection rule at the pooled signal: an agent concedes bad news with unconditional probability equal to its inverse, and

both agents disclose fully once the prior is high enough. With asymmetric priors (Section 6), each agent discloses more when his rival is stronger, and when one agent is much stronger, the favorite reveals nothing while the underdog discloses fully.

The second set of results, in Section 4, concerns the firm's utility with equal priors. Stronger agents may disclose less, since an agent's own strength can outweigh his rival's, yet under every well S-shaped rule the firm is better off facing them; when the density of the noise is convex instead (Subsection 7.2), agents reveal good news, and stronger agents can hurt the firm. Within the class, the best rule for the firm is generated by i.i.d. uniform shocks and sits exactly at its edge. It beats the noiseless benchmark once agents are good with probability above roughly 21%, and, with binary types, delivers the first best once they are more likely good than bad; a continuum of types changes this (Subsection 7.1). At low priors, wide noise barely separates the pooled signal from the revealed one: as the prior vanishes, the rule attains half of the first best, against two thirds for the noiseless benchmark.

More noise is not always better, though. Scaling the uniform shocks traces out the trade-off (Subsection 4.3): more noise leaves disclosure unchanged and only blurs selection, while less noise lowers her utility and pushes the equilibrium toward the noiseless one once agents stop disclosing two signals. The firm thus does best at the least noise that keeps disclosures binary, which lies at or beyond the edge of the class, so a moderately noisy judge beats both a perfectly sharp and an excessively blurry one.

The third set of results, in Section 5, asks when the firm can attain her first best with any rule. With binary types and equal priors, she needs full disclosure and a rule that always picks a revealed good agent over a revealed bad one. If she cannot tell whether the pool is strong or weak – whether both priors lie above or below one half – only one rule depending on the gap between signals, as noisy evaluation does, can deliver both, and it is linear, which leaves agents indifferent among all disclosures: robustness

to the prior costs uniqueness. If she knows which, a rich family of rules makes full disclosure the unique equilibrium. With i.i.d. shocks, this fails for a weak pool: full disclosure is never an equilibrium once an agent is more likely bad than good.

Section 7 collects extensions. Beyond the well S-shaped class, the equilibrium stays unique, with at most two signals per agent, under a weaker condition that normal noise meets down to a quarter of the variance that well S-shapedness requires (Subsection 7.3); with less noise, agents randomize over more signals. With a continuum of types (Subsection 7.1), each agent still reveals his type below a threshold and pools all types above it, but under a convex prior CDF, the noiseless rule attains the first best, which no noisy rule can, and under a concave one, the ranking depends on how dispersed the pool is. Finally, a head start for one agent (Subsection 7.5) keeps the evaluation accurate but breaks anonymity, induces both agents to concede bad news, and outperforms every well S-shaped rule for sufficiently low priors.

1.1 Literature review

This paper contributes to the Bayesian persuasion literature (Kamenica and Gentzkow, 2011; Kamenica, 2019), in particular to its strand on competing senders. Most results of Gentzkow and Kamenica (2017) require a Blackwell-connected information environment, which fails when each sender can only disclose information about himself, as here, and their comparison with collusion is vacuous, since the agents' game is constant-sum. Each agent's best reply solves a single-sender problem with an S-shaped value function, so it is an upper censorship as characterized by Kolotilin et al. (2022), whose result also gives the equilibrium under a general prior (Subsection 7.1.1).

The closest paper is Boleslavsky and Cotton (2015), part of the literature on grading policies (Ostrovsky and Schwarz, 2010; Chan et al., 2007; Popov and Bernhardt, 2013; Bar et al., 2012): two schools design grades, and an evaluator hires the graduate with

the higher posterior. Under this noiseless rule, the equilibrium is unique, posteriors are uniform on an interval, possibly with atoms at 0 and 1, and neither school ever discloses fully. Disclosure increases with the number of senders, both in this setting, where it becomes full in the limit (Au and Kawai, 2020), and when senders also compete in prices (Hwang et al., 2026). In search markets, whether competition raises informativeness depends on what sellers observe and how buyers search (Board and Lu, 2018; Au and Whitmeyer, 2023; He and Li, 2023).

An imperfect evaluation can also benefit the evaluator by strengthening investment incentives, as strategic grading does in Boleslavsky and Cotton (2015) and a noisy signal of universities' relative quality does in Boleslavsky and Li (2026). Here the gain comes through information instead: a suitably noisy evaluation makes disclosures more informative. Closer in spirit, an organization that can fund only one of two proposals induces more informative evidence than an unconstrained one and is better off unless both are ex-ante strong (Boleslavsky and Cotton, 2018), and moderate attention costs can make a rationally inattentive consumer select the better of two disclosed products more often than when attention is free (Jain and Whitmeyer, 2026). In both, the gain comes from a constraint on the receiver; I instead treat the evaluation error as a primitive of the selection rule, whose shape determines the equilibrium and how much information the firm extracts.

My normative results also relate to the idea that a receiver can extract more information by committing to use it less, through random payoff adjustments (Tsakas et al., 2021) or, when agents manipulate their data at a cost, through a less responsive allocation rule, a test that passes some noncompliant agents, or a score that underweights some features (Frankel and Kartik, 2022; Perez-Richet and Skreta, 2022; Ball, 2025). Extra noise in an already noisy channel can benefit the sender and sometimes both players (Tsakas and Tsakas, 2021), and coarse grades can inform the public better by

inducing participation (Harbaugh and Rasmusen, 2018). Here disclosure is costless and flexible, and noise works only through competition, by changing the curvature of each agent’s value function. Randomized choice also serves other purposes: breaking ties in school choice (Abdulkadiroğlu and Sönmez, 2003; Erdil and Ergin, 2008), saving on costly signaling (Hoppe et al., 2009), economizing on verification (Border and Sobel, 1987), deterring gaming (Ederer et al., 2018), and keeping improper reasons out of selection (Stone, 2011; Fang and Casadevall, 2016).

The agents’ game is also a contest in which each agent chooses a distribution with a fixed mean, and noise gives it a difference-form success function (Lazear and Rosen, 1981; Che and Gale, 2000). When contestants choose the riskiness of their performance, limiting competition helps a principal select the better one (Hvide and Kristiansen, 2003; Fang and Noe, 2022), and a noisy selection rule is a new instrument with the same effect. In effort contests, an imperfectly accurate contest can also be optimal (Wang, 2010; Morgan et al., 2022; Letina et al., 2023), and the shape of the noise matters (Drugov and Ryvkin, 2020); see Fu and Wu (2019) for a survey.

2. MODEL

Agents. There are 2 agents and the firm (from here on she). Each agent’s type θ_i is drawn independently from $\{0, 1\}$, and let p_i be the probability of $\theta_i = 1$. From here on, I refer to p_i as a *prior*. For notational simplicity, if $p_1 = p_2$, I refer to them simply as p .

Information and Timing. Agents play a simultaneous-move game in which, before the realization of his type, each agent i commits to a signal structure. Then, after the realization of each agent’s type θ_i , the firm receives a signal s_i according to that structure. Since the prior is binary, I can re-identify the reported signal with its posterior mean $s_i := \mathbb{E}[\theta_i | s_i]$, and hence WLOG agent i chooses a distribution of posterior means

G_i (from here on disclosure). Let $\mathcal{G}(p_i)$ be the set of feasible disclosures for a prior p_i . Since the prior is binary, $\mathcal{G}(p_i) := \{G_i : \text{supp}(G_i) \subseteq [0, 1], \int_0^1 s_i dG_i(s_i) = p_i\}$, see Gentzkow and Kamenica (2016).

Selection. Agents select signal structures knowing the firm's *selection rule*: a function $v : [-1, 1] \mapsto [0, 1]$, such that $v(x) - 0.5$ is odd, increasing and continuous on $[-1, 1]$. If the firm observes signals s_1, s_2 , the respective probabilities of agents 1, 2 being selected are $(v(s_1 - s_2), v(s_2 - s_1))$, and since $v(x) - 0.5$ is odd, $v(s_1 - s_2) + v(s_2 - s_1) \equiv 1$ for all s_1, s_2 , so that the firm always selects exactly one agent. The oddness of $v(x) - 0.5$ also means that the selection rule treats the agents anonymously; Subsection 7.5 discusses biased rules. The selection rule is assumed to be exogenously given to the firm for most of the sections; however, I endogenize the choice of $v(x)$ in Section 5.

Noisy evaluation. Such selection rules arise naturally when the firm evaluates the agents with noise. Suppose that, instead of observing s_i perfectly, the firm observes agent i 's score $s_i + \varepsilon_i$, where ε_i is a random shock, independent of s_i, s_j , reflecting the idiosyncrasies of whoever is judging – a particular committee member's mood, the order in which proposals were read, an unmodeled personal rapport. Taking the scores at face value, the firm selects the agent with the higher score. If $v(x)$ is the CDF of a continuous $\xi := \varepsilon_j - \varepsilon_i$, restricted to $[-1, 1]$, she selects agent i with probability

$$\mathbb{P}(s_i + \varepsilon_i > s_j + \varepsilon_j) = \mathbb{P}(\xi < s_i - s_j) = v(s_i - s_j).$$

If ξ is also symmetric¹, $v(x)$ satisfies the assumptions on the selection rule. Conversely, every selection rule is generated in this way, for example by $\varepsilon_j = -\varepsilon_i = \xi/2$, where ξ is symmetric with CDF $v(x)$ on $[-1, 1]$; these shocks are exchangeable but not

¹A sufficient condition for this is the exchangeability of $\varepsilon_i, \varepsilon_j$.

independent, and Corollary 6 shows that independence is restrictive. While I usually give $v(x)$ explicitly, I sometimes describe it implicitly, saying that $v(x)$ is *generated by* a distribution of $(\varepsilon_i, \varepsilon_j)$ that satisfies these assumptions.

Example 1. If $v(x)$ is generated by ε_i i.i.d. Gumbel, normal or Cauchy, then $v(x)$ is the CDF of a logistic, normal or Cauchy distribution, respectively.

Payoffs. WLOG, each agent receives 1 if selected and 0 otherwise. If the firm selects agent i , she receives θ_i , so her ex-ante expected utility equals the reported signal s_i . The shocks ε_i that may generate the selection rule do not enter the firm's utility.

Agents' Equilibrium. Since θ_i is unobservable to agent i , he should simply maximize the ex-ante probability of being selected when choosing a signal structure $G_i(s_i)$. Hence, an equilibrium of the agents' informational game is a tuple (G_1^*, G_2^*) , where

$$G_i^* \in \arg \max_{G_i \in \mathcal{G}(p_i)} \int_0^1 \int_0^1 v(s_i - s_j) dG_j^*(s_j) dG_i(s_i).$$

For any (p_1, p_2) and selection rule $v(x)$, an equilibrium of this game always exists, because the following properties are satisfied. First, the game is constant-sum, since $v(x) - 0.5$ is odd; second, payoffs are linear and continuous in G_i , since $v(x)$ is continuous; third, $\mathcal{G}(p_i)$ is a compact convex subset of a topological vector space (see Proposition 1 in Kleiner et al. (2021)). Hence, Sion's minimax theorem ensures the existence of an equilibrium. Finally, all equilibria are pairs of max-minimizer strategies for both agents and form a product set.

Interpretation. The model is deliberately abstract and says nothing about a price: agent i 's type θ_i is simply the value the firm realizes from picking i , with no wage, term sheet or other transfer attached to that choice. This fits any setting in which a decision-maker

allocates one indivisible prize between two competing options and cannot pay more to secure a stronger one. My running example is **internal capital budgeting**: a firm's capital committee (*the firm*) must fund exactly one of two project proposals this cycle, and each project's sponsor (*agent i*) controls how much hard evidence – the scope of a pilot, the rigor of a feasibility study, which numbers are led with – to place in front of the committee. The same primitives fit a **grant panel** choosing between two proposals, a **startup accelerator** deciding which of two founders gets its last slot this cycle, and a **procurement office** running a technical, price-free evaluation of two bids. They also cover grading: a university chooses a grading policy for a graduate who competes for a single place or award and is evaluated by an admissions office or a scholarship trust. I use *the firm* and *agent* as generic labels and switch between these readings wherever a concrete example helps. The binary prior simplifies the analysis and sharpens the results; Subsection 7.1 relaxes it.

Under any of these readings, the disclosure $G_i(s_i)$ is the design of the hard information about the project, applicant or student – the scope and pre-registration of a pilot study, the depth of a due-diligence file, a school's grading policy – summarized by the distribution of the posterior s_i that the agent is good. Equivalently, as in Boleslavsky and Cotton (2015), each type generates a more primitive random report – a grade, a pilot outcome, a due-diligence finding – whose distribution satisfies the monotone likelihood ratio property, so that, by Bayes' rule, the posterior increases in the report.

The selection rule $v(x)$ translates the two disclosed signals into the ex-ante probability of being picked when the firm selects the agent with the higher score. The shocks ε_i that generate $v(x)$ are imperfections of the kind documented in Section 1: idiosyncrasies of the evaluator that are uncorrelated with the substance of the disclosure and with the firm's realized payoff, but still move her decision. An agent may also have outcome-relevant characteristics that are uncorrelated with his signal

and influence the firm's decision, but I set these aside for two reasons. First, for an agent it makes no difference whether the shock affects the firm's payoff, as long as it shapes the selection rule in the same way. Second, comparing the firm's utility across selection rules is easier to interpret when the shock is pure evaluation noise.

3. WELL S-SHAPED SELECTION RULES

In this section, I study the agents' game under the additional assumption that the selection rule $v(x)$ has a concave density with a unique mode. This is sufficient for an equilibrium with binary support strategies to exist for all $p \in (0, 1)$, and for it to be unique when the priors of the agents are equal. I then study the symmetric equilibrium: how disclosures change with the priors, when full disclosure is an equilibrium, and which characteristic of $v(x)$ drives the equilibrium.

3.1 Preliminary definitions and lemmas

I begin with the key definition of this and the following sections.

Definition 1. A function $v : [-1, 1] \rightarrow [0, 1]$ is called *well S-shaped* if $v(x) - 0.5$ is odd and increasing, and $v'(x)$ is concave on $[-1, 1]$ and uniquely maximized at $x = 0$.

Since $v(x) - 0.5$ is odd, $v'(x)$ is even, so a concave $v'(x)$ is always maximized at $x = 0$. Requiring this maximum to be unique only rules out densities that are flat around 0, including the linear selection rule. Without this requirement, the equilibrium may fail to be unique: if $v(x)$ is linear on some $[-b, b]$, then for priors (p, p) with $p \leq \frac{b}{2}$, both $(\{p\}; \{p\})$ and $(\{0, 2p\}; \{0, 2p\})$ are equilibria. If $v(x)$ is generated by $\varepsilon_i, \varepsilon_j$ as in Section 2, well S-shapedness means that ξ has a concave density on $[-1, 1]$ with a unique mode, which holds for the distributions of Example 1 if their tails are wide enough.

Example 2. For each of the three classes² from Example 1, there is a threshold $\bar{V}$ s.t. $v(x)$ is well S-shaped if $\text{Var}[\varepsilon_i] \geq \bar{V}$. So is the rule generated by i.i.d. $\varepsilon_i \sim U[0, u]$ with $u \geq 1$, although its triangular density is linear on $[0, 1]$.

All results for equal priors hold on the whole well S-shaped class, which must include rules such as the one generated by $U[0, 1]$ shocks, the best one for the firm (Theorem 2). Where uniqueness must extend beyond equal binary priors, I additionally require that $v'(x)$ is *not linear near 0*, i.e., not linear on any interval $[0, \varepsilon]$, so that the density is not linear next to its mode. This does not change the shape of value functions, but makes every best reply unique (Lemma 1), which pins down the equilibrium for asymmetric priors (Proposition 6) and for non-binary priors (Proposition 9); without it, asymmetric priors can support multiple equilibria (Remarks 2, 4).

Let $\bar{v}(s_i, G_j) := \int_0^1 v(s_i - s_j) dG_j(s_j)$, often written $\bar{v}(s_i)$, be the *value function* of agent i . By choosing a disclosure G_i , agent i solves a Bayesian persuasion problem as in Kamenica and Gentzkow (2011):

$$\max_{G_i \in \mathcal{G}(p_i)} \int_0^1 \bar{v}(s_i, G_j) dG_i(s_i) \quad (\text{BP})$$

I call G_i an *optimal disclosure* for $(\bar{v}(s_i), p_i)$ if it solves (BP), and *the optimal disclosure* if, in addition, no other G_i solves it.

To find an optimal disclosure, I use the *concavification* $\bar{v}^{cv}(s_i)$ of $\bar{v}(s_i)$ for a fixed G_j , the pointwise lowest concave function majorizing $\bar{v}(s_i)$. Then G_i is an optimal disclosure for $\bar{v}(s_i)$ if and only if it is optimal for $\bar{v}^{cv}(s_i)$ and its support is contained in the *concavification support* $\text{supp}(\bar{v}^{cv}) := \{s_i : \bar{v}(s_i) = \bar{v}^{cv}(s_i)\}$. For the concave $\bar{v}^{cv}(s_i)$, this is simple: if it is linear on some interval around p_i , any $G_i \in \mathcal{G}(p_i)$ supported in that interval is optimal; otherwise, the optimal disclosure has $\text{supp}(G_i) = \{p_i\}$ (no disclosure), see Kamenica and Gentzkow (2011).

²Cauchy shocks have no finite variance; for them, $\text{Var}[\varepsilon_i]$ is the squared scale parameter throughout.

A well S-shaped $v(x)$ helps because if $v'(x)$ is concave on $[-1, 1]$, then $\bar{v}'(s_i)$ is concave on $[0, 1]$ for every G_j , which gives $\bar{v}(s_i)$ the following shape.

Definition 2. $\bar{v}(s)$ is called *S-shaped* on $[a, b]$, if there exists some $c \in [a, b]$ s.t. $\bar{v}(s)$ is convex on $[a, c]$ and concave on $[c, b]$.

This is the standard definition of an S-shaped curve. As $v(x)$ is continuously differentiable, so is $\bar{v}(s_i)$, and then $\bar{v}(s_i)$ is S-shaped if and only if $\bar{v}'(s_i)$ is quasi-concave. Since $\bar{v}'(s_i)$ is even concave, $\bar{v}(s_i)$ is S-shaped for every well S-shaped $v(x)$ and every G_j . As quasi-concave functions are not closed under addition, there is no simple extension of the well S-shaped class for which $\bar{v}(s_i)$ is always S-shaped. Characterizing the equilibrium, however, needs this shape only against the rival's equilibrium disclosures. Using this, Proposition 12 goes beyond the class: for all priors, the equilibrium stays unique, with at most two signals per agent, and for normal noise this holds down to a quarter of the variance threshold in Example 2.


So under a well S-shaped $v(x)$, an agent's marginal gain from a higher signal (posterior mean) is concave, and against any disclosure G_j , $\bar{v}(s_i)$ has at most two regions of different convexity. Hence its concavification consists of at most two parts, which strongly restricts the equilibrium disclosures, characterized in Lemma 1 (ii).

Lemma 1. *If $v(x)$ is well S-shaped, then*

- (i) $\bar{v}(s_i, G_j)$ is S-shaped on $[0, 1]$ for all $p_j \in (0, 1)$ and $G_j \in \mathcal{G}(p_j)$.
- (ii) *An optimal disclosure for $(\bar{v}(s_i), p_i)$ is $\{0, a\}$ if $p_i < a$ and $\{p_i\}$ otherwise, where $a = \max \arg \max_{s_i \in [0, 1]} slp(s_i)$, $slp(s_i) := \frac{\bar{v}(s_i) - \bar{v}(0)}{s_i}$ and $slp(0) := \bar{v}'(0)$. If $v'(x)$ is not linear near 0, it is the optimal disclosure.*

Remark 1. I identify a disclosure G_i with its support whenever $|supp(G_i)| \leq 2$.

Figure 1: Illustration of Lemma 1 (ii)



This is not the image of $\bar{v}(s_i)$, but of $\bar{v}(s_i) - bs_i$ for some $b > 0$ for clarity. Since $\text{cav}(f(x) + bx) = \text{cav}(f(x)) + bx$ for all x, b , this does not influence the optimal disclosure.

Lemma 1 thus restricts an agent's optimal disclosure to a very limited class, the *bad news* disclosures: the agent reveals the low type with some probability and otherwise sends a pooled signal, with full ($\{0, 1\}$) and no ($\{p_i\}$) disclosure as the edge cases. Binary support itself is not special: two-point disclosures are the pure strategies of an agent, every other disclosure is their mixture, and against any continuous value function some disclosure with at most two support points is a best reply. What a well S-shaped $v(x)$ adds is that, unless no disclosure is optimal, such a best reply has 0 in its support, and that it is unique if $v'(x)$ is not linear near 0. Moreover, the point a depends only on the rival's disclosure, and the agent's own prior only determines whether he pools at p_i or plays $\{0, a\}$.

The first claim was shown above. For the second, the concavification of an S-shaped $\bar{v}(s_i)$ coincides with the secant from $\bar{v}(0)$ to $\bar{v}(a)$ on $[0, a]$ and with $\bar{v}(s_i)$ on $[a, 1]$. Hence $\{0, a\}$ is an optimal disclosure if $a > p_i$, and $\{p_i\}$ if $a \leq p_i$, and if $v'(x)$ is not linear near 0, these are **the** optimal disclosures. Figure 1 illustrates this claim.

By the Bayesian constraint, positive scaling of $v(x)$ and adding a linear function to it

do not change the concavification support, and hence do not change equilibria for any (p_1, p_2) . I represent each such equivalence class of selection rules by the normalization

$$\tilde{v}(x) := \frac{v(x) - 0.5 - x \cdot v'(0)}{|v(1) - 0.5 - v'(0)|}$$

It satisfies $\tilde{v}(0) = \tilde{v}'(0) = 0$ and $\tilde{v}(1) = -1$, which simplifies the exposition, and it is well defined for every well S-shaped $v(x)$: as $v'(x)$ is uniquely maximized at $x = 0$, $v(1) - 0.5 = \int_0^1 v'(x)dx < v'(0)$.

The key characteristic of $\tilde{v}(x)$ is its elasticity $\mathcal{E}_{\tilde{v}}(x) = \frac{x\tilde{v}'(x)}{\tilde{v}(x)}$. Since $\tilde{v}(0) = 0$, $\mathcal{E}_{\tilde{v}}(a)$ is the ratio of the slopes of the tangent to $\tilde{v}(x)$ at $x = a$ and of the secant from 0 to a , so it measures how curved $\tilde{v}(x)$ is on $[0, a]$. Hence it governs the curvature of $\bar{v}(s_i)$ and the location of the support point a from Lemma 1, as the results below show.

3.2 Main equilibrium results

3.2.1 Full disclosure

A natural first question is whether the firm can ever get the first best information, full disclosure by both agents, in equilibrium. Under a well S-shaped $v(x)$, the answer is simple: both priors must be high enough.

Definition 3. For a given well S-shaped $v(x)$, let $\bar{p} := 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)} = 1 - \frac{1}{|\tilde{v}'(1)|}$.

Proposition 1. *Let $v(x)$ be well S-shaped. Then both agents playing full disclosure is an equilibrium if and only if $p_1, p_2 \geq \bar{p}$, and it is the unique equilibrium if, in addition, $v'(x)$ is not linear near 0, or $p_1, p_2 > \bar{p}$, or $p_1 = p_2$.*

Competing with a sufficiently good rival induces an agent to disclose fully, and the proof of the condition $p_1, p_2 \geq \bar{p}$ is built on exactly that. If agent j plays $\{0, 1\}$, the

normalized value function of agent i is

$$\bar{v}(s_i) = (1 - p_j)\tilde{v}(s_i) + p_j\tilde{v}(s_i - 1) = (1 - p_j)\tilde{v}(s_i) - p_j\tilde{v}(1 - s_i).$$

By Lemma 1, it is S-shaped, so $\{0, 1\}$ is optimal if and only if the secant from 0 to 1 is weakly flatter than the tangent at $x = 1$ (see Figure 1b):

$$\bar{v}(1) - \bar{v}(0) \leq \bar{v}'(1) \Leftrightarrow \tilde{v}(1) \leq (1 - p_j)\tilde{v}'(1) \Leftrightarrow p_j \geq 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(1)}.$$

For uniqueness, as the game is constant-sum, all equilibria form a product set, so any disclosure played in equilibrium is a best reply to full disclosure. An S-shaped $\bar{v}(s_i)$ can touch its secant from 0 to 1 at some $s \in (0, 1)$ only by coinciding with it on $[s, 1]$, which non-linearity of $v'(x)$ near 0, or a tangent at 1 strictly steeper than the secant ($p_j > \bar{p}$), rules out, and then full disclosure is the only best reply. For $p_1 = p_2 = \bar{p}$, the best reply need not be unique, but uniqueness follows from Theorem 1.

Remark 2. Without the additional conditions, uniqueness may fail. For $v(x) = 0.5 + x - 0.5x|x|$, with $\bar{p} = 0.5$, and $(p_1, p_2) = (0.75, 0.5)$, agent 1's value function against $\{0, 1\}$ is linear, and $\{0, 1\}$ remains a best reply to any $G_1 \in \mathcal{G}(0.75)$, since $\mathbb{E}_{G_1}[s_1^2] \geq 0.5$. So $(G_1; \{0, 1\})$ is an equilibrium for every such G_1 .

Proposition 1 contrasts with the noiseless benchmark³ of Boleslavsky and Cotton (2015), in which neither school ever uses a fully informative grading policy. This is not a feature of well S-shaped rules alone: Section 5 shows that full disclosure can be reached by selection rules far beyond this class.

Since $\tilde{v}'(x)$ is concave with $\tilde{v}'(0) = 0$, and $\tilde{v}(x)$ is the (oriented) area under it, $\mathcal{E}_{\tilde{v}}(x) \geq 2$ for every well S-shaped $v(x)$, and $\mathcal{E}_{\tilde{v}}(1) = 2$ only if $\tilde{v}(x) = -x^2$ on $[0, 1]$

³The noiseless rule $v(x) = 0.5 + 0.5 \cdot \text{sign}(x)$ is not a selection rule in the sense of Section 2, as it is discontinuous at 0; Appendix C and Subsection 7.1.2 give its equilibria.

(Lemma 10 in Appendix A.1). Hence $\bar{p} \geq 0.5$, with equality only in that case, so Proposition 1 gives the following.

Corollary 1. *For any well S-shaped $v(x)$, $(\{0, 1\}; \{0, 1\})$ is never an equilibrium if $p_1 < 0.5$ or $p_2 < 0.5$, and, unless $\tilde{v}(x) = -x^2$ on $[0, 1]$, also if $p_1 = 0.5$ or $p_2 = 0.5$.*

In other words, unless $\tilde{v}(x) = -x^2$ on $[0, 1]$, in which case $\bar{p} = 0.5$, full disclosure requires both agents to be more likely good than bad.

3.2.2 Symmetric equilibrium

For equal priors, Lemma 1 and Proposition 1 characterize the equilibrium on the well S-shaped class. Unless it is full disclosure, it is a bad news equilibrium where the probability of conceding bad news ($s_i = 0$) is inversely proportional to the elasticity of $\tilde{v}(x)$ at the pooled signal a .

Theorem 1. *Let $v(x)$ be well S-shaped and $p_1 = p_2 = p$. If $p \geq \bar{p}$, the unique equilibrium is $(\{0, 1\}; \{0, 1\})$. If $p < \bar{p}$, the unique equilibrium is $(\{0, a\}; \{0, a\})$, where a solves $p = a(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)})$, and $\mathbb{P}(s_i = 0) = \frac{1}{\mathcal{E}_{\tilde{v}}(a)}$.*

The intuition is simple. Against $\{0, a\}$, an agent's normalized value function is

$$\bar{v}(s_i) = \mathbb{P}(s_i = 0)\tilde{v}(s_i) + \mathbb{P}(s_i = a)\tilde{v}(s_i - a) = \frac{a-p}{a}\tilde{v}(s_i) + \frac{p}{a}\tilde{v}(s_i - a),$$

where the second term $\tilde{v}(s_i - a) = -\tilde{v}(a - s_i)$ reflects $\tilde{v}(s_i)$ on $[0, a]$. For $\{0, a\}$ to be a best reply, the secant from $\bar{v}(0)$ to $\bar{v}(a)$ must be tangent at $s_i = a$. Since $\tilde{v}'(0) = 0$, the second term has zero derivative at a , so $\bar{v}'(a) = \frac{a-p}{a}\tilde{v}'(a)$, while the secant has the slope $\frac{\tilde{v}(a)}{a}$, the same for both terms. Hence, in equilibrium,

$$\frac{a-p}{a}\tilde{v}'(a) = \frac{\tilde{v}(a)}{a} \Leftrightarrow \frac{a-p}{a} = \frac{1}{\mathcal{E}_{\tilde{v}}(a)} \Leftrightarrow p = a \left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}\right).$$

Such a is unique, continuous and strictly increasing in p : the pooled signal rises with the prior. Indeed, $p = a - \frac{\tilde{v}(a)}{\tilde{v}'(a)}$ is the point where the tangent to $\tilde{v}(x)$ at $x = a$, $y = \tilde{v}(a) + (x - a)\tilde{v}'(a)$, crosses the x -axis. Since $\tilde{v}(x)$ is strictly concave and decreasing on $[0, 1]$ with $\tilde{v}(0) = 0$, this point is continuous and strictly increasing in a , and so is its inverse $a(p)$. As $\mathbb{P}(s_i = 0) = \frac{a-p}{a}$, this also gives the inverse elasticity result.

Uniqueness needs one more step, since the best reply to $\{0, a\}$ need not be unique. As all equilibria form a symmetric product set, a disclosure G is played in equilibrium only if it is a best reply to $\{0, a\}$ and $\{0, a\}$ is a best reply to G . The value function above touches the secant from $\bar{v}(0)$ to $\bar{v}(a)$ either only at 0 and a , so that $\{0, a\}$ is the unique best reply, or on all of $[0, a]$. The latter happens only if $\tilde{v}(x) = -cx^2$ on $[0, a]$, where $c > 0$ as $v'(x)$ is uniquely maximized at 0, and then $a = 2p$ and every G with support in $[0, 2p]$ is a best reply to $\{0, 2p\}$. But against any such $G \neq \{0, 2p\}$, the value function lies strictly above the secant from 0 to $2p$ just below $2p$, so $\{0, 2p\}$ is not a best reply to G . This step, and with it uniqueness, fails for $c = 0$, a density flat around 0.

The concavity of the density makes the agents concede bad news: under the mirror-image, reverse S-shaped rules of Subsection 7.2, they reveal good news instead.

3.2.3 No disclosure

No disclosure is the other edge case of a bad news disclosure, and, unlike full disclosure, it is never played by both agents.

Lemma 2. *Let $v(x)$ be well S-shaped. Then $(\{p_1\}; \{p_2\})$ is never an equilibrium.*

Against no disclosure, an agent's value function $v(s_i - p_j)$ is convex on $[0, p_j]$ and concave on $[p_j, 1]$, so pooling is a best reply only for the agent with the strictly higher prior. Still, no disclosure can be approached in the limit: for $\tilde{v}_n(x) = -x^n$ on $[0, 1]$, Theorem 1 gives $a = \frac{np}{n-1} \rightarrow p$. The next lemma gives a sufficient condition.

Lemma 3. Let $p \in (0, 1)$, let $v_n(x)$ be well S-shaped for all $n \in \mathbb{N}$, and let $(\{0, a_n\}; \{0, a_n\})$ be the equilibrium for v_n and priors (p, p) . If $\mathcal{E}_n(x) \rightarrow \infty$ uniformly on $(0, 1)$, then $a_n \rightarrow p$.

3.3 Informativeness of the disclosure

This subsection studies how the prior and the selection rule affect the informativeness of disclosure in the symmetric bad news equilibrium, which is unique for every well S-shaped $v(x)$ by Theorem 1.

3.3.1 Informativeness and change in prior

For a fixed p , a higher a means a more informative bad news disclosure. In equilibrium, a rises with p , but this does not imply more informative disclosure: if a rises only a little relative to p , the disclosure may become less informative. To compare disclosures for different a and p , I represent them as experiments $\pi_i : \{0, 1\} \rightarrow [0, 1]$ and use the Blackwell order (see Blackwell (1953)).

A bad news experiment sends $s_i = 0$ or $s_i = a$ if $\theta_i = 0$, and always $s_i = a$ if $\theta_i = 1$, so it is characterized by the conditional probability of revealing the bad type, $\frac{\mathbb{P}(s_i=0)}{\mathbb{P}(\theta_i=0)}$. A higher probability gives a more informative experiment, since a lower one is obtained by garbling. Hence, I say that agent i discloses more for p_{i1} than for p_{i2} if $\frac{\mathbb{P}(s_{i1}=0)}{1-p_{i1}} > \frac{\mathbb{P}(s_{i2}=0)}{1-p_{i2}}$ for the corresponding equilibrium strategies G_{i1}, G_{i2} . By Theorem 1, this measure depends on a single variable:

$$\mathbb{P}(s_i = 0) = \frac{1}{\mathcal{E}_{\bar{v}}(a)} \text{ and } p = a \left(1 - \frac{1}{\mathcal{E}_{\bar{v}}(a)}\right) \text{ hence } \frac{\mathbb{P}(s_i = 0)}{1-p} = \frac{1}{1 + (\mathcal{E}_{\bar{v}}(a) - 1)(1-a)}.$$

This gives the following lemma.

Lemma 4. Let $v(x)$ be well S-shaped and $(\{0, a\}; \{0, a\})$ the equilibrium for $p \in (0, \bar{p})$. Then disclosure increases in the prior at p if and only if $(\mathcal{E}_{\bar{v}}(x) - 1)(1 - x)$ decreases at $x = a$.

Corollary 2. *Let $v(x)$ be well S-shaped with $\mathcal{E}_{\tilde{v}}(x)$ non-increasing. Then in equilibrium agents disclose more for p_2 than for p_1 whenever $0 < p_1 < p_2 \leq \bar{p}$.*

Some algebra shows that Corollary 2 applies to the natural examples.

Example 3. For each of the three classes from Example 1 and every $\text{Var}[\varepsilon_i] \geq \bar{V}$, $\mathcal{E}_{\tilde{v}}(x)$ is non-increasing, so in equilibrium agents disclose more as p increases.

However, the elasticity of a well S-shaped rule can vary flexibly, and ex-ante stronger agents may then disclose less.

Example 4. There exists a well S-shaped $v(x)$ with $\tilde{v}(x) = -\frac{x^2}{16(1-x)^2}$ on $[0, 0.5]$ s.t. in equilibrium agents disclose less for p_2 than for p_1 if $0 < p_1 < p_2 \leq 0.375$.

3.3.2 Informativeness and change in selection rule

Next, fix the prior and change the selection rule. If $\tilde{v}(x)$ has a constant elasticity, a higher elasticity lowers $\mathbb{P}(s_i = 0)$ for a fixed p , and hence a , making the signal less informative. The next corollary extends this argument to any two selection rules.

Corollary 3. *Let $v_e(x), v_n(x)$ be well S-shaped, let $p_1 = p_2 = p \in (0, \min(\bar{p}_e, \bar{p}_n))$, and let $(\{0, a_i\}; \{0, a_i\})$ be the equilibrium for v_i , $i \in \{e, n\}$. If $\tilde{v}_e(x)$ is more elastic than $\tilde{v}_n(x)$, i.e. $\mathcal{E}_{\tilde{v}_e}(x) > \mathcal{E}_{\tilde{v}_n}(x)$ for all $x \in (0, 1)$, then $a_e < a_n$.*

So a more elastic selection rule suppresses disclosure, and within the well S-shaped class, the most informative disclosure is induced by the rules with $\mathcal{E}_{\tilde{v}}(x) = 2$, i.e. $\tilde{v}(x) = -x^2$ on $[0, 1]$. For the natural examples, this reads intuitively in terms of noise.

Example 5. For each of the three classes from Example 1, $\mathcal{E}_{\tilde{v}}(x)$ increases in $\text{Var}[\varepsilon_i]$ on $[\bar{V}, +\infty)$ for every $x \in (0, 1)$, so noisier shocks lead to less disclosure in equilibrium.

4. UTILITY OF THE FIRM

This section studies the firm's utility under a well S-shaped selection rule: whether it increases in the prior in the symmetric equilibrium, which rule in the class maximizes it, and, outside the class, the optimal level of noise.

4.1 Monotonicity with a change in the prior

It is natural to expect the firm to benefit from ex-ante stronger agents. But stronger agents do not always disclose more, and this effect could outweigh the higher prior.

For a well S-shaped $v(x)$ and priors (p, p) , let (G, G) be the unique equilibrium and $u_F(p) := \mathbb{E}_G[s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)]$ the firm's expected utility.

Proposition 2. *Let $v(x)$ be well S-shaped. Then $u_F(p)$ is increasing in p .*

So, despite possibly decreasing disclosure, a firm that uses a well S-shaped selection rule is never harmed by ex-ante stronger agents. The result is not immediate: the equilibrium disclosures for different priors need not be ranked by first-order stochastic dominance or in the Blackwell order, and under a noisy selection rule the firm's payoff need not be monotone or convex in the signals. However, the change from p to $p' > p$ splits into two steps within the bad news class. First, keeping the prior p , the pooled signal rises from $a(p)$ to $a(p')$: this spread reveals the bad type more often and widens the gap between the signals, so the firm selects the worse agent less often. Second, keeping the pooled signal $a(p')$, the prior rises to p' , which moves mass from 0 to $a(p')$, and on $\{0, a\}$ the firm's payoff is monotone. Both steps benefit the firm. This logic relies on the pooled signal moving away from the revealed one as p increases. Under the reverse S-shaped selection rules of Subsection 7.2, it moves toward it, and there exist $p_1 < p_2$ s.t. $u_F(p_1) > u_F(p_2)$ in the unique equilibrium.

A different question is how often noise makes the firm hire the agent with the lower

posterior. For $p < \bar{p}$, posteriors differ with probability $\frac{2}{\mathcal{E}_{\bar{v}}(a)}(1 - \frac{1}{\mathcal{E}_{\bar{v}}(a)})$, and the firm then hires the agent with $s_i = 0$ with probability $1 - v(a)$. If $\mathcal{E}_{\bar{v}}(x)$ is non-decreasing, the first factor does not increase in p , as $\frac{1}{\mathcal{E}_{\bar{v}}(a)} \leq \frac{1}{2}$, and the second decreases, so such reversals become rarer. Under the condition of Corollary 2, which makes disclosure increase in p , the two factors move in opposite directions, and either can dominate. For the three classes from Example 1, which satisfy it by Example 3, a direct computation shows that reversals still become rarer, but for $v(x) = 0.5 + 0.6x - 0.1e^{5-\frac{5}{x}}$ on $[0, 1]$, the probability of a reversal increases in p for $p < 0.36$.

4.2 Maximal utility in the well S-shaped class

I now depart from an exogenous $v(x)$: for a fixed p , the firm chooses the selection rule to maximize $u_F(p)$, for instance, by choosing between committee structures, evaluation protocols or review panels. By changing the biases that shape her selection rule, she changes the equilibrium disclosures and her utility.

Two questions arise. First, what is the best equilibrium utility within the well S-shaped class for each p , and which rules attain it? Second, how does it compare with the noiseless selection rule? The answer to the latter is not obvious: in the noiseless equilibrium of Boleslavsky and Cotton (2015), an agent with some probability uses a uniformly distributed posterior, which is less informative than the binary disclosure under a well S-shaped rule, so noise trades selection precision for more information.

The natural examples show an important pattern.

Example 6. For each of the three classes from Example 1, $u_F(p)$ decreases in $\text{Var}[\varepsilon_i]$ on $[\bar{V}, +\infty)$ for every $p \in (0, 1)$, so noisier shocks lower the firm's equilibrium utility.

So within each of these classes, the best rule lies at the “edge” of the well S-shaped class: the lowest level of noise that keeps $v(x)$ well S-shaped. The next theorem gives a tight upper bound on the firm's utility over the whole class and the rules that attain it.

Theorem 2. Let $v(x)$ be well S-shaped. Then

$$(i) \ u_F(p) \leq u_F^{ws}(p) = \begin{cases} p + 2p^2(1-p), & p \leq 0.5 \\ 2p - p^2, & p \geq 0.5 \end{cases}.$$

(ii) If $p \geq 0.5$, then $u_F(p) = u_F^{ws}(p)$ if and only if $p \geq \bar{p}$ and $v(1) = 1$.

(iii) If $p < 0.5$, then $u_F(p) = u_F^{ws}(p)$ if and only if $v(x) = 0.5 + x - 0.5x|x|$.

Corollary 4. Let $v(x) = 0.5 + x - 0.5x|x|$. Then

(i) $u_F(p) = u_F^{ws}(p)$ for all $p \in (0, 1)$;

(ii) $u_F(p)$ is strictly higher than under the noiseless rule if and only if $p > \frac{3-\sqrt{3}}{6} \approx 0.21$.




Figure 2: Losses from the best well S-shaped selection rules

So the selection rule $v^*(x) = 0.5 + x - 0.5x|x|$ is optimal within the well S-shaped class for every p , and uniquely so for $p < 0.5$. It has a triangular density and is generated by two independent $\varepsilon_i \sim U[0, 1]$. Its optimality has a simple source. In the equilibrium $(\{0, a\}; \{0, a\})$, the firm's utility is $p + \mathbb{P}(s_1 \neq s_2) \cdot a \cdot (v(a) - 0.5)$: she obtains p from a random pick and gains only when the signals differ, by their gap

times her excess probability of selecting the higher one. This gain rises with the pooled signal a and, at a given a , with the accuracy $v(a)$, and $v^*(x)$ is best on both counts. It induces the highest pooled signal, $a = \min(2p, 1)$ (Lemma 11), and since a concave density on $[0, 1]$ declines no faster, in proportion, than the triangular density $1 - x$, and its mass there is at most 0.5, $v^*(x)$ is also the most accurate well S-shaped rule at every gap: $v(x) \leq v^*(x)$ on $[0, 1]$ (Lemma 12). So $v^*(x)$ lies at the far edge of the class, as in the natural examples, and there the firm does not trade sharper selection against more disclosure; this trade-off arises only beyond the edge (Subsection 4.3).

Figure 2 shows the share of the first best utility $u_F^*(p) = 2p - p^2$ (Section 5) attained by $v^*(x)$ and by the noiseless rule (Appendix C). By Corollary 4, $v^*(x)$ outperforms the noiseless rule if and only if $p > \frac{3-\sqrt{3}}{6} \approx 0.21$, and attains the first best for $p \geq 0.5$, so other well S-shaped rules may beat the noiseless rule as well. The reason is that noise can induce more disclosure, which outweighs the cost of sometimes selecting a bad agent over a good one, especially for high priors, where it induces full disclosure.

4.3 The optimal level of noise

Theorem 2 leaves open what happens beyond the edge of the well S-shaped class, where the noise is too small for the density of the shock difference to be concave. The selection rules generated by i.i.d. $\varepsilon_i \sim U[0, u]$, which have a triangular density on $[-u, u]$, cover both sides of this edge. For $u \geq 1$, $v(x)$ is well S-shaped with $\tilde{v}(x) = -x^2$ on $[0, 1]$, and $u = 1$ gives $v^*(x)$; for $u < 1$, $v'(x)$ is not concave on $[0, 1]$, and as $u \rightarrow 0$, $v(x)$ converges pointwise to the noiseless selection rule. For every u and p , a symmetric equilibrium can be constructed explicitly (Proposition 13), and it has binary support if and only if $u \geq 2 \min(p, 1 - p)$; for smaller u , each agent randomizes over more than two posteriors.⁴ The construction is technical and left to Subsection 7.4; the next

⁴For $u < 1$, I do not prove uniqueness, but the equilibria of the discretized game converge to this one.

proposition concerns only the informativeness of disclosure and the firm's utility.

Proposition 3. *Let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$, $u \in (0, 1]$, and let $u_F(p, u)$ be the firm's utility in the equilibrium (G_u, G_u) constructed in Proposition 13.*

- (i) *If $p \leq 0.5$, G_u is strictly increasing in u in the Blackwell order on $(0, 2p]$, and $G_u = \{0, 2p\}$ for $u \in [2p, 1]$. If $p > 0.5$, $G_u = \{0, 1\}$ for $u \in [2 - 2p, 1]$.*
- (ii) *If $p \leq 0.5$, $u_F(p, u)$ is uniquely maximized at $u = 2p$ and strictly decreasing on $[2p, 1]$. If $p > 0.5$, $u_F(p, u)$ equals the first best $u_F^*(p) = 2p - p^2$ if and only if $u \in [2 - 2p, 1]$.*
- (iii) *As $u \rightarrow 0$, G_u converges weakly to the equilibrium disclosure for the noiseless rule.*

The proposition considers only $u \leq 1$: for $u \geq 1$, $v(x)$ is well S-shaped with $\tilde{v}(x) = -x^2$ on $[0, 1]$, so the equilibrium disclosure does not change with u , while $v(x) = 1 - \frac{(u-x)^2}{2u^2}$ decreases in u on $(0, 1]$, and so does the firm's utility.

Part (i) shows that for $p \leq 0.5$, less noise only suppresses disclosure. For $p > 0.5$ and $u < 2 - 2p$, this ranking breaks down. For instance, for $p = \frac{2}{3}$, G_u puts $(\frac{1}{5}, \frac{1}{5}, \frac{3}{5})$ on $\{0, \frac{1}{3}, 1\}$ for $u = \frac{1}{3}$ and $(\frac{4}{21}, \frac{4}{21}, \frac{13}{21})$ on $\{0, \frac{1}{4}, 1\}$ for $u = \frac{1}{2}$. The latter reveals the good type more often but the bad type less often, and since a mean-preserving spread on $[0, 1]$ cannot lower the probabilities of 0 and 1, neither is more informative. The same holds for any two $u \in [1 - p, 2 - 2p]$ if $p \geq \frac{5}{9}$ (see the proof of Proposition 3). Still, on this range, the probability of revealing the good type and the variance increase in u .

Part (ii) shows that the firm's utility is maximized at the least noise at which agents play a binary disclosure, $\{0, 2p\}$ or $\{0, 1\}$, and less noise, which breaks the binary structure, strictly lowers it. So the optimal level of noise is intermediate: it trades off disclosure by the agents against precision of selection by the firm. Moreover, at least within this family, the firm never benefits from non-binary disclosures, so going far beyond the well S-shaped class may be unnecessary.

Lastly, part (iii) shows that there is no discontinuity at zero noise: as $u \rightarrow 0$, $v(x)$ converges pointwise to $0.5 + 0.5 \cdot \text{sign}(x)$, and the equilibrium to the noiseless one derived in Boleslavsky and Cotton (2015) (see Appendix C). Simulations show the same for logistic and normal noise.

5. DESIGNING THE SELECTION RULE

The previous sections studied equilibria for a given well S-shaped selection rule; this section takes the normative side. Subsection 4.2 found the best well S-shaped rule, which outperforms the noiseless rule for sufficiently high priors and attains the first best for all priors above 0.5. The next question is which $v(x)$ maximizes the firm's utility if she can choose any selection rule, and which properties of $v(x)$ are necessary or sufficient for optimality. I start with the main definitions.

I say that $v(x)$ is *revelation-accurate* if $v(1) = 1$, i.e. whenever agent i is revealed to be good and agent j bad ($s_i = 1, s_j = 0$), the firm selects agent i with probability 1. This is a simple necessary condition for optimality: if $v(1) < 1$, an affine modification of $v(x)$ leads to the same equilibria but gives the firm a strictly higher utility.

I also say that $v(x)$ *implements a full disclosure equilibrium (uniquely) for (p_1, p_2)* if, for these priors, full disclosure by both agents is an (the unique) equilibrium, and *for p* if it does so for the symmetric priors (p, p) .

The firm's maximal equilibrium utility⁵ under a given selection rule and her first best utility, both for symmetric priors, are defined as follows.

Definition 4. For a selection rule $v(x)$ and priors (p, p) , let E be the set of all equilibria. Then the maximal expected utility of the firm among all equilibria is

$$\bar{u}_F(p) := \max_{(G_1^*, G_2^*) \in E} \mathbb{E}_{G_1^*, G_2^*} [s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)].$$

⁵The maximum is well defined: by the continuity of $v(x)$, each agent's set of max-minimizers is compact, and so is the set of all equilibria.

Definition 5. Let $\mathcal{V}$ be the set of all selection rules. Then the first best expected utility of the firm is $u_F^*(p) := \sup_{G_1, G_2 \in \mathcal{G}(p), v \in \mathcal{V}} \mathbb{E}_{G_1, G_2}[s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)]$.

The first result shows that, for symmetric agents, a full disclosure equilibrium and revelation accuracy are not only sufficient, but also necessary for the first best.

Lemma 5. *For every selection rule $v(x)$, $\bar{u}_F(p) \leq u_F^*(p) = 2p - p^2$, with equality if and only if $v(x)$ implements a full disclosure equilibrium for p and is revelation-accurate.*

The proof is simple. For priors (p, p) ,

$$u_F^*(p) = \max_{G_1, G_2 \in \mathcal{G}(p)} \mathbb{E}_{G_1, G_2}[\max(s_1, s_2)].$$

As $\max(s_1, s_2)$ is convex and piecewise linear, the maximum is attained if and only if at least one agent plays full disclosure. As all equilibria form a symmetric product set, if full disclosure and some other disclosure form an equilibrium, so do two full disclosures. Since $s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) \leq \max(s_1, s_2)$, $\bar{u}_F(p) < u_F^*(p)$ if $v(x)$ does not implement a full disclosure equilibrium, and in a full disclosure equilibrium, the firm gets the first best if and only if $v(x)$ is revelation-accurate. The result generally fails for $p_1 \neq p_2$: the set of equilibria is then not symmetric and may contain only equilibria in which exactly one agent discloses fully.

For every $p \in (0, 1)$, the selection rules that implement a full disclosure equilibrium form a non-empty class, described by an infinite system of linear inequalities:

$$(1 - p)\hat{v}(x) - p\hat{v}(1 - x) \leq (x - p)\hat{v}(1) \quad \forall x \in (0, 1), \quad \text{where } \hat{v}(x) := v(x) - 0.5.$$

Except for $p = 0.5$, this class has no simple structure, and the assumption that the firm knows the priors exactly is restrictive. Instead, suppose that the firm only knows that the priors lie in some exogenous set S . This robust approach leads to sharper results.

First, although a full disclosure equilibrium is not necessary for the first best under asymmetric priors, this approach covers this case too.

Lemma 6. *Let $S \subseteq (0, 1)$. A selection rule $v(x)$ implements a full disclosure equilibrium for all $(p_1, p_2) \in S^2$ if and only if it does so for all $p \in S$.*

The *only if* direction is straightforward; as for *if*, suppose that $v(x)$ implements a full disclosure equilibrium for some $p \in S$, and let agent j with prior $p_j = p$ play full disclosure. Agent i 's value function does not depend on his own prior, and since full disclosure is a best reply for $p_i = p$, its concavification is the secant from 0 to 1, so full disclosure is a best reply for every $p_i \in (0, 1)$. Hence, if $v(x)$ implements a full disclosure equilibrium for all $p \in S$, then it does so for all $(p_1, p_2) \in S^2$.

Second, robustly implementing full disclosure strongly restricts $v(x)$.

Proposition 4. *Let a selection rule $v(x)$ implement a full disclosure equilibrium for all $(p_1, p_2) \in S^2$, where $\inf S < 0.5 < \sup S$. Then*

- (i) *$v(x)$ implements a full disclosure equilibrium for all $p \in (0, 1)$;*
- (ii) *$v(x) = 0.5 + ax$, where $a \in [0, 0.5]$.*

Corollary 5. *Let $v(x)$ be a selection rule with $\bar{u}_F(p) = u_F^*(p)$ for $p \in \{p_1, p_2\}$, where $p_1 < 0.5 < p_2$. Then $v(x) = 0.5 + 0.5x$.*

The proof of Proposition 4 is simple. By Lemma 6, the system of inequalities above holds for every $p \in S$. Adding it at x and at $1 - x$ with weights $1 - p$ and p gives $(1 - 2p)(\hat{v}(x) - x\hat{v}(1)) \leq 0$ for all $x \in (0, 1)$. So a prior in S below 0.5 rules out any upward deviation of $\hat{v}(x)$ from the linear function $x\hat{v}(1)$, and a prior in S above 0.5 rules out any downward one. Hence, $v(x) = 0.5 + x(v(1) - 0.5)$ is linear, so it implements full disclosure for all priors, and any $a = v(1) - 0.5 \in [0, 0.5]$ is valid.

So a firm that is unsure about the priors can robustly attain her first best only with $v(x) = 0.5 + 0.5x$. But this rule has a crucial issue. Since $v(x)$, and hence $\bar{v}(s_i)$, is linear, every pair of disclosures is an equilibrium, and the firm need not get the first best: robustness comes at the cost of complete indifference.

This tension is specific to rules that depend only on the difference between the signals, as those generated by additive evaluation noise do. A more general anonymous rule avoids it: if the firm selects agent i with probability $0.5 + 0.5(s_i^2 - s_j^2)$, each agent maximizes $\mathbb{E}[s_i^2]$, so full disclosure is strictly dominant for every prior, and as this rule always selects the good type over the bad one, the firm attains the first best for all priors in the unique equilibrium. I do not pursue such rules further, since additive noise cannot generate them.

The issue disappears under slightly less robustness. Suppose that the firm knows whether both agents are more or less likely to be good than bad. The rules that implement a full disclosure equilibrium in either case are described by the deviation of $\hat{v}(x)$ from its secant from 0 to 1.

Definition 6. Let $v(x)$ be a selection rule, let $a := \hat{v}(1)$ and $g(x) := \hat{v}(x) - ax$ on $[0, 1]$. Then $v(x)$ is (strictly) *positively* [*negatively*] *balanced* if $g(x) = g(1 - x)$ for all $x \in [0, 1]$ and $g(x) \geq (>)0$ [$g(x) \leq (<)0$] for all $x \in (0, 1)$.

The symmetry of $g(x)$ alone is exactly the condition for a full disclosure equilibrium at $p_1 = p_2 = 0.5$, as shown after Proposition 5; the sign of $g(x)$ then determines the half-interval of priors on which full disclosure extends. For a selection rule, $a \in [0, 0.5]$ and $g(0) = g(1) = 0$ hold automatically, and for a symmetric $g(x)$, the monotonicity of $v(x)$ is equivalent to $|g(x) - g(y)| \leq a|x - y|$.

Proposition 5. Let $v(x)$ be a selection rule.

- (i) $v(x)$ implements a full disclosure equilibrium $\forall (p_1, p_2) \in [0.5, 1]^2$ [$\forall (p_1, p_2) \in (0, 0.5]^2$] if and only if it is positively [*negatively*] balanced.

(ii) If $v(x)$ is strictly positively [negatively] balanced, it implements a full disclosure equilibrium uniquely $\forall (p_1, p_2) \in (0.5, 1)^2$ [$\forall (p_1, p_2) \in (0, 0.5)^2$].

Part (i) gives necessary and sufficient conditions for implementing full disclosure for all priors on a half-interval. They impose a specific structure: $v(x)$ can deviate from the universal implementation $v(x) = 0.5 + ax$ ($v(1) = 1$ again implies $a = 0.5$) only by a function that is symmetric around 0.5 on $[0, 1]$, non-negative for strong pools and non-positive for weak ones.

The proof is again simple: the conditions hold for all $p \in (0, 0.5]$ if and only if they hold for $p = 0$ and $p = 0.5$, and the latter gives the symmetry. If agent j with $p = 0.5$ discloses fully, agent i 's normalized value function is odd around $s_i = 0.5$:

$$\bar{v}(s_i) = 0.5\hat{v}(s_i) - 0.5\hat{v}(1 - s_i) = -\bar{v}(1 - s_i) \quad \forall s_i \in [0, 1].$$

Hence $\{0, 1\}$ is a best reply only if $\bar{v}(s_i)$ is linear:

$$\bar{v}(s_i) = 0.5\hat{v}(s_i) - 0.5\hat{v}(1 - s_i) = (s_i - 0.5)\hat{v}(1) \Leftrightarrow \hat{v}(s_i) - s_i\hat{v}(1) = \hat{v}(1 - s_i) - (1 - s_i)\hat{v}(1).$$

That is, the deviation $g(s_i)$ of $\hat{v}(s_i)$ from the secant from 0 to 1 must be symmetric around 0.5: $g(s_i) = g(1 - s_i)$ for all $s_i \in [0, 1]$.

Part (ii) gives a sufficient condition for full disclosure to be the unique equilibrium. As all equilibria form a product set, it suffices that $\{0, 1\}$ is each agent's unique best reply to itself. This is also why $p = 0.5$ is excluded: there, any $v(x)$ that implements full disclosure makes $\bar{v}(s_i)$ linear, so the best reply is not unique. The condition is not necessary, but since it is only a slight strengthening of (i), the room for other rules with a unique full disclosure equilibrium is small, and checking uniqueness for them requires showing that no disclosure supported on $\{s_i : g(s_i) = 0\}$ is a max-minimizer.

A firm facing a weak pool therefore needs a negatively balanced rule, and evaluation

noise of the kind in Section 2 cannot deliver one: by the next result, which extends Corollary 1 beyond the well S-shaped class, i.i.d. shocks rule out full disclosure as soon as one prior is below 0.5.

Corollary 6. *Let $v(x)$ be a selection rule with $v'(0) > v'(1)$. Then $(\{0, 1\}; \{0, 1\})$ is never an equilibrium if $p_1 < 0.5$ or $p_2 < 0.5$. In particular, this holds for every $v(x)$ generated by i.i.d. shocks with a square-integrable density.*

Remark 3. Like $\tilde{v}(x)$, $g(x)$ removes a linear part of $v(x)$, but it subtracts the secant from 0 to 1 rather than the tangent at 0. So, for every selection rule, equilibria depend on $v(x)$ only through $g(x)$, and revelation accuracy only through a . A well S-shaped $v(x)$ is positively balanced if and only if $\tilde{v}(x) = -x^2$ on $[0, 1]$.

6. ASYMMETRIC PRIORS

This section characterizes the equilibrium for asymmetric priors under a well S-shaped selection rule with $v'(x)$ not linear near 0. Without loss of generality let $p_2 \leq p_1$. By Lemma 1, every best reply is unique and takes one of three forms: no disclosure $\{p_i\}$, full disclosure $\{0, 1\}$, or bad news $\{0, c\}$ with $p_i < c < 1$. In general, an agent's best reply depends on his rival's whole disclosure, but in the following example it depends only on the rival's prior, which makes the equilibrium explicit.

Example 7. Let $\tilde{v}(x) = -x^3$. Agent i 's best reply is $\{0, \min(\frac{3p_i}{2}, 1)\}$, or $\{p_i\}$ if $p_i \geq \frac{3p_j}{2}$. So for $p_2 \leq p_1$, the unique equilibrium is $(\{0, 1\}; \{0, 1\})$ if $p_2 \geq \frac{2}{3}$, and otherwise

- (i) $(\{p_1\}; \{0, 1\})$ if $p_1 \geq \max(\frac{3p_2}{2}, \frac{2}{3})$;
- (ii) $(\{p_1\}; \{0, \frac{3p_1}{2}\})$ if $\frac{3p_2}{2} \leq p_1 < \frac{2}{3}$;
- (iii) $(\{0, \frac{3p_2}{2}\}; \{0, 1\})$ if $\frac{2}{3} \leq p_1 < \frac{3p_2}{2}$;
- (iv) $(\{0, \frac{3p_2}{2}\}; \{0, \frac{3p_1}{2}\})$ if $p_1 < \min(\frac{3p_2}{2}, \frac{2}{3})$.

For every such selection rule, the same regions arise, with boundaries given by the following objects; the last column gives them for Example 7.

	definition	Example 7
$\bar{p}$	$1 - \frac{1}{\varepsilon_{\bar{v}}(1)}$: full disclosure for $p_1, p_2 \geq \bar{p}$	$\frac{2}{3}$
$b(p_1)$	agent 2's best reply to $\{p_1\}$ is $\{0, b(p_1)\}$ if $p_2 < b(p_1)$, else $\{p_2\}$	$\min(\frac{3p_1}{2}, 1)$
p^*	smallest p_1 with $b(p_1) = 1$	$\frac{2}{3}$
$a(p_2)$	agent 1's best reply to $\{0, 1\}$ is $\{0, a(p_2)\}$ if $p_1 < a(p_2)$, else $\{p_1\}$	$\frac{3p_2}{2}$
$\check{p}_2(p_1)$	largest p_2 for which $\{p_1\}$ is agent 1's best reply to $\{0, b(p_1)\}$	$\frac{2p_1}{3}$
$\hat{p}_1(p_2)$	smallest p_1 for which $\{0, 1\}$ is agent 2's best reply to $\{0, a(p_2)\}$	$\frac{2}{3}$

Proposition 6. *Let $v(x)$ be well S-shaped with $v'(x)$ not linear near 0, and let $p_1, p_2 \in (0, 1)$ with $p_2 \leq p_1$. Then the equilibrium is unique, and it is $(\{0, 1\}; \{0, 1\})$ if $p_2 \geq \bar{p}$, and otherwise*

- (i) $(\{p_1\}; \{0, 1\})$ if $p_1 \geq \max\{p^*, a(p_2)\}$;
- (ii) $(\{p_1\}; \{0, b(p_1)\})$ if $p_1 < p^*$ and $p_2 \leq \check{p}_2(p_1)$;
- (iii) $(\{0, a(p_2)\}; \{0, 1\})$ if $\hat{p}_1(p_2) \leq p_1 < a(p_2)$;
- (iv) $(\{0, a\}; \{0, b\})$ with $p_1 < a \leq b < 1$ otherwise, where $a = b$ if and only if $p_1 = p_2$.


The equilibrium strategies are continuous in (p_1, p_2) .

Figure 3 depicts these regions. In Example 7, regions (i)–(iv) are exactly the cases listed, and $\bar{p} = p^*$: it is the knife-edge case in panel (c), between $\bar{p} < p^*$ and $\bar{p} > p^*$ in panels (a) and (b). The boundary $a(\cdot)$ is monotone, but $\hat{p}_1(\cdot)$ and $\check{p}_2(\cdot)$ need not be: in panel (d), neither is monotone. Non-linearity of $v'(x)$ near 0 cannot be dropped from Proposition 6: without it, the equilibrium need not be unique, and disclosures need not take these forms (Remark 4).

Remark 4. Let $v(x) = v^*(x)$, for which $v'(x)$ is linear on $[0, 1]$, and $(p_1, p_2) = (0.4, 0.3)$. Then agent 2 plays $\{0, 0.6\}$ in every equilibrium, and agent 1's value function against it

is linear on $[0, 0.6]$. So agent 1's equilibrium disclosures are exactly those with support in $[0, 0.6]$ and $\mathbb{E}[s_1^2] = 0.18$, which keep $\{0, 0.6\}$ a best reply of agent 2, such as $\{0, 0.45\}$, $\{0.3, 0.6\}$ and their mixtures.

Figure 3: Equilibria for asymmetric priors, with $\tilde{v}(x)$ given on $[0, 1]$



Colors are for the corresponding equilibrium type: black – $(\{0, 1\}; \{0, 1\})$; red – $(\{p_1\}; \{0, 1\})$; green – $(\{0, a\}; \{0, b\})$; blue – $(\{p_1\}; \{0, b\})$; yellow – $(\{0, a\}; \{0, 1\})$.

Every pair of forms except no disclosure by both agents (Lemma 2) is played for some priors, and the agent with the lower prior never plays no disclosure. For a fixed $v(x)$, however, not every such pair is played: full disclosure is a best reply to $\{0, c\}$ only if $c \geq p^*$, so $(\{0, c\}; \{0, 1\})$ is an equilibrium only if $c \geq p^*$. In regions (ii) and (iii), and in Example 7 also in region (iv), an agent's bad news disclosure depends only on his rival's prior. More generally, it is the rival who forces an agent to disclose.

Proposition 7. *Let $v(x)$ be as in Proposition 6. Then the equilibrium disclosure of agent i is non-decreasing in p_j , and strictly increasing whenever agent i plays $\{0, a\}$ with $p_i < a < 1$.*

An agent's own prior usually has the opposite effect: in Example 7, his high signal does not depend on p_i , so he discloses less as p_i increases. This effect need not be monotone, though, as the next example shows.

Example 8. Let $v(x)$ be well S-shaped with $\tilde{v}(x) = -x^{2.5}$ on $[0, 1]$, and let $p_j = 0.6$. Then agent i 's equilibrium disclosure is non-monotone in p_i .

Region (i) is a favorite–underdog pattern: when p_1 is high and p_2 is low, the favorite reveals nothing, while the underdog gambles on full disclosure. Against full disclosure, agent 1's value function $(1 - p_2)v(s_1) + p_2v(s_1 - 1)$ is close to $v(s_1)$ for low p_2 , so it is concave except near 0, and pooling at p_1 is optimal. Against no disclosure, agent 2's value function $v(s_2 - p_1)$ is convex on $[0, p_1]$ and concave on $[p_1, 1]$, and for high p_1 , its concavification is supported on $\{0, 1\}$ despite the concave part. So an agent who is nearly sure to win plays safe, while one who is likely to lose gambles; under a well reverse S-shaped selection rule, the roles are reversed (Subsection 7.2).

For the firm, the monotonicity of Proposition 2 extends to asymmetric priors everywhere except in region (iv) of Proposition 6.

Proposition 8. *Let $v(x)$ be as in Proposition 6 and $p_2 \leq p_1$. In the interior of the full disclosure region and of each of the regions (i)–(iii), both partial derivatives of u_F are positive.*

Example 9. Let $v(x) = 0.5 + (20x - x^{20})/38$ on $[0, 1]$, so that $\tilde{v}(x) = -x^{20}$ and $v(1) = 1$, and let $p_1 = 0.924$. Then u_F decreases in p_2 on $[0.839, 0.873]$, where the equilibrium is in region (iv): once agent 2 stops disclosing fully, his high signal b decreases in p_2 .

For the agents, this question is trivial under equal priors, as each is selected with probability 0.5. Under asymmetric priors, since the game is constant-sum, each agent's equilibrium payoff $u_i(p_1, p_2) = \mathbb{E}_{G_1, G_2}[v(s_i - s_j)]$ is well defined for any selection rule, whether or not the equilibrium is unique.

Lemma 7. *For any selection rule, $u_i(p_1, p_2)$ is non-decreasing in p_i and non-increasing in p_j . If $v(x)$ is strictly increasing, in particular if it is well S-shaped, $u_i(p_1, p_2)$ is strictly increasing in p_i and strictly decreasing in p_j .*

7. DISCUSSION

This section collects extensions and results excluded from the main text. Subsection 7.1 relaxes the binary prior. Subsections 7.2–7.4 leave the well S-shaped class: Subsection 7.2 studies its mirror image, Subsection 7.3 describes equilibria around its boundary, and Subsection 7.4 constructs the equilibrium for the triangular selection rules of Subsection 4.3. Subsection 7.5 relaxes the anonymity of the selection rule.

7.1 Non-binary priors

Every result up to this point assumes a binary prior. That assumption buys tractability, but it is not innocuous: the two-point distribution on $\{0, 1\}$ is the *largest* element in the convex order among all distributions on $[0, 1]$ with mean p , so it gives agents the richest possible set of feasible disclosures. This subsection relaxes it.

7.1.1 Censorship equilibrium

Under a well S-shaped rule, an agent's value function is S-shaped against every disclosure of his rival, so if the prior has a strictly positive density, Theorem 1 of Kolotilin et al. (2022) applies: the optimal disclosure reveals types below some θ^* and pools all types above it into one signal. The same censorship logic applies to a binary prior in the quantile space. Let $F^{-1}(x)$ be its quantile function:

$$F^{-1}(x) = \begin{cases} 0, & x \in [0, 1-p) \\ 1, & x \in [1-p, 1] \end{cases}.$$

Then a censorship with quantile $q < 1-p$ reveals $\theta_i = 0$ with probability q (below the quantile) and otherwise sends the pooled signal $s_i = \frac{(1-p-q) \cdot 0 + p \cdot 1}{1-p-q+p} = \frac{p}{1-q}$ (above the quantile), while a censorship with quantile $q \geq 1-p$ is full disclosure.

With a positive density $f(x)$, the equilibrium is again a symmetric upper censorship:

Proposition 9. *Suppose that $v(x)$ is well S-shaped and $F(x)$ has a strictly positive density $f(x)$ on $[0, 1]$. Then $(G(x), G(x))$ is an equilibrium, unique if $v'(x)$ is not linear near 0, where*

$$G(x) = \begin{cases} F(x) & , x \leq \theta^* \\ F(\theta^*) & , x \in (\theta^*, m^*) \text{ with } m^* = \mathbb{E}[\theta | \theta \geq \theta^*] \text{ and where } \theta^* \in (0, 1) \text{ solves} \\ 1 & , x \geq m^* \end{cases}$$

$$\int_0^{\theta^*} [\tilde{v}(m^* - s) - \tilde{v}(\theta^* - s) - (m^* - \theta^*)\tilde{v}'(m^* - s)]f(s)ds + \tilde{v}(m^* - \theta^*)(1 - F(\theta^*)) = 0 \quad (1)$$

For the best well S-shaped rule from Corollary 4, the equation collapses entirely.

Corollary 7. *Let $v(x) = 0.5 + x - 0.5x|x|$. Then θ^* in Proposition 9 solves $F(\theta^*) = 0.5$.*

So this rule induces median censorship as an equilibrium, whatever the prior is.

Step 2 of the proof of Proposition 9 does not use the density, so the median censorship remains an equilibrium whenever F is continuous at its median, and in quantile terms the bad news equilibrium $\{0, 2p\}$ under a binary prior is a median censorship too.

7.1.2 The noiseless benchmark

With a binary prior the only restriction on a disclosure G is that its mean equal p , but with a general prior the feasible set becomes smaller: by Gentzkow and Kamenica (2016), G is attainable if and only if it is a mean-preserving contraction of F , i.e. $\mathbb{E}_G[\phi] \leq \mathbb{E}_F[\phi]$ for every convex ϕ . This substantially changes the noiseless benchmark and the interpretation of Subsection 4.2. Here I count the noiseless rule $v(x) = 0.5 + 0.5 \cdot \text{sign}(x)$ among selection rules, with $\underline{\theta} = \min \text{supp}(F)$, $\bar{\theta} = \max \text{supp}(F)$.

Proposition 10. *Let F be atomless. Then the noiseless selection rule implements a full disclosure equilibrium if and only if F is convex on $[\underline{\theta}, \bar{\theta}]$, and then it is the unique equilibrium.*

Both directions rest on the fact that, against full disclosure, the deviator's value function is F itself. If F is convex, then $\mathbb{E}_G[F] \leq \mathbb{E}_F[F] = 0.5$ for every feasible G , so no deviation beats full disclosure; otherwise some interval of types can be pooled into its mean at a profit, and if F has a density, these are types where it decreases.

With a general prior, the first best is $\mathbb{E}[\max(\theta_1, \theta_2)]$, and the comparison in Subsection 4.2 reverses sharply for convex priors.

Corollary 8. *Let F be atomless. A selection rule attains the first best in some equilibrium if and only if it is the noiseless selection rule and F is convex on $[\underline{\theta}, \bar{\theta}]$.*

The first best requires both agents to disclose fully and the firm never to select the worse of two distinct types. A well S-shaped rule always fails the second requirement, while under the noiseless rule the first one holds exactly when F is convex.

For concave priors the equilibrium structure is also simple: a suitable uniform distribution is always feasible, precisely because the density is decreasing.

Proposition 11. *Let F be concave on its support $[\underline{\theta}, \bar{\theta}]$, possibly with an atom at $\underline{\theta}$, and with mean p . Then both agents playing $G = U[\underline{\theta}, 2p - \underline{\theta}]$ is the unique equilibrium of the noiseless game, and the firm's equilibrium utility is $u_F = \frac{4p - \underline{\theta}}{3}$.*

Under the noiseless selection rule, the firm's payoff depends on the prior only through its mean and its lower support point, so, for a given lower support point, it does not grow as the pool of agents becomes more dispersed. The first best does, since $\max(\theta_1, \theta_2)$ is convex, and so may the payoff under a noisy selection rule. Hence, for concave priors, the comparison from Subsection 4.2 depends on how dispersed F is.

Example 10. Let $p = \frac{1}{3}$ and $\underline{\theta} = 0$, so that the noiseless selection rule gives $u_F = \frac{4}{9} \approx 0.444$ under every concave F . Under $v^*(x)$ and the median censorship of Corollary 7, the firm gets $u_F \approx 0.399$ for $F(x) = 2x - x^2$ and $u_F \approx 0.457$ for $F(x) = \frac{1+2x}{3}$.

So the noiseless selection rule is better under the first prior and worse under the second, which is a mean-preserving spread of the first. The binary prior of Sections 3–6 is the extreme case: it is the most dispersed prior with mean p .

When $\text{supp}(F) = [0, 1]$, a convex F has $p \geq 0.5$ and a concave F has $p \leq 0.5$, so convex and concave priors describe strong and weak pools in the sense of Section 5, although most priors are neither.

Overall, the binary prior is harmless for the equilibrium characterization, but not for the comparison with the noiseless rule: the advantage of noise disappears for convex priors and, for concave ones, depends on how dispersed the pool is.

7.2 Reverse S-shaped selection rules

The well S-shaped class has a mirror image, in which the density of the selection rule is convex rather than concave. A concave density puts an agent's marginal gain from a higher signal near a tie, as with an urge to win, whereas a convex density puts it at large margins, as with an urge to surpass.

Definition 7. A function $v : [-1, 1] \rightarrow [0, 1]$ is called *well reverse S-shaped* if $v(x) - 0.5$ is odd and increasing, and $v'(x)$ is convex on $[-1, 1]$ and uniquely minimized at $x = 0$.

The following lemma transfers the equilibrium analysis to this class.

Lemma 8. (i) For every well reverse S-shaped $v_2(x)$, there is a well S-shaped $v_1(x)$ with $\tilde{v}_2(x) = -\tilde{v}_1(x)$.

(ii) Let $\tilde{v}_2(x) = -\tilde{v}_1(x)$. If $(G_1(s_1), G_2(s_2))$ is an equilibrium for $v_1(x)$ and priors (p_1, p_2) , then $(\check{G}_1, \check{G}_2)$ is an equilibrium for $v_2(x)$ and priors $(1 - p_1, 1 - p_2)$, where $\check{G}_i$ is the distribution of $1 - s_i$ for $s_i \sim G_i$.

Hence, for equal priors, the unique equilibrium of a well reverse S-shaped $v(x)$ is $(\{b, 1\}; \{b, 1\})$, where b increases in p analogously to Theorem 1: each agent reveals good news rather than conceding bad news, and full disclosure is reached for $p \leq \frac{1}{\varepsilon_{\bar{v}}(1)}$, mirroring Proposition 1. In particular, such rules can deliver the first best for $p < 0.5$: on $[0, 1]$, $v(x) = 0.5 + (0.5 - \eta)x + \eta x|x|$ with $\eta \in (0, 0.5]$ equals $0.5 + 0.5x - \eta x(1 - x)$, so it is revelation-accurate and strictly negatively balanced (Proposition 5).

The guarantee of Proposition 2, however, fails for many such rules.

Example 11. For $v(x) = 0.5 + 0.5x^3$, $u_F(p)$ is non-monotone in p .

For these rules, disclosure often decreases in p unless the elasticity increases sharply, mirroring Lemma 4 and Corollary 2, and since $v(x)$ is convex on $[0, 1]$, this effect can outweigh the increase in the prior. Asymmetric equilibria are flipped too: under a well reverse S-shaped rule with $v'(x)$ not linear near 0, the agent with the high prior discloses fully and the other discloses nothing, the opposite of the pattern in Section 6.

7.3 Around the boundary of the well S-shaped class

This subsection describes what changes as the selection rule leaves the well S-shaped class: under a weaker condition, the equilibrium keeps the forms of Section 6, whereas

further away equilibrium disclosures may have non-binary support.

7.3.1 Binary support equilibrium

Although no simple extension of the well S-shaped class keeps $\bar{v}(s_i)$ S-shaped against every G_j (Section 3), the equilibrium forms of Lemma 1 and Proposition 6 survive outside it. Their proofs use the shape of $v(x)$ only through the shape of agent i 's value function against the disclosures that agent j plays in equilibrium, $\{p_j\}$ and $\{0, c\}$. Against $\{0, c\}$, the sign of $\bar{v}''(s_i)$ on $(0, c)$ depends on $w(s_i)$ and $w(c - s_i)$, where $w(x) := -v''(x)$, so only pairs of points with sum at most 1 matter.

Proposition 12. *Let $v(x)$ be a selection rule with $v'(x)$ differentiable on $(0, 1]$, $w(x) > 0$ on $(0, 1]$, and $\frac{w(c-x)}{w(x)}$ strictly decreasing in x on $(0, c)$ for every $c \in (0, 1]$. Then for all priors, the equilibrium is unique, and each agent i plays $\{p_i\}$ or $\{0, a_i\}$ with $a_i \in (p_i, 1]$.*

These conditions hold for every well S-shaped $v(x)$ with strictly concave and differentiable $v'(x)$, since $w(x)$ is then strictly increasing, but they do not require $v'(x)$ to be concave: $w(x)$ may decrease near $x = 1$ if it increases fast enough near 0. For $\xi \sim N(0, \sigma^2)$, $w(x)$ is proportional to $xe^{-x^2/(2\sigma^2)}$, and the conditions hold if and only if $\sigma \geq \frac{1}{2}$, whereas $v(x)$ is well S-shaped only for $\sigma \geq 1$; in terms of Example 2, the threshold drops from $\text{Var}[\varepsilon_i] \geq \frac{1}{2}$ to $\text{Var}[\varepsilon_i] \geq \frac{1}{8}$. This bound is nearly tight: in simulations, the equilibrium has binary support for all priors for $\sigma = 0.48$, but for $\sigma = 0.46$ and $(p_1, p_2) = (0.62, 0.48)$, agent 1 discloses three signals in every equilibrium.

7.3.2 Non-binary support equilibrium

Outside the well S-shaped class, equilibrium disclosures need not have binary support, and their structure varies considerably with the selection rule.

Lemma 9. *Let $v(x)$ be real-analytic on $[-1, 1]$ with $v''(x) > 0$ for all $x \in [-1, 0)$. Then for all $p_i, p_j \in (0, 1)$ and $G_j \in \mathcal{G}(p_j)$, every optimal disclosure for $(\bar{v}(s_i, G_j), p_i)$ has finite support.*

These conditions hold for the selection rules generated by i.i.d. Gumbel, normal or Cauchy shocks of any variance, so for these natural examples, equilibrium disclosures are discrete. By the identity theorem, an analytic $\bar{v}(s_i)$ coincides with a line either on a finite set or on all of $[0, 1]$, and the latter is ruled out by the convexity of $v(x)$ on $[-1, 0]$. Still, simulations⁶ show that the number of support points grows as the noise decreases, mirroring the classical result for games on the unit square with bell-shaped kernels (see Karlin (1959)), in which optimal strategies are finitely supported and the number of support points grows without bound as the kernel sharpens.

In contrast, a selection rule that is not analytic can induce an equilibrium disclosure with a continuous part, already for exponential shocks.

Example 12. Let $v(x)$ be generated by i.i.d. $\varepsilon_i \sim \text{Exp}(\lambda)$, i.e. the Laplace CDF $v(x) = 0.5 + 0.5 \cdot \text{sign}(x) \left(1 - e^{-\lambda|x|}\right)$. Then for all $p \leq 0.5$, both agents playing

$$G = \frac{1}{2(1 + \lambda p)} \left(\delta_{\{0\}} + \delta_{\{2p\}} \right) + \frac{\lambda p}{1 + \lambda p} U[0, 2p]$$

is a symmetric equilibrium.

In general, finding such equilibria explicitly is hard. Whenever an equilibrium disclosure of agent j contains an interval $[a, b]$ in its support, $\bar{v}(s_j)$ must be linear on $[a, b]$, and this linearity is a functional equation for the equilibrium disclosure G_i^* :

$$\bar{v}'(s_j) = \int_0^1 v'(s_j - s_i) dG_i^*(s_i) = \text{const} \quad \forall s_j \in [a, b].$$

This is a Fredholm integral equation of the first kind with a difference kernel on a finite interval. Such equations have no general explicit solution: the general theory reduces them to auxiliary equations (see Sakhnovich (2015)), and closed-form solutions are known only for special kernels, such as kernels with a rational Fourier transform, as in

⁶For $p = 0.3$ and $\xi \sim N(0, \sigma^2)$, the equilibrium G_i^* has 5 support points for $\sigma = 0.1$ and 9 for $\sigma = 0.05$.

Example 12, or piecewise polynomial kernels (see Polyanin and Manzhirov (2008)), which include the triangular selection rules studied in the next subsection.

7.4 Triangular selection rules

This subsection constructs the symmetric equilibrium used in Subsection 4.3 for the selection rules generated by i.i.d. $\varepsilon_i \sim U[0, u]$ with $u < 1$. Since $v(x)$ is piecewise quadratic, $v'(x)$ is piecewise linear, so the functional equation from Subsection 7.3 reduces to a difference equation, and one can find $G_i(s_i)$ s.t. $\bar{v}(s_j)$ is linear on $[a, b]$.

Definition 8. Say χ is a proper u -partition of the interval $[a, b]$ if its CDF is given by:

$$G(x) = \begin{cases} 0, & x < a \\ \frac{i+1}{k+1}, & x \in [a + i \cdot u, b - (k - 1 - i) \cdot u), i \in \{0, \dots, k - 1\} \\ \frac{i+1}{k}, & x \in [b - (k - 1 - i) \cdot u, a + (i + 1) \cdot u), i \in \{0, \dots, k - 2\} \\ 1, & x \geq b \end{cases}, \text{ with } k = \lceil \frac{b-a}{u} \rceil$$

Remark 5. This G solves the functional equation $G(x - u) + G(x + u) = 2G(x)$, i.e. $\mathbb{P}(x < \chi \leq x + u) = \mathbb{P}(x - u < \chi \leq x)$, for all $x \in [a, b)$, subject to $\text{supp}(\chi) \subseteq [a, b]$.

The proper u -partition of $[a, b]$ is the unique $G_i(s_i)$ supported in $[a, b]$ that makes $\bar{v}(s_j) = \int_0^1 v(s_j - s_i) dG_i(s_i)$ linear on $[a, b]$. Both agents playing it is often not an equilibrium, since $\bar{v}(s_j)$ beyond $[a, b]$ also matters, but it is the key ingredient: the next proposition constructs a symmetric equilibrium for all (u, p) from mixtures of proper u -partitions and, possibly, $\delta_{\{1\}}$.

Proposition 13. Let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$. Then a symmetric equilibrium disclosure strategy is:

- (i) If $p \leq 0.5$, disclosure is a proper u -partition of $[0, 2p]$.
- (ii) If $p > 0.5$ and $u \geq 2 - 2p$, disclosure is $\{0, 1\}$.

(iii) Otherwise, disclosure mixes a proper u -partition of $[0, a]$ for some a with a mass point at 1 and (possibly) with a proper u -partition of $[0, 1 - u]$.

So binary support survives a slight violation of well S-shapedness for extreme priors, $p \leq 0.5u$ or $p \geq 1 - 0.5u$, but fails for all $p \in (0.5u, 1 - 0.5u)$. The last case, whose exact equations are in Appendix B, shows how complex equilibria can be for general $v(x)$: as the signal 1 must be treated separately, an equilibrium may require a three-way mixture of two proper u -partitions and a mass point.

7.5 Biased selection rules

Since $v(x) - 0.5$ is odd, every selection rule in this paper treats the agents anonymously: exchanging their signals exchanges their selection probabilities. Anonymity is natural when the agents are ex-ante identical, and it is often required by equal-treatment norms, but it rules out a simple alternative to noise – favoring one of the agents. Head starts of this kind are a standard instrument of contest design (Kirkegaard, 2012), and a designer who wants to select a high-ability contestant can benefit from biasing the contest even between ex-ante identical contestants (Kawamura and Moreno de Barreda, 2014). To compare this instrument with noise, suppose that the firm uses the noiseless rule with a head start $b \in (0, 1)$ for agent 1: agent 1 is selected if and only if $s_1 + b \geq s_2$, so that ties are broken in his favor.⁷ I consider symmetric priors $p \leq 0.5$, since for $p > 0.5$ the well S-shaped rule from Corollary 4 already attains the first best.

Proposition 14. *Let $p_1 = p_2 = p \leq 0.5$, and let agent 1 be selected if and only if $s_1 + b \geq s_2$.*

- (i) *If $b \leq 1 - \sqrt{2p}$, let $M = L$, $c = \frac{2p}{L(2b+L)}$ and $d = \frac{1}{b+L}$, where $L = p + \sqrt{p^2 + 2pb}$.*
- (ii) *If $1 - \sqrt{2p} < b < 1 - p$, let $M = 1 - b$, $c = \frac{2p}{1-b^2}$ and $d = \frac{2(1-b-p)}{1-b^2}$.*

⁷Tie-breaking matters only in case (ii) of Proposition 14: with ties broken equally, no exact equilibrium exists, but ε -equilibria with agent 1's atom just above $1 - b$ converge to it, as does the firm's utility.




Figure 4: Equilibrium disclosures under the biased noiseless rule for $p = b = 0.2$

(iii) If $b \geq 1 - p$, agent 1 is selected with probability one in every equilibrium, and $u_F = p$.

In cases (i) and (ii), the unique equilibrium is: agent 1 sends $s_1 = 0$ with probability bd , $s_1 = M$ with probability $1 - d(b + M)$, which is zero in case (i), and otherwise draws s_1 from $U[0, M]$; agent 2 sends $s_2 = 0$ with probability $1 - cM$ and otherwise draws s_2 from $U[b, b + M]$. The firm's utility is $u_F = p + cdM \left(\frac{M^2}{6} + bM + b^2 \right)$.

The equilibrium keeps the structure of the noiseless equilibrium of Boleslavsky and Cotton (2015): on the range where the agents compete, the signal of agent 1 and the handicapped signal of agent 2, $s_2 - b$, are both uniformly distributed on $[0, M]$, so that each agent's value function is linear on the support of his disclosure. What the head start changes is what lies outside this range, see Figure 4. Agent 2 never sends a signal in $(0, b]$, since such a signal never wins, and concedes bad news instead. In turn, agent 1 concedes bad news too, since the signal $s_1 = 0$ still wins against every concession of agent 2. So, in contrast to the noiseless rule, under which neither agent ever reveals the bad type for $p \leq 0.5$, both agents now reveal it with positive probability. This additional information comes at the cost of selecting agent 1 whenever $s_1 < s_2 \leq s_1 + b$.

Corollary 9. Let $p \leq 0.5$.

- (i) A small head start strictly improves on the noiseless rule: the right derivative of u_F with respect to b at $b = 0$ is positive.
- (ii) As $p \rightarrow 0$, the share of the first best $\frac{u_F}{u_F^*}$ under the optimal head start converges to 1, whereas under every well S-shaped rule it is at most $\frac{u_F^{ws}}{u_F^*}$, which converges to 0.5.

Hence, the information gain dominates for small head starts, and for sufficiently low priors, a biased noiseless rule outperforms every well S-shaped selection rule, and even the best selection rule within the triangular family of Proposition 3, whose share of the first best converges to 0.75. This ranking reverses as p approaches 0.5, where the best noisy selection rules attain the first best and the biased rule does not. Both instruments are weakly dominated by revelation-accurate negatively balanced rules, which attain the first best for all $p < 0.5$ by Proposition 5 but, by Corollary 6, cannot be generated by i.i.d. shocks. The two instruments work through different channels: noise changes the curvature of the agents' value functions, whereas a head start keeps them linear and instead makes a range of signals useless for one of the agents. Whether combining a head start with noise improves further on both is left for future research.

REFERENCES

- Abdulkadiroğlu, A. and T. Sönmez (2003). School choice: a mechanism design approach. *American Economic Review* 93(3), 729–747.
- Au, P. H. and K. Kawai (2020). Competitive information disclosure by multiple senders. *Games and Economic Behavior* 119, 56–78.
- Au, P. H. and M. Whitmeyer (2023). Attraction versus persuasion: Information provision in search markets. *Journal of Political Economy* 131(1), 202–245.
- Ball, I. (2025). Scoring strategic agents. *American Economic Journal: Microeconomics* 17(1), 97–129.

- Bar, T., V. Kadiyali, and A. Zussman (2012). Putting grades in context. *Journal of Labor Economics* 30(2), 445–478.
- Blackwell, D. (1953). Equivalent comparisons of experiments. *The Annals of Mathematical Statistics* 24(2), 265–272.
- Board, S. and J. Lu (2018). Competitive information disclosure in search markets. *Journal of Political Economy* 126(5), 1965–2010.
- Boleslavsky, R. and C. Cotton (2015). Grading standards and education quality. *American Economic Journal: Microeconomics* 7(2), 248–279.
- Boleslavsky, R. and C. Cotton (2018). Limited capacity in project selection: Competition through evidence production. *Economic Theory* 65(2), 385–421.
- Boleslavsky, R. and C. Li (2026). Evaluating university graduates: Prestige versus merit. *Journal of Economics & Management Strategy*. Forthcoming.
- Border, K. C. and J. Sobel (1987). Samurai accountant: a theory of auditing and plunder. *Review of Economic Studies* 54(4), 525–540.
- Chan, W., H. Li, and W. Suen (2007). A signaling theory of grade inflation. *International Economic Review* 48(3), 1065–1090.
- Che, Y.-K. and I. Gale (2000). Difference-form contests and the robustness of all-pay auctions. *Games and Economic Behavior* 30(1), 22–43.
- Drugov, M. and D. Ryvkin (2020). How noise affects effort in tournaments. *Journal of Economic Theory* 188, 105065.
- Ederer, F., R. Holden, and M. Meyer (2018). Gaming and strategic opacity in incentive provision. *RAND Journal of Economics* 49(4), 819–854.

- Erdil, A. and H. Ergin (2008). What's the matter with tie-breaking? Improving efficiency in school choice. *American Economic Review* 98(3), 669–689.
- Fang, D. and T. Noe (2022). Less competition, more meritocracy? *Journal of Labor Economics* 40(3), 669–701.
- Fang, F. C. and A. Casadevall (2016). Research funding: the case for a modified lottery. *mBio* 7(2), e00422–16.
- Frankel, A. and N. Kartik (2022). Improving information from manipulable data. *Journal of the European Economic Association* 20(1), 79–115.
- Fu, Q. and Z. Wu (2019). Contests: Theory and topics. In *Oxford Research Encyclopedia of Economics and Finance*. Oxford University Press.
- Gentzkow, M. and E. Kamenica (2016). A Rothschild-Stiglitz approach to Bayesian persuasion. *American Economic Review* 106(5), 597–601.
- Gentzkow, M. and E. Kamenica (2017). Competition in persuasion. *The Review of Economic Studies* 84(1), 300–322.
- Harbaugh, R. and E. Rasmusen (2018). Coarse grades: Informing the public by withholding information. *American Economic Journal: Microeconomics* 10(1), 210–235.
- He, W. and J. Li (2023). Competitive information disclosure in random search markets. *Games and Economic Behavior* 140, 132–153.
- Heyard, R., M. Ott, G. Salanti, and M. Egger (2022). Rethinking the funding line at the Swiss National Science Foundation: Bayesian ranking and lottery. *Statistics and Public Policy* 9(1), 110–121.
- Hoppe, H. C., B. Moldovanu, and A. Sela (2009). The theory of assortative matching based on costly signals. *Review of Economic Studies* 76(1), 253–281.

- Hvide, H. K. and E. G. Kristiansen (2003). Risk taking in selection contests. *Games and Economic Behavior* 42(1), 172–179.
- Hwang, I., K. Kim, and R. Boleslavsky (2026). Competitive advertising and pricing. *The Review of Economic Studies*. Forthcoming.
- Jain, V. and M. Whitmeyer (2026). Competitive disclosure of information to a rationally inattentive agent. *Journal of Economic Behavior & Organization* 242, 107410.
- Kamenica, E. (2019). Bayesian persuasion and information design. *Annual Review of Economics* 11, 249–272.
- Kamenica, E. and M. Gentzkow (2011). Bayesian persuasion. *American Economic Review* 101(6), 2590–2615.
- Karlin, S. (1959). *Mathematical Methods and Theory in Games, Programming, and Economics*, Vol. II: *The Theory of Infinite Games*. Reading, MA: Addison-Wesley.
- Kawamura, K. and I. Moreno de Barreda (2014). Biasing selection contests with ex-ante identical agents. *Economics Letters* 123(2), 240–243.
- Kirkegaard, R. (2012). Favoritism in asymmetric contests: Head starts and handicaps. *Games and Economic Behavior* 76(1), 226–248.
- Kleiner, A., B. Moldovanu, and P. Strack (2021). Extreme points and majorization: Economic applications. *Econometrica* 89(4), 1557–1593.
- Kolotilin, A., T. Mylovannov, and A. Zapechelnyuk (2022). Censorship as optimal persuasion. *Theoretical Economics* 17(2), 561–585.
- Lazear, E. P. and S. Rosen (1981). Rank-order tournaments as optimum labor contracts. *Journal of Political Economy* 89(5), 841–864.

- Letina, I., S. Liu, and N. Netzer (2023). Optimal contest design: Tuning the heat. *Journal of Economic Theory* 213, 105616.
- Morgan, J., J. Tumlinson, and F. Várdy (2022). The limits of meritocracy. *Journal of Economic Theory* 201, 105414.
- Niculescu, C. P. and L.-E. Persson (2003). Old and new on the Hermite–Hadamard inequality. *Real Analysis Exchange* 29(2), 663–685.
- Ostrovsky, M. and M. Schwarz (2010). Information disclosure and unraveling in matching markets. *American Economic Journal: Microeconomics* 2(2), 34–63.
- Perez-Richet, E. and V. Skreta (2022). Test design under falsification. *Econometrica* 90(3), 1109–1142.
- Pier, E. L., M. Brauer, A. Filut, A. Kaatz, J. Raclaw, M. J. Nathan, C. E. Ford, and M. Carnes (2018). Low agreement among reviewers evaluating the same NIH grant applications. *Proceedings of the National Academy of Sciences* 115(12), 2952–2957.
- Polyanin, A. D. and A. V. Manzhirov (2008). *Handbook of Integral Equations* (Second ed.). Boca Raton, FL: Chapman & Hall/CRC.
- Popov, S. V. and D. Bernhardt (2013). University competition, grading standards, and grade inflation. *Economic Inquiry* 51(3), 1764–1778.
- Prendergast, C. and R. H. Topel (1996). Favoritism in organizations. *Journal of Political Economy* 104(5), 958–978.
- Röbbecke, M. and D. Simon (2023). Riskante Forschung und teilrandomisierte Begutachtungsverfahren: Neue Wege der Förderlinie “Experiment!” der VolkswagenStiftung. *Beiträge zur Hochschulforschung* 45(2), 8–30.

- Sakhnovich, L. A. (2015). *Integral Equations with Difference Kernels on Finite Intervals* (Second ed.), Volume 84 of *Operator Theory: Advances and Applications*. Cham: Birkhäuser.
- Scharfstein, D. S. and J. C. Stein (2000). The dark side of internal capital markets: Divisional rent-seeking and inefficient investment. *The Journal of Finance* 55(6), 2537–2564.
- Stone, P. (2011). *The Luck of the Draw: The Role of Lotteries in Decision Making*. New York: Oxford University Press.
- Tsakas, E. and N. Tsakas (2021). Noisy persuasion. *Games and Economic Behavior* 130, 44–61.
- Tsakas, E., N. Tsakas, and D. Xefteris (2021). Resisting persuasion. *Economic Theory* 72(3), 723–742.
- Wang, Z. (2010). The optimal accuracy level in asymmetric contests. *The B.E. Journal of Theoretical Economics* 10(1), Article 13.

A. PROOFS OF SECTIONS 3–5

A.1 Properties of well S-shaped selection rules

Throughout the proofs, I use the following properties of a well S-shaped $v(x)$.

- 1) *Regularity*. Since $v'(x)$ is concave, it is continuous on $(-1, 1)$ and has one-sided derivatives there; I set $v'(\pm 1)$ equal to its one-sided limits. I write $v''(x)$ and $\tilde{v}''(x)$ for either one-sided derivative, and every statement involving them holds for both.
- 2) *Shape*. Since $v(x)$ is increasing, $v'(1) \geq 0$, and since $v'(x)$ is even, concave and uniquely maximized at 0, $v'(x) \geq (1 - |x|)v'(0) + |x|v'(1) > v'(1) \geq 0$ on $(-1, 1)$, so $v(x)$ is strictly

increasing. For $0 \leq x < y \leq 1$, the same argument gives $v'(x) \geq \frac{y-x}{y}v'(0) + \frac{x}{y}v'(y) > v'(y)$, so $v'(x)$ is strictly decreasing on $[0, 1]$. Since $\tilde{v}'(x) = \frac{v'(x)-v'(0)}{|v(1)-0.5-v'(0)|}$, it is even, concave, and strictly decreasing on $[0, 1]$ with $\tilde{v}'(0) = 0$. So $\tilde{v}(x)$ is negative, strictly decreasing and strictly concave on $(0, 1]$. By the concavity of $\tilde{v}'(x)$, $\tilde{v}''(x) \leq \frac{\tilde{v}'(x)}{x} < 0$ on $(0, 1)$, and $-\tilde{v}''(x)$ is non-decreasing on $(0, 1)$.

3) *Elasticity*. The following lemma bounds the elasticity of $\tilde{v}(x)$ from below.

Lemma 10. *Let $v(x)$ be well S-shaped. Then $\mathcal{E}_{\tilde{v}}(x) \geq 2$ for all $x \in (0, 1]$, with equality at x only if $\tilde{v}'(t)$ is linear on $[0, x]$. Hence $\frac{\tilde{v}(x)}{x^2}$ is non-increasing on $(0, 1]$, so $\tilde{v}(x) \geq -x^2$ on $[0, 1]$; if $v'(x)$ is not linear near 0, the ratio is strictly decreasing.*

Proof. Let $x \in (0, 1]$ and $\phi := -\tilde{v}'$, which is convex on $[0, x]$ with $\phi(0) = 0$ by 2), so $\phi(t) \leq \frac{t}{x}\phi(x)$ for $t \in [0, x]$. Hence $-\tilde{v}(x) = \int_0^x \phi(t)dt \leq \frac{x\phi(x)}{2}$, i.e., $\mathcal{E}_{\tilde{v}}(x) \geq 2$, with equality only if ϕ is linear on $[0, x]$. Finally, $(\frac{\tilde{v}(x)}{x^2})' = \frac{\tilde{v}'(x)(\mathcal{E}_{\tilde{v}}(x)-2)}{x^3} \leq 0$, with strict inequality if $v'(x)$ is not linear near 0, and $\tilde{v}(1) = -1$. $\square$

4) *Non-linearity near 0*. If $v'(x)$ is not linear near 0, then it is not affine on any interval whose closure contains 0: on an interval with 0 in its interior, an affine $v'(x)$ would be constant, contradicting the unique maximum at 0, and an affine piece ending at 0 gives, by evenness, one starting at 0. Hence, for every $G \in \mathcal{G}(p)$, $\bar{v}'_G(s) := \int_0^1 v'(s-t)dG(t)$ is not affine on any interval $[\alpha, \beta]$ with $G([\alpha, \beta]) > 0$. Indeed, $\bar{v}'_G$ is a mixture of the concave functions $v'(s-t)$, so it is affine on $[\alpha, \beta]$ only if $v'(x)$ is affine on $[\alpha-t, \beta-t]$ for G -almost every t , and the closure of this interval contains 0 whenever $t \in [\alpha, \beta]$.

A.2 Proof of Lemma 1

(i) As shown in the text, $\bar{v}'(s_i) = \int_0^1 v'(s_i - s_j)dG_j(s_j)$ is concave on $[0, 1]$. For a maximizer c of $\bar{v}'$, $\bar{v}$ is convex on $[0, c]$ and concave on $[c, 1]$.

(ii) With $slp(0) = \bar{v}'(0)$, slp is continuous on $[0, 1]$, so a exists. For $s \in (0, 1]$, differentiating gives $slp'(s) = \frac{N(s)}{s^2}$, where

$$N(s) := s\bar{v}'(s) - \bar{v}(s) + \bar{v}(0) = \int_0^s (\bar{v}'(s) - \bar{v}'(t)) dt.$$

By (i), $N(s) \geq 0$ on $[0, c]$, and $N(s)$ is non-increasing on $[c, 1]$, since for $c \leq s_1 < s_2 \leq 1$,

$$N(s_2) - N(s_1) = s_1 (\bar{v}'(s_2) - \bar{v}'(s_1)) + \int_{s_1}^{s_2} (\bar{v}'(s_2) - \bar{v}'(t)) dt \leq 0.$$

Hence slp is non-decreasing on $[0, c]$ and changes its monotonicity on $[c, 1]$ at most once, from non-decreasing to non-increasing. So $a \geq c$, and $\bar{v}$ is concave on $[a, 1]$. Let $l(s) := \bar{v}(0) + s \cdot slp(a)$. Then $l \geq \bar{v}$ on $[0, a]$ with equality at 0 and a , and if $a < 1$, $N(a) = 0$, i.e., l is tangent to $\bar{v}$ at a . Therefore $\bar{v}^{cv}$ equals l on $[0, a]$ and $\bar{v}$ on $[a, 1]$. By Kamenica and Gentzkow (2011), a disclosure is optimal if and only if it is supported where $\bar{v}$ touches a line supporting $\bar{v}^{cv}$ at p_i . So $\{0, a\}$ is optimal if $p_i < a$, and $\{p_i\}$ is optimal if $p_i \geq a$, where $\bar{v}^{cv}(p_i) = \bar{v}(p_i)$.

If $v'(x)$ is not linear near 0, the concave $\bar{v}'$ is strictly increasing on $[0, z_1]$, constant on $[z_1, z_2]$ and strictly decreasing on $[z_2, 1]$ for some $z_1 \leq z_2$, and $z_1 > 0$ unless $z_2 = 0$: if $\bar{v}'$ were constant on $[0, z_2]$ with $z_2 > 0$, then $G_j([0, z_2]) = 0$ by property 4) in Appendix A.1, so $\bar{v}''(s) = \int_{z_2}^1 v''(s - s_j) dG_j(s_j) > 0$ on $(0, z_2)$, as $v'' > 0$ on $[-1, 0)$ by property 2), a contradiction. Hence $N > 0$ on $(0, z_2]$ and N is strictly decreasing on $[z_2, 1]$, so $\bar{v} < l$ on $(0, a)$, and $\bar{v}$ is strictly concave on $[a, 1]$. Hence the supporting line at p_i touches $\bar{v}$ only on $\{0, a\}$ if $p_i \leq a$ and only at p_i if $p_i > a$, so the optimum is unique.

A.3 Proof of Proposition 1

Let $\bar{v}(s_i)$ denote the value function of agent i for $\tilde{v}(x)$ rather than $v(x)$, which does not change best replies. If agent j plays $\{0, 1\}$, then $\bar{v}(s_i) = (1 - p_j)\tilde{v}(s_i) - p_j\tilde{v}(1 - s_i)$,

which is S-shaped by Lemma 1, with $\bar{v}(1) - \bar{v}(0) = \tilde{v}(1) = -1$ and $\bar{v}'(1) = (1 - p_j)\tilde{v}'(1)$. By Lemma 1, $\{0, 1\}$ is a best reply if and only if $a = 1$, i.e., if and only if $\bar{v}$ lies below its secant $L(s) := \bar{v}(0) - s$ on $[0, 1]$. This implies $\bar{v}'(1) \geq -1$. Conversely, if $a < 1$, then $\bar{v}$ is concave on $[a, 1]$ by the proof of Lemma 1, so $\bar{v}(a) \leq \bar{v}(1) - (1 - a)\bar{v}'(1)$, while $slp(a) > slp(1)$ gives $\bar{v}(a) > \bar{v}(0) - a$; together, $\bar{v}'(1) < -1$. Hence

$$\{0, 1\} \text{ is a best reply} \Leftrightarrow (1 - p_j)\tilde{v}'(1) \geq -1 \Leftrightarrow p_j \geq 1 - \frac{1}{\mathcal{E}_{\bar{v}}(1)} = \bar{p},$$

and $(\{0, 1\}; \{0, 1\})$ is an equilibrium if and only if $p_1, p_2 \geq \bar{p}$.

For uniqueness, since all equilibria form a product set, it suffices that $\{0, 1\}$ is the unique best reply to $\{0, 1\}$ for both agents. Another best reply exists only if $D := \bar{v} - L \leq 0$ vanishes at some $s_0 \in (0, 1)$. Since D is convex on $[0, c]$ and concave on $[c, 1]$ for some c , it then vanishes on $[s_0, 1]$: a convex D with an interior maximum at $s_0 < c$ vanishes on $[0, c]$, and a concave D that vanishes at both ends of $[\max(s_0, c), 1]$ is non-negative there. Hence $\bar{v}'(1) = -1$, i.e., $p_j = \bar{p}$, and $\bar{v}$ is affine on $[s_0, 1]$, which property 4) in Appendix A.1 rules out if $v'(x)$ is not linear near 0, as $\{0, 1\}$ puts mass at 1. This proves uniqueness if $v'(x)$ is not linear near 0 or $p_1, p_2 > \bar{p}$. For $p_1 = p_2 = \bar{p}$, the uniqueness follows from the uniqueness part of the proof of Theorem 1, which covers $p = \bar{p}$.

A.4 Proof of Theorem 1

If $p \geq \bar{p}$, full disclosure is an equilibrium by Proposition 1, and it is unique for $p > \bar{p}$ by Proposition 1 and for $p = \bar{p}$ by the uniqueness part of this proof below. So let $p < \bar{p}$, and suppose agent j plays a disclosure $\{0, a\}$. Then for i we have

$$\bar{v}(s_i) = \frac{a - p}{a}\tilde{v}(s_i) + \frac{p}{a}\tilde{v}(s_i - a).$$

Then, for $\{0, a\}$ to be a best reply, it should be true that the slope of the secant is equal to the slope of the tangent:

$$\bar{v}'(a) = \frac{a-p}{a} \tilde{v}'(a) = \frac{a-p}{a} \frac{\tilde{v}(a)}{a} + \frac{p}{a} \frac{\tilde{v}(a)}{a},$$

hence $(a-p)\tilde{v}'(a) = \tilde{v}(a)$.

Then, since $\tilde{v}'(a) < 0 \forall a \in (0, 1]$, a should be a solution of the equation $p = a - \frac{\tilde{v}(a)}{\tilde{v}'(a)}$. The RHS is the zero of the tangent $T_a(x) := \tilde{v}(a) + (x-a)\tilde{v}'(a)$, which is strictly decreasing in x , and it lies in $[0, a)$. For $a_1 < a_2$ and all $x \leq a_1$,

$$T_{a_2}(x) - T_{a_1}(x) = \int_{a_1}^{a_2} (\tilde{v}'(t) - \tilde{v}'(a_2)) dt + (a_1 - x) (\tilde{v}'(a_1) - \tilde{v}'(a_2)) > 0,$$

since $\tilde{v}'(x)$ is strictly decreasing on $[0, 1]$. So T_{a_2} is positive at the zero of T_{a_1} , and its own zero lies strictly to the right. Hence the RHS is strictly increasing in a , and the solution is unique and monotone in p , if it exists.

To prove existence, notice that the RHS is continuous in a and its value at $a = 1$ is $\bar{p}$. By the definition of $\mathcal{E}_{\tilde{v}}(x)$ and Lemma 10,

$$a - \frac{\tilde{v}(a)}{\tilde{v}'(a)} = a \left(1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)} \right) \in \left[\frac{a}{2}, a \right),$$

hence the RHS converges to 0 as $a \rightarrow 0_+$. So $\forall p \in (0, \bar{p})$ there always exists a unique monotone solution to this equation, which is exactly the equation from the statement. Moreover, as the inverse of a continuous and strictly increasing function, $a(p)$ is continuous and strictly increasing.

Best reply. Let $p \in (0, \bar{p}]$, where for $p = \bar{p}$, $a = 1$ satisfies $(a-p)\tilde{v}'(a) = \tilde{v}(a)$ by the definition of $\bar{p}$. Let $l(s)$ be the secant from $(0, \bar{v}(0))$ to $(a, \bar{v}(a))$ and $d(s) := l(s) - \bar{v}(s)$,

so that $d(0) = d(a) = 0$, and $d'(a) = 0$ by this equation. As $\tilde{v}'(x)$ is even,

$$d'(s) = \frac{\tilde{v}(a)}{a} - \frac{a-p}{a}\tilde{v}'(s) - \frac{p}{a}\tilde{v}'(a-s),$$

which is convex on $[0, a]$, since $\tilde{v}'(x)$ is concave. By the condition above, $d'(0) = \frac{\tilde{v}(a)}{a} - \frac{p}{a}\tilde{v}'(a) = \frac{a-2p}{a}\tilde{v}'(a) \geq 0$, since $\tilde{v}'(a) < 0$ and $\frac{p}{a} = 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)} \in [\frac{1}{2}, 1)$ by Lemma 10, i.e., $p < a \leq 2p$. So $d' \geq 0$ on some $[0, z]$ and $d' \leq 0$ on $[z, a]$, and hence $d \geq 0$ on $[0, a]$. For $s \in (a, 1]$, both s and $s - a$ lie in $(0, 1]$, where $\tilde{v}'(x)$ is strictly decreasing, so

$$\bar{v}'(s) = \frac{a-p}{a}\tilde{v}'(s) + \frac{p}{a}\tilde{v}'(s-a) < \frac{a-p}{a}\tilde{v}'(a) + \frac{p}{a}\tilde{v}'(0) = \bar{v}'(a),$$

and $\bar{v}(s)$ lies strictly below its tangent $l(s)$ on $(a, 1]$. Hence $l(s)$ supports $\bar{v}(s)$ on $[0, 1]$ and touches it at 0 and $a > p$, so $\{0, a\}$ is a best reply to itself, and every best reply to $\{0, a\}$ is supported on $\{s \in [0, a] : d(s) = 0\}$.

Uniqueness. By the constant-sum property, all equilibria form a symmetric product set $E \times E$ with $\{0, a\} \in E$. Hence, if $G \in E$, then both $(G, \{0, a\})$ and $(\{0, a\}, G)$ are equilibria, i.e., G is a best reply to $\{0, a\}$, and $\{0, a\}$ is a best reply to G .

Step 1: best replies to $\{0, a\}$. Suppose that $d(c) = 0$ for some $c \in (0, a)$. Then c minimizes d on $[0, a]$, so $d'(c) = 0$. By convexity, $d'(s) \leq 0$ on $[c, a]$, and since $\int_c^a d'(s)ds = d(a) - d(c) = 0$, $d' \equiv 0$ on $[c, a]$. Again by convexity, $d'(s) \geq 0$ on $[0, c]$, and since $d(0) = d(c) = 0$, $d' \equiv 0$ on $[0, c]$. Hence $\frac{a-p}{a}\tilde{v}'(s) + \frac{p}{a}\tilde{v}'(a-s)$ is constant on $[0, a]$. A sum of two concave functions is affine only if both are affine, so $\tilde{v}'(s)$ is affine on $[0, a]$, and as $\tilde{v}'(0) = 0 > \tilde{v}'(a)$, $\tilde{v}'(s) = -Ks$ on $[0, a]$ for some $K > 0$. Then

$$\frac{a-p}{a}\tilde{v}'(s) + \frac{p}{a}\tilde{v}'(a-s) = -pK - \frac{K(a-2p)}{a}s$$

is constant only if $a = 2p$.

So, either $d(s) > 0$ on $(0, a)$, in which case $\{0, a\}$ is the only best reply to $\{0, a\}$, and the equilibrium is unique, or $\tilde{v}(x) = -\frac{K}{2}x^2$ on $[0, a]$ and $a = 2p$.

Step 2: the quadratic case. Let $\tilde{v}(x) = -\frac{K}{2}x|x|$ on $[-2p, 2p]$ and $a = 2p$, so that $d \equiv 0$ on $[0, 2p]$ and every G with support in $[0, 2p]$ and mean p is a best reply to $\{0, 2p\}$. Take such a $G \neq \{0, 2p\}$ and let $w(s) := \int_0^{2p} \tilde{v}(s-t)dG(t)$ be the normalized value function against it. For $s \in [0, 2p]$, $w'(s) = -K \int_0^{2p} |s-t|dG(t)$, hence $w'(2p) = -Kp$, while $w(0) = \frac{K}{2}\mathbb{E}_G[t^2]$ and $w(2p) = -\frac{K}{2}\mathbb{E}_G[(2p-t)^2] = -\frac{K}{2}\mathbb{E}_G[t^2]$. So the slope of the secant from $(0, w(0))$ to $(2p, w(2p))$ exceeds $w'(2p)$ by

$$-\frac{K\mathbb{E}_G[t^2]}{2p} + Kp = \frac{K}{2p} \left(2p^2 - \mathbb{E}_G[t^2] \right) > 0,$$

since $t^2 \leq 2pt$ on $[0, 2p]$, with equality only at $t \in \{0, 2p\}$. Hence $w(s)$ lies strictly above the secant for $s < 2p$ close enough to $2p$, so, for such $s > p$, the disclosure $\{0, s\}$ yields a strictly higher payoff against G than $\{0, 2p\}$. Thus $\{0, 2p\}$ is not a best reply to G , $G \notin E$, and the equilibrium is unique.

Let me also introduce a bound on a that is used in later proofs.

Lemma 11. *Let $v(x)$ be well S-shaped and $p \in (0, \bar{p})$. In the symmetric equilibrium $(\{0, a\}; \{0, a\})$, $p < a \leq 2p$, and $a < 2p$ if $v'(x)$ is not linear near 0.*

Proof. Since $\frac{p}{a} = 1 - \frac{1}{\mathcal{E}_{\tilde{v}}(a)}$, and $\mathcal{E}_{\tilde{v}}(a) \geq 2$ by Lemma 10, with strict inequality if $v'(x)$ is not linear near 0, $p < a \leq 2p$, and $a < 2p$ in the latter case. $\square$

A.5 Proof of Lemma 2

Against $\{p_j\}$, $\bar{v}(s) = v(s - p_j)$, so $\bar{v}'(s) = v'(s - p_j)$ is strictly increasing on $[0, p_j]$, since $v'(x)$ is even and strictly decreasing on $[0, 1]$ by property 2) in Appendix A.1. As in the proof of Lemma 1, let $slp(s) := \frac{\bar{v}(s) - \bar{v}(0)}{s}$, so that $slp'(s) = \frac{N(s)}{s^2}$ with $N(s) = \int_0^s (\bar{v}'(s) - \bar{v}'(t))dt$. Then $N > 0$ on $(0, p_j]$, and, by continuity, slp is strictly

increasing on $(0, s]$ for some $s \in (p_j, 1]$. If $p_i \leq p_j$, the disclosure $\{0, s\}$ with mean p_i yields $\bar{v}(0) + p_i slp(s) > \bar{v}(0) + p_i slp(p_i) = \bar{v}(p_i)$, so $\{p_i\}$ is not a best reply. Hence $\{p_i\}$ is a best reply only if $p_i > p_j$, which cannot hold for both agents.

A.6 Proof of Lemma 3

Since $\mathcal{E}_n(x) \rightarrow \infty$ uniformly on $(0, 1)$ and $\mathcal{E}_n(x)$ is continuous on $(0, 1]$, $\varepsilon_n := \sup_{x \in (0, 1]} \frac{1}{\mathcal{E}_n(x)} \rightarrow 0$. Hence $p < \bar{p}_n = 1 - \frac{1}{\mathcal{E}_n(1)}$ for large n , and then, by Theorem 1, $a_n = \frac{p}{1 - 1/\mathcal{E}_n(a_n)} \in \left[p, \frac{p}{1 - \varepsilon_n} \right]$, so $a_n \rightarrow p$.

A.7 Proof of Lemma 4

By Theorem 1, $\frac{\mathbb{P}(s_i=0)}{1-p} = \frac{1}{1 + (\mathcal{E}_{\bar{v}}(a) - 1)(1-a)}$, where, by its proof, $a(p)$ is continuous and strictly increasing, so it maps $(p, p + \varepsilon)$ and $(p - \varepsilon, p)$ onto a right and a left neighbourhood of $a(p)$. Therefore agents disclose more for all $p_h \in (p, p + \varepsilon)$ and less for all $p_l \in (p - \varepsilon, p)$ if and only if $(\mathcal{E}_{\bar{v}}(x) - 1)(1 - x)$ is decreasing at $x = a$.

A.8 Proof of Proposition 2

By Theorem 1 and Proposition 1, the equilibrium is $(\{0, a\}; \{0, a\})$ with $a = a(p) > p$, where $a(p)$ is non-decreasing on $(0, 1)$, as $a(p) = 1$ for $p \geq \bar{p}$. When both agents play $\{0, a\}$ with prior $p < a$, the firm gets

$$U(p, a) := \frac{p^2}{a} + \frac{2p(a-p)}{a}v(a) = p + p\left(1 - \frac{p}{a}\right)(2v(a) - 1).$$

For a fixed p , U is non-decreasing in a , as $1 - \frac{p}{a}$ and $2v(a) - 1$ are non-negative and non-decreasing. For a fixed a , $\partial_p U = 1 + \left(1 - \frac{2p}{a}\right)(2v(a) - 1) > 2(1 - v(a)) \geq 0$, as $1 - \frac{2p}{a} > -1$ and $0 < 2v(a) - 1 \leq 1$. Hence, for $p_1 < p_2$,

$$u_F(p_1) = U(p_1, a(p_1)) \leq U(p_1, a(p_2)) < U(p_2, a(p_2)) = u_F(p_2).$$

A.9 Proof of Theorem 2

Lemma 12. *Let $v(x)$ be well S-shaped and $\hat{v}(x) := v(x) - 0.5$. Then $\hat{v}(x) \leq x(2-x)\hat{v}(1) \leq x - \frac{x^2}{2}$ for all $x \in [0, 1]$, and if $\hat{v}(x) = x - \frac{x^2}{2}$ for some $x \in (0, 1)$, then $v = v^*$.*

Proof. For $0 \leq t < s < 1$, s is a convex combination of t and 1 with weight $\frac{1-s}{1-t}$ on t , so the concavity of $v'(x)$ and $v'(1) \geq 0$ give $v'(s) \geq \frac{1-s}{1-t}v'(t)$. Hence $r(t) := \frac{v'(t)}{1-t}$ is non-decreasing on $[0, 1)$: the density $v'(t)$ has a monotone likelihood ratio relative to the triangular density $1-t$. Let $T(x) := \int_0^x (1-t)dt = x - \frac{x^2}{2}$. For $x \in (0, 1)$,

$$\frac{\hat{v}(x)}{T(x)} = \frac{\int_0^x r(t)(1-t)dt}{\int_0^x (1-t)dt} \leq r(x) \leq \frac{\int_x^1 r(t)(1-t)dt}{\int_x^1 (1-t)dt} = \frac{\hat{v}(1) - \hat{v}(x)}{T(1) - T(x)},$$

so $\hat{v}(x) \leq \frac{T(x)}{T(1)}\hat{v}(1) = x(2-x)\hat{v}(1) \leq x - \frac{x^2}{2}$, as $\hat{v}(1) \leq 0.5$. If $\hat{v}(x) = x - \frac{x^2}{2}$ for some $x \in (0, 1)$, then $\hat{v}(1) = 0.5$ and both inequalities in the display are equalities, so r , which is continuous on $[0, 1)$ by property 1) in Appendix A.1, is constant there and equal to $2\hat{v}(1) = 1$. Hence $v'(t) = 1-t$ on $[0, 1]$, and $v = v^*$, as $v'(x)$ is even. $\square$

By Theorem 1 and Proposition 1, the equilibrium is $(\{0, a\}; \{0, a\})$, where $a = 1$ if and only if $p \geq \bar{p}$, and $p < a \leq a^* := \min(2p, 1)$ by Lemma 11 and $\bar{p} \geq 0.5$ (Lemma 10). As in the proof of Proposition 2, $u_F(p) = p + 2p(1 - \frac{p}{a})\hat{v}(a)$, so, by Lemma 12,

$$u_F(p) \leq p + 2p\left(1 - \frac{p}{a}\right)\left(a - \frac{a^2}{2}\right) \leq p + 2p\left(1 - \frac{p}{a^*}\right)\left(a^* - \frac{a^{*2}}{2}\right) = u_F^{ws}(p),$$

where the second inequality holds as both factors are positive and strictly increasing in a on $(p, 1]$. This proves (i), and equality holds if and only if $a = a^*$ and $\hat{v}(a) = a - \frac{a^2}{2}$.

(ii) If $p \geq 0.5$, then $a^* = 1$, so equality means $a = 1$, i.e., $p \geq \bar{p}$, and $v(1) = 1$.

(iii) If $p < 0.5$, then $a^* = 2p \in (0, 1)$, so equality requires $\hat{v}(2p) = 2p - 2p^2$, hence $v = v^*$ by Lemma 12. Conversely, for v^* , $\tilde{v}(x) = -x^2$ on $[0, 1]$, so $a = 2p$ by Theorem 1,

and both inequalities hold with equality.

A.10 Proof of Corollary 4

(ii) By the Fact in Appendix C, the utility of the firm under the noiseless selection rule is $\frac{4p}{3}$ for $p \leq 0.5$ and $1 - \frac{(1-p)^2(4p-1)}{3p^2}$ for $p > 0.5$. For $p \leq 0.5$, by (i),

$$p + 2p^2(1-p) - \frac{4p}{3} = -\frac{p}{3} (6p^2 - 6p + 1) > 0 \Leftrightarrow p \in \left(\frac{3 - \sqrt{3}}{6}, \frac{3 + \sqrt{3}}{6} \right).$$

For $p > 0.5$, by (i),

$$2p - p^2 - 1 + \frac{(1-p)^2(4p-1)}{3p^2} = \frac{(1-p)^2(4p-1-3p^2)}{3p^2} = \frac{(1-p)^3(3p-1)}{3p^2} > 0.$$

A.11 Proof of Proposition 3

Throughout, G_u is the equilibrium disclosure constructed in Proposition 13, and proper u -partitions are as defined in Subsection 7.4.

(i) Let $p \leq 0.5$, $u < 2p$, $k = \lceil \frac{2p}{u} \rceil \geq 2$ and $r = 2p - (k-1)u \in (0, u]$. The proper u -partition of $[0, 2p]$ has atoms iu with masses $\frac{k-i}{k(k+1)}$ and atoms $iu + r = 2p - (k-1-i)u$ with masses $\frac{i+1}{k(k+1)}$, $i = 0, \dots, k-1$. For fixed k , as u increases, iu moves at rate i and $iu + r$ at rate $-(k-1-i)$, so $\Psi_u(t) := \int_0^t G_u(x)dx$ satisfies $\partial_u \Psi_u(t) = -M(t)$, where $M(t)$ sums rate times mass over the atoms below t . The pair $\{iu, iu + r\}$ contributes $\frac{2i+1-k}{k(k+1)}$ and the atom iu alone $\frac{i(k-i)}{k(k+1)}$, hence $M(t) = 0$ for $t \in (ju, ju + r]$ and $M(t) = \frac{(j+1)(j+1-k)}{k(k+1)} < 0$ for $t \in (ju + r, (j+1)u]$, $j \leq k-2$. So $\Psi_u(t)$ is non-decreasing in u , and strictly increasing for t in a set of positive measure, i.e. G_u is a strict mean-preserving spread of $G_{u'}$ for $u' < u$ with the same k . At $u = \frac{2p}{k}$, $r = u$ and G_u is the uniform distribution on $\{0, \frac{2p}{k}, \dots, 2p\}$, which is also its limit as $u \uparrow \frac{2p}{k}$, where $k+1$ is used and $r \rightarrow 0$; so G_u is continuous in u and the ranking extends to $(0, 2p]$. By

Proposition 13, $G_u = \{0, \min(2p, 1)\}$ if $u \geq 2p$ and $p \leq 0.5$, or $u \geq 2 - 2p$ and $p > 0.5$.

For the example after Proposition 3, let $p \geq \frac{5}{9}$ and $u \in [1 - p, 2 - 2p)$. Then $k = 1$ in case (iii) of the full statement in the proof of Proposition 13, and $a = 2u - \frac{u^2}{1-p} \leq 1 - u$, as $3u - \frac{u^2}{1-p}$ is maximized at $u = \frac{3(1-p)}{2}$, where it equals $\frac{9(1-p)}{4} \leq 1$. The proper u -partition of $[0, a]$ is $\{0, a\}$ with equal probabilities, so G_u puts probabilities $\frac{1-p}{2-a}$, $\frac{1-p}{2-a}$ and $\frac{2p-a}{2-a}$ on $\{0, a, 1\}$. As $\frac{da}{du} = 2 - \frac{2u}{1-p} < 0$ for $u > 1 - p$, a higher u lowers the probability of 0 and raises that of 1, so no two of these disclosures are ranked in the Blackwell order.

(ii) Since $v(-x) = 1 - v(x)$, with $d = s_1 - s_2$,

$$u_F = \mathbb{E}[s_1 v(d) + s_2 v(-d)] = \mathbb{E}[\max(s_1, s_2)] - \mathbb{E}[|d|(1 - v(|d|))],$$

and $v(x) = 1 - \frac{(u-x)^2}{2u^2}$ for $x \in [0, u]$. Let $p \leq 0.5$. For $u \geq 2p$, $G_u = \{0, 2p\}$ and

$$u_F = \frac{p}{2} + p v(2p) = \frac{3p}{2} - \frac{p(u - 2p)^2}{2u^2},$$

which equals $\frac{3p}{2}$ at $u = 2p$ and is strictly decreasing on $[2p, 1]$. For $u < 2p$, $u_F \leq \mathbb{E}[\max(s_1, s_2)] < \frac{3p}{2}$: for $s_2 \in [0, 2p]$, the convex function $\max(\cdot, s_2)$ lies below its chord on $[0, 2p]$, strictly on $(0, 2p)$ if $s_2 \in (0, 2p)$, so $\mathbb{E}_{s_1}[\max(s_1, s_2)] \leq p + \frac{s_2}{2}$, strictly if $s_2 \in (0, 2p)$, as G_u has mass in $(0, 2p)$; averaging over s_2 gives the bound $\frac{3p}{2}$, which is strict for the same reason.

Let $p > 0.5$. For $u \in [2 - 2p, 1]$, $G_u = \{0, 1\}$ and $v(1) = 1$, so $u_F = u_F^*(p)$. For $u < 2 - 2p$, $v(x)$ does not implement a full disclosure equilibrium by the proof of Proposition 13 (ii), so $u_F < u_F^*(p)$ by Lemma 5.

(iii) The proper u -partition of $[0, a]$ with $k = \lceil \frac{a}{u} \rceil$ puts mass $\frac{1}{k}$ on each pair $\{iu, iu + r\} \subset [iu, (i+1)u]$, $i = 0, \dots, k-1$, so it converges weakly to $U[0, a]$ as $u \rightarrow 0$. If $p \leq 0.5$, G_u is the proper u -partition of $[0, 2p]$ and converges to $U[0, 2p]$. If $p > 0.5$ and $u < \min(2p - 1, 2 - 2p)$, then with k from Proposition 13 (iii), we have

$a \in ((k-1)u, ku]$ and $ku < 2-2p \leq (k+1)u$, so $a \in (2-2p-2u, 2-2p)$ and $a < 1-u$. Hence case (iii) applies, $a \rightarrow 2-2p$, the weight $\frac{2(1-p)}{2-a} \rightarrow \frac{1-p}{p}$, and G_u converges to the $(\frac{1-p}{p}, \frac{2p-1}{p})$ mix of $U[0, 2-2p]$ and $\delta_{\{1\}}$ — exactly the distributions from Appendix C.

A.12 Proof of Lemma 5

Since $v(-x) = 1 - v(x)$, for $s_1 \geq s_2$,

$$\begin{aligned} s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) &= s_2 + (s_1 - s_2)v(s_1 - s_2) \leq s_2 + (s_1 - s_2)v(1) \\ &= v(1) \max(s_1, s_2) + (1 - v(1)) \min(s_1, s_2) \leq \max(s_1, s_2), \end{aligned}$$

where the first inequality holds as $v(x)$ is increasing and $s_1 - s_2 \leq 1$, and the second as $v(1) \leq 1$, strictly if $s_1 \neq s_2$ and $v(1) < 1$. Hence $u_F^*(p) \leq \max_{G_1, G_2 \in \mathcal{G}(p)} \mathbb{E}[\max(s_1, s_2)]$. For $s_2 \in [0, 1]$, $\max(s, s_2)$ lies below its secant $s_2 + s(1 - s_2)$ on $[0, 1]$, strictly on $(0, 1)$ if $s_2 \in (0, 1)$. Hence $\mathbb{E}_{G_1}[\max(s_1, s_2)] \leq s_2 + p(1 - s_2)$, with equality for all s_2 if G_1 is full disclosure, and for $s_2 \in (0, 1)$ only then. Averaging over s_2 , $\mathbb{E}[\max(s_1, s_2)] \leq 2p - p^2$, with equality if and only if at least one agent plays full disclosure. If both agents play full disclosure and $v(1) = 1$, then $s_1 - s_2 \in \{-1, 0, 1\}$ and both inequalities above are equalities, so $u_F^*(p) = 2p - p^2$.

By definition, $\bar{u}_F(p) \leq u_F^*(p)$. Let (G_1^*, G_2^*) be an equilibrium with $\mathbb{E}[s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1)] = u_F^*(p)$. Then all inequalities above are equalities, so some agent plays full disclosure, and since then $s_1 \neq s_2$ with positive probability, $v(1) = 1$. As all equilibria form a symmetric product set, $(\{0, 1\}; \{0, 1\})$ is then an equilibrium too. Conversely, if $(\{0, 1\}; \{0, 1\})$ is an equilibrium and $v(1) = 1$, then $s_1 - s_2 \in \{-1, 0, 1\}$ and both inequalities are equalities, so $\bar{u}_F(p) = u_F^*(p)$.

A.13 Proof of Lemma 6

Let agent j with prior p_j play full disclosure. Then $\bar{v}(s_i) - 0.5 = (1 - p_j)\hat{v}(s_i) - p_j\hat{v}(1 - s_i)$ does not depend on p_i , and full disclosure is a best reply of agent i for any $p_i \in (0, 1)$ if and only if $\bar{v}$ lies below its secant from 0 to 1:

$$(1 - p_j)\hat{v}(x) - p_j\hat{v}(1 - x) \leq \hat{v}(1)(x - p_j) \quad \forall x \in (0, 1).$$

Hence $(\{0, 1\}; \{0, 1\})$ is an equilibrium for (p_1, p_2) if and only if this condition holds for $p_j \in \{p_1, p_2\}$, so it is an equilibrium for all $(p_1, p_2) \in S^2$ if and only if the condition holds for all $p_j \in S$, i.e., if and only if it is an equilibrium for all (p, p) , $p \in S$.

A.14 Proof of Proposition 4

Let $a := \hat{v}(1)$; since $v(x) \leq 1$ is increasing, $a \in [0, 0.5]$. By the proof of Lemma 6, $v(x)$ implements a full disclosure equilibrium for p if and only if $(1 - p)\hat{v}(x) - p\hat{v}(1 - x) \leq a(x - p)$ for all $x \in (0, 1)$. Applying it also at $1 - x$ and summing the two inequalities with weights $1 - p$ and p gives $(1 - 2p)(\hat{v}(x) - ax) \leq 0$. Taking $p \in S$ below and above 0.5 yields $\hat{v}(x) = ax$, which proves (ii); then the condition holds with equality for all $p \in (0, 1)$, which proves (i).

A.15 Proof of Proposition 5

(i) Let a and $g(x)$ be as in Definition 6. Then the condition $(1 - p)\hat{v}(x) - p\hat{v}(1 - x) \leq a(x - p) \quad \forall x \in (0, 1)$ from the proof of Lemma 6 simplifies to

$$(1 - p)g(x) - pg(1 - x) \leq 0 \quad \forall x \in (0, 1).$$

Since this condition is linear in p , it holds $\forall p \in [0.5, 1)$ if and only if it holds for

$p = 0.5$ and $p = 1$; the latter means $g(1 - x) \geq 0 \forall x \in (0, 1)$.

$p = 0.5$ implies $0.5g(x) - 0.5g(1 - x) \leq 0 \forall x \in (0, 1)$. Replacing x with $1 - x$, we get that $0.5g(x) - 0.5g(1 - x) \geq 0 \forall x \in (0, 1)$, hence $0.5g(x) = 0.5g(1 - x) \geq 0 \forall x \in (0, 1)$, and since by construction $g(0) = g(1) = 0$, $g(x + 0.5)$ is even on $[-0.5, 0.5]$.

Finally, since $\hat{v}(x)$ should be increasing, on the one hand, for $x < y$, $g(x) + ax \leq g(y) + ay$; on the other hand, for $1 - y < 1 - x$,

$$g(1 - y) + a(1 - y) \leq g(1 - x) + a(1 - x) \Leftrightarrow g(y) - ay \leq g(x) - ax,$$

hence $|g(x) - g(y)| \leq a(y - x) = a|x - y|$, and the converse is immediate.

The proof for $p \in (0, 0.5]$ is done analogously.

(ii) Since all equilibria form a product set, if full disclosure is each agent's unique best reply to full disclosure, it is the unique equilibrium.

The uniqueness of the best reply is equivalent to $(1 - p)\hat{v}(x) - p\hat{v}(1 - x) < a(x - p) \forall x \in (0, 1)$, and it can be rewritten as $(1 - 2p)g(x) < 0 \forall x \in (0, 1)$, if $g(x + 0.5)$ is even. Hence, if $g(x) < 0 \forall x \in (0, 1)$, this condition holds $\forall p < 0.5$, and vice versa.

A.16 Proof of Corollary 6

Suppose that agent j plays $\{0, 1\}$. Then $\bar{v}(s_i) = (1 - p_j)v(s_i) + p_jv(s_i - 1)$, and $\{0, 1\}$ is a best reply of agent i only if the secant from 0 to 1 lies weakly above $\bar{v}(s_i)$ on $[0, 1]$, which requires $\bar{v}'(0) \leq \bar{v}(1) - \bar{v}(0) \leq \bar{v}'(1)$. Since $v'(x)$ is even, $\bar{v}'(0) = (1 - p_j)v'(0) + p_jv'(1)$ and $\bar{v}'(1) = (1 - p_j)v'(1) + p_jv'(0)$, so this requires $(2p_j - 1)(v'(0) - v'(1)) \geq 0$, i.e., $p_j \geq 0.5$. Hence in a full disclosure equilibrium both priors are at least 0.5.

If $v(x)$ is generated by i.i.d. shocks with a square-integrable density f , then $v'(x) = \int f(t)f(t + x)dt$ is continuous, and by the Cauchy-Schwarz inequality,

$$v'(1) = \int f(t)f(t + 1)dt \leq \int f^2(t)dt = v'(0),$$

with equality only if $f(t + 1) = f(t)$ almost everywhere, which is impossible for a probability density. Hence $v'(0) > v'(1)$.

B. PROOFS OF SECTIONS 6 AND 7

B.1 Proof of Proposition 6

If $p_2 \geq \bar{p}$, the claim follows from Proposition 1, so let $p_2 < \bar{p}$.

Preliminaries. For a disclosure of agent j let $N(s) := s\bar{v}'(s) - \bar{v}(s) + \bar{v}(0)$, so that $slp'(s) = N(s)/s^2$. By the proof of Lemma 1 (ii), $N > 0$ on $(0, a)$ and $N < 0$ on $(a, 1]$, with a from Lemma 1. Hence agent i plays $\{p_i\}$ if and only if $N(p_i) \leq 0$, and $\{0, 1\}$ if and only if $N(1) \geq 0$. For $y, c \in [0, 1]$, let

$$A(y) := \tilde{v}(y) - y\tilde{v}'(y) = - \int_0^y x\tilde{v}''(x)dx,$$

$$B(y; c) := y\tilde{v}'(c - y) + \tilde{v}(c - y) - \tilde{v}(c) = - \int_0^y x\tilde{v}''(c - x)dx.$$

Since $-\tilde{v}'' > 0$ on $(0, 1]$, $A(y) > 0$, and $B(y; c) > 0$ for $0 < y \leq c$. If agent j plays $\{0, c\}$, so that $\mathbb{P}(s_j = c) = q := \frac{p_j}{c}$, then $\bar{v}(s) = (1 - q)\tilde{v}(s) + q\tilde{v}(s - c)$ and

$$N(s) = -(1 - q)A(s) + qB(s; c); \tag{2}$$

no disclosure is the case $c = p_j$, $q = 1$.

Step 1: $b(\cdot)$ and p^ .* Against $\{p_1\}$, $N(s) = B(s; p_1)$. Hence agent 2 never plays $\{p_2\}$, as $p_2 \leq p_1$, and plays $\{0, b(p_1)\}$, where $b(p_1)$ solves $B(b; p_1) = 0$, i.e. $b\tilde{v}'(b - p_1) = \tilde{v}(b - p_1) + \tilde{v}(p_1)$, or $b(p_1) = 1$ if $B(1; p_1) \geq 0$. With $b = p_1 + \tilde{b}$ this equation reads

$$\varphi(\tilde{b}) := \tilde{v}(\tilde{b}) - (p_1 + \tilde{b})\tilde{v}'(\tilde{b}) + \tilde{v}(p_1) = 0,$$

where $\varphi'(\tilde{b}) = -(p_1 + \tilde{b})\tilde{v}''(\tilde{b}) > 0$, $\varphi(0) = \tilde{v}(p_1) < 0 < 2A(p_1) = \varphi(p_1)$ and $\partial_{p_1}\varphi = \tilde{v}'(p_1) - \tilde{v}'(\tilde{b}) < 0$ for $\tilde{b} < p_1$. So the root is unique, $p_1 < b(p_1) < 2p_1$, and b is continuous, strictly increasing while $b < 1$. Let p^* be the smallest p_1 with $b(p_1) = 1$, solving $\tilde{v}(1 - p^*) + \tilde{v}(p^*) = \tilde{v}'(1 - p^*)$; it exists and $p^* \in (\frac{1}{2}, 1)$, since $b(\frac{1}{2}) < 1$ and $B(1; 1) = 1 > 0$. Thus against $\{x\}$ the other agent plays $\{0, 1\}$ if and only if $B(1; x) \geq 0$, i.e. $x \geq p^*$, as b is non-decreasing.

Step 2: $a(\cdot)$. Against $\{0, 1\}$, by (2),

$$N(s) = (A(s) + B(s; 1))(p_2 - \Phi(s)), \quad \Phi := \frac{A}{A + B(\cdot; 1)}.$$

Φ is strictly increasing: writing $w = -\tilde{v}''$,

$$\partial_y B(y; 1) A(y) - B(y; 1) A'(y) = y \int_0^y x [w(1 - y)w(x) - w(1 - x)w(y)] dx < 0,$$

since $w(x) \leq w(y)$ and $w(1 - y) \leq w(1 - x)$ for $x < y$, and $w(x) < w(y)$ for x close to 0, as $v'(x)$ is not linear near 0. Moreover, $\Phi(1) = \frac{|\tilde{v}'(1)| - 1}{|\tilde{v}'(1)|} = \bar{p}$. Hence agent 1 plays $\{p_1\}$ if $p_1 \geq a(p_2)$ and $\{0, a(p_2)\}$ otherwise, where $a(p_2) = 0$ if $p_2 \leq \Phi(0^+)$, $a(p_2) = \Phi^{-1}(p_2)$ if $\Phi(0^+) < p_2 < \bar{p}$, and $a(p_2) = 1$ if $p_2 \geq \bar{p}$; a is continuous and non-decreasing, since Φ is continuous and strictly increasing.

Step 3: $\check{p}_2(\cdot)$ and $\hat{p}_1(\cdot)$. By (2), against $\{0, b(p_1)\}$, agent 1 plays $\{p_1\}$ if and only if

$$p_2 \leq \check{p}_2(p_1) := \frac{b(p_1)A(p_1)}{A(p_1) + B(p_1; b(p_1))},$$

and against $\{0, a\}$ with $a > p_1$, agent 2 plays $\{0, 1\}$ if and only if $-(a - p_1)A(1) + p_1B(1; a) \geq 0$. By Step 1, $B(1; a) \geq 0$ if and only if $a \geq p^*$, so for $a < p^*$ this fails, and for $a \geq p^*$ it reads $p_1 \geq \frac{aA(1)}{A(1) + B(1; a)}$. For $a = a(p_2) \geq p^*$ denote this bound by $\hat{p}_1(p_2)$.

Step 4: cases (i)–(iii). A pair of disclosures is an equilibrium if and only if each is

a best reply to the other. By Steps 1 and 2, $(\{p_1\}; \{0, 1\})$ is an equilibrium if and only if $p_1 \geq p^*$ and $p_1 \geq a(p_2)$; by Steps 1 and 3, $(\{p_1\}; \{0, b\})$ with $b < 1$ is an equilibrium if and only if $b = b(p_1)$, $p_1 < p^*$ and $p_2 \leq \check{p}_2(p_1)$; by Steps 2 and 3, $(\{0, a\}; \{0, 1\})$ is an equilibrium if and only if $a = a(p_2) \geq p^*$ and $\hat{p}_1(p_2) \leq p_1 < a(p_2)$. An equilibrium exists by the minimax theorem, and as the game is constant-sum, all equilibria form a product set. By Lemma 1 every best reply is unique, hence so is the equilibrium.

Step 5: case (iv). At $p_1 = p_2 = p$ the game is symmetric, so the unique equilibrium is symmetric and none of the forms in (i)–(iii) is an equilibrium. By Step 4 this gives: $a(p) > p$ if $p \geq p^*$; $\check{p}_2(p) < p$ if $p < p^*$; and $\hat{p}_1(p) > p$ if $a(p) \geq p^*$ and $p < \bar{p}$, since then $p < a(p)$. Now let $p_2 \leq p_1$ and suppose that none of (i)–(iii) holds. By Lemma 1 each agent plays $\{p_i\}$, $\{0, 1\}$ or $\{0, c\}$ with $p_i < c < 1$, and by Lemma 2 not both play no disclosure. Apart from (i)–(iv), this leaves the following forms, and each requires, by Steps 1–3 with the roles of the agents exchanged (the functions b , a , $\check{p}_2$ and $\hat{p}_1$ stay the same, only their arguments change), a condition incompatible with $p_2 \leq p_1$:

- $(\{0, b(p_2)\}; \{p_2\})$ requires $p_2 < p^*$ and $p_1 \leq \check{p}_2(p_2) < p_2$;
- $(\{0, 1\}; \{p_2\})$ requires $p_2 \geq p^*$ and $p_2 \geq a(p_1) \geq a(p_2) > p_2$;
- $(\{0, 1\}; \{0, a(p_1)\})$ with $a(p_1) < 1$ requires $a(p_1) \geq p^*$, $p_1 < \bar{p}$ and $p_2 \geq \hat{p}_1(p_1) > p_1$;
- $(\{0, 1\}; \{0, 1\})$ requires $p_2 \geq \bar{p}$ by Proposition 1.

Hence the equilibrium is $(\{0, a\}; \{0, b\})$ with $p_1 < a < 1$ and $p_2 < b < 1$.

Step 6: $a \leq b$ and continuity. The correspondence $p \mapsto \mathcal{G}(p)$ is continuous: upper hemicontinuous since the mean is weakly continuous, and lower hemicontinuous since any $G \in \mathcal{G}(p)$ is approximated by mixtures of G with δ_0 or δ_1 . As payoffs are continuous, the equilibrium correspondence has a closed graph, and being single-valued with values in a compact set, it is continuous. If $a = b = c$ in (iv), then, since $B(c; c) = -\tilde{v}(c)$, the indifference conditions $N(c) = 0$ of the two agents, with $q = \frac{p_2}{c}$ and

$q = \frac{p_1}{c}$, read $\frac{c}{p_2} - 1 = -\frac{\tilde{v}(c)}{A(c)} = \frac{c}{p_1} - 1$, so $p_1 = p_2$; conversely $a = b$ on the diagonal by symmetry. On the convex set $U = \{0 < p_2 < p_1 < 1, p_2 < \bar{p}\}$, let R be the set where (iv) holds. R is open in U , as the strategies in (iv) are not weak limits of $\{0, 1\}$ or of point masses, and $a - b \neq 0$ is continuous on R . Since U is connected and (i) holds for p_1 close to 1, each connected component of R has a boundary point in $U \setminus R$. There one of (i)–(iii) holds, and continuity gives $(a, b) \rightarrow (p_1, 1), (p_1, b(p_1))$ or $(a(p_2), 1)$ respectively, with $a(p_2) < 1$ as $p_2 < \bar{p}$. Each limit has $a < b$, hence $a < b$ on R .

B.2 Proof of Proposition 7

Describe a disclosure of agent i by its high signal $c_i \in [p_i, 1]$, where $c_i = p_i$ stands for $\{p_i\}$. For a fixed p_i , $\frac{\mathbb{P}(s_i=0)}{1-p_i} = \frac{c_i-p_i}{c_i(1-p_i)}$ is increasing in c_i , so it suffices to show that c_i is non-decreasing in p_j . By Lemma 1, the best reply to c_j is $c_i = \max\{r(c_j, \frac{p_i}{c_j}), p_i\}$, where $r(c, q)$ is the point at which N from (2) changes sign; it is continuous in (c, q) . With $w = -\tilde{v}''$, which is odd and non-decreasing,

$$\partial_q N(s) = A(s) + B(s; c) = \int_0^s x [w(x) + w(c - x)] dx > 0,$$

since $w(x) + w(c - x)$ is at least $w(x) > 0$ if $x \leq c$ and equals $w(x) - w(x - c) \geq 0$ if $x > c$. Hence $r(c, q)$ is non-decreasing in q , and strictly increasing where $r(c, q) \in (0, 1)$, and $r(c, 1) = b(c)$ from Step 1 of the proof of Proposition 6.

Let $\rho(a) := r(a, \frac{p_1}{a})$, so that agent 2's best reply to a is $\max\{\rho(a), p_2\}$, and let $\Gamma(a) := \max\{r(c, \frac{p_2}{c}), p_1\}$ with $c = \max\{\rho(a), p_2\}$, for $a \in [p_1, 1]$. Γ is continuous and its fixed points are exactly the equilibria, so by Step 4 of the proof of Proposition 6 it has a unique fixed point a^* ; since $\Gamma(p_1) \geq p_1$, $\Gamma(a) > a$ on $[p_1, a^*)$. As p_2 increases to p'_2 , $\Gamma(a)$ does not decrease for any a : if $\rho(a) \geq p'_2$, only the weight $\frac{p_2}{\rho(a)}$ increases; if $\rho(a) \leq p_2$, agent 2 pools and $b(p_2) \leq b(p'_2)$; and if $p_2 < \rho(a) < p'_2$, then $r(\rho, \frac{p_2}{\rho}) \leq r(\rho, 1) = b(\rho) \leq b(p'_2)$. Hence the fixed point does not decrease, and if

$p_1 < a^* < 1$, $\Gamma(a^*)$ strictly increases, so the fixed point strictly increases. The argument does not use $p_2 \leq p_1$, so it applies to both agents.

B.3 Proof of Proposition 8

Since $v(-x) = 1 - v(x)$, $s_1 v(s_1 - s_2) + s_2 v(s_2 - s_1) = \frac{s_1 + s_2}{2} + H(s_1 - s_2)$, where $H(x) := x \hat{v}(x)$ is even, so the firm's expected utility is

$$u_F = \frac{p_1 + p_2}{2} + \mathbb{E}[H(s_1 - s_2)].$$

For $x \in (0, 1]$, $0 < h(x) := H'(x) = \hat{v}(x) + x v'(x) \leq 2\hat{v}(x) \leq 2x$, since $v'(x)$ is decreasing on $[0, 1]$ and $v'(0) \leq 1$, as $0.5 \geq \hat{v}(1) \geq \frac{v'(0)}{2}$ by the concavity of $v'(x)$. Also, $H(x) \leq \frac{x}{2}$.

Full disclosure. $u_F = \frac{p_1 + p_2}{2} + H(1)(p_1 + p_2 - 2p_1 p_2)$, so $\partial_{p_1} u_F = 0.5 + H(1)(1 - 2p_2) > 0.5 - H(1) \geq 0$, and similarly for p_2 .

Region (i). $u_F = \frac{p_1 + p_2}{2} + (1 - p_2)H(p_1) + p_2 H(1 - p_1)$, so $\partial_{p_2} u_F = 0.5 - H(p_1) + H(1 - p_1) > 0$ and, as $p_2 \leq p_1$ and $h(x) \leq 2x$,

$$\partial_{p_1} u_F = 0.5 + (1 - p_2)h(p_1) - p_2 h(1 - p_1) > 0.5 - 2p_1(1 - p_1) \geq 0.$$

Region (ii). Let $b = b(p_1)$ and $y := b - p_1$. By Step 1 of the proof of Proposition 6, $0 < y < p_1$ and $y' = -\frac{\partial_{p_1} \varphi}{\varphi'} > 0$. Then

$$u_F = \frac{p_1 + p_2}{2} + H(p_1) - \frac{p_2}{b}(H(p_1) - H(y)),$$

where $0 < H(p_1) - H(y) \leq \frac{p_1}{2} < \frac{b}{2}$. Hence $\partial_{p_2} u_F > 0$ and

$$\partial_{p_1} u_F = 0.5 + \left(1 - \frac{p_2}{b}\right)h(p_1) + \frac{p_2}{b}h(y)y' + \frac{p_2(1 + y')}{b^2}(H(p_1) - H(y)) > 0.$$

Region (iii). Let $a = a(p_2)$, which is non-decreasing by Step 2 of the proof of

Proposition 6, with $p_2 \leq p_1 < a$ and $a \geq p^* > 0.5$. Then

$$u_F = \frac{p_1 + p_2}{2} + p_2 H(1) + \frac{p_1}{a} K, \quad \text{where} \quad K := (1 - p_2)H(a) + p_2(H(1 - a) - H(1)),$$

so $\partial_{p_1} u_F = 0.5 + \frac{K}{a} \geq 0.5 - \frac{p_2 H(1)}{a} > 0$ and, with $K_a := \partial_a K$,

$$\partial_{p_2} u_F = 0.5 + H(1) + \frac{p_1}{a} (H(1 - a) - H(1) - H(a)) + \frac{p_1 a'}{a^2} (a K_a - K).$$

The first three terms sum to at least $0.5 - H(a) > 0$, and

$$a K_a - K = (1 - p_2) a^2 v'(a) + p_2 (\hat{v}(1) - \hat{v}(1 - a) - a(1 - a) v'(1 - a)) \geq 0,$$

since, by the concavity of $v'(x)$ and $a \geq 0.5$,

$$\hat{v}(1) - \hat{v}(1 - a) = \int_{1-a}^1 v'(x) dx \geq \frac{a}{2} v'(1 - a) \geq a(1 - a) v'(1 - a).$$

Example 9. Both disclosures solve the indifference conditions $N = 0$ from (2), which I solved numerically. For instance, for $p_2 = 0.84, 0.86, 0.873$, $(a, b) \approx (0.9375, 0.9996)$, $(0.9406, 0.9925)$ and $(0.9435, 0.9876)$, and $u_F \approx 0.96145, 0.96092$ and 0.96076 .

B.4 Proof of Lemma 7

Write $u_i(p_i, p_j)$ for agent i 's equilibrium payoff when his prior is p_i and agent j 's prior is p_j . For $G_j \in \mathcal{G}(p_j)$, let $\bar{v}_{G_j}(x)$ be agent i 's value function against G_j and $\bar{v}_{G_j}^{cv}$ its concavification on $[0, 1]$. By Kamenica and Gentzkow (2011), $\bar{v}_{G_j}^{cv}(p_i) = \max_{H \in \mathcal{G}(p_i)} \mathbb{E}_H[\bar{v}_{G_j}]$ is agent i 's best payoff against G_j , where the maximum is attained as $\bar{v}_{G_j}$ is continuous.

The game is constant-sum, so agent j 's equilibrium disclosure G_j^* minimizes i 's payoff:

$$u_i(p_i, p_j) = \min_{G_j \in \mathcal{G}(p_j)} \bar{v}_{G_j}^{cv}(p_i) = \bar{v}_{G_j^*}^{cv}(p_i).$$

Step 1: each $\bar{v}_{G_j}^{cv}$ is non-decreasing on $[0, 1]$, and strictly increasing if $v(x)$ is. As $v(x)$ is increasing, so is $\bar{v}_{G_j}$, hence $\bar{v}_{G_j} \leq \bar{v}_{G_j}(1)$ on $[0, 1]$. The constant $\bar{v}_{G_j}(1)$ is then a concave majorant of $\bar{v}_{G_j}$, so $\bar{v}_{G_j}^{cv} \leq \bar{v}_{G_j}(1)$, while $\bar{v}_{G_j}^{cv}(1) = \bar{v}_{G_j}(1)$, since $\{1\}$ is the only disclosure with mean 1. So the concave function $\bar{v}_{G_j}^{cv}$ attains its maximum at $x = 1$. For $0 \leq x < y \leq 1$, write $y = \lambda x + (1 - \lambda) \cdot 1$ with $\lambda = \frac{1-y}{1-x} \in [0, 1)$; by concavity,

$$\bar{v}_{G_j}^{cv}(y) \geq \lambda \bar{v}_{G_j}^{cv}(x) + (1 - \lambda) \bar{v}_{G_j}^{cv}(1) \geq \bar{v}_{G_j}^{cv}(x).$$

If $v(x)$ is strictly increasing, so is $\bar{v}_{G_j}$. Then every $H \in \mathcal{G}(x)$ with $x < 1$ puts positive mass on $[0, 1)$, where $\bar{v}_{G_j} < \bar{v}_{G_j}(1)$, so $\bar{v}_{G_j}^{cv}(x) = \max_{H \in \mathcal{G}(x)} \mathbb{E}_H[\bar{v}_{G_j}] < \bar{v}_{G_j}(1) = \bar{v}_{G_j}^{cv}(1)$, and the last inequality is strict.

Step 2: monotonicity of u_i . Let $p_i < p'_i$, and let G_j^* be agent j 's equilibrium disclosure for priors (p'_i, p_j) . By Step 1,

$$u_i(p_i, p_j) \leq \bar{v}_{G_j^*}^{cv}(p_i) \leq (<) \bar{v}_{G_j^*}^{cv}(p'_i) = u_i(p'_i, p_j).$$

Finally, since $v(x) + v(-x) = 1$, $u_i(p_i, p_j) = 1 - u_j(p_j, p_i)$, so, by the above applied to agent j , u_i is non-increasing (strictly decreasing) in p_j .

B.5 Proof of Proposition 9

For $t \in (0, 1)$, let G_t be the upper censorship with cutoff t , which pools the types above t at $m(t) := \mathbb{E}[\theta | \theta \geq t] > t$, and let

$$w_t(x) := \int_0^t \tilde{v}(x-s)f(s)ds + (1-F(t))\tilde{v}(x-m(t))$$

be the normalized value function against G_t , which has the same best replies as the value function itself. Let L_t be the tangent to w_t at $m(t)$. Since $\tilde{v}$ is odd with $\tilde{v}'(0) = 0$, the left-hand side of (1) at $\theta^* = t$ equals $E(t) := L_t(t) - w_t(t)$. So (1) is the first-order condition of Kolotilin et al. (2022) for the cutoff of an upper censorship against G_t , evaluated at the cutoff t itself; Steps 2 and 3 verify directly that its solutions are equilibria, which avoids any regularity condition on f .

Step 1: (1) has a solution. E is continuous on $(0, 1)$, and $E(t) \rightarrow \tilde{v}(\mathbb{E}[\theta]) < 0$ as $t \rightarrow 0$. Let $t \geq \frac{3}{4}$ and $m = m(t)$. For $s \leq t$, the integrand in (1) is non-negative by the concavity of $\tilde{v}$ on $[0, 1]$, and for $s \leq \frac{1}{2}$ it is at least $\frac{\kappa}{2}(m-t)^2$, where $\kappa := -\tilde{v}'(\frac{1}{4}) > 0$, since $\tilde{v}''(x) \leq \frac{\tilde{v}'(x)}{x} \leq -\kappa$ on $[\frac{1}{4}, 1]$ by property 2) in Appendix A.1. As $\tilde{v}(m-t) \geq -(m-t)^2$ by Lemma 10, the last term of $E(t)$ is at least $-(m-t)^2(1-F(t))$, so

$$E(t) \geq (m-t)^2 \left(\frac{\kappa}{2} F\left(\frac{1}{2}\right) - (1-F(t)) \right) > 0$$

for t close to 1. Hence E has a zero in $(0, 1)$.

Step 2: every solution gives an equilibrium. Let $E(t) = 0$, $m = m(t)$, $w = w_t$, $L = L_t$ and $D := L - w$, so that $D(t) = D(m) = D'(m) = 0$ and $D' = w'(m) - w'$, where $w'(x) = \int \tilde{v}'(x-s)dG_t(s)$ is concave, as a mixture of concave functions. If $w'(x) > w'(m)$ for some $x > m$, then $w' < w'(m)$ on $[0, m)$ by concavity, so D is strictly increasing on $[0, m]$, contradicting $D(t) = D(m)$. Hence D is non-decreasing, and

non-negative, on $[m, 1]$. The set of $x \leq m$ with $w'(x) \geq w'(m)$ is an interval $[\alpha, m]$, so D is non-decreasing on $[t, \alpha]$ and non-increasing on $[\max(\alpha, t), m]$, and $D(t) = D(m) = 0$ gives $D \geq 0$ on $[t, m]$. If $\alpha > t$, then $w'(t) < w'(\alpha)$, so w' is increasing on $[0, t]$ and $w'(t) < w'(m)$. If $\alpha \leq t$, then $D = 0$ on $[t, m]$, so w' is constant on $[t, m]$, hence maximal there and non-decreasing on $[0, t]$. In both cases

$$p(x) := \begin{cases} w(x), & x \leq t, \\ L(x), & x \geq t, \end{cases}$$

is convex, $p \geq w$, and $p = w$ on $[0, t] \cup \{m\} \supseteq \text{supp}(G_t)$. Since p is affine on $[t, 1]$ and G_t pools $[t, 1]$ at its conditional mean, $\int p \, dG_t = \int p \, dF$. So every mean-preserving contraction H of F satisfies

$$\int w \, dH \leq \int p \, dH \leq \int p \, dF = \int p \, dG_t = \int w \, dG_t,$$

i.e., G_t is a best reply to itself, and (G_t, G_t) is an equilibrium.

Step 3: uniqueness. Now suppose that $v'(x)$ is not linear near 0. By property 4) in Appendix A.1, w' is affine on no interval that G_t charges. Hence $\alpha > t$ in Step 2, as otherwise w' is constant on $[t, m]$; $D > 0$ on $(t, 1] \setminus \{m\}$, as otherwise D , and hence $w' - w'(m)$, vanishes on an interval with endpoint m ; and w' is strictly increasing on $[0, t]$, so p is strictly convex on $[0, t]$ with a kink at t . Let H be a best reply to G_t , so that both inequalities above are equalities. The first one gives $\text{supp}(H) \subseteq [0, t] \cup \{m\}$. Writing $p(x) = p(0) + p'(0)x + \int_0^1 (x - y)^+ dp'(y)$, the second one gives $\int_0^1 (\Psi_F(y) - \Psi_H(y)) dp'(y) = 0$, where $\Psi_K(y) := \int_0^y K(x) dx \leq \Psi_F(y)$ for every mean-preserving contraction K of F . As dp' charges every subinterval of $[0, t]$, $\Psi_H = \Psi_F$ on $[0, t]$, so $H = F$ on $[0, t)$. The remaining mass $1 - F(t)$ of H lies on $\{t, m\}$ with mean m , so it is all at m : $H = G_t$. So G_t is the unique best reply to itself. As

all equilibria form a product set, every equilibrium strategy is a best reply to G_t , so (G_t, G_t) is the unique equilibrium, and by Step 2, (1) has no other solution.

B.6 Proof of Corollary 7

Here $\tilde{v}(x) = -x|x|$, so for $0 \leq s \leq \theta^* \leq m^*$, the integrand in (1) is

$$-(m^* - s)^2 + (\theta^* - s)^2 + 2(m^* - \theta^*)(m^* - s) = (m^* - \theta^*)^2,$$

and (1) becomes $(m^* - \theta^*)^2(2F(\theta^*) - 1) = 0$. Since $m^* > \theta^*$, $F(\theta^*) = 0.5$.

The same computation, with $dF(s)$ in place of $f(s)ds$, applies to any prior F that is continuous at a median θ^* , and Step 2 of the proof of Proposition 9 does not use the density of F . Hence, for every such prior, the median censorship is a symmetric equilibrium under $v(x) = 0.5 + x - 0.5x|x|$.

B.7 A convexity lemma

The following lemma extends the converse of the Hermite–Hadamard inequality (see Niculescu and Persson (2003)), which is the case of a uniform K .

Lemma 13. *Let a CDF K and a function h be continuous on an interval J , with K strictly increasing. If $h(\mathbb{E}_K[\theta|\theta \in J']) \leq \mathbb{E}_K[h(\theta)|\theta \in J']$ for every interval $J' \subseteq J$, then h is convex.*

Proof. Let $x < z$ in J , let ℓ be the secant of h through x and z , and suppose that $g := h - \ell$ has a positive maximum on $[x, z]$, first attained at $c \in (x, z)$. For $\delta > 0$ small enough that $\mathbb{E}_K[\theta|\theta \in [c - \delta, z]] > c$, there is $\delta' \in (0, z - c)$ such that $J' := [c - \delta, c + \delta']$ satisfies $\mathbb{E}_K[\theta|\theta \in J'] = c$, as K is continuous and strictly increasing. Since $g < g(c)$ on $[c - \delta, c)$, which has positive K -mass, and ℓ is affine, $\mathbb{E}_K[h(\theta)|\theta \in J'] = \ell(c) + \mathbb{E}_K[g(\theta)|\theta \in J'] < h(c)$, a contradiction. Hence $h \leq \ell$ on $[x, z]$. $\square$

B.8 Proof of Proposition 10

Let $I = [\underline{\theta}, \bar{\theta}]$. For a CDF K let $\Psi_K(y) := \int_{\underline{\theta}}^y K(x)dx$, so that a disclosure G is feasible if and only if $\Psi_G \leq \Psi_F$ on I with equality at $\bar{\theta}$, and for convex ϕ and feasible G ,

$$\int \phi dF - \int \phi dG = \int (\Psi_F(y) - \Psi_G(y)) d\phi'(y) \geq 0.$$

Against an atomless disclosure H , the payoff of a signal s is $H(s)$. In particular, against full disclosure, a deviation to G yields $\mathbb{E}_G[F]$, and full disclosure yields $\mathbb{E}_F[F] = 0.5$.

Sufficiency. If F is convex on I , the display with $\phi = F$ gives $\mathbb{E}_G[F] \leq \mathbb{E}_F[F] = 0.5$ for every feasible G , so (F, F) is an equilibrium.

Necessity. Pooling the types in an interval $J \subseteq I$ into their mean m_J is feasible and changes the deviator's payoff by $\mathbb{P}(\theta \in J)(F(m_J) - \mathbb{E}_F[F(\theta)|\theta \in J])$, so in a full disclosure equilibrium $F(m_J) \leq \mathbb{E}_F[F(\theta)|\theta \in J]$ for every J . First, F is strictly increasing on I : otherwise $F = \gamma \in (0, 1)$ on a maximal interval $[L, R] \subset I$ with $L < R$, and for $J = \{\theta : \gamma - r \leq F(\theta) \leq \gamma + \frac{r}{2}\}$, $\mathbb{E}_F[F(\theta)|\theta \in J] = \gamma - \frac{r}{4}$, while $m_J \rightarrow \frac{2L+R}{3} > L$ as $r \rightarrow 0$, so $F(m_J) \geq \gamma$ for small r . Then F is convex on I by Lemma 13 with $K = h = F$.

Uniqueness. Let F be convex, so that it is strictly increasing on I , and let (F, H) be an equilibrium. As all equilibria form a symmetric product set, (H, H) and (H, F) are equilibria too, and each agent gets 0.5 in all of them. Against H , the payoff of a signal s is $H(s^-) + \frac{1}{2}(H(s) - H(s^-))$. First, H is atomless. It has no atom at $\underline{\theta}$ or $\bar{\theta}$, as $\Psi_H \leq \Psi_F$ with equality at $\bar{\theta}$ and F is atomless. If H had an atom $a > 0$ at $m \in (\underline{\theta}, \bar{\theta})$, then $\Psi_H(m) < \Psi_F(m)$, since otherwise $\Psi_F - \Psi_H \geq 0$ would have a minimum at m with left derivative $F(m) - H(m^-)$ above its right derivative $F(m) - H(m)$. So replacing the atom by masses $\frac{2a}{3}$ at $m + \varepsilon$ and $\frac{a}{3}$ at $m - 2\varepsilon$ is feasible for small ε , as $\Psi_F - \Psi_H > 0$ near m , and against H it raises the payoff by $a^2(\frac{2}{3} - \frac{1}{2}) + o(1) > 0$, contradicting that H is a best reply to H . Second, H is convex on I : as F is a best reply to H , pooling the

types in an interval $J \subseteq I$ does not pay, i.e., $H(m_J) \leq \mathbb{E}_F[H(\theta)|\theta \in J]$, and Lemma 13 applies with $K = F$ and $h = H$. Finally, as F and H are convex, $\int F dH \leq \int F dF$ and $\int H dH \leq \int H dF$, and all four integrals equal 0.5. By the identity above, $\Psi_F = \Psi_H$ on the supports of dF' and dH' , and there $F = H$, since each such point in $(\underline{\theta}, \bar{\theta})$ is a minimum of the differentiable function $\Psi_F - \Psi_H \geq 0$, while $F = H$ at $\underline{\theta}$ and $\bar{\theta}$. On each open interval off these supports, F and H are affine and agree at the endpoints. Hence $H = F$, and (F, F) is the unique equilibrium.

B.9 Proof of Corollary 8

Let $d = s_1 - s_2$. For any selection rule and feasible G_1, G_2 , since $v(-x) = 1 - v(x)$,

$$u_F = \mathbb{E}[\max(s_1, s_2)] - \mathbb{E}[|d|(1 - v(|d|))] \leq \mathbb{E}[\max(s_1, s_2)] \leq \mathbb{E}[\max(\theta_1, \theta_2)],$$

where, as $\max(s, s') = s' + (s - s')^+$, the last gap equals $\mathbb{E}_F[(\Psi_F - \Psi_{G_2})(\theta)] + \mathbb{E}_{G_2}[(\Psi_F - \Psi_{G_1})(s)]$, with Ψ as in the proof of Proposition 10. The first term vanishes only if $\Psi_{G_2} = \Psi_F$ on $\text{supp}(F)$. Then $G_2 = F$: each point of $\text{supp}(F)$ in $(\underline{\theta}, \bar{\theta})$ is a minimum of $\Psi_F - \Psi_{G_2} \geq 0$, so $G_2 = F$ there, G_2 has no atom at $\underline{\theta}$, and on each gap of $\text{supp}(F)$, G_2 is constant, as it equals F at both ends. Likewise, the second term then vanishes only if $G_1 = F$. So the first best requires full disclosure by both agents, and then also $|d|(1 - v(|d|)) = 0$ almost surely. As F is atomless, $0 < |d| < \varepsilon$ with positive probability for every $\varepsilon > 0$, so $v(x) = 1$ for arbitrarily small $x > 0$, i.e., $v(x)$ is the noiseless selection rule, as it is increasing and $v(x) \leq 1$. By Proposition 10, (F, F) is then an equilibrium if and only if F is convex, and it attains the first best. Finally, a well S-shaped $v(x)$ is strictly increasing, so it is not the noiseless rule.

B.10 Proof of Proposition 11

Let $c = 2p - \underline{\theta}$ and $G = U[\underline{\theta}, c]$. As F is concave, it lies above the uniform CDF on $[\underline{\theta}, \bar{\theta}]$, so $p \leq \frac{\underline{\theta} + \bar{\theta}}{2}$, i.e. $c \leq \bar{\theta}$. Moreover, $F - G$ is concave on $[\underline{\theta}, c]$ and non-negative at $\underline{\theta}$, while $F - G = F - 1 \leq 0$ on $[c, \bar{\theta}]$, so $F - G$ changes sign at most once, from positive to negative. Since F and G have the same mean, G is a mean-preserving contraction of F by the single-crossing condition, i.e. it is feasible. Against G , $\bar{v}(s) = G(s)$, which is concave on $[\underline{\theta}, \bar{\theta}]$, so by Jensen's inequality every feasible H gives $\mathbb{E}_H[G] \leq G(p) = 0.5 = \mathbb{E}_G[G]$, and (G, G) is an equilibrium. Finally,

$$u_F = \mathbb{E}[\max(s_1, s_2)] = \underline{\theta} + \frac{2}{3}(c - \underline{\theta}) = \frac{4p - \underline{\theta}}{3}.$$

For uniqueness, let (G, H) be an equilibrium, so that (H, H) and (H, G) are equilibria too. As in the proof of Proposition 10, H has no atom in $(\underline{\theta}, \bar{\theta}]$, and an atom $a > 0$ at $\underline{\theta}$ would not be a best reply to H either: pooling it with a small mass ε of H above $\underline{\theta}$ raises the payoff per unit of the atom from $\frac{a}{2}$ to at least a , at a total cost of at most $\varepsilon < \frac{a^2}{2}$. So H is atomless. As H is a best reply to G and $G(s) = \min\{\frac{s - \underline{\theta}}{c - \underline{\theta}}, 1\}$ is concave, $\mathbb{E}_H[G] \leq G(p) = 0.5$ with equality only if $\text{supp}(H) \subseteq [\underline{\theta}, c]$. As G is a best reply to H and every mean-preserving contraction of G is feasible, pooling the types of G in any interval $J \subseteq [\underline{\theta}, c]$ does not pay, so H is convex on $[\underline{\theta}, c]$ by Lemma 13 with $K = G$. Hence $H \leq G$ on $[\underline{\theta}, c]$, as G is the secant of H there, and $\int_{\underline{\theta}}^c (G - H)dx = \mathbb{E}_H[s] - \mathbb{E}_G[s] = 0$, so $H = G$.

B.11 Proof of Lemma 8

(i) Consider $v_1(x) = 0.5 + 0.5x - a\tilde{v}_2(x)$. The convexity of $\tilde{v}_2'(x)$ and the uniqueness of its minimum at 0 are simply inverted into the concavity of $v_1'(x)$ and the uniqueness of its maximum at 0, and $v_1(x) - 0.5$ is trivially odd and increasing for sufficiently small

a , since $\tilde{v}'_2(x)$ is bounded. So $v_1(x)$ is well S-shaped for such a , while

$$\tilde{v}_1(x) = \frac{0.5x - a\tilde{v}_2(x) - x(0.5 - a\tilde{v}'_2(0))}{|0.5 - a - (0.5 - a\tilde{v}'_2(0))|} = -\tilde{v}_2(x).$$

(ii) Since a normalization changes payoffs by a positive factor and terms independent of the disclosures, $v_k(x)$ and $\tilde{v}_k(x)$ have the same equilibria, $k \in \{1, 2\}$. The map $G \mapsto \check{G}$, where $\check{G}$ is the distribution of $1 - s$ for $s \sim G$, is a bijection between $\mathcal{G}(p)$ and $\mathcal{G}(1 - p)$. Since $\tilde{v}_1(x)$ is odd, $\tilde{v}_2((1 - s_i) - (1 - s_j)) = -\tilde{v}_1(s_j - s_i) = \tilde{v}_1(s_i - s_j)$, so each agent's payoff from $(\check{H}_1, \check{H}_2)$ under $\tilde{v}_2$ equals that from (H_1, H_2) under $\tilde{v}_1$ for all H_1, H_2 . Hence $(\check{G}_1, \check{G}_2)$ is an equilibrium for $\tilde{v}_2$ and priors $(1 - p_1, 1 - p_2)$.

B.12 Proof of Proposition 12

Let $G_j = \{0, c\}$ with $c \in (p_j, 1]$, so that $\mathbb{P}(s_j = c) = q := \frac{p_j}{c} \in (0, 1)$. Since $v''(x)$ is odd, $\bar{v}''(s) = (1 - q)v''(s) + qv''(s - c) = -(1 - q)w(s) - qw(s - c) < 0$ on $(c, 1]$, and on $(0, c)$,

$$\bar{v}''(s) = -(1 - q)w(s) + qw(c - s) = w(s) \left(q \frac{w(c - s)}{w(s)} - (1 - q) \right),$$

which, as the ratio is strictly decreasing, is first positive and then negative, with at most one zero. If $G_j = \{p_j\}$, then $\bar{v}''(s) = v''(s - p_j)$ is positive on $[0, p_j)$ and negative on $(p_j, 1]$. In both cases, there is $c^* \in [0, 1]$ such that $\bar{v}$ is strictly convex on $[0, c^*]$ and strictly concave on $[c^*, 1]$, so, as in the proof of Lemma 1, agent i 's best reply is unique and is either $\{p_i\}$ or $\{0, a_i\}$ with $a_i \in (p_i, 1]$.

Now restrict each agent i to these disclosures, indexed by $a_i \in [p_i, 1]$, where $a_i = p_i$ stands for $\{p_i\}$. Payoffs are continuous in (a_1, a_2) , and, by the above, the best reply within this family is unique and is also a best reply among all disclosures. By the maximum theorem, each agent's best reply is continuous in the other agent's index, so Brouwer's theorem gives a pair of mutual best replies, which is an equilibrium. Since

all equilibria form a product set and each agent's best reply to the other's equilibrium disclosure is unique, this equilibrium is unique.

B.13 Proof of Lemma 9

Since $v(x)$ is real-analytic on $[-1, 1]$, it extends analytically to $\{z \in \mathbb{C} : \text{dist}(z, [-1, 1]) < \delta\}$ for some $\delta > 0$, hence $\bar{v}(s_i) = \int_0^1 v(s_i - s_j) dG_j(s_j)$ extends analytically to $\{z \in \mathbb{C} : \text{dist}(z, [0, 1]) < \delta\}$. Let $l(s_i)$ be a line supporting $\bar{v}^{cv}(s_i)$ at p_i . Then $l(s_i) \geq \bar{v}(s_i)$ on $[0, 1]$, and every optimal disclosure is supported on $\{s_i \in [0, 1] : \bar{v}(s_i) = l(s_i)\}$. If this set were infinite, it would have an accumulation point in $[0, 1]$, and by the identity theorem $\bar{v}(s_i) = l(s_i)$ on $[0, 1]$, so $\bar{v}''(0) = 0$. However, $\bar{v}''(0) = \int_0^1 v''(-s_j) dG_j(s_j) > 0$, since $v''(-s_j) > 0$ for $s_j \in (0, 1]$ and G_j puts positive mass on $(0, 1]$ as $p_j > 0$.

B.14 Proof of Proposition 13

Full statement: let $v(x)$ be the CDF of a triangular distribution on $[-u, u]$; then there exists a symmetric equilibrium in which each agent's disclosure G is:

- (i) Proper u -partition of $[0, 2p]$ if $p \leq 0.5$
- (ii) $\{0, 1\}$, if $p > 0.5, u \geq 2 - 2p$.
- (iii) $(\frac{2(1-p)}{2-a}, \frac{2p-a}{2-a})$ mix of the proper u -partition of $[0, a]$ with $\delta_{\{1\}}$, if (i) $p > 0.5, u < 2 - 2p$, and (ii) $a \leq 1 - u$, where $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}, k = \lceil \frac{2-2p}{u} \rceil - 1$.
- (iv) If, in the solution of the equations in (iii), $a > 1 - u$, the true equilibrium G is constructed as follows: let $k = \lceil \frac{1}{u} \rceil$, and let $\tilde{G}$ be a $(\frac{k-1}{k}, \frac{1}{k})$ mix of the proper u -partition of $[0, 1 - u]$ and $\delta_{\{1\}}$. Then let G be the

$$\left(\frac{(2p-a)k}{(k-1)(1-u)+2-ka}, \frac{(k-1)(1-u)+2-2pk}{(k-1)(1-u)+2-ka} \right)$$

mix of $\tilde{G}$ and the proper u -partition of $[0, a]$, where $a \in (1 - u, (k - 1)u)$ is the unique solution of $(1 - a)\bar{v}(0) + a\bar{v}(1) = \bar{v}(a)$.

If $u \geq 1$, then $v(x)$ is quadratic and $\tilde{v}(x) = -x^2$ on $[0, 1]$; hence I refer the reader to Proposition 1 and Theorem 1. Consider now $u < 1$, where $v(x)$ is the CDF of the triangular distribution on $[-u, u]$:

$$v(x) = \begin{cases} 0, & x < -u, \\ \frac{(u+x)^2}{2u^2}, & x \in [-u, 0), \\ 1 - \frac{(u-x)^2}{2u^2}, & x \in [0, u), \\ 1, & x \geq u, \end{cases} \quad v'(x) = \begin{cases} 0, & x < -u, \\ \frac{u+x}{u^2}, & x \in [-u, 0), \\ \frac{u-x}{u^2}, & x \in [0, u), \\ 0, & x \geq u, \end{cases}$$

i.e. $v'(x) = \frac{\max(u-|x|, 0)}{u^2}$.

Here I use the continuity of $v'(x)$, which makes $\bar{v}'(x)$ continuous.

(i) Suppose that $p \leq 0.5, u < 1$, and let $G(x)$ be the CDF of the proper u -partition χ

on $[0, 2p]$ with $k = \lceil \frac{2p}{u} \rceil$. Let's analyze $\bar{v}'(x) = \int_0^1 v'(x-s)dG(s) = \int_0^{2p} v'(x-s)dG(s)$:

$$\begin{aligned}
\int_0^{2p} v'(x-s)dG(s) &= \sum_{i=0}^{k-1} v'(x-iu)\mathbb{P}(\chi = iu) + \sum_{i=0}^{k-1} v'(x-2p+iu)\mathbb{P}(\chi = 2p-iu) \\
&= \sum_{i=0}^{k-1} v'(x-iu) \frac{k-i}{k(k+1)} + \sum_{i=0}^{k-1} v'(x-2p+iu) \frac{k-i}{k(k+1)} \\
&= \sum_{i=0}^{k-1} \frac{\max(u-|x-iu|, 0)}{u^2} \frac{k-i}{k(k+1)} \\
&\quad + \sum_{i=0}^{k-1} \frac{\max(u-|x-2p+iu|, 0)}{u^2} \frac{k-i}{k(k+1)} \\
&= \underbrace{\sum_{i=0}^k \frac{\max(u-|x-iu|, 0)}{u^2} \frac{k-i}{k(k+1)}}_{S_L} \\
&\quad + \underbrace{\sum_{i=0}^k \frac{\max(u-|x-2p+iu|, 0)}{u^2} \frac{k-i}{k(k+1)}}_{S_R}.
\end{aligned}$$

Consider S_L and say that $a_i = \max(u-|x-iu|, 0)$ is *active* if $a_i > 0$. Then, as long as $x \in (iu, (i+1)u)$, only a_i and a_{i+1} are active, so

$$S_L = \frac{(u-(x-iu))(k-i) + (u-((i+1)u-x))(k-i-1)}{u^2 k(k+1)} = \frac{ku-x}{u^2 k(k+1)}.$$

By the continuity of S_L at $x = iu$, $S_L = \frac{ku-x}{u^2 k(k+1)}$ on $[0, ku]$, and hence on $[0, 2p]$, since $ku \geq 2p$. Analogously, $S_R = \frac{ku+x-2p}{u^2 k(k+1)}$ on $[0, 2p]$, hence $\bar{v}'(x) = \frac{2ku-2p}{u^2 k(k+1)}$ on $[0, 2p]$.

For $x > 2p$, $S_L = \frac{(ku-x)^+}{u^2 k(k+1)}$ and $S_R = \frac{k(u-x+2p)^+}{u^2 k(k+1)}$.

$$\bar{v}'(x) = \begin{cases} \frac{2ku-2p}{u^2k(k+1)}, & x \in [0, 2p], \\ \frac{2ku-x(1+k)+2kp}{u^2k(k+1)}, & x \in (2p, ku], \\ \frac{k(u-x+2p)}{u^2k(k+1)}, & x \in (ku, 2p+u], \\ 0, & x \geq 2p+u. \end{cases}$$

$\bar{v}'(x)$ is continuous, decreasing on $[2p, 1]$ and constant on $[0, 2p]$, hence $\bar{v}(x)$ is concave on $[0, 1]$, and linear on $[0, 2p]$, so indeed a pair of proper u -partitions of $[0, 2p]$ is an equilibrium. Also, $\bar{v}'(x)$ is convex on $[2p, 1]$.

(ii) Suppose that $p > 0.5, 2 - 2p \leq u < 1$; then $\bar{v}(x) = (1 - p)v(x) - pv(1 - x)$. Then

$$\bar{v}'(x) = \frac{(1 - p) \max(u - x, 0) + p \max(u + x - 1, 0)}{u^2},$$

where the numerator is a piecewise linear convex function, hence $\bar{v}(x)$ is concave and then convex, and we need to check that $\bar{v}'(0) \leq \bar{v}(1) - \bar{v}(0)$.

$\bar{v}'(0) = \frac{1-p}{u}$, $\bar{v}(1) - \bar{v}(0) = v(1) - v(0) = 0.5$, so $\{0, 1\}$ is a best reply iff $2 - 2p \leq u$.

(iii) Suppose $p > 0.5, u < 2 - 2p$, and $a \leq 1 - u$, where $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}$ for $k = \lceil \frac{2-2p}{u} \rceil - 1$. Then consider the $(\frac{2(1-p)}{2-a}, 1 - \frac{2(1-p)}{2-a})$ mix of the proper u -partition of $[0, a]$ with $\delta_{\{1\}}$; notice that this is a proper G , since the proper u -partition of $[0, a]$ has a mean $0.5a$, and $\frac{2(1-p)}{2-a} \cdot 0.5a + 1 - \frac{2(1-p)}{2-a} = p$.

First, $\lceil \frac{2-2p}{u} \rceil - 1 = \lceil \frac{a}{u} \rceil$, so k is the same in the proper u -partition of $[0, a]$:

$$\left\lceil \frac{a}{u} \right\rceil = \left\lceil 2k - \frac{k(k+1)u}{2(1-p)} \right\rceil = k,$$

since $k < \frac{2(1-p)}{u} \leq k + 1$, hence $k \leq \frac{k(k+1)u}{2(1-p)} < k + 1$.

Since $\bar{v}(x)$ is linear in the strategy of the other agent,

$$\bar{v}'(x) = \frac{2(1-p)}{2-a} S(x) + \frac{2p-a}{2-a} \cdot \frac{\max(u+x-1, 0)}{u^2},$$

where $S(x)$ is $\bar{v}'(x)$ from (i) against the proper u -partition χ of $[0, a]$. Hence $\bar{v}(x)$ is linear on $[0, a]$ with slope $\bar{v}'(0) = \frac{2(1-p)}{2-a} \cdot \frac{2ku-a}{u^2 k(k+1)}$. Since $v(1-s) = 1$ for $s \leq a \leq 1-u$,

$$\bar{v}(1) - \bar{v}(0) = \frac{2(1-p)}{2-a} \mathbb{E}[v(\chi)] + \frac{2p-a}{2(2-a)}, \quad \mathbb{E}[v(\chi)] = 1 - \frac{1}{2(k+1)} - \frac{(ku-a)^2}{2u^2 k(k+1)},$$

and the equation $\bar{v}(1) - \bar{v}(0) = \bar{v}'(0)$ reduces to a linear equation in a with the root $a = 2ku - \frac{k(k+1)u^2}{2(1-p)}$. Hence $\int_a^1 (\bar{v}'(x) - \bar{v}'(a)) dx = 0$.

On $[a, 1]$ both summands of $\bar{v}'(x)$ are convex: the first by (i), the second as a maximum of linear functions. Hence $\bar{v}'(x) - \bar{v}'(a)$ is convex on $[a, 1]$, vanishes at a and integrates to zero, so $\exists c : \bar{v}'(x) \leq \bar{v}'(a) \forall x \in [a, c]$ and $\bar{v}'(x) \geq \bar{v}'(a) \forall x \in [c, 1]$. Therefore $\bar{v}(x) \leq \bar{v}(a) + (x-a)\bar{v}'(a)$ on $[a, 1]$, with equality at $x = 1$, so $[0, a] \cup \{1\}$ is the concavification support, and (G, G) is an equilibrium.

(iv) Notice that G is a feasible distribution, since the mean of $\tilde{G}$ is $\frac{k-1}{k} \frac{1-u}{2} + \frac{1}{k} = \frac{Q}{2k}$ with $Q := (k-1)(1-u) + 2$, the mean of the proper u -partition of $[0, a]$ is $\frac{a}{2}$, and $\beta := \frac{(2p-a)k}{Q-ka}$ solves $\beta \frac{Q}{2k} + (1-\beta) \frac{a}{2} = p$.

Suppose that a from (iii) satisfies $a > 1-u$. I first show that if the other agent plays the $(\frac{k-1}{k}, \frac{1}{k})$ mix of the proper u -partition of $[0, 1-u]$ and $\delta_{\{1\}}$, where $k = \lceil \frac{1}{u} \rceil$, then

$\bar{v}(x)$ is linear on $[0, (k-1)u]$. Again, consider $\bar{v}'(x)$, noting that $\lceil \frac{1-u}{u} \rceil = k-1$:

$$\begin{aligned} \bar{v}'(x) = \frac{k-1}{k} & \left[\underbrace{\sum_{i=0}^{k-1} \frac{\max(u - |x - iu|, 0)}{u^2} \frac{k-1-i}{k(k-1)}}_{S_L} \right. \\ & \left. + \underbrace{\sum_{i=0}^{k-1} \frac{\max(u - |x - 1 + (i+1)u|, 0)}{u^2} \frac{k-1-i}{k(k-1)}}_{S_R} \right] + \frac{1}{k} \frac{\max(u - |x - 1|, 0)}{u^2}. \end{aligned}$$

By the proof of (i), $\bar{v}'(x)$ is constant on $[0, 1-u]$ (since the last term is zero on that interval) and equal to $\frac{k-1}{k} \frac{2(k-1)u - (1-u)}{u^2 k(k-1)} = \frac{(2k-1)u-1}{u^2 k^2}$. On $[1-u, (k-1)u]$, the only active terms are a_{k-2} in S_L , a_0 in S_R and the last term.

$$\begin{aligned} \bar{v}'(x) &= \frac{k-1}{k} \left[\frac{((k-1)u - x)}{u^2} \frac{1}{k(k-1)} + \frac{1-x}{u^2} \frac{k-1}{k(k-1)} \right] + \frac{1}{k} \frac{x+u-1}{u^2} \\ &= \frac{((k-1)u - x) + (k-1)(1-x) + k(x+u-1)}{k^2 u^2} = \frac{(2k-1)u-1}{k^2 u^2}. \end{aligned}$$

Finally, on $[(k-1)u, 1]$, only a_0 in S_R and the last term are active.

$$\bar{v}'(x) = \frac{k-1}{k} \frac{1-x}{u^2} \frac{k-1}{k(k-1)} + \frac{1}{k} \frac{x+u-1}{u^2} = \frac{(k-1)(1-x) + k(x+u-1)}{k^2 u^2} = \frac{x+ku-1}{k^2 u^2}.$$

So against $\tilde{G}$, $\bar{v}'(x)$ is constant on $[0, (k-1)u]$ and increasing linear on $[(k-1)u, 1]$, while against the proper u -partition of $[0, a]$ it is constant on $[0, a]$ and convex on $[a, 1]$ by (i). Hence, if the other agent plays G , $\bar{v}(x)$ is linear on $[0, a]$ and $\bar{v}'(x)$ is convex on $[a, 1]$, and as in (iii), if a solves $(1-a)\bar{v}(0) + a\bar{v}(1) = \bar{v}(a)$, then $[0, a] \cup \{1\}$ is the concavification support, and (G, G) is an equilibrium.

It remains to show that such an $a \in (1-u, (k-1)u)$ exists and is unique, and that $\beta \in (0, 1)$. Let a_3 and $k_3 = \lceil \frac{2-2p}{u} \rceil - 1$ be a and k from (iii). Since $a_3 \leq k_3 u$ and

$k_3 \leq k-1$, $a_3 > 1-u > (k-2)u$ requires $k_3 = k-1$, and then $a_3 > 1-u$ is equivalent to $p < p_u := 1 - \frac{k(k-1)u^2}{2((2k-1)u-1)}$. Let χ be the proper u -partition of $[0, a]$ and, for a disclosure H of the other agent, let s_H be the slope of $\bar{v}(x)$ on $[0, a]$ and $C_H := \bar{v}(1) - \bar{v}(0)$, so that the condition on a is $s_G = C_G$. A direct computation gives

$$s_{\tilde{G}} - C_{\tilde{G}} = -\frac{((k-1)u-1)^2}{2k^2u^2}, \quad \frac{d}{da}(s_{\chi} - C_{\chi}) = -\frac{1-a}{(k-1)u^2},$$

and $s_{\chi} = \frac{2(k-1)u-a}{u^2k(k-1)}$ by (i). Hence

$$\Lambda(a) := (Q - ka)(s_G - C_G) = -(2p - a)\frac{((k-1)u-1)^2}{2ku^2} + (Q - 2kp)(s_{\chi} - C_{\chi})$$

has the sign of $s_G - C_G$, as $Q - ka \geq (k+1)(1 - (k-1)u) > 0$, and $\Lambda(a)$ is quadratic in a with the leading coefficient $\frac{Q-2kp}{2(k-1)u^2}$. Since $p < p_u$ and $\frac{Q}{2k} - p_u = \frac{(k-1)((k-1)u-1)^2}{2k((2k-1)u-1)} > 0$, $2kp < Q$ and Λ is strictly convex. Moreover,

$$\begin{aligned} \Lambda(1-u) &= \frac{(1+u)((2k-1)u-1)}{k(k-1)u^2}(p_u - p) > 0, \\ \Lambda((k-1)u) &= -\frac{(k+1)(2p-1)((k-1)u-1)^2}{2ku^2} < 0, \end{aligned}$$

so Λ has a unique root in $(1-u, (k-1)u)$. And $a < 1 < 2p$ and $2kp < Q$ give $\beta \in (0, 1)$.

B.15 Proof of Proposition 14

Let $X = s_1$ and $Y = s_2$. The value functions are $\bar{v}_1(x) = \mathbb{P}(Y \leq x + b)$ and $\bar{v}_2(y) = \mathbb{P}(X + b < y)$. In cases (i) and (ii), let M , c and d be as in the statement.

Equilibrium. Against Y , $\bar{v}_1(x) = 1 - cM + cx$ on $[0, M]$ and $\bar{v}_1(x) = 1$ on $[M, 1]$, so $\bar{v}_1$ is concave and linear on $[0, M] \supseteq \text{supp}(X)$, and X is a best reply. Against X , $\bar{v}_2(y) = 0$ on $[0, b]$ and $\bar{v}_2(y) = bd + d(y - b) = dy$ on $(b, b + M]$, where in case (ii) the atom of X at $1-b$ is not beaten by $y = 1$. In case (i), $d(b+L) = 1$, so $\bar{v}_2(y) = 1 \leq dy$ on $[b+L, 1]$. Hence

$\bar{v}_2$ lies weakly below the line dy and coincides with it on $\{0\} \cup (b, b + M] \supseteq \text{supp}(Y)$, so Y is a best reply. By construction, both disclosures have mean p , and are well defined: in case (i), $1 - cL = 1 - \frac{2p}{2b+L} \geq 0$ as $L \geq 2p$, and $b + L \leq 1$ if and only if $b \leq 1 - \sqrt{2p}$; in case (ii), $1 - \frac{2p}{1+b} \geq 0$ as $p \leq 0.5$, and $d \in [0, 1]$ if and only if $1 - \sqrt{2p} \leq b \leq 1 - p$.

Uniqueness. As the game is constant-sum, all equilibria form a product set. Let X' be an equilibrium disclosure of agent 1. Then Y is a best reply to X' , and since $\text{supp}(Y) \supseteq \{0\} \cup [b, b + M]$, $\mathbb{P}(X' < y - b) = \kappa y$ on $(b, b + M]$ for some κ , i.e. X' has an atom κb at 0 and density κ on $(0, M)$. Since X' is also a best reply to Y and $p < M$, $\text{supp}(X') \subseteq [0, M]$, and the mass and the mean of X' pin down κ and its atom at M , hence $X' = X$. Let Y' be an equilibrium disclosure of agent 2. Since Y' is a best reply to X , it puts no mass outside $\{0\} \cup (b, b + M]$, and since X is a best reply to Y' and $\text{supp}(X) \supseteq [0, M]$, $\mathbb{P}(Y' \leq x + b)$ is linear on $[0, M]$. Hence Y' has an atom at 0 and constant density on $(b, b + M]$, both pinned down by its mass and mean, so $Y' = Y$.

Utility of the firm. The firm receives X unless $Y > X + b$, so

$$\begin{aligned} u_F &= p + \mathbb{E}[(Y - X)\mathbb{1}\{Y > X + b\}] = p + c \int_b^{b+M} \int_{[0, y-b)} (y - x) dX dy \\ &= p + cd \int_0^M \left(\frac{t^2}{2} + 2bt + b^2 \right) dt, \end{aligned}$$

which gives the expression in the statement. Finally, if $b \geq 1 - p$, agent 1 is selected with probability one by playing no disclosure, since $p + b \geq 1 \geq s_2$, so he is selected with probability one in every equilibrium, and $u_F = \mathbb{E}[s_1] = p$.

B.16 Proof of Corollary 9

(i) If $p < 0.5$, case (i) of Proposition 14 applies for small b , and

$$u_F = p + \frac{2p}{(2b + L)(b + L)} \left(\frac{L^2}{6} + bL + b^2 \right).$$

At $b = 0$, $L = 2p$ and $\frac{\partial L}{\partial b} = 1$, which gives $\frac{\partial u_F}{\partial b} = 0.5$. If $p = 0.5$, case (ii) of Proposition 14 applies for all $b > 0$, and

$$u_F = p + \frac{2p(1-b-p)(1+4b+b^2)}{3(1-b)(1+b)^2},$$

which gives $\frac{\partial u_F}{\partial b} = \frac{1}{6}$ at $b = 0$.

(ii) For $b = 1 - \sqrt{2p}$, $L = 1 - b = \sqrt{2p}$, and

$$u_F = p + \frac{2p}{2 - \sqrt{2p}} \left(\frac{p}{3} + (1 - \sqrt{2p})\sqrt{2p} + (1 - \sqrt{2p})^2 \right) = 2p + o(p),$$

while $u_F^*(p) = 2p - p^2$, so the share of the first best under the optimal head start converges to 1. For every well S-shaped selection rule, Theorem 2 gives $\frac{u_F}{u_F^*} \leq \frac{1+2p(1-p)}{2-p} \rightarrow 0.5$.

C. EQUATIONS FOR THE NOISELESS SELECTION RULE

Fact (Boleslavsky and Cotton (2015) Lemma 3). *The unique symmetric equilibrium for $v(x) = 0.5 + 0.5 \cdot \text{sign}(x)$ has the form (G, G) , where for $p \geq 0.5$*

$$G(x) = \begin{cases} \frac{x}{2p}, & x \leq 2 - 2p, \\ \frac{1}{p} - 1, & 2 - 2p \leq x < 1, \\ 1, & x = 1, \end{cases}$$

while for $p \leq 0.5$

$$G(x) = \begin{cases} \frac{x}{2p}, & x \leq 2p, \\ 1, & x \in [2p, 1]. \end{cases}$$

Hence the firm's equilibrium utility is

$$u_F^0(p) = \mathbb{E}_G[\max(s_1, s_2)] = \begin{cases} \frac{4p}{3}, & 0 < p \leq 0.5, \\ 1 - \frac{(1-p)^2(4p-1)}{3p^2}, & 0.5 < p < 1. \end{cases}$$