

Nonequilibrium Dynamics and Superadiabatic Fluxon Motion for Reversible Computing

Karpur Shukla

Department of Physics, Carnegie Mellon University

Department of Electrical and Computer Engineering, Brown University

Presented at the Center for Computing Research, Sandia National Laboratories

Date: February 13, 2020



e-mail: kshukla@alumni.cmu.edu
karpur_shukla@brown.edu



Outline

- Resource theory of quantum thermodynamics.
- Modelling generalized reversible computing in quantum open systems.
- State space geometric properties of open systems and impact on generalized reversible computing.
- Application to the dissipation delay product.
- Other work: reversible computing via superconducting circuits and superadiabatic fluxon motion.

Resource Theory

- Resource theory^[1]: all free states $\hat{\rho} \in \mathcal{D}(\mathcal{H})$ and free operations $\Phi \in \text{Aut}(\mathcal{D}(\mathcal{H}))$ which can be implemented with no energetic cost or dissipation, with conditions.
 - Ex.: Resource theory of bipartite entanglement. Free states: separable states. Free operations: local unitaries & classical communication (LOCC). Necessary & sufficient condition: $|\psi\rangle\langle\psi|_{AB} \mapsto |\phi\rangle\langle\phi|_{AB}$ iff $S(\psi_A) \geq S(\phi_A)$.
- Resource theory of quantum thermodynamics (RTQT):
 - Free states for nontrivial $\beta\hat{H}$: equilibrium thermal states $\{\hat{\rho}_G\}$.
 - Free operations for nontrivial $\beta\hat{H}$: all operations that preserve $\{\hat{\rho}_G\}$. *Thermal operations*.
 - Condition for $\hat{\rho} \mapsto \hat{\sigma}$ under thermal operations: *thermomajorization*.

[1] – N. H. Y. Ng and M. P. Woods, in *Thermodynamics in the Quantum Regime*, edited by F. Binder *et al.* (Springer Nature, Cham, 2018); see also arXiv:1805.09564.

Thermal Operations & Thermomajorization

- Start: system S coupled to environment E . E starts in thermal state: $\hat{\rho}_{E,G}$.
- Thermal operations: all quantum channels $\mathcal{E}(\hat{\rho}_S): \hat{\rho}_S \mapsto \text{Tr}_E \left\{ \hat{U}_{SE} (\hat{\rho}_S \otimes \hat{\rho}_{E,G}) \hat{U}_{SE}^\dagger \right\}$.
 - Energy-conserving unitary dynamics: global unitary evolution \hat{U}_{SE} across SE .
 - Require $[\hat{U}_{SE}, \hat{H}_S + \hat{H}_E] = 0$ at all times.
 - Maps thermal states to thermal states (but not necessarily the same one).
- Thermomajorization: necessary & sufficient for $\mathcal{E}(\hat{\rho})$ to map $\hat{\rho}_S \mapsto \hat{\sigma}_S$. For eigenvalues $\lambda_i(\hat{\rho})$ of $\hat{\rho}$ with corresponding energies $E_{i,\hat{\rho}}$, require:

$$\left(\sum_i e^{-\beta E_{i,\hat{\rho}}} , \sum_i \lambda_i(\hat{\rho})^\downarrow \right) \geq \left(\sum_i e^{-\beta E_{i,\hat{\sigma}}} , \sum_i \lambda_i(\hat{\sigma})^\downarrow \right)$$

- $\lambda_i(\hat{\rho})^\downarrow$: order eigenvalues of $\hat{\rho}$ by decreasing value.

Nonequilibrium Landauer Limit

- Using RTQT, can get an explicit nonequilibrium Landauer bound^[13].

$$\beta \langle Q \rangle \geq -\ln \left[\text{Tr}_S \text{Tr}_E \left\{ (\hat{\rho}_S(0) \otimes \mathbb{1}_E) \hat{U}_{SE}^\dagger (\mathbb{1}_S \otimes \hat{\rho}_{E,\beta}) \hat{U}_{SE} \right\} \right]$$

- Ejection of information in correlated bits^[3]: loss of prior correlations to environment.
 - Ejection of *uncorrelated* bits to the environment does *not* contribute to change in entropy.
- Dissipation bound depends on subsystem non-unitarity of \hat{U}_{SE} .
 - Measure of non-unitarity^[4] on E : $\mathcal{N}_E := \left\| \sum_{jk} \lambda_k \langle s_k | \hat{U}_{SE} | s_j \rangle \langle s_j | \hat{U}_{SE} | s_k \rangle - \mathbb{1}_E \right\|$.
 - $\mathcal{N}_E = 0$: no lower bound on dissipation.
 - $\langle s_a | \hat{U}_{SE} | s_b \rangle$: environment Kraus operators. Maps $\mathcal{D}(\mathcal{H}_E)$ to itself via operator algebra.

[2] – J. Goold, M. Paternostro, and K. Modi, Phys. Rev. Lett. 114, 060602 (2015).

[3] – M. Frank, arXiv:1806.10183.

[4] – G. Guarnieri *et al.*, New J. Phys. 19, 103038 (2017).

Modelling Reversible Computational Processes

- Standard^[5] method: N density matrices encoding N -ary alphabet.
 - Bijection between physical and computational states.
 - Generalized classical reversible computing: more physical states than computational states.
- Need to generalize to GRC, thermal operations, and nonequilibrium dynamics.
 - Want physical state \rightarrow computational state mapping to always commute with $\hat{U}_{SE(MW)}$. The same physical state always maps to the same computational state.
 - Can permit long-lived quantum coherences between physical states within an equiv. class.
 - Want to represent these as thermal operations, possibly within a larger open system.

(System) Kraus Operators

- ∞ -simal evolution for density matrices in open (noneq.) quantum systems:

$$\mathcal{E}_{dt}[\hat{\rho}_S(t)] = \hat{\rho}_S(t + dt) = \sum_{jk} \langle e_k | \hat{U}_{SE}(dt) | e_j \rangle \hat{\rho}_S(t) \langle e_j | \hat{U}_{SE}(dt) | e_k \rangle$$

- $\hat{E}_{jk} := \sqrt{\lambda_j} \langle e_k | \hat{U}_{SE} | e_j \rangle$: (system) Kraus operators over environment eigenstates $\{|e_i\rangle\}$.
- Any map of the form $\hat{\rho}_S(t) \mapsto \sum_{jk} \hat{E}_{jk}(t) \hat{\rho}_S(0) \hat{E}_{jk}^\dagger(t)$ is a *CPTP map*.
- $\hat{E}_{jk} \in \text{Op}(\mathcal{H}_S)$: takes $\mathcal{D}(\mathcal{H}_S)$ to itself, but expression depends on $\{|e_i\rangle\}$.
 - Done via algebra operation $\text{Op}(\mathcal{H}_S) \times \text{Op}(\mathcal{H}_S) \rightarrow \text{Op}(\mathcal{H}_S)$. $\hat{E}_{jk} \notin L_1(\text{Op}(\mathcal{H}_S))$.

The GKSL Superoperator

- GKSL superop.: time evolution generator at lowest order. $\mathcal{E}_{dt} = \mathcal{I} + dt \mathcal{L} + \dots$, and $\mathcal{L} := \lim_{dt \rightarrow 0} (\mathcal{E}_{dt} - \mathcal{I})/dt$.
- Kraus operators give evolution of $\hat{\rho}$ under quantum jump:

$$\hat{E}_0 := \mathbb{1}_S - i\hat{H}_S dt - \frac{1}{2} \sum_{j,k>0} \hat{F}_{jk}^\dagger \hat{F}_{jk}; \quad \hat{E}_{j,k>0} = \sqrt{\kappa_{jk} dt} \hat{F}_{jk}$$

- GKSL superoperator in terms of jump operators $\{\hat{F}_{jk}\}$:

$$\mathcal{L}[\hat{\rho}_S] := \frac{d\hat{\rho}_S}{dt} = -i[\hat{H}_S, \hat{\rho}_S] + \frac{1}{2} \sum_{j,k>0} \kappa_{jk} \left(2\hat{F}_{jk} \hat{\rho}_S \hat{F}_{jk}^\dagger - \left\{ \hat{F}_{jk}^\dagger \hat{F}_{jk}, \hat{\rho}_S \right\} \right)$$

Spectrum of the GKSL Superoperator


- Formal solution to GKSL equation: $\hat{\rho}_S(t) = e^{t\mathcal{L}} \hat{\rho}_S(0)$.
 - Examine spectral decomposition of \mathcal{L} via Choi-Jamiołkowski isomorphism: mapping of operators $\mathbb{F}^{n \times m}$ to vector space \mathbb{F}^{nm} , employing Hilbert-Schmidt inner product.
 - $|a\rangle\langle b| \mapsto |a\rangle \otimes |b\rangle$; $\hat{A} \mapsto |\hat{A}\rangle\rangle$. Inner product $\langle\langle \hat{A} | \hat{B} \rangle\rangle = \text{Tr}\{\hat{A}^\dagger \hat{B}\}$.
- Steady state solutions: defined as $|\hat{\rho}_{SS}\rangle\rangle := \lim_{t \rightarrow \infty} e^{t\mathcal{L}} |\hat{\rho}_S(0)\rangle\rangle$.
- System with *multiple* steady states: assuming \mathcal{L} is unitarily diagonalizable, have:

$$|\hat{\rho}_S(t)\rangle\rangle = e^{t\mathcal{L}} |\hat{\rho}_S(0)\rangle\rangle = \sum_a e^{t\lambda_a} |\hat{x}_a\rangle\rangle \langle\langle \hat{y}_a | \hat{\rho}_S(0) \rangle\rangle$$

- $|\hat{x}_a\rangle\rangle$, $\langle\langle \hat{y}_a |$, λ_a are right eigenstates, left eigenstates, and eigenvalues of \mathcal{L} .
- $\Re\{\lambda_a\} < 0$: damped states. $\Re\{\lambda_a\} = 0$: steady states. ($\Re\{\lambda_a\} > 0$ blows up.)


Four-Corners Decomposition

- Multiple steady states form^[6] *asymptotic subspace* $As(\mathcal{H})$ of nonzero steady states.
 - Right eigenvectors of \mathcal{L} with pure imaginary eigenvalues.
 - *Corresponding* left eigenvectors are system conserved currents.
 - All initial states have components along these states.
 - $As(\mathcal{H})$ can have further nontrivial dynamics.
- *Four-corners decomposition*: decomposition of \mathcal{L} in terms of $As(\mathcal{H})$ and conserved currents.




Example Types of Asymptotic Subspaces

- Decoherent subspace: decoherence between steady states in dynamics purely within $\text{As}(\mathcal{H})$.
 - Decoherence-free subspace (DFS): no further decoherence.
- Noiseless subsystem (NS): $\text{As}(\mathcal{H})$ is tensor product of DFS subspace and subspace spanned by different steady state.
- von Neumann algebra: $\text{As}(\mathcal{H})$ is a direct sum of DFS or NS blocks.
 - von Neumann algebra: C^* -algebra which also includes all \hat{A} that satisfy $\langle \phi | \hat{A}_n | \psi \rangle \rightarrow \langle \phi | \hat{A} | \psi \rangle$ for $|\phi\rangle, |\psi\rangle \in \mathcal{H}$ and $\{\hat{A}_n\} \in L_1(\mathcal{H})$.



Generalized Reversible Computing & Four Corners

- Classical reversible computing: surjective map from physical to computational states, equivalence classes.
 - All states *within* a class must have same noncomputational entropy: related by a unitary transformation.
 - We permit intra-class coherences: each class can be a DFS.
 - Each class in a given computational scheme must have same computational entropy: each DFS block has same dimension.
 - \therefore quantum channels performing classical RC can be represented by the von Neumann algebra.
- Quantum GRC: permit inter-class coherences.



Berry-Wilczek-Zee Connection

- Time evolution in adiabatic approximation, expressed in parameter space: $t \mapsto \lambda^\mu \in \mathbb{R}^n$.
- Nondegenerate eigenspaces: Berry phase & connection given by:

$$\phi = \oint_C d\lambda^\mu A_\mu; \quad A_\mu = i \langle n(\lambda) | \partial_\mu | n(\lambda) \rangle$$

- A_μ as U(1) gauge theory: under $|n(t)\rangle \mapsto e^{i\xi} |n(t)\rangle$, have $A_\mu \mapsto A_\mu - \partial_\mu \xi$.
- N-fold degenerate eigenspace: Berry-Wilczek-Zee connection is U(N) matrix. Eigenvalues of *Wilson loop* are generalizations of Berry phase:

$$W = \text{Tr } \mathcal{P} \exp \left\{ \oint_C d\lambda^\mu A_{ab;\mu} \right\}; \quad A_{\mu;ab} = i \langle n_a(\lambda) | \partial_\mu | n_b(\lambda) \rangle$$

- $A_{\mu;ab}$ as U(N) gauge theory: under $|n_b(t)\rangle \mapsto \sum_a U_{ab} |n_a(t)\rangle$, have $A_\mu \mapsto iU^{-1} \partial_\mu U - U^{-1} A_\mu U$.

Berry-Wilczek-Zee to Operator Space

- Systems with multiple steady states: expect geometric effects to arise, especially in quantities not dependent on adiabatic path in parameter space.
- Basis for $\text{Op}(\mathcal{H}_S)$: $\{|N_{ab}\rangle\rangle\} := |n_a\rangle\langle n_b|$. BWZ connection and Wilson loops in decoherence-free subspace:

$$\mathcal{W} = \text{Tr } \mathcal{P} \exp \left\{ \oint_C d\lambda^\mu \mathcal{A}_{\mu; abcd} \right\}; \quad \mathcal{A}_{\mu; abcd} = \langle\langle N_{cd} | \partial_\mu | N_{ab} \rangle\rangle$$


- Noiseless subsystem: $\text{As}(\mathcal{H})$ is tensor product of DFS subspace and subspace spanned by different steady state. \mathcal{W} and $\mathcal{A}_{\mu; abcd}$ can mix these!
- von Neumann algebra: $\mathcal{A}_{\mu; abcd}$ is distinct on each block. No mixing between directly summed blocks.

Dissipation Delay Product

- Major quantity of interest for characterizing efficiency of reversible operations.
 - Goal: characterizing *general* (protocol-based, device-independent) efficiency properties of reversible operations.
- Dissipation-delay product: product of energy dissipation incurred by transition process and delay of process.
 - For perfectly unitary time evolution, dissipation is zero.
 - Want to quantify energy dissipation for reversible operation. Dissipation as a function of delay.
 - Geometric approach: both dissipation (via this approach to calculating bound on average dissipation) and delay (preliminarily, can see this from quantum speed limit^[7]) will depend on geometry of states.

Dissipation Length for Single Steady States

- *Thermodynamic* dissipation length: minimal dissipation amount for processes.
 - Construct manifold for set of all possible Gibbs states. (*Not* the same manifold!)
 - Geodesics: minimal dissipation process.
- Derived^[9] for GKSL dynamics with a *single* steady state.
 - Current status: extending perturbation theory^[10] of steady states for \mathcal{L} with slowly varying parameters to multiple steady states.



[8] – M. Scandi and M. Peranau-Llobet, Quantum 3, 197 (2019). Image from here.

[9] – V. Cavina, A. Mari, and V. Giovannetti, Phys. Rev. Lett. 119, 050601 (2017).

Superadiabaticity / Shortcuts to Adiabaticity (STA)

- Goal: more quickly reach the final results of adiabatic time evolution of \hat{H} .
 - Ideally, want to reproduce full spectrum (states and their respective densities) that we'd have gotten from adiabatic evolution.
 - Alternately, want to find all of the invariants of the adiabatic motion.
 - *No* excitation (perturbations) allowed in final \hat{H} (relative to the time-evolved form of the unperturbed initial \hat{H} , but such perturbations permitted along the way.
 - Mapping $\{|\psi_i\rangle\} \rightarrow \{|\psi_f\rangle\}$ should be state independent: holds for *any* possible $|\psi_i\rangle$.

STA Techniques

- Counterdiabatic driving (CD): add perturbations, not necessarily adiabatic, to $\hat{H}_0(t)$. Dynamics should result in same spectral decomposition at end.

$$\hat{H}_{CD} = i\hbar \sum_n \{|\partial_t n\rangle\langle n| - \langle n|\partial_t n\rangle \cdot |n\rangle\langle n|\}$$

- Invariant-based engineering (IE): construct dynamical invariants of the full Hamiltonian from 1st integrals of motion. examine spectral decomposition of this invariant.
 - Lewis-Riesenfeld invariant has $d_t \hat{I} = 0$, so its eigenvalues are time independent and time evolved expectation values of its eigenvectors are constant. Calculate \hat{H} from time evolution.

$$d\hat{I}/dt = \partial\hat{I}/\partial t + i[\hat{H}, \hat{I}]/\hbar = 0$$

Quantum-Classical Correspondence for STA

- In parameter (Berry) space, define^[11] $\hat{\xi} := i\hbar \sum_n \{ |\partial_\mu n\rangle \langle n| - \langle n | \partial_\mu n \rangle \cdot |n\rangle \langle n| \}$. Full \hat{H} is then:

$$\hat{H} = \hat{H}_0 + \hat{H}_{CD} = \hat{H}_0 + i\hbar \sum_n \{ |\partial_\mu n\rangle \langle n| - \langle n | \partial_\mu n \rangle \cdot |n\rangle \langle n| \} = \hat{H}_0 + \frac{dR_\mu}{dt} \cdot \hat{\xi}$$

- $\hat{\xi}$ satisfies (and is specified by):

$$[\hat{\xi}, \hat{H}_0] = i\hbar \left(\partial_\mu \hat{H}_0 - \sum_n |n\rangle \langle n | \partial_\mu \hat{H}_0 | n \rangle \langle n| \right); \quad \langle n | \hat{\xi} | n \rangle = 0$$

- Correspondence: commutator \rightarrow Poisson bracket. For a classical H_0 , microcanonical average $\langle \alpha \rangle_\mu$ of classical observable α on constant energy shells given by:

$$\langle \alpha \rangle_\mu = \frac{\int d^{3n}p \, d^{3n}q \, \alpha \, \delta(E - H_0)}{\partial_E \int d^{3n}p \, d^{3n}q \, \theta(E - H_0)}; \quad \{\xi, H_0\} = \nabla H_0 - \langle \nabla H_0 \rangle_\mu; \quad \langle \xi \rangle_\mu = 0$$

Classical STA and Completely Integrable Systems

- Procedure^[12]: for Hamiltonian H_0 that depends on parameters R , determine ξ that satisfies

$$\frac{\partial H_0}{\partial \vec{R}} = \{\xi, H_0\} + \frac{dE_0}{d\vec{R}}$$

- Goal for classical STA (and invariant-based engineering): finding invariants.
 - Classical STA: ξ is an invariant used to solve a classical PDE.
 - Invariant-based engineering: Lewis-Riesenfeld invariants used to find a convenient eigenbasis for \hat{H} .
- Lewis-Riesenfeld equation $\partial \hat{I} / \partial t + i[\hat{H}, \hat{I}] / \hbar = 0$ can be directly related to soliton theory: all soliton equations can be reformulated as $dL/dt + [P, L] = 0$, with L determined from the PDE's symmetry group.

Quantum and Classical STA of KdV Equation

- Korteweg-de Vries equation: $6 \cdot \partial u / \partial R^\mu - 6u \cdot \partial u / \partial x + \partial u^3 / \partial x^3 = 0$
- Quantum STA developed by Lewis-Riesenfeld invariants: $i\hbar \partial \hat{H}_0 / \partial R^\mu = [\hat{\xi}, \hat{H}_0]$
 - Solution^[12]: for $\hat{H}_0 = \hat{p}^2 + u$, have $\hat{\xi} = -2\hat{p}/3 - (\hat{p}u + u\hat{p})/2$
- Classical STA: $\partial H_0 / \partial R^\mu = \{\xi, \hat{H}_0\}$
 - Solution^[12]: for $\hat{H}_0 = p^2 + u$, have $\xi = -2p/3 - pu$
 - Relies on solving a classical PDE for ξ .
- Notice that these provide somewhat different results. Can't just naively go from commutator to Poisson!
- Next step: apply this technique to sine Gordon equation.

Conclusions and Next Steps

- To develop reversible computing and minimal dissipation properties, can embed framework of generalized reversible computing within GKSL dynamics.
- Current work: extending RC to a proper open system / nonequilibrium quantum thermodynamic footing. (Sandia Tracking #1068410)
 - Gives us fundamental quantum thermo results “for free”; e.g. thermodynamic length for multiple SSs.
 - Can give us a device-independent, protocol-based metric for efficiency of reversible operations!
- Current work: shortcut to adiabaticity for fluxons.
 - Relies intrinsically on infinite degrees of freedom (complete integrability) of sine-Gordon equation.

Landauer's Limit

- Landauer's limit^[13]: one bit of information lost in computational process dissipates $\Delta E \geq k_B T \ln 2$ of energy as heat.
 - Dissipation due to increase in entropy: $\Delta S \geq k_B \ln 2$. Links information and physics!
 - Ejection of information in correlated bits^[14]: loss of prior correlations to environment.
 - Ejection of *uncorrelated* bits to the environment does *not* contribute to change in entropy.
- No-hiding theorem^[14]: information can't be destroyed.
 - Moves from system S to environment E . Global unitary evolution over $\mathcal{H}_S \otimes \mathcal{H}_E$.
 - Information lost from original system can't remain in SE correlations.
- Conventional computing: *entirely* irreversible. (Ex.: clearing memory.)

[13] – R. Landauer, IBM J. Res. Dev. 5, 163 (1961).

[3] – M. Frank, arXiv:1806.10183.

[14] – S. Braunstein and A. K. Pati, Phys. Rev. Lett. 98, 080502 (2007).

Practical Motivation for Reversible Computing

- Dennard scaling: power density of transistors remains constant as transistor size decreases.
- Moore's law: number of transistors in given integrated circuit area doubles (approximately) every 18 months.
 - Dennard: ended by 2008^[15]. Moore: ending now^[16].
- *Main* challenge: energy efficiency! Reversible computing: avoids major source of energy dissipation.
 - Almost all quantum computing is reversible, but classical reversible computing is the *lower bound* case on dissipation.

[15] – B. Deng *et al.*, ACM Trans. Arch. Code Opt. 15, 8:1 (2018).

[16] – H. Khan, D. Hounshell, and E. Fuchs, Nature Electronics 1, 14 (2018).

Classical Computing as a Lower Dissipative Bound

- Information processing expressed as a thermal operation^[17]. Dissipation:

$$\Delta E_Q \geq k_B T (S(\hat{\rho}_S) - S(\hat{\varrho}_S)) + S \left(\hat{U}_{SME} (\hat{\rho}_S \otimes \hat{\rho}_M \otimes \hat{\rho}_E) \hat{U}_{SME}^\dagger \parallel \hat{\varrho}_S \otimes \hat{\rho}_M \otimes \hat{\rho}_E \right)$$

- System S coupled to environment E and catalyst M ; same as splitting E into M and E .
- Channel: $\mathcal{E}(\hat{\rho}_S): \hat{\rho}_S \mapsto \hat{\varrho}_S := \text{Tr}_M \text{Tr}_E \left\{ \hat{U}_{SME} (\hat{\rho}_S \otimes \hat{\rho}_M \otimes \hat{\rho}_E) \hat{U}_{SME}^\dagger \right\}$.
- First term: information cost of classical IP. Second term: quantum IP.
- Classical IP is a *lower* dissipative bound! Quantum IP can be equal at best.
 - Classical IP: signal states correspond to orthogonal quantum states.
 - Pure unitaries and single input & output operations match classical IP dissipation bound.

Resource Theory & Reversible Computing


- Key issue: IP operations usually involve a catalyst machine, which sometimes needs to be reset. Reset destroys correlations (mutual info.): dissipation!
 - This is also the source of dissipation in irreversible (classical) computing.
- Thermal operations *can* provide a framework for quantum RC with dissipation arbitrarily close to classical RC.
 - Same idea as classical RC: preserve correlations. Rigorously proved^[18]: examine the transition $\hat{\rho}_S \otimes \hat{\rho}_M \otimes |w\rangle\langle w| \mapsto \hat{\varrho}_{SM} \otimes |w - \Delta\rangle\langle w - \Delta|$ for SME coupled to work bit $|w\rangle$. Resetting condition: $\text{Tr}_S \hat{\varrho}_{SM} = \hat{\rho}_M$.
 - $S(\hat{\rho}_S \| \text{Tr}_S \hat{\varrho}_{SM})$ can be arbitrarily close to zero: arbitrarily close to classical IP bound.

Thermal bath


$$\hat{\rho}_E, \hat{H}_E$$

Machine


Machine



Energy-preserving transformation



Correlation




System



System

Generalized Reversible Computing

- Reversible computing: reversibly transform computational states, instead of destructively overwriting them.
- Computational states c : states representing computational information.
 - $\{c\}$ partitions $\{\hat{\rho}\}$ in equiv. classes. All $\{\hat{\rho}\}$ in same class are linked by unitary transform: have same entropy.
- GRC: Bijections on the probability-1 subset of $\{c\}$.
- Lower dissipative bound on classical RC can be zero!
 - Quantum RC: bounded only by mutual entropy term.



Quantum Geometric Tensor from GKSL Dynamics

- Distance in parameter space:

$$ds^2 = \|\psi(\lambda + d\lambda) - \psi(\lambda)\|^2 = \langle \partial_\mu \psi | \partial_\nu \psi \rangle d\lambda^\mu d\lambda^\nu = (\gamma_{\mu\nu} + i\sigma_{\mu\nu}) d\lambda^\mu d\lambda^\nu$$

- $\sigma_{\mu\nu}$: Berry curvature. $\sigma_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.
- $g_{\mu\nu} = \gamma_{\mu\nu} - A_\mu \wedge A_\nu$: quantum geometric tensor. Metric on the manifold of states.
- Interest is in quantum geometric tensor induced by GKSL dynamics.
 - Wilczek-Zee-Berry connection: $\mathcal{A}_{\mu;abcd} = \langle\langle N_{cd} | \partial_\mu | N_{ab} \rangle\rangle = \text{Tr}\{\hat{N}_{cd} \partial_\mu \hat{N}_{ab}\}$.
 - Induced QGT for vN algebra: $\mathcal{Q}_{\alpha\beta;abcd} = \partial_\alpha \mathcal{A}_{\beta;abcd} + \mathcal{A}_{\alpha;abcd} \wedge \mathcal{A}_{\beta;abcd} - \langle\langle \hat{\rho}_{cd} | \partial_\alpha \partial_\beta | \hat{\rho}_{ab} \rangle\rangle$.

More STA Techniques

- Variational counterdiabatic driving: same as CD, except use variational method in quantum to find perturbative term.
- Lie algebraic invariant-based engineering: same as IE, except construct dynamical invariants based on Lie algebra of underlying symmetries.
- Scaling laws: same as IE, except construct invariant from overarching scaling laws of \hat{H} .
- Variational method: construct approximate dynamics using Lagrangian and calculate via saddle-point approximation.
- Fast forward: same as scaling law approach, but also requires renormalization.