

NOTES ON



COMMUTATIVE ALGEBRA



Lecture Notes & Problem Discussions

SAYAN KAR

Last updated
July 30, 2026

ABSTRACT

These are lecture notes for a first course in Commutative Algebra, covering rings, ideals, and modules, together with the accompanying problem-set discussions worked out alongside the lectures. The standard references for this material are [\[AM69\]](#), [\[Bly90\]](#), and [\[DF04\]](#).

CONTENTS

1	Lecture Notes	5
1.1	Lecture 1	5
1.2	Lecture 2	7
1.3	Lecture 3	11
1.4	Lecture 4	13
2	Problem Discussions	17
2.1	Assignment 0	17

CHAPTER 1

LECTURE NOTES

1.1 Lecture 1

Convention. Rings are commutative with 1.

Definition 1.1. Let R, S be rings. A *morphism* φ from R to S is a map $\varphi : R \rightarrow S$ such that

- (1) $\varphi(r + s) = \varphi(r) + \varphi(s)$,
- (2) $\varphi(rs) = \varphi(r)\varphi(s)$,
- (3) $\varphi(1) = 1$.

Convention. A subring R must contain 1.

Question. What is $R[X]$?

X is not an element of R . Consider

$$S = \{f : \mathbb{N} \rightarrow R \mid f(r) = 0 \text{ for all but finitely many } r\},$$

with the usual addition and multiplication in sequences (Cauchy product). Then S is a ring with identity

$$1_S = (1, 0, 0, \dots, 0).$$

Define $\varphi : R \rightarrow S$ by $\varphi(a) = (a, 0, \dots, 0)$. Then φ is a ring homomorphism, for sure.

Define $X = (0, 1, 0, \dots)$ and observe $X^2 = (0, 0, 1, 0, \dots)$. A polynomial P over R can then be identified as an element of S , say

$$P = (P_0, P_1, \dots, P_n, 0, \dots) = \varphi(P_0) \cdot 1_S + \varphi(P_1) \cdot X + \dots + \varphi(P_n) \cdot X^n.$$

Definition 1.2. Let R be a ring. A non-zero, non-unit element $p \in R$ is *prime* if

$$p \mid ab \implies p \mid a \text{ or } p \mid b$$

(include the elementwise definition: for $a, b \in R$, $a \mid b$ if $\exists r \in R$ such that $b = r \cdot a$).

Definition 1.3. Let R be a ring. A proper ideal P of R is called a *prime ideal* if

$$ab \in P \implies a \in P \text{ or } b \in P.$$

Definition 1.4. Let R be a ring. $I \subseteq R$ is called an *ideal* if there exists a ring S and a morphism $\varphi : R \rightarrow S$ such that $I = \ker(\varphi)$.

Remark 1.5. This immediately gives the elementwise definition of ideals.

$$\mathcal{N}(R) = \{\text{all nilpotent elements of } R\} \rightsquigarrow \text{the nil-radical of } R.$$

Theorem 1.6.

$$\mathcal{N}(R) = \bigcap_{P \in \text{Spec } R} P, \quad \text{where } \text{Spec } R = \text{all prime ideals of } R.$$

Proof. Let $x \in \mathcal{N}(R)$. Then for any $P \in \text{Spec } R$, there exists $n_P \in \mathbb{N}$ such that

$$x^{n_P} = 0 \implies 0 \in P \implies x^{n_P} \in P \implies x \in P.$$

Now let $x \notin \mathcal{N}(R)$, and define

$$S = \{I \subseteq R \text{ ideal} \mid x^n \notin I \text{ for all } n\}.$$

$S \neq \emptyset$, as $\langle 0 \rangle \in S$, and S is a poset (ordered by inclusion). By Zorn's Lemma, let M be a maximal element of S .

Claim. M is a prime ideal.

Proof of claim. Let $a \notin M$, $b \notin M$. Now,

$$M + \langle a \rangle, M + \langle b \rangle \notin S \implies \exists m, n \in \mathbb{N} \text{ such that}$$

$$x^m \in M + \langle a \rangle \text{ and } x^n \in M + \langle b \rangle \implies x^{m+n} \in M + \langle ab \rangle.$$

If $ab \in M$, then $x^{m+n} \in M$, a contradiction. Hence $ab \notin M$, so M is prime. □

Since $x \notin M$ and $M \in \text{Spec } R$, we conclude $x \notin \bigcap_{P \in \text{Spec } R} P$, completing the proof. □

Question. What are the units of $R[X]$?

If R is an integral domain, then they are precisely the units of R .

Proposition 1.7. $f(X) = a_n X^n + \cdots + a_1 X + a_0$ is a unit if a_0 is a unit and the a_i 's are nilpotent.

Proof. Let $P \in \text{Spec } R$. We have a natural morphism $\varphi : R[X] \rightarrow (R/P)[X]$ given by

$$\varphi(b_m X^m + \cdots + b_1 X + b_0) = \overline{b_m} X^m + \cdots + \overline{b_1} X + \overline{b_0}, \quad (\overline{b_i} = b_i \bmod P).$$

Then $\varphi(f)$ is a unit in $(R/P)[X]$, and since R/P is an integral domain, this forces

$$\overline{a_i} \equiv \overline{0} \pmod{P} \text{ for all } i \in [n], \quad \text{and} \quad \overline{a_0} \text{ is a unit of } R/P.$$

That is, $a_0 \notin P$ and $a_1, \dots, a_n \in P$. As $P \in \text{Spec } R$ was arbitrary,

$$a_0 \notin P \text{ and } a_i \in P \quad \forall i \in [n], \quad \text{for all } P \in \text{Spec } R.$$

The other direction is trivial. □

1.2 Lecture 2

Definition 1.8. Let R be a ring and let $(M, +)$ be an abelian group. A *scalar multiplication*

$$R \times M \longrightarrow M, \quad (r, m) \longmapsto rm,$$

makes M into an R -module if the following hold for all $r, s \in R$ and $m, m_1, m_2 \in M$:

- (i) $(r + s)(m) = rm + sm$,
- (ii) $r(m_1 + m_2) = rm_1 + rm_2$,
- (iii) $r(sm) = rsm = s(rm)$,
- (iv) $1 \cdot m = m$.

Definition 1.9. If M, N are R -modules, a map $\varphi : M \rightarrow N$ is called an R -module morphism if

- (i) $\varphi(m_1 + m_2) = \varphi(m_1) + \varphi(m_2)$,
- (ii) $\varphi(rm) = r\varphi(m)$.

Definition 1.10. Let R be a ring and S a nonempty set. A *free R -module on S* is a pair (F, f) , where F is an R -module and $f : S \rightarrow F$ is a set-theoretic map, such that: for any R -module M and any set-theoretic map $g : S \rightarrow M$, there exists a unique R -module morphism $h : F \rightarrow M$

such that

$$\begin{array}{ccc} S & \xrightarrow{g} & M \\ f \downarrow & \nearrow h & \\ F & & \end{array}$$

commutes, i.e. $h \circ f = g$.

Proposition 1.11. Let (F, f) be a free R -module on S . Then:

- (1) f is injective;
- (2) $\text{Im}(f)$ generates F as an R -module.

Proof of (1). Let $x, y \in S$ with $x \neq y$. **Goal:** $f(x) \neq f(y)$.

Take any R -module M with at least two elements, and define $g : S \rightarrow M$ such that $g(x) \neq g(y)$. By definition, there exists a (unique) morphism $h : F \rightarrow M$ such that $h \circ f = g$. Then

$$(h \circ f)(x) = g(x) \neq g(y) = (h \circ f)(y) \implies f(x) \neq f(y). \quad \square$$

Definition 1.12. Let M be an R -module and $\emptyset \neq A \subseteq M$. The *submodule of M generated by A* is, equivalently:

- (1) the smallest submodule of M containing A ;
- (2) the intersection of all submodules of M containing A ;
- (3) the set of all finite R -linear combinations of elements of A .

Proof of (2). Let A be the submodule of F generated by $\text{Im}(f)$. Since $f(s) \in \text{Im}(f) \subseteq A$ for every $s \in S$, f factors as $f = i_A \circ f^*$ for a set-theoretic map $f^* : S \rightarrow A$ and the inclusion $i_A : A \hookrightarrow F$:

$$\begin{array}{ccc} S & \xrightarrow{f^*} & A \\ & \searrow f & \downarrow i_A \\ & & F \end{array}$$

As (F, f) is free on S and A is an R -module, there is a unique morphism $h : F \rightarrow A$ such that $h \circ f = f^*$:

$$\begin{array}{ccc} S & \xrightarrow{f^*} & A \\ f \downarrow & \nearrow h & \\ F & & \end{array}$$

It is enough to show i_A is onto. We compute

$$(i_A \circ h) \circ f = i_A \circ (h \circ f) = i_A \circ f^* = f.$$

So $i_A \circ h$ and id_F both satisfy $(-) \circ f = f$; by uniqueness in the definition of freeness (applied with $M = F, g = f$), $i_A \circ h = \text{id}_F$. Hence i_A is onto, i.e. $A = F$, so $\text{Im}(f)$ generates F . $\square$

Theorem 1.13: Uniqueness of free modules. Let (F, f) be a free R -module on a set S . Then (F', f') is also a free R -module on S if and only if there exists a unique R -module isomorphism $j : F \rightarrow F'$ such that $j \circ f = f'$.

Proof. Suppose (F', f') is also free on S . Since (F, f) is free (applied to $M = F', g = f'$), there is a morphism $h : F \rightarrow F'$ with $h \circ f = f'$:

$$\begin{array}{ccc} S & \xrightarrow{f'} & F' \\ f \downarrow & \nearrow h & \\ F & & \end{array}$$

Likewise, since (F', f') is free (applied to $M = F, g = f$), there is a morphism $h' : F' \rightarrow F$ with $h' \circ f' = f$:

$$\begin{array}{ccc} S & \xrightarrow{f} & F \\ f' \downarrow & \nearrow h' & \\ F' & & \end{array}$$

Then $(h' \circ h) \circ f = h' \circ (h \circ f) = h' \circ f' = f$. So $h' \circ h$ and id_F both satisfy $(-) \circ f = f$; by uniqueness (with $M = F, g = f$), $h' \circ h = \text{id}_F$. Similarly $h \circ h' = \text{id}_{F'}$. So $j := h$ is an isomorphism with $j \circ f = f'$.

Converse. Suppose there is an isomorphism $j : F \rightarrow F'$ with $j \circ f = f'$. To show (F', f') is free on S , take any R -module M and set-theoretic map $g : S \rightarrow M$; we must produce a unique $h' : F' \rightarrow M$ with $h' \circ f' = g$:

$$\begin{array}{ccc} S & \xrightarrow{g} & M \\ f' \downarrow & \nearrow h' & \\ F' & & \end{array}$$

Since (F, f) is free, there is a unique $h : F \rightarrow M$ with $h \circ f = g$. From $j \circ f = f'$ we get $j^{-1} \circ f' = f$, so

$$(h \circ j^{-1}) \circ f' = h \circ (j^{-1} \circ f') = h \circ f = g,$$

i.e. $h' := h \circ j^{-1}$ works:

$$\begin{array}{ccc} S & \xrightarrow{g} & M \\ f' \downarrow & \nearrow f & \uparrow h \\ F' & \xrightarrow{j^{-1}} & F \end{array}$$

Uniqueness. Suppose $t : F' \rightarrow M$ also satisfies $t \circ f' = g$. Then $t \circ j : F \rightarrow M$ satisfies $(t \circ$

$j) \circ f = t \circ (j \circ f) = t \circ f' = g$, so by uniqueness of h (from freeness of (F, f)), $t \circ j = h$, hence $t = h \circ j^{-1} = h'$. $\square$

Theorem 1.14: Existence of free R -modules. For any ring R and nonempty set S , a free R -module on S exists.

Proof. Define

$$F = \{\theta : S \rightarrow R \mid \theta(s) = 0 \text{ for almost all } s \in S\}.$$

For $\theta_1, \theta_2 \in F$ and $\lambda \in R$, define

$$(\theta_1 + \theta_2)(s) = \theta_1(s) + \theta_2(s), \quad (\lambda\theta)(s) = \lambda\theta(s).$$

One checks that F is an R -module. Define $f : S \rightarrow F$ by $f(s) = \delta_s$, where $\delta_s(t) = 1$ if $t = s$ and 0 otherwise.

Existence of h . Let M be an R -module and $g : S \rightarrow M$ a set-theoretic map. Define $h : F \rightarrow M$ by

$$h(\theta) = \sum_{s \in S} \theta(s) g(s),$$

a finite sum for every $\theta \in F$. For any $t \in S$,

$$(h \circ f)(t) = h(f(t)) = h(\delta_t) = \delta_t(t) g(t) = g(t),$$

so $h \circ f = g$:

$$\begin{array}{ccc} S & \xrightarrow{g} & M \\ f \downarrow & \nearrow h & \\ F & & \end{array}$$

Uniqueness of h . For $\theta \in F$ and $t \in S$,

$$\theta(t) = \theta(t) \cdot 1_R = \sum_{s \in S} \theta(s) \delta_s(t) = \left(\sum_{s \in S} \theta(s) \delta_s \right) (t) \quad \forall t \in S,$$

so $\theta = \sum_{s \in S} \theta(s) \delta_s$. Now let $h' : F \rightarrow M$ be any R -module morphism with $h' \circ f = g$. Then for $\theta \in F$,

$$h'(\theta) = h' \left(\sum_{s \in S} \theta(s) \delta_s \right) = \sum_{s \in S} \theta(s) h'(\delta_s) = \sum_{s \in S} \theta(s) h'(f(s)) = \sum_{s \in S} \theta(s) g(s) = h(\theta).$$

Hence $h = h'$. $\square$

1.3 Lecture 3

Theorem 1.15. Let R be a ring such that every prime ideal is principal. Then every ideal is principal.

Proof. Let $S = \{\text{all non-principal ideals of } R\}$. For the sake of contradiction (FTSOC), suppose $S \neq \emptyset$. By Zorn's Lemma, let I be a maximal element of S .

Since I is not principal, I is not prime by hypothesis, so there exist $a, b \in R$ with $a \notin I, b \notin I$, but $ab \in I$.

Let $J = \langle a \rangle + I \supsetneq I$; since I is maximal among non-principal ideals, $J \notin S$, so J is principal, say $J = \langle c \rangle$.

Also let $K = \{r \in R \mid ra \in I\} \supsetneq I$ (since $b \in K \setminus I$); again $K \notin S$, so $K = \langle d \rangle$ for some d .

$I \subseteq \langle cd \rangle$. Let $x \in I$. Since $I \subseteq J = \langle c \rangle$, there is $\lambda \in R$ with $\lambda c = x$. Now $\lambda a \in I$, so $\lambda \in K = \langle d \rangle$, and hence $x \in \langle cd \rangle$. Thus $I \subseteq \langle cd \rangle$.

$\langle cd \rangle \subseteq I$. There exist $u \in R, v \in I$ such that $c = ua + v$ (since c generates $J = \langle a \rangle + I$). Also $da \in I$ (since d generates K). Then

$$cd = (ua + v)d = u(ad) + vd \in I,$$

since $ad = da \in I$ and $v \in I$. Hence $\langle cd \rangle \subseteq I$.

Together, $I = \langle cd \rangle$, contradicting $I \in S$. So $S = \emptyset$, i.e. every ideal of R is principal. $\square$

Theorem 1.16. Let R be a ring such that every prime ideal is finitely generated. Then every ideal of R is finitely generated.

Proof. Let $A = \{I \text{ ideal of } R \mid I \text{ is not finitely generated}\}$. By Zorn's Lemma, let M be a maximal element of A .

Claim: M is a maximal ideal. Suppose not, so there is an ideal N with $M \subsetneq N \subsetneq R$. Since N is maximal (hence prime), it is finitely generated by hypothesis: there exist $a_1, \dots, a_n \in R$ such that $N = \langle a_1, \dots, a_n \rangle$.

Since $M \subsetneq N$, some $a_i \notin M$. Then $M + \langle a_i \rangle \supsetneq M$, so by maximality of M in A , $M + \langle a_i \rangle \notin A$; hence there exist $b_1, \dots, b_k \in R$ such that

$$M + \langle a_i \rangle = \langle b_1, \dots, b_k \rangle.$$

$\square$

Lemma 1.17. Let I, J_1, J_2 be ideals of R such that $I \subseteq J_1 \cup J_2$. Then either $I \subseteq J_1$ or $I \subseteq J_2$.

Proof. FTSOC, suppose $I \setminus J_1 \neq \emptyset$ and $I \setminus J_2 \neq \emptyset$. Let $x \in I \setminus J_1$ and $y \in I \setminus J_2$; since $I \subseteq J_1 \cup J_2$, this forces $x \in J_2$ and $y \in J_1$. Also $x + y \in I \subseteq J_1 \cup J_2$, so $x + y \in J_1$ or $x + y \in J_2$; WLOG $x + y \in J_1$. Since $y \in J_1$, $x = (x + y) - y \in J_1$, contradicting $x \notin J_1$. $\square$

Theorem 1.18: Prime Avoidance. Let $I, J_1, \dots, J_n$ be ideals of R such that

- (1) $I \subseteq J_1 \cup \dots \cup J_n$,
- (2) $J_3, \dots, J_n \in \text{Spec } R$.

Then $I \subseteq J_i$ for some i .

Proof. We induct on n . The case $n = 2$ needs no primality hypothesis and is the lemma above; assume $n \geq 3$ and the result for any collection of $n - 1$ such ideals.

We prove the contrapositive: if $I \not\subseteq J_i$ for every i , then $I \not\subseteq \bigcup_{i=1}^n J_i$.

Fix i . Apply the induction hypothesis to I together with the $n - 1$ ideals $\{J_j\}_{j \neq i}$ (which still has all but at most two members prime): if $I \subseteq \bigcup_{j \neq i} J_j$, the induction hypothesis would give $I \subseteq J_j$ for some $j \neq i$, contradicting our assumption. So $I \not\subseteq \bigcup_{j \neq i} J_j$, and we may choose

$$z_i \in I \setminus \bigcup_{j \neq i} J_j.$$

Since $I \subseteq \bigcup_{i=1}^n J_i$ by hypothesis (1), and $z_i \notin J_j$ for every $j \neq i$, we must have $z_i \in J_i$, for every i .

Set

$$z := z_1 z_2 \cdots z_{n-1} + z_n \in I$$

(a sum of an element of I and a product of elements of I , hence itself in I). We show $z \notin J_i$ for every i , which gives $I \not\subseteq \bigcup_{i=1}^n J_i$, as desired.

- If $i \in \{1, \dots, n - 1\}$: since $z_i \in J_i$ is one of the factors of $z_1 \cdots z_{n-1}$, this product lies in J_i . If $z \in J_i$, then $z_n = z - z_1 \cdots z_{n-1} \in J_i$, contradicting $z_n \notin J_i$ (as $i \neq n$).
- If $i = n$: if $z \in J_n$, then since $z_n \in J_n$ as well, $z_1 \cdots z_{n-1} = z - z_n \in J_n$. As J_n is prime, some factor $z_k \in J_n$ for a $k \in \{1, \dots, n - 1\}$, contradicting $z_k \notin J_n$ (as $k \neq n$).

This completes the induction. □

Proposition 1.19. An element of $\mathbb{Q}(i)$ lies in $\mathbb{Z}[i]$ if and only if it is the root of a monic polynomial over $\mathbb{Z}$.

Proof. Let $a + bi \in \mathbb{Q}(i)$. Then $a + bi$ is a root of

$$P(X) = X^2 - 2aX + (a^2 + b^2) \in \mathbb{Q}[X].$$

Claim: $2a \in \mathbb{Z}$ and $a^2 + b^2 \in \mathbb{Z}$ together imply $a, b \in \mathbb{Z}$.

Since $2a \in \mathbb{Z}$, $(2a)^2 \in \mathbb{Z}$. Since $a^2 + b^2 \in \mathbb{Z}$, $4(a^2 + b^2) = (2a)^2 + (2b)^2 \in \mathbb{Z}$, so $(2b)^2 = 4(a^2 + b^2) - (2a)^2 \in \mathbb{Z}$. As $2b \in \mathbb{Q}$ has integer square, $2b \in \mathbb{Z}$.

Now $(2a)^2 + (2b)^2 = 4(a^2 + b^2) \equiv 0 \pmod{4}$. Since the square of any integer is $\equiv 0 \pmod{4}$ if it

is even, and $\equiv 1 \pmod{4}$ if it is odd, the only way for the sum to be $\equiv 0 \pmod{4}$ is

$$(2a)^2 \equiv (2b)^2 \equiv 0 \pmod{4},$$

forcing $2a, 2b$ to both be even, i.e. $a, b \in \mathbb{Z}$.

Since $a + bi \in \mathbb{Z}[i]$ exactly when $a, b \in \mathbb{Z}$, and $a + bi$ is a root of a monic polynomial over $\mathbb{Z}$ exactly when $2a, a^2 + b^2 \in \mathbb{Z}$ (the coefficients of $P(X)$), the claim gives the equivalence. $\square$

Definition 1.20. Let $R \hookrightarrow S$ be rings. An element $\alpha \in S$ is called *integral* over R if α is a root of a monic polynomial

$$X^n + a_{n-1}X^{n-1} + \cdots + a_0$$

over R (i.e. with $a_0, \dots, a_{n-1} \in R$).

1.4 Lecture 4

Definition 1.21. Let R be a subring of S , and let $U \subseteq S$ be a subset. Define

$$R[U] := \text{the smallest subring of } S \text{ containing } R \text{ and } U.$$

Remark 1.22. For $U, V \subseteq S$,

$$R[U][V] = R[U \cup V] = R[V][U].$$

In particular, if $V = \{u_1, \dots, u_n\}$, then

$$R[u_1, \dots, u_n] = R[u_1][u_2] \cdots [u_n].$$

Proposition 1.23.

$$R[u] = \{a_0 + a_1u + \cdots + a_nu^n \mid a_i \in R, n \in \mathbb{N}_{\geq 0}\}.$$

Proof. Let $T = \{a_0 + a_1u + \cdots + a_nu^n \mid a_i \in R, n \in \mathbb{N}_{\geq 0}\}$.

T is a **subring** of S . It contains 1 (take $n = 0, a_0 = 1$). It is closed under addition: given two such expressions, pad the shorter one with zero coefficients to a common degree and add coefficientwise, which stays in R since R is a ring. It is closed under multiplication: expanding a product of two such expressions and collecting powers of u , each resulting coefficient is a finite sum of products of elements of R , hence lies in R .

$R[u] \subseteq T$. Since $R \subseteq T$ (take $n = 0$) and $u \in T$ (take $n = 1, a_0 = 0, a_1 = 1$), and T is a subring of S containing R and u , while $R[u]$ is by definition the *smallest* such subring, we get $R[u] \subseteq T$.

$T \subseteq R[u]$. Since $R[u]$ is a subring containing R and u , it is closed under addition and multiplication. So for any $a_0, \dots, a_n \in R \subseteq R[u]$, each $a_iu^i \in R[u]$, and hence $a_0 + a_1u + \cdots + a_nu^n \in R[u]$.

Combining both containments, $R[u] = T$. □

Example 1.24. The map $\mathbb{Z}[X] \rightarrow \mathbb{Z}[i]$, $X \mapsto i$, is a surjective ring morphism with kernel $\langle X^2 + 1 \rangle$.

Proposition 1.25. Let R, S be rings and $\varphi : R \rightarrow S$ be a morphism. Let $\alpha \in S$ be an element. Then φ can be uniquely extended to a morphism $\tilde{\varphi} : R[X] \rightarrow S$ such that $\tilde{\varphi}|_R = \varphi$ and $\tilde{\varphi}(X) = \alpha$.

Proof sketch. Define $\tilde{\varphi} : R[X] \rightarrow S$ by

$$\tilde{\varphi}(a_0 + a_1X + \cdots + a_nX^n) = \varphi(a_0) + \varphi(a_1)\alpha + \cdots + \varphi(a_n)\alpha^n,$$

and $\tilde{\varphi}(A \cdot B) = \tilde{\varphi}(A) \cdot \tilde{\varphi}(B)$. □

Let R be a subring of S , and let $u \in S$. Then there is a unique morphism $\varphi : R[X] \rightarrow S$ such that $\varphi|_R = \text{id}_R$ and $\varphi(X) = u$ (apply the above with $\varphi = \text{id}_R$, $\alpha = u$). Then $\text{Im}(\varphi) = R[u]$; letting $I = \ker \varphi$,

$$R[X]/I \cong R[u].$$

Notation. $I \cap R = \langle 0 \rangle$.

Conversely, let I be an ideal of $R[X]$ such that $I \cap R = \langle 0 \rangle$. We have the natural projection map

$$R[X] \longrightarrow R[X]/I.$$

Since $I \cap R = 0$, $R \hookrightarrow R[X]/I$, so we can identify $a \in R$ with its image $a + I$.

Question. What is the image of $R[X]$ in $R[X]/I$? (It will be determined by $\text{Im}(R)$ and that of X .)

Under the projection, $X \mapsto X + I$. Writing $u = X + I$,

$$R[X]/I \cong R[u].$$

Definition 1.26. Let $R \hookrightarrow S$. S is said to be an *integral extension* of R if each $\alpha \in S$ is integral over R .

Notation. (Recall.) Let R be a commutative ring with 1, and let $A \in M_n(R)$. Then

$$A \cdot \text{adj}(A) = \text{adj}(A) \cdot A = \det(A) \cdot I_n.$$

Theorem 1.27. Let $R \hookrightarrow S$ and $b \in S$. Then

$$b \text{ is integral over } R \iff R[b] \text{ is a finitely generated } R\text{-module}.$$

Proof. ($\Rightarrow$) Assume b is integral over R ; then there is a monic polynomial $f(X) \in R[X]$ such that

$f(b) = 0$. Let $\deg f = n \geq 1$. Take any $g(b) \in R[b]$. By the division algorithm,

$$g(X) = q(X)f(X) + r(X), \quad \deg r < n,$$

so $g(b) = r(b)$. Hence $R[b]$ is generated (as an R -module) by $\{1, b, \dots, b^{n-1}\}$.

($\Leftarrow$) Conversely, let $R[b]$ be a finitely generated R -module, say $R[b] = \langle w_1, \dots, w_n \rangle$. Then for each $i \in [n]$,

$$bw_i = a_{i1}w_1 + \dots + a_{in}w_n, \quad a_{ij} \in R,$$

i.e. $(bI - A)\vec{w} = 0$, where $A = (a_{ij})$. Multiplying by $\text{adj}(bI - A)$,

$$\det(bI - A) w_i = 0 \quad \forall i \in [n].$$

Since $1 \in R[b] = \langle w_1, \dots, w_n \rangle$, we get $\det(bI - A) = 0$, which is a monic equation in b (of degree n). Hence b is integral over R . $\square$

Theorem 1.28. Let $R \hookrightarrow S$, and let $b_1, b_2, \dots, b_n \in S$. Then each b_i is integral over R if and only if $R[b_1, \dots, b_n]$ is a finitely generated R -module.

Proof. Induct on n . The case $n = 1$ is the previous theorem.

For $n > 1$: assume $b_1, \dots, b_n$ are integral over R . In particular b_n is integral over R , hence also integral over $R[b_1, \dots, b_{n-1}]$; by the previous theorem (applied over the base ring $R[b_1, \dots, b_{n-1}]$), $R[b_1, \dots, b_{n-1}][b_n]$ is a finitely generated $R[b_1, \dots, b_{n-1}]$ -module, which (combined with the induction hypothesis that $R[b_1, \dots, b_{n-1}]$ is a finitely generated R -module) makes $R[b_1, \dots, b_n]$ a finitely generated R -module.

The converse is the same as the $n = 1$ case. $\square$

Exercise 1.29. Let $R \hookrightarrow S$, and let $a, b \in S$ be integral over R . Then $a \pm b$ and ab are integral over R .

CHAPTER 2

PROBLEM DISCUSSIONS

2.1 Assignment 0

Convention. By a ring, we mean a commutative ring with 1. Also, we exclude the zero ring.

1. Let I be an ideal of R and $\text{rad}(I) := \{a \in R \mid a^n \in I \text{ for some } n \geq 1\}$. Prove the following:

(a) $\text{rad}(\text{rad}(I)) = \text{rad}(I)$

Solution. The containment $\text{rad}(I) \subseteq \text{rad}(\text{rad}(I))$ is immediate, since $I \subseteq \text{rad}(I)$ and $\text{rad}(-)$ is monotone.

Conversely, let $x \in \text{rad}(\text{rad}(I))$. Then $x^n \in \text{rad}(I)$ for some n , so $(x^n)^m = x^{nm} \in I$ for some m . Hence $x \in \text{rad}(I)$. $\square$

(b) $\text{rad}(IJ) = \text{rad}(I \cap J) = \text{rad}(I) \cap \text{rad}(J)$

Solution. We show $\text{rad}(I) \cap \text{rad}(J) \subseteq \text{rad}(IJ) \subseteq \text{rad}(I \cap J) \subseteq \text{rad}(I) \cap \text{rad}(J)$, giving all three sets equal.

- $\text{rad}(I) \cap \text{rad}(J) \subseteq \text{rad}(IJ)$: if $x \in \text{rad}(I) \cap \text{rad}(J)$, then $x^n \in I$ and $x^m \in J$ for some n, m , so $x^{n+m} = x^n x^m \in IJ$, i.e. $x \in \text{rad}(IJ)$.
- $\text{rad}(IJ) \subseteq \text{rad}(I \cap J)$: since $IJ \subseteq I \cap J$, this follows from monotonicity of rad .
- $\text{rad}(I \cap J) \subseteq \text{rad}(I) \cap \text{rad}(J)$: since $I \cap J \subseteq I$ and $I \cap J \subseteq J$, monotonicity gives $\text{rad}(I \cap J) \subseteq \text{rad}(I)$ and $\text{rad}(I \cap J) \subseteq \text{rad}(J)$.

$\square$

(c) $\text{rad}(I) = R$ if and only if $I = R$

Solution. $(\Rightarrow)$ If $\text{rad}(I) = R$, then for every $a \in R$ there is $n \in \mathbb{N}$ with $a^n \in I$. Taking $a = 1$: $1^n = 1 \in I$, so $I = R$.

$(\Leftarrow)$ Trivial: if $I = R$ then $\text{rad}(I) = R$ as well. $\square$

(d) $\text{rad}(I + J) = \text{rad}(\text{rad}(I) + \text{rad}(J))$

Solution. ($\subseteq$) Let $x \in \text{rad}(I + J)$. Then $x^k = u + v$ for some $k \in \mathbb{N}$, $u \in I$, $v \in J$. Since $u \in I \subseteq \text{rad}(I)$ and $v \in J \subseteq \text{rad}(J)$, $x^k \in \text{rad}(I) + \text{rad}(J)$, so $x \in \text{rad}(\text{rad}(I) + \text{rad}(J))$.

($\supseteq$) Since $I \subseteq I + J$, monotonicity gives $\text{rad}(I) \subseteq \text{rad}(I + J)$; similarly $\text{rad}(J) \subseteq \text{rad}(I + J)$. As $\text{rad}(I + J)$ is an ideal, $\text{rad}(I) + \text{rad}(J) \subseteq \text{rad}(I + J)$. Applying $\text{rad}(-)$ and using part (a),

$$\text{rad}(\text{rad}(I) + \text{rad}(J)) \subseteq \text{rad}(\text{rad}(I + J)) = \text{rad}(I + J).$$

□

(e) If $P \in \text{Spec}(R)$, then $\text{rad}(P^n) = P$ for all $n > 0$

Solution. First, $\text{rad}(P) = P$: the containment $P \subseteq \text{rad}(P)$ always holds, and $\text{rad}(P) \subseteq P$ follows since P is prime (if $x^n \in P$, primality forces $x \in P$).

Now induct on n . The base case $n = 1$ is $\text{rad}(P) = P$, shown above. For the inductive step, using part (b) with $I = P^{n-1}$, $J = P$:

$$\text{rad}(P^n) = \text{rad}(P^{n-1} \cdot P) = \text{rad}(P^{n-1}) \cap \text{rad}(P) = P \cap P = P,$$

using the induction hypothesis $\text{rad}(P^{n-1}) = P$.

□

2. Let $R[X]$ be the polynomial ring over a ring R . Prove that the nilradical of $R[X]$ is equal to the Jacobson radical of $R[X]$.

Solution. Since every maximal ideal is prime, intersecting over the (larger) family of prime ideals gives a smaller or equal intersection than intersecting over maximal ideals alone; hence, for any ring, $\mathcal{N}(R[X]) \subseteq J(R[X])$.

For the reverse containment, we use the identity $\mathcal{N}(R[X]) = (\mathcal{N}(R))[X]$ (proof deferred). Take $f(X) \in J(R[X])$. By the standard characterization of the Jacobson radical, $1 - g(X)f(X)$ is a unit for every $g(X) \in R[X]$; taking $g(X) = -X$,

$$1 + Xf(X) \text{ is a unit in } R[X].$$

By the Lecture 1 proposition on units of $R[X]$, since the constant term of $1 + Xf(X)$ is 1 (a unit), all remaining coefficients of $1 + Xf(X)$ – i.e. all coefficients of $f(X)$ – are nilpotent. Hence $f(X) \in (\mathcal{N}(R))[X] = \mathcal{N}(R[X])$.

Therefore $J(R[X]) \subseteq \mathcal{N}(R[X])$, and combined with the reverse containment above, $\mathcal{N}(R[X]) = J(R[X])$. □

3. Let $R[X]$ be the polynomial ring over a ring R . Let $J \subset R$ be an ideal. Define

$$J[X] := \{a_0 + a_1X + \cdots + a_nX^n \mid a_i \in J \forall i\}.$$

Prove that $J[X]$ is an ideal of $R[X]$. Give an example of an ideal of $R[X]$ which is not of this form. Prove that the rings $R[X]/J[X]$ and $(R/J)[X]$ are isomorphic.

Solution. That $J[X]$ is an ideal of $R[X]$ is a direct verification (closed under addition and absorbs multiplication by $R[X]$, coefficientwise, since J is an ideal of R).

Example of an ideal not of this form. Let

$$I = \{p(X) \in R[X] \mid \exists q(X) \in R[X] \text{ such that } p(X) = X \cdot q(X)\} = \langle X \rangle,$$

the ideal of polynomials with zero constant term. This is not of the form $J[X]$ for any proper ideal $J \subsetneq R$: an element of $J[X]$ has *every* coefficient in J , whereas I places no constraint on coefficients beyond the constant term.

Isomorphism. Define $\varphi : R[X]/J[X] \rightarrow (R/J)[X]$ by

$$\varphi(r(X) + J[X]) = \sum_{i=0}^{\deg r} (r_i + J) X^i,$$

i.e. reduce each coefficient of $r(X)$ modulo J . This is induced by the coefficientwise reduction map $R[X] \rightarrow (R/J)[X]$, which is a surjective ring morphism with kernel exactly $J[X]$; by the First Isomorphism Theorem, φ is a well-defined ring isomorphism. $\square$

4. A commutative ring is called a *local ring* if it has only one maximal ideal. Let R be a local ring with unique maximal ideal $\mathfrak{m}$. Prove that any element $a \in R \setminus \mathfrak{m}$ is a unit in R . Conversely, let R be a ring such that the set of non-units forms an ideal M . Prove that R is local with maximal ideal M .

Solution. $(\Rightarrow)$ FTSOC, suppose $a \in R \setminus \mathfrak{m}$ is not a unit. Then $\langle a \rangle \neq R$, i.e. $1 \notin \langle a \rangle$. Let

$$A = \{I \subseteq R \text{ an ideal} \mid a \in I \text{ but } 1 \notin I\}.$$

Since $\langle a \rangle \in A$, $A \neq \emptyset$. By Zorn's Lemma, let M be a maximal element of A (ordered by inclusion). One checks M is a maximal ideal of R . Since R is local, $M = \mathfrak{m}$, and since $a \in M$, we get $a \in \mathfrak{m}$ – contradicting $a \notin \mathfrak{m}$. Hence a must be a unit.

(Converse) Suppose $M = \{x \in R \mid x \text{ is not a unit}\}$ is an ideal. Every proper ideal I of R consists entirely of non-units (else I would contain a unit, forcing $I = R$), so $I \subseteq M$. In particular M itself is a proper ideal (as $1 \notin M$), and since every proper ideal is contained in M , M is the unique maximal ideal of R . Hence R is local with maximal ideal M . $\square$

5. An element $e \in R$ is called an *idempotent* if $e^2 = e$. Let R be a local ring. Prove that the only idempotents of R are 0 and 1.

Solution. **Lemma.** If R is a local ring, then for every $a \in R$, either a or $1 - a$ is a unit.

Proof. Suppose neither a nor $1 - a$ is a unit. Then $a \in \mathfrak{m}$ and $1 - a \in \mathfrak{m}$ (the unique maximal ideal), so $1 = a + (1 - a) \in \mathfrak{m}$, forcing $\mathfrak{m} = R$ – a contradiction.

Now let $e \in R$ be idempotent: $e^2 = e$, i.e. $e(e - 1) = 0$. By the lemma, at least one of $e, 1 - e$ is a unit; multiplying $e(e - 1) = 0$ by its inverse forces the other factor to vanish. Hence $e = 0$ or $e = 1$. $\square$

6. Let $R = \mathbb{Z}[i]$. Prove that the ideal $P = \langle 1 + i \rangle$ is a prime ideal of R . Prove that $\langle 2 \rangle = P^2$ in R .
7. Prove that the ring $\mathbb{Z}[i]/\langle 2 + i \rangle$ is isomorphic to the ring $\mathbb{Z}/5\mathbb{Z}$.
8. Let R be a ring and $a, b \in R$ be two *comaximal* elements, i.e., there exist $x, y \in R$ such that $xa + yb = 1$. Prove that for any $m, n \in \mathbb{N}$, one has a^m and b^n are also comaximal (i.e., there exist $\lambda, \mu \in R$ such that $\lambda a^m + \mu b^n = 1$).
9. Let R be an integral domain. Let $a, b \in R$ be such that their gcd exists. If d, d' are both gcds of a, b , prove that d and d' are associates.
10. Let R be an integral domain such that $\gcd(a, b)$ exists for any $a, b \in R \setminus \{0\}$. Then prove that $\text{lcm}(a, b)$ exists for any $a, b \in R \setminus \{0\}$. (Hint: First prove that $\gcd(ma, mb) = m \gcd(a, b)$.)
11. Let R be an integral domain such that $\text{lcm}(a, b)$ exists for any $a, b \in R \setminus \{0\}$. Then prove that $\gcd(a, b)$ exists for any $a, b \in R \setminus \{0\}$.
12. Let R be an integral domain such that $\gcd(a, b)$ exists for any $a, b \in R \setminus \{0\}$. If $\gcd(a, b) = 1$ and $\gcd(a, c) = 1$, then prove that $\gcd(a, bc) = 1$.
13. Let $R = \mathbb{Z}[\sqrt{-5}]$. Verify the following:

(a) There are two distinct (i.e. non-associate) factorizations of 6 into irreducibles in R :

$$6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5}).$$

(b) The ideals $\langle 2, 1 \pm \sqrt{-5} \rangle, \langle 3, 1 \pm \sqrt{-5} \rangle$ are all prime ideals.

(c) The following factorization of ideals into prime ideals holds:

- $\langle 2 \rangle = \langle 2, 1 + \sqrt{-5} \rangle^2$,
- $\langle 3 \rangle = \langle 3, 1 + \sqrt{-5} \rangle \langle 3, 1 - \sqrt{-5} \rangle$,
- $\langle 1 + \sqrt{-5} \rangle = \langle 2, 1 + \sqrt{-5} \rangle \langle 3, 1 + \sqrt{-5} \rangle$,
- $\langle 1 - \sqrt{-5} \rangle = \langle 2, 1 + \sqrt{-5} \rangle \langle 3, 1 - \sqrt{-5} \rangle$.

(d) Uniqueness of factorization is restored in (a) at the level of ideals.

(e) The ideal $\langle 2, 1 + \sqrt{-5} \rangle$ is not a principal ideal.

14. Show that $\langle 2, 1 + \sqrt{-17} \rangle$ is not a principal ideal in $\mathbb{Z}[\sqrt{-17}]$. Is this a prime ideal?

15. Let R be an ID. Assume that:

(a) any two $a, b \in R \setminus \{0\}$ have gcd, say, d , and there are $x, y \in R$ such that $d = ax + by$;

- (b) if $a_1, a_2, \dots$ are nonzero elements of R such that a_{i+1} divides a_i for $i = 1, 2, \dots$, then there is $N > 0$ such that $a_n = u_n a_N$ for some unit $u_n \in R$ for all $n \geq N$.

Show that R is a PID.

16. Prove that $R[X]/\langle X^3 + X \rangle$ is a product of fields.
17. Let $m, n \in \mathbb{Z}$ be two nonzero elements and $d = \gcd(m, n)$ in $\mathbb{Z}$. Prove that d is also a gcd of m, n in $\mathbb{Z}[i]$.
18. Let $m, n \in \mathbb{Z}$ be two nonzero elements and let m divide n in $\mathbb{Z}[i]$. Does m also divide n in $\mathbb{Z}$?
19. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of R -modules. Let M', M'' be both finitely generated. Show that M is finitely generated.
20. From Dummit–Foote: Exercise 10.1: 5, 8, 9, 10, 11, 15.
21. From Dummit–Foote: Exercise 10.2: 3, 4, 5, 6, 7, 9, 10, 11, 12, 13.
22. An R -module is called *simple* if it is not the zero module and it has no proper submodule.
 - (a) Prove that an R -module M is simple if and only if it is isomorphic (as an R -module) to $R/\mathfrak{m}$, where $\mathfrak{m}$ is a maximal ideal of R . Show also that M is generated by any non-zero element.
 - (b) Let M, N be simple R -modules. Prove that a module morphism $\phi : M \rightarrow N$ is either the zero morphism or an isomorphism. Conclude that if M is simple, $\text{Hom}_R(M, M)$ is a division ring.
23. Let M be an additive abelian group and $\text{End}(M)$ be the set of all group morphisms from M to M .
 - (a) Show that $\text{End}(M)$ is a ring with identity (where addition is defined pointwise, and multiplication is composition of maps).
 - (b) Show that M is an $\text{End}(M)$ -module under the action:

$$(f, m) \mapsto fm := f(m).$$
 - (c) Let R be a ring and $\phi : R \rightarrow \text{End}(M)$ be a ring morphism. Show that M is an R -module under the action:

$$(r, m) \mapsto rm := \phi(r)(m).$$
24. (Converse of 4(c)): Let R be a ring and M be an R -module. Show that there is a ring morphism $\phi : R \rightarrow \text{End}(M)$.
25. Let M be an R -module and A, B, C be its submodules such that

$$A \subseteq B, \quad A + C = B + C, \quad A \cap C = B \cap C.$$

Prove that $A = B$.

26. Let $f : M \rightarrow N$ be an R -module morphism. Let A be a submodule of M . Prove that $f^{-1}(f(A)) = A + \text{Ker}(f)$. Now let B be a submodule of N . Prove that $f(f^{-1}(B)) = B \cap \text{Im}(f)$. Also, show that

$$f(A \cap f^{-1}(B)) = f(A) \cap B.$$

27. Let R be a commutative ring. Show that a map $f : R \times R \rightarrow R$ is an R -module morphism if and only if $f(x, y) = \alpha x + \beta y$ for some $\alpha, \beta \in R$.
28. Let $f : M \rightarrow N$ be an R -module morphism. Show that f can be expressed as the composition of three R -module morphisms (for appropriate R -modules): $f = g \circ h \circ k$, where g is surjective, h is an isomorphism, and k is injective.
29. Let R be an integral domain and $a \in R$ be non-zero. Let n be any positive integer. Prove that Ra^n / Ra^{n+1} is isomorphic to R/Ra (as R -modules).

BIBLIOGRAPHY

- [AM69] M. F. Atiyah and I. G. Macdonald. *Introduction to Commutative Algebra*. Addison-Wesley, 1969.
- [Bly90] T. S. Blyth. *Module Theory: An Approach to Linear Algebra*. Clarendon Press, Oxford University Press, Oxford, 2nd edition, 1990.
- [DF04] David S. Dummit and Richard M. Foote. *Abstract Algebra*. John Wiley & Sons, Hoboken, NJ, 3rd edition, 2004.