
INTRODUCTION TO MODEL CATEGORIES

INCOMPLETE

Karshit Joshi

Math and Physics Student
Chennai Mathematical Institute
9th September 2026

Contents

1 Preliminaries	3
2 Derived Functors	4
2.1 Homotopical Categories and Derived Functors	4
2.2 Enriched Category Theory	10
3 Weak Factorisation Systems	13
4 Model Categories	13

1 Preliminaries

This is an exposition written by me to take notes of what I read about Model Categories and certain ideas that I think are very useful while reading Model Categories. Any mistakes made in this exposition is purely mine and please report them to [karshitkjoshi@cmi.ac.in]. I am mostly using "Homotopical Algebra" by Daniel Quillen[1], "Model Categories" by Mark Hovey[2] and "Categorical Homotopy Theory" by Emily Riehl[3] as my primary resources to read about the topic from, and will try my best to mention any other resources that I use along the way in the bibliography. I am grateful to Chennai Mathematical Institute to expose me to the resources and wonderful people that helped me during this writing.

I am very grateful to Prof. Aditya Karnataki, who encouraged me to write this exposition and introducing me to writing exposition which led me to find an appreciation for writing mathematical expositions.

2 Derived Functors

2.1 Homotopical Categories and Derived Functors

An important question has always been in different settings and even in category theory, "Are these two objects isomorphic ?", while the answer to the question depends on the what we consider an isomorphism. Fixing the context, it could perhaps be easier or difficult to differentiate between objects.

A good strategy in our toolset has always been to classify objects "upto some property". Note that the creation of such classification creates a new category where the context is changed to something which is much coarser than the current context that we are trying to work in. This often helps us to focus ourselves towards a particular way to differentiate objects rather than having the vague goal of differentiating objects with nothing ahead of us. Common examples of this phenomenon include Homotopy Groups, Homology Groups, Euler Characteristic, Betti Numbers, etc. We shall now try to build this notion in category theory for abstract categories.

Suppose we start with a category $\mathcal{C}$. We have some property $\mathcal{P}$ along which we can test our objects to see if they are "equivalent" or not. If identifying $\mathcal{P}$ on 2 different objects tells us that they have the same property. Our desire is to consider those objects to be the same. A natural way to store all of this data of identification is in a category, with the purpose that we will inherit all the data naturally by doing a construction on $\mathcal{C}$.

We are now clear that we must perform construction on $\mathcal{C}$ to create a category which characterizes the data that we get from $\mathcal{P}$. There are naturally 2 ways to do this.¹

- Consider the objects of $\mathcal{C}$, consider the equivalence relation imposed by $\mathcal{P}$ on the collection of object of $\mathcal{C}$. We quotient by the equivalence relation. We also forget all the maps between objects which were identified. The residue will turn out to be a category with a few more natural steps. This can be thought of as skeleton category of localisation given by the second point.
- Consider the maps of $\mathcal{C}$, formally introduce the inverses of maps between objects, "which characterize the equivalence of $\mathcal{P}$ ". This just naturally forms a category once again after completing all possible compositions with the new maps.²

Both the above ways are useful in different ways that we will explore throughout the span of this text. Notice the difference, that the first construction forgets how things are actually equivalent whereas the second does not due to the existence of all the maps which "characterize the equivalence of $\mathcal{P}$ ". This especially comes into the picture if you view the first construction as the manifestation of "existence of an equivalence" rather than "equivalent

¹A person who already has a vague idea about infinity categories might realize that the 2 above localisations can be unified under a kind of construction that can be done in infinity categories called n-truncation. This allows to truncate all data from n and above to get rid of the precise witnesses which exhibit the homotopy between (n-1) morphisms and replace them with just existence statements

²There is an interesting question for keen observers. There are 2 ways to achieve this. (1) Introduce the "abstract" formal inverses of all maps we consider weak equivalences (2) Create an equivalence relation between maps and quotient by the equivalence relation which will force certain "concrete" maps to become formal inverses of other "concrete" maps. When exactly are these constructions equivalent/isomorphic ? In some sense, the answer for why they would be equivalent is that having a left inverse upto some homotopy can make the inverse unique upto homotopy which we can test if all such inverses have a characterisation in the base category or not. I still need to know more about this though.

with the witness map”, wherein the second construction is much more suited to the latter. This often comes into play when you are considering n -truncations of an infinity category which are the objects we are truly trying to study indirectly via model categories. Due to the previous reason, this also plays a significant role in homotopy type theory.[Note that the first construction is also technically inline with Leibniz equivalence of identibles which says that ”if two objects have the same interactions, they are the same”.i.e. Isomorphisms should not exist as objects, instead all isomorphisms must be equalities. This plays a major role in homotopy type theory.]

Since the second construction retains much more information from the ambient category. It is natural to call it the appropriate ”localisation” of the category. The terminology of localisation is inherited from ring theory, wherein we localize a ring against a prime ideal which requires to invert all elements outside the prime ideal.

We now turn our focus to actually selecting the maps of the category $\mathcal{C}$, to be our equivalences which ”we believe” characterizes some property $\mathcal{P}$. Here is a subtle but important point that we ”choose” a property to classify objects against. In the context of abstract categories, the selection of maps in the category is in direct correspondence to the abstract property $\mathcal{P}$. Hence, the selection of property is equivalent to selection of which maps to invert. Hence, the structure we obtain is a deliberate construction, not a natural structure to consider, i.e. The structure is a choice, not a natural byproduct of $\mathcal{C}$.

It is also natural to expect that all isomorphisms are also a part of the collection of maps we invert mainly because, if $\mathcal{C}$ cannot distinguish between 2 objects, they must possess the same set of properties at all levels [Note that this is essentially the application of Yoneda Lemma]. Hence, this helps us to arrive at our first definition.

Definition 2.1. A **homotopical category** is a category $\mathcal{C}$ equipped with a wide* subcategory $\mathcal{W}$ such that, for any composable triple of arrows,i.e. f, g, h . If hg and gf are in $\mathcal{W}$, so are f, g, h , and hgf .

The arrows in $\mathcal{W}$ are called equivalences. The property above concerning the maps is called the 2-of-6 property.[abbrieviation of ”2 out of 6”] (*) By a wide subcategory, we mean a subcategory which contains all identities of the category $\mathcal{C}$ and hence by 2-of-6 property, all isomorphisms of $\mathcal{C}$ aswell.

There is a weaker condition that is also often considered instead of the 2-of-6 property. It is called 2-of-3 property.

Definition 2.2. 2-of-3 Property occurs if any 2 of the 3 possible composition of 2 composable maps are in $\mathcal{W}$. Then $\mathcal{W}$ contains all 3.

A small caveat in the above 2 definitions is that $\mathcal{W}$ must also contain all the identities for each object in $\mathcal{C}$ which has a clear enough reason to be included.

Now that we have concretely defined what is the data that we would like to work with. We now approach the problem of carrying out the construction of the localisation of the category as we previously mentioned. The following explicit construction is due to Pierre Gabriel and Michel Zisman. Its morphisms are equivalence classes of finite zig-zags (i.e.compositions which maps from $\mathcal{W}$ in the string with finite length) with arrows in $\mathcal{W}$ to be inverted with

the same name but dual of its original counterpart subjected to the following relations.

Definition 2.3. The localisation category/homotopy category $\text{Ho}\mathcal{C}$ of a homotopical category $(\mathcal{C}, \mathcal{W})$ is the formal localisation of $\mathcal{C}$ at the subcategory $\mathcal{W}$. The explicit construction of the homotopy category is as follows:

- Objects of $\text{Ho}\mathcal{C}$ are the same as $\mathcal{C}$.
- All strings of compositions which contain $\xleftarrow{\mathcal{W}} \xrightarrow{\mathcal{W}}$ or $\xrightarrow{\mathcal{W}} \xleftarrow{\mathcal{W}}$ or identities pointing forward or backwards as a part of the string are equivalent to strings of compositions with the earlier substrings removed. This induces the natural equivalence relation of strings of composable arrows.

One can simply notice that there is a natural functor $\mathcal{C} \xrightarrow{L} \text{Ho}\mathcal{C}$. This functor also possesses the universal property that any other functor which inverts all morphisms in $\mathcal{W}$ (not necessarily only morphisms in $\mathcal{W}$) factors through L .

An important point is that not all morphisms of $\mathcal{C}$ that become isomorphisms should be present in the subcategory $\mathcal{W}$. When this does not happen, we call the homotopical category $\mathcal{C}$ to be saturated [i.e. all morphisms whose inverse exists in $\text{Ho}\mathcal{C}$ are also present in $\mathcal{W}$].

Another important point is that performing this construction on a locally small category does not ensure that it will remain a locally small category. This is simply due to the reason that the collection of objects form a class and each object has its respective equivalence class. There could be a lot of choices of a single map about which object to passthrough for each object's equivalence class. Considering the possibility that the equivalence class of an object may itself form a class. The collection of maps between any two maps can form a class. For this reason, there exists another richer localisation than the one we came up with, which is inspired from the localisation of rings, called "Calculus of Fractions" due to Pierre Gabriel, Pierre Zisman. However, I plan to delve into it just yet.

Therefore, one of the problems that the model category construction solves is to keep the localisation of the category to be locally small.

We now try to study functors between Homotopical Categories which can extend to maps between the respective homotopy categories under suitable conditions on the functor itself. The condition on the functor is also natural, in the sense that it must send equivalences to equivalences ,i.e. it preserves all equivalences. We call such functors, "Homotopical Functors", by the universal property of localisation, it induces a functor between the respective homotopy categories.

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow L_{\mathcal{C}} & & \downarrow L_{\mathcal{D}} \\ \text{Ho}\mathcal{C} & \xrightarrow{F'} & \text{Ho}\mathcal{D} \end{array}$$

It is a natural desire to study functors which are not homotopical. However, when that happens notice that $L_{\mathcal{C}}$ and $L_{\mathcal{D}} \circ F$ form a span diagram, which hints us towards looking at left and right Kan Extensions along the appropriate direction carrying the intuition that we are trying to find the best functor which approximates F while losing the least amount

of information while doing it. This naturally leads to the following definition.

Definition 2.4. A **total left derived functor** LF of a functor F between Homotopical Categories $\mathcal{C}$ and $\mathcal{D}$ is a right Kan extension $Ran_{L_{\mathcal{C}}}(L_{\mathcal{D}} \circ F)$ [We shall also sometimes refer the $Ran_{L_{\mathcal{C}}}(L_{\mathcal{D}} \circ F) \circ L_{\mathcal{C}}$ as the total left derived functor with an equivalent universal property]

This also alludes to being related to derived functors in the homological algebra sense. We are eventually heading towards that direction. Now we introduce the definition of a left derived functor which won't just yet make any sense but its better to go through the definition first and then realise it.

Definition 2.5. A **left derived functor** of $F : \mathcal{C} \rightarrow \mathcal{D}$ is a homotopical functor $\mathcal{L}F : \mathcal{M} \rightarrow \mathcal{N}$ along with a natural transformation $\tau : \mathcal{L}F \Rightarrow F$ such that $L_{\mathcal{D}}\tau : \tau \circ \mathcal{L}F \Rightarrow L_{\mathcal{D}} \circ F$. the last operation is also better known as "whiskering" ,i.e. left derived functor is a direct lift of the total left derived functor along the localisation functor.

The total left derived functor is essentially nothing but the induced functor between the homotopy categories from the functor between the homotopical categories. In classical notion, this will only occur when the functor respects the weak equivalences that we have chosen ,i.e. the functor is well-defined on the isomorphism classes of objects equivalently F is not a homotopy functor. The main problem is that most functors that we are aware of ,i.e. $Hom(-, A)$, $- \otimes A$ do not respect the quasi-isomorphisms in the category chain complexes. This tells you that a total left derived functor. Now, borrowing the intuition for a Kan Extension, One can realise that this is precisely the problem that we would like to solve which is why the goal is to find the total left derived functor ,i.e. the best approximation of F . We now introduce some more definitions which are directly motivated from the topology.


Definition 2.6. A **left deformation** on a homotopical category $\mathcal{C}$ consists of an endofunctor $\mathcal{Q}$ together with a natural "weak equivalence", not natural isomorphism instead its a natural isomorphism as a homotopical endofunctor on the homotopy category $q : \mathcal{Q} \Leftrightarrow 1$.

We are already tip-toeing into the deep waters of model categories, as we go into the following example which considers the weak factorisation system. If $\mathcal{M}$ is a model category with a functorial cofibration-trivial fibration factorisation, then the action of this factorisation on arrows of the form $\phi \rightarrow x$ produces a functor $\mathcal{Q}$ whose image lands in the subcategory of cofibrant objects with natural weak equivalence $q_x : \mathcal{Q}x \sim x$. I just copied the previous example verbatim from Riehl's book but I believe this is an important example to look at.


In the above example, we use a weak factorisation system to break any map into a cofibration and weakly equivalent fibration which allows us to find an object q_x for each x for which $\phi \rightarrow x$ is a cofibration and it is equivalent to x which will constitute a larger functor $q : x \sim q_x$ sending the category to its subcategory of cofibrant objects/replacements. As it will soon turn out that such replacements are of extreme importance to create what

are homotopy limit/colimits which are of grave importance to work in the homotopy category, which in most cases does not even have all limits and colimits. A reasonable question to ask is that why q_x is called a cofibrant object/replacement even though its weak equivalence with x is a fibration. There is a very important reasoning behind this. It is for, How exactly do we deal with extension/lifting problems ?


Many times in mathematics, again and once again you must encounter this precise diagram.



When does exactly such a lift/extension [dotted arrow] exist such that both the triangles commute ? Often times, a lifting or extension problem can be precisely phrased in this way given the existence of an initial and a terminal object.



Lifting Problem



Extension Problem

And it is due to the prominence of these problems which are solved by cofibrant objects because the property of $\phi \rightarrow C$ being a cofibration is precisely that it lifts all maps against any fibration $B \rightarrow D$. This solves the lifting problem for any object C because as long as $B \rightarrow D$ is a fibration, in the homotopy category I can replace C with q_C and lift any map $q_C \rightarrow D$ without any worries. A dual situation will occur when weak factorisation system is trivial-cofibrations and fibrations. Wherein you will be replacing objects with fibrant replacements. This is precisely why a Weak Factorisation System is ingrained into the definition of a Model Category because this allows us to have some level of control over the maps in the homotopy category [while obviously increasing a lot of conditions on the original category like i.e. there existing a weak factorisation system in the first place] and evidently enough, this is enough for us to get limits and colimits of diagrams in the homotopy category which in turn allows us to actually compute legible objects which are significant to us. But given a significant idea of how we are going to solve our problems, we leave this here for the reader to digest this situation as this is one of the major parts of model categories which build the bigger picture. Instead we try to bring back our focus onto realising the

current definitions with our knowledge from Homological Algebra.³

We know that the category of left R -modules have the following property: for any R -module A , there exists a projective module P together with an epimorphism $P \twoheadrightarrow A$. This way, repeating the procedure we get the **projective resolution**, an acyclic (except in degree zero) bounded-below chain complex of projectives that maps quasi-isomorphically to the chain complex concentrated in degree 0 at module A . Making a specific natural choice of projective resolution (free resolution), gives us a left deformation for the homotopical category of the homotopical category of bounded-below chain complexes where the weak equivalences are the quasi-isomorphisms. Dually, any R -module embeds into an injective module, repeating that gives us injective resolutions. Similarly, one can arrive at right deformation on the category of bounded-below cochain complexes.

Any additive functor F from the category of R -modules to category of S -modules induces obvious functors between the respective chain complex categories. These functors preserve the chain homotopy equivalences by very easy verification. But, they do not preserve quasi-isomorphisms which are the actual weak-equivalences that we would like to look at because they are completely faithful to homology/cohomology groups. Although, for projective resolutions, any chain complex made of projective objects, a quasi-isomorphism entails a chain homotopy equivalence which is preserved by $F_\bullet$. The dual versions of the above statements also follows correctly. This also fits into the previous picture that we are essentially taking a cofibrant replacement in the category of chain complexes and since F preserves the quasi-isomorphisms in the subcategory of projective chain complexes. We can get a total derived functor between the respective homotopy category chain complexes.

The classically regarded as the left derived functor of $F : \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$ can now be viewed as below using what we have realised up until now:

$$\mathbf{Mod}_R \xrightarrow{dego} \mathbf{Ch}_{\geq 0}(R) \rightarrow \mathbf{Ch}_{\geq 0}(S) \xrightarrow{H_0} \mathbf{Mod}_S$$

which takes the cofibrant replacement of an R -module A , applies $F_\bullet$, and then computes the 0th homology. Please note that it is very possible that the natural deformation functors does not exist for your abelian category, in which case also this procedure will pass through just by knowing that there is a weak factorisation system on the category. Also note that if you do not have such a natural deformation functor then you will not get a derived functor but instead a total derived functor due to the reason that a person now must make a choice of cofibrant replacement for any object rather than there being a natural choice to pick for each choice.

At this point I expect the reader to understand that why are does one care about fibrations, cofibrations, weak equivalences, weak factorisation systems, fibrant and cofibrant replacements, which are some of the most important objects in understanding/dealing with model categories. Now, it is an adequate point to take a look into homotopy (co)limits and how are they exactly constructed while expecting the reader to have realised the importance of knowing how to reliably construct limits/colimits in the homotopy category and why will they be very useful. Now we shall go into a bit of enriched category theory as it is useful in model categories framework, calculating homotopy (co)limits aswell as understanding models of infinity categories.

³I insist the reader to atleast read the definition of a Weak Factorisation System at this point in time.

2.2 Enriched Category Theory

The core idea behind enriched category theory is that to try to localise a category, you must invert maps which in turn force you to deal with the hom-sets or in many cases hom-objects which now have more structure because our desire to localise. Therefore, to keep track of data as it behaves in the ambient category but also be able to see the consequences of localisation as play around with hom-objects which is precisely what enriching a category is all about, replacing hom-sets with hom-objects in some other category D .

Precisely speaking, we are going make our hom-objects precisely to be spaces/topological spaces which carry the information of homotopies and higher homotopies in the terms of paths, paths between paths, paths between paths between paths, up until infinity (i.e. simplicial set, which can also be realised to be a CW-complex).

Definition 2.7. The base category over which we enrich must be a **symmetric monoidal category** $(V, \times, \star)$. Here V is an ordinary category. $\times : V \times V \rightarrow V$ is a bifunctor and $\star$ is the unit object. It is also equipped with the following natural transformations.

$$v \times w \cong w \times v, u \times (v \times w) \cong (u \times v) \times w, \star \times v \cong v \cong v \times \star$$

We also do not have to worry about the associativity holding at higher levels.

The monoidal product is supposed to serve the composition operation as well as the unit is supposed to be the initial object in the category V to point out the identity map for each endo-object for each object.

Definition 2.8. A V -category $\underline{D}$ consists of

- a collection of objects, i.e. $x, y, z \in \underline{D}$
- for each pair $x, y \in \underline{D}$, a hom-object $\underline{D}(x, y) \in V$.
- for each $x \in \underline{D}$, a morphism $id_x : \star \rightarrow \underline{D}(x, x) \in V$.
- for each triple $x, y, z \in \underline{D}$, a morphism $\circ : \underline{D}(y, z) \times \underline{D}(x, y) \rightarrow \underline{D}(x, z)$ in V .

such that the following diagrams commute for all $x, y, z, w \in \underline{D}$:

$$\begin{array}{ccc}
 \underline{D}(zw, y) \times \underline{D}(y, z) \times \underline{D}(x, y) & \xrightarrow{1 \times \circ} & \underline{D}(z, w) \times \underline{D}(x, z) \\
 \circ \times 1 \downarrow & & \downarrow \circ \\
 \underline{D}(y, w) \times \underline{D}(x, y) & \xrightarrow{\circ} & \underline{D}(x, w) \\
 \\
 \underline{D}(x, y) \times \star & \xrightarrow{1 \times id_x} & \underline{D}(x, y) \times \underline{D}(x, x) \\
 \searrow \cong & & \swarrow \circ \\
 & \underline{D}(x, y) & \\
 \\
 \underline{D}(x, y) \times \star & \xrightarrow{id_y \times 1} & \underline{D}(y, y) \times \underline{D}(x, y) \\
 \searrow \cong & & \swarrow \circ \\
 & \underline{D}(x, y) &
 \end{array}$$

The goal here is to take as much advantage of the base category structure as possible. And it turns out that in many nice categories, currying is possible and there is a parametrized adjunction between the $\times$ and the "internal hom". This alludes to the fact that such an adjunction is only possible in a category when the hom-objects of the base category lie within itself ,i.e. the base category is enriched over itself. This is a necessary but also sufficient condition for currying. We obviously cannot let go of such a nice structure, therefore we are going to introduce a definition for it.

Definition 2.9. We call a monoidal category **closed** if each functor $- \times d : \mathcal{D} \rightarrow \mathcal{D}$ admits a right adjoint $\underline{\mathcal{D}}(d, -)$, the right adjoints in this family of parameterized adjunctions assemble in a unique way into a bifunctor.

$$\underline{\mathcal{D}}(-, -) : \mathcal{D}^{op} \times \mathcal{D} \rightarrow \mathcal{D}$$

such that there exists natural isomorphisms for all $u, v, w \in \mathcal{V}$

$$\mathcal{D}(u \times v, w) \cong \mathcal{D}(u, \underline{\mathcal{D}}(v, w))$$

what follows are the categories which are also called **closed symmetric monoidal categories**, which are natural to understand by now. In the special case that the monoidal product is the cartesian product(i.e. the categorical product) on $\mathcal{V}$, one says that $\mathcal{V}$ is **cartesian closed**. We now introduce a construction which is very useful later on in topology. For any cartesian closed symmetric monoidal category $(\mathcal{V}, \times, \star)$, which we also suppose to be complete and cocomplete. Consider the slice category under $\star$, the terminal object of $\mathcal{V}$. It should be instantly clear what the construction is supposed to be. Let this category be $\mathcal{V}_\star$, otherwise known as the category of **based objects** in $\mathcal{V}$. We define the product to be the **smash product** which is the pushout square of the following kind.

$$\begin{array}{ccc} v \sqcup w & \longrightarrow & v \times w \\ \downarrow & & \downarrow \\ \star & \longrightarrow & v \wedge w \end{array}$$

This is the natural way to define a product on this particular slice category because here we are just quotienting the edges along $\star$ in the product to a single point which can be viewed using the pushout formula or actually calculating the colimit of the diagram. This construction is associative upto isomorphism (it would be associative if the underlying monoidal category is strictly associative). Since $\mathcal{V}$ is cartesian closed, the functors $v \times -$ preserve colimits. This may allow you to consider based spaces where you might not have looked for before. At this point the point of enriched categories should be clear. We would like to makes paths in the mapping space of two objects to create homotopies and create paths between paths to create higher homotopies. This should now be clear that for homotopy theory we would like to enrich our categories over some kind of simplicial objects, perhaps simplicial sets. Although before moving into simplicial categories we shall take a look at something else before.

There is a natural way to define a "good" functor between symmetric monoidal category, i.e. it preserves the symmetric monoidal structure in the sense that we describe below.

Definition 2.10. A functor $F : \mathcal{V} \rightarrow \mathcal{U}$ between symmetric monoidal categories $(\mathcal{V}, \times, \star)$ and $(\mathcal{U}, \otimes, \mathbb{I})$ is **lax monoidal** if there exist associative and unital natural transformations

$$\mathbb{I} \rightarrow F(\star) \text{ and } Fv \otimes Fv' \rightarrow F(v \times v')$$

One would have expected that the above natural transformation is a natural isomorphism instead. While that is a completely valid way to think, infact there is indeed a definition specifically for them aswell but the above definition also serves the purpose of change of base functors which are of great importance while learning enriched category theory. We shall mention the other variants of the definition down below.

- We call a functor **strong monoidal** if the above natural transformations become natural isomorphisms.
- We call a functor **strict monoidal** if the above natural transformations just become equalities or identities.
- We call a functor **oplax monoidal** if the direction of natural transformations in the definition of lax monoidal functors is reversed.

We now delve into the usage of lax monoidal functors that we mentioned before, Change of Base functors.

Proposition 2.11. In the presence of any lax monoidal functor $F : \mathcal{V} \rightarrow \mathcal{U}$, any $\mathcal{V}$ – category has an associated $\mathcal{U}$ – category with the same objects.

The proof really just follows simply from trying to find the object in $\mathcal{U}$ for any string of operations in $\mathcal{V}$ using the natural transformations given by the lax monoidal functor which essentially completes the proof by using the coherence conditions on $\mathcal{V}$ first and then $\mathcal{U}$. This can be used to lose the enrichment structure of any $\mathcal{V}$ – category, i.e. change the base of the category from $\mathcal{V}$ to **Set**. Using the proposition for this specific case, we get what is called "losing" the enrichment for an enriched category. The lax monoidal functor that we use to achieve this is the $\mathcal{V}(\star, -)$.

Definition 2.12. The **underlying category** $\mathcal{C}_0$ of a $\mathcal{V}$ – category $\underline{\mathcal{C}}$ has the same objects as before but has the hom-sets as

$$\mathcal{C}_0(x, y) = \mathcal{V}(\star, \underline{\mathcal{C}}(x, y))$$

One can now reasonably think about closed monoidal categories, when you lose the enrichment on a self-enriched category, you would just get the same category back with hom-sets instead of hom-objects assuming the category was locally-small. We shall now talk about enriched functors and enriched natural transformations a little bit before going into the meat of the situation that is the simplicial category.

Definition 2.13. A $\mathcal{V}$ -functor $F : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{D}}$ between $\mathcal{V}$ -categories is given by an object map between the enriched categories and a map between the base categories of the following form

$$\underline{\mathcal{C}}(x, y) \xrightarrow{F_{x,y}} \underline{\mathcal{D}}(Fx, Fy)$$

I hope this gives you the suitable style in which enriched category theory is defined. You can go on to define enriched natural transformations, enriched Yoneda lemma, enriched adjunctions and so on but we shall stop here and move onto a much more important topic of Simplicial Categories ,i.e. categories enriched over $(\mathbf{SSet}, \times, \star)$.

Any simplicial category $\underline{\mathcal{C}}$ gives a simplicial object in $\mathbf{Cat}$. Any simplicial category gives us a simplicial object in the $\mathbf{Cat}$, i.e. the enrichment yields a structure such that, therefore at each level you can define a C_n which contains the same objects as in the unenriched category and all n-simplicies(string of arrows with n maps) morphisms between the objects. We may denote this simplicial object as $\mathcal{C}_\bullet$. Under this understanding, one can expect that C_0 is the underlying category itself.

3 Weak Factorisation Systems

4 Model Categories

References

- [1] D. Quillen, “Homotopical algebra,” 1965.
- [2] M. Hovey, “Model categories,” 1999.
- [3] E. Riehl, “Categorical homotopy theory,” 2014.