
INTRODUCTION TO MODEL CATEGORIES

10th September 2026

Karshit Joshi

Math and Physics Student
Chennai Mathematical Institute
9th September 2026

Contents

1	Preliminaries	3
2	Derived Functors	4
2.1	Homotopical Categories and Derived Functors	4
3	Weak Factorisation Systems	7
4	Model Categories	7

1 Preliminaries

This is an exposition written by me to take notes of what I read about Model Categories and certain ideas that I think are very useful while reading Model Categories. Any mistakes made in this exposition is purely mine and please report them to [karshitkjoshi@cmi.ac.in]. I am mostly using "Homotopical Algebra" by Daniel Quillen[1], "Model Categories" by Mark Hovey[2] and "Categorical Homotopy Theory" by Emily Riehl[3] as my primary resources to read about the topic from, and will try my best to mention any other resources that I use along the way in the bibliography. I am grateful to Chennai Mathematical Institute to expose me to the resources and wonderful people that helped me during this writing.

I am very grateful to Prof. Aditya Karnataki, who encouraged me to write this exposition and introducing me to writing exposition which led me to find an appreciation for writing expositions. I am also grateful to Bhanu Kiran Sandepudi and Aleksandr Zakharov for giving me their free time at various occasions to discuss questions that I came across while writing.

2 Derived Functors

2.1 Homotopical Categories and Derived Functors

An important question has always been in different settings and even in category theory, "Are these two objects isomorphic?", while the answer to the question depends on the what we consider an isomorphism. Fixing the context, it could perhaps be easier or difficult to differentiate between objects.

A good strategy in our toolset has always been to classify objects "upto some property". Note that the creation of such classification creates a new category where the context is changed to something which is much coarser than the current context that we are trying to work in. This often helps us to focus ourselves towards a particular way to differentiate objects rather than having the vague goal of differentiating objects with nothing ahead of us. Common examples of this phenomenon include Homotopy Groups, Homology Groups, Euler Characteristic, Betti Numbers, etc. We shall now try to build this notion in category theory for abstract categories.

Suppose we start with a category $\mathcal{C}$. We have some property $\mathcal{P}$ along which we can test our objects to see if they are "equivalent" or not. If identifying $\mathcal{P}$ on 2 different objects tells us that they have the same property. Our desire is to consider those objects to be the same. A natural way to store all of this data of identification is in a category, with the purpose that we will inherit all the data naturally by doing a construction on $\mathcal{C}$.

We are now clear that we must perform construction on $\mathcal{C}$ to create a category which characterizes the data that we get from $\mathcal{P}$. There are naturally 2 ways to do this.

- Consider the objects of $\mathcal{C}$, consider the equivalence relation imposed by $\mathcal{P}$ on the collection of object of $\mathcal{C}$. We quotient by the equivalence relation. We also forget all the maps between objects which were identified. The residue will turn out to be a category with a few more natural steps. This can be thought of as skeleton category of localisation given by the second point.
- Consider the maps of $\mathcal{C}$, formally introduce the inverses of maps between objects, "which characterize the equivalence of $\mathcal{P}$ ". This just naturally forms a category once again after completing all possible compositions with the new maps.

Both the above ways are useful in different ways that we will explore throughout the span of this text. Notice the difference, that the first construction forgets how things are actually equivalent whereas the second does not due to the existence of all the maps which "characterize the equivalence of $\mathcal{P}$ ". This especially comes into the picture if you view the first construction as the manifestation of "existence of an equivalence" rather than "equivalent with the witness map", wherein the second construction is much more suited to the latter. This often comes into play when you are considering n-truncations of an infinity category which are the objects we are truly trying to study indirectly via model categories. Due to the previous reason, this also plays a significant role in homotopy type theory. [Note that the first construction is also technically inline with Leibniz equivalence of identibles which says that "if two objects have the same interactions, they are the same" i.e. Isomorphisms should not exist as objects, instead all isomorphisms must be equalities. This plays a major role in homotopy type theory.]

Since the second construction retains much more information from the ambient category. It is natural to call it the appropriate "localisation" of the category. The terminology of localisation is inherited from ring theory, wherein we localize a ring against a prime ideal which requires to invert all elements outside the prime ideal.

We now turn our focus to actually selecting the maps of the category $\mathcal{C}$, to be our equivalences which "we believe" characterizes some property $\mathcal{P}$. Here is a subtle but important point that we "choose" a property to classify objects against. In the context of abstract categories, the selection of maps in the category is in direct correspondence to the abstract property $\mathcal{P}$. Hence, the selection of property is equivalent to selection of which maps to invert. Hence, the structure we obtain is a deliberate construction, not a natural structure to consider, i.e. The structure is a choice, not a natural byproduct of $\mathcal{C}$.

It is also natural to expect that all isomorphisms are also a part of the collection of maps we invert mainly because, if $\mathcal{C}$ cannot distinguish between 2 objects, they must possess the same set of properties at all levels [Note that this is essentially the application of Yoneda Lemma]. Hence, this helps us to arrive at our first definition.

Definition 2.1. A **homotopical category** is a category $\mathcal{C}$ equipped with a wide* subcategory $\mathcal{W}$ such that, for any composable triple of arrows, i.e. f, g, h . If hg and gf are in $\mathcal{W}$, so are f, g, h , and hgf .

The arrows in $\mathcal{W}$ are called equivalences. The property above concerning the maps is called the 2-of-6 property. [abbreviation of "2 out of 6"] (*) By a wide subcategory, we mean a subcategory which contains all identities of the category $\mathcal{C}$ and hence by 2-of-6 property, all isomorphisms of $\mathcal{C}$ as well.

There is a weaker condition that is also often considered instead of the 2-of-6 property. It is called 2-of-3 property.

Definition 2.2. 2-of-3 Property occurs if any 2 of the 3 possible composition of 2 composable maps are in $\mathcal{W}$. Then $\mathcal{W}$ contains all 3.

A small caveat in the above 2 definitions is that $\mathcal{W}$ must also contain all the identities for each object in $\mathcal{C}$ which has a clear enough reason to be included.

Now that we have concretely defined what is the data that we would like to work with. We now approach the problem of carrying out the construction of the localisation of the category as we previously mentioned. The following explicit construction is due to Pierre Gabriel and Michel Zisman. Its morphisms are equivalence classes of finite zig-zags (i.e. compositions which maps from $\mathcal{W}$ in the string with finite length) with arrows in $\mathcal{W}$ to be inverted with the same name but dual of its original counterpart subjected to the following relations.

Definition 2.3. The localisation category/homotopy category $\text{Ho}\mathcal{C}$ of a homotopical category $(\mathcal{C}, \mathcal{W})$ is the formal localisation of $\mathcal{C}$ at the subcategory $\mathcal{W}$. The explicit construction of the homotopy category is as follows:

- Objects of $\text{Ho}\mathcal{C}$ are the same as $\mathcal{C}$.
- All strings of compositions which contain $\xleftarrow{\mathcal{W}} \xrightarrow{\mathcal{W}}$ or $\xrightarrow{\mathcal{W}} \xleftarrow{\mathcal{W}}$ or identities pointing forward or backwards as a part of the string are equivalent to strings of compositions with the earlier substrings removed. This induces the natural equivalence relation of strings of composable arrows.

One can simply notice that there is a natural functor $\mathcal{C} \xrightarrow{L} \text{Ho}\mathcal{C}$. This functor also possesses the universal property that any other functor which inverts all morphisms in $\mathcal{W}$ (not necessarily only morphisms in $\mathcal{W}$) factors through L .

An important point is that not all morphisms of $\mathcal{C}$ that become isomorphisms should be present in the subcategory $\mathcal{W}$. When this does not happen, we call the homotopical category $\mathcal{C}$ to be saturated [i.e. all morphisms whose inverse exists in $\text{Ho}\mathcal{C}$ are also present in $\mathcal{W}$].

Another important point is that performing this construction on a locally small category does not ensure that it will remain a locally small category. This is simply due to the reason that the collection of objects form a class and each object has its respective equivalence class. There could be a lot of choices of a single map about which object to passthrough for each object's equivalence class. Considering the possibility that the equivalence class of an object may itself form a class. The collection of maps between any two maps can form a class. For this reason, there exists another richer localisation than the one we came up with, which is inspired from the localisation of rings, called "Calculus of Fractions" due to Pierre Gabriel, Pierre Zisman. However, I plan to delve into it just yet.

Therefore, one of the problems that the model category construction solves is to keep the localisation of the category to be locally small.

We now try to study functors between Homotopical Categories which can extend to maps between the respective homotopy categories under suitable conditions on the functor itself. The condition on the functor is also natural, in the sense that it must send equivalences to equivalences ,i.e. it preserves all equivalences. We call such functors, "Homotopical Functors", by the universal property of localisation, it induces a functor between the respective homotopy categories.

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow L_{\mathcal{C}} & & \downarrow L_{\mathcal{D}} \\ \text{Ho}\mathcal{C} & \xrightarrow{F'} & \text{Ho}\mathcal{D} \end{array}$$

It is a natural desire to study functors which are not homotopical. However, when that happens notice that $L_{\mathcal{C}}$ and $L_{\mathcal{D}} \circ F$ form a span diagram, which hints us towards looking at left and right Kan Extensions along the appropriate direction carrying the intuition that we are trying to find the best functor which approximates F while losing the least amount of information while doing it. This naturally leads to the following definition.

Definition 2.4. A **total left derived functor** LF of a functor F between Homotopical Categories $\mathcal{C}$ and $\mathcal{D}$ is a right Kan extension $Ran_{L_{\mathcal{C}}}(L_{\mathcal{D}} \circ F)$:

This also alludes to being related to derived functors in the homological algebra sense.

3 Weak Factorisation Systems

4 Model Categories

References

- [1] D. Quillen, “Homotopical algebra,” 1965.
- [2] M. Hovey, “Model categories,” 1999.
- [3] E. Riehl, “Categorical homotopy theory,” 2014.