## theoretical exercise 2

# Pattern Recognition (2018)

Prof. Dr. Hauke Schramm

Marc Engelmann Lasse B. Petersen Jan N. Steeg

due on 30.11.2018

### Exercise T-2.1

#### **Problem**

Consider a sample space X comprising three possible outcomes  $X = v_1, v_2, v_3$ .

We define the events

$$E = \{v_1, v_2\}$$
$$F = \{v_1, v_3\}$$

and denote by  $E^c$  the complement of E.

Compute  $P(F|E^c)$ , the conditional probability of F given  $E^c$ , using the conditional probability formula for the events A and B:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

#### Solution

$$P(F|E^{c}) = \frac{P(F \cap E^{c})}{P(E^{c})}$$

$$= \frac{P(F) - P(F \cap E)}{1 - P(E)}$$

$$= \frac{P(\{v_{1}\} \cup \{v_{3}\}) - P((\{v_{1}\} \cup \{v_{3}\}) \cap (\{v_{1}\} \cup \{v_{2}\}))}{1 - P(\{v_{1}\} \cup \{v_{2}\})}$$

$$= \frac{P(\{v_{3}\}) - P(\{v_{2}\} \cup \{v_{3}\})}{1 - P(\{v_{1}\} \cup \{v_{2}\})}$$

$$= \frac{P(v_{3}) - (P(v_{2}) + P(v_{3}))}{1 - (p(v_{1}) + p(v_{2}))}$$
(1)

Exercise T-2.2

#### **Problem**

The Census Bureau has estimated the following survival probabilities for men:

- 1. probability that a man lives at least 70 years: 80 %
- 2. probability that a man lives at least 80 years: 50 %

What is the conditional probability that a man lives at least 80 years given that he has just celebrated his 70th birthday?

#### Solution

Given probabilities:

$$p(70) = 0.8$$
: (Chance a man lives at least 70 years)  $p(80) = 0.5$ : (Chance a man lives at least 80 years)

Calculation of p(80|70):

$$p(70|80) = 1$$
 (no man can live 80 years if he died before living 70 years)  
$$p(80|70) = \frac{p(70|80) \cdot p(80)}{p(70)} = \frac{1 \cdot 0.5}{0.8} = 0.625$$

Solution: The conditional probability that a man lives at least 80 years given that he has just celebrated his 70th birthday is at 62.5%.