

Gierer-Meinhart equations

PS5_GM_equation.mlx

Joe Howard, modified from Hugo Bowne-Anderson 2014

April 8, 2024

In this script, we solve the Gierer-Meinhardt equations 12a/b or 15a/b from (Kybernetik, 12:30-39, 1972) for a variety of parameters.

The partial differential equations are the dimensionless G-M system:

$$\frac{\partial}{\partial t} a(x, t) = \frac{a^2}{h + h_0} - a + \sigma + D \frac{\partial^2}{\partial x^2} a$$

$$\frac{\partial}{\partial t} h(x, t) = \omega(a^2 - h) + \frac{\partial^2}{\partial x^2} h$$

%Notes:

- (1) the derivatives are partial derivatives (t is time, x is space);
- (2) t , x , a and h are all dimensionless;
- (4) D is ratio of diffusion constants: activator/inhibitor and is < 1
- (5) ω is a reaction term
- (6) σ is a baseline synthesis term for a (assumed small, 0.01).
- 7) h_0 is a small positive term (0.1); it keeps the first term in Eqn 1 finite

```
clear;  
close all;
```

GM_solve_record is the PDE solver. It takes 3 main parameters:

$P(1) = D$

$P(2) = \omega$

$P(3) = \sigma$ (chosen to be close to zero);

There are also three other constants specified below:

tmax = amount of time that we run the system;

delt = time increment;

L = length (spatial) of the system;

and there is h_0 , which is set at 0.1 in the function

GM_solve_record outputs the concentrations of activator and inhibitor over the time course and plots the solution as surfaces for $a(x, t)$ and $h(x, t)$ and as an animation.

We consider the phase plane of the G-M system with x-axis given by D (ratio of diffusion constants: activator/inhibitor) and y-axis ω (the reaction

% term) while keeping everything else constant.

Question 1.

$$Q1) \frac{\partial A}{\partial t} = k_a \frac{A^2}{H+H_0} - \omega_a A + \Sigma + D_a \frac{\partial^2 A}{\partial x^2} \quad \text{--- ①}$$

$$\frac{\partial H}{\partial t} = k_h A^2 - \omega_h H + D_h \frac{\partial^2 H}{\partial x^2}$$

[using: $T = \alpha t \Rightarrow dT = \alpha dt$
 $x = \beta x \Rightarrow dx = \beta x$
 $A = \chi a \Rightarrow dA = \chi \cdot da$
 $H = \delta h \Rightarrow dH = \delta \cdot dh$

$$\begin{aligned} \frac{\partial a}{\partial t} &= \frac{\alpha}{\chi} \left(k_a \cdot \frac{\chi^2 a^2}{\delta(h+h_0)} - \omega_a \chi a + \Sigma + D_a \frac{\chi^2}{\beta^2} \cdot \frac{\partial a^2}{\partial x^2} \right) \\ &= \frac{k_a \chi \alpha}{\delta} \cdot \frac{a^2}{h+h_0} - \omega_a \chi a + \Sigma + \frac{D_a \chi \alpha}{\beta^2} \cdot \frac{\partial a^2}{\partial x^2} \end{aligned}$$

$$\begin{aligned} \frac{\partial h}{\partial t} &= \frac{\alpha}{\delta} \left(k_h \chi^2 a^2 - \omega_h \delta h + \frac{D_h \delta^2}{\beta^2} \cdot \frac{\partial h^2}{\partial x^2} \right) \\ &= \frac{k_h \alpha \chi^2}{\delta} a^2 - \omega_h \alpha h + \frac{D_h \alpha \delta}{\beta^2} \cdot \frac{\partial h^2}{\partial x^2} \end{aligned}$$

Let $\chi = 1/\omega_a \rightarrow \omega_h = \omega \omega_a = \frac{\omega}{\chi}$
 $\frac{\beta^2}{\alpha \delta} = D_h \rightarrow D_a = D D_h = \frac{D \beta^2}{\alpha \delta}$
 $\frac{\delta}{\chi \alpha} = k_a$

$$\frac{\partial a}{\partial t} = \frac{a^2}{h+h_0} - a + \Sigma + \frac{D \chi}{\delta} \cdot \frac{\partial a^2}{\partial x^2}$$

$$\frac{\partial h}{\partial t} = \frac{k_h \chi}{k_a} a^2 - \omega \cdot \frac{\alpha}{\chi} h + \frac{\partial h^2}{\partial x^2}$$

Question 2.

$$Q2) h_0 \ll 1, \sigma = 0$$

$$\text{Also, } \frac{\partial^2 a}{\partial x^2} = 0; \quad \frac{\partial^2 h}{\partial x^2} = 0$$

$$\text{Setting } \frac{\partial a}{\partial t} = \frac{\partial h}{\partial t} = 0,$$

$$\omega(a^2 - h) = 0$$

$$\text{So, } a^2 = h$$

$$\text{Now, } \frac{a^2}{h} - a + \sigma = 0$$

$$\Rightarrow \boxed{a = 1} \quad \because \sigma = 0$$

$$\text{and } \boxed{h = 1}$$

Question 3.

Q3) $a = a_0 \sin kx$
 $h = h_0 \sin kx$

So, $\frac{\partial^2 a}{\partial x^2} = -a_0 k^2 \sin kx = -k^2 a$

$\frac{\partial^2 h}{\partial x^2} = -h_0 k^2 \sin kx = -k^2 h$

The dynamical system is then:

$$\frac{\partial a}{\partial t} = \frac{a^2}{h+h_0} - a(1+Dk^2)$$

$$\frac{\partial h}{\partial t} = \omega(a^2-h) - k^2 h$$

Now, the Jacobian,

$$J = \begin{pmatrix} \frac{2a}{h} - 1 - Dk^2 & -\frac{a^2}{h^2} \\ 2a\omega & -\omega - k^2 \end{pmatrix}$$

$$J(1,1) = \begin{pmatrix} 1-Dk^2 & -1 \\ 2\omega & -\omega-k^2 \end{pmatrix}$$

(i) When spatial gradient is 0,

$$J(1,1) = \begin{pmatrix} 1 & -1 \\ 2\omega & -\omega \end{pmatrix}$$

$$\text{tr}_J = 1-\omega$$

$$\det_J = \omega$$

For a stable fixed pt.,

$\text{tr} < 0$ and $\det > 0$

$$1-\omega < 0 \quad 4\omega > 0$$

$$\omega > 1$$

So we need: $\omega > 1$

(iii) $\omega = \frac{\omega_h}{\omega_a} > 1$

This means that degradation rate of the inhibitor (H) must be higher than that of the activator (A) for the fixed point to be stable.

Question 4.

Q4) From 3(i),

$$J = \begin{pmatrix} \frac{2a}{h} - 1 - Dk^2 & -\frac{a^2}{h^2} \\ 2a\omega & -\omega - k^2 \end{pmatrix}$$

$$\begin{aligned} \text{So, } \det J &= \left(\frac{2a}{h} - 1 - Dk^2 \right) (-\omega - k^2) + \frac{2a^2\omega}{h^2} \\ &= Dk^4 + \left(\omega D - \frac{2a}{h} + 1 \right) k^2 + \left(1 - \frac{2a}{h} \right) \omega \end{aligned}$$

$$\begin{aligned} \frac{\partial(\det J)}{\partial(k^2)} &= \frac{\partial(\det J)}{\partial k} \cdot \frac{1}{2k} \\ &= 2Dk^2 + \omega D - \frac{2a}{h} + 1 \end{aligned}$$



Setting $\det' J = 0$,


$$k^2 = \frac{a}{Dh} - \frac{\omega}{2} - \frac{1}{2D}$$

At the fixed pt. (1,1),

$$k^2 = \frac{1}{2D} - \frac{\omega}{2}$$

```
tmax = 100;      %maximum time (can be changed from 50 to 500 if necessary)
delt = 1;        %time interval
L = 20;          %length
% P = [0.05 ; 2 ; 0.01 ];      %Spatial pattern
% P = [0.4 ; 2 ; 0.01 ];      %Homogenous in time and space
% P = [0.4 ; 1; 0.01];        %temporal oscillation
P = [0.4 ; 0.5; 0.01];        %trivial solution
soln = GM_solve_record_JH(P , tmax , delt, L, 'test_pattern1.avi');
```





(i) The wavelength from the plot is ~ 3.25 units. Based on the prediction from question 4, it is ~ 2.09 units.

(ii) The steady fixed point occurs at $(a, h) = (0.75, 0.575)$. The prediction from question 2 was $(1, 1)$.

(iii)

(iv) The fixed point occurs at $(0, 0)$. The Jacobian evaluated at $(0, 0)$ shows that the fixed point is stable (see below).

$$Q5) \quad J = \begin{pmatrix} \frac{2a}{h} - 1 - Dk^2 & -\frac{a^2}{h^2} \\ 2a\omega & -\omega - k^2 \end{pmatrix}$$

$$J(0,0) = \begin{pmatrix} -1 - Dk^2 & 0 \\ 0 & -\omega - k^2 \end{pmatrix}$$

$$\begin{aligned} \tau_f &= -1 - Dk^2 - \omega - k^2 \\ &= -1 - (0.4)k^2 - 0.5 - k^2 \\ &= -(1.5 + 1.4k^2) \\ &< 0 \end{aligned}$$

So the fixed pt is stable.

Phase diagram


Alter $P(1) = D$ and $P(2) = \omega$ to move around the phase plane & draw (by hand or by computer) a phase diagram for this Gierer-Meinhardt system with D on the x-axis and ω on the y-axis.

```
tmax = 200;
ctr = 1;
for x1 = 0.05:0.1:0.5
    for x2 = 0.05:0.5:5
        P = [x1; x2; 0.01]; %trivial solution
        disp(P);
        sol = GM_solve_record_JH(P, tmax, delt, L, 'test_pattern2.avi');


        t = linspace(0,tmax,tmax/delt);
        x = linspace(0,L,200); %the mesh on which we solve (200 space
points)
        figure(100+ctr);
        for n = 2:2:length(t)
            set(gca, 'FontSize', 18, 'LineWidth', 1); %<- Set properties
            plot( x , sol(n,:,1), 'LineWidth',3);
            hold on
            plot( x , sol(n,:,2), 'r', 'LineWidth',3);
            hold off
            legend('activator', 'inhibitor', 'Location', 'SouthEast');
            title(strcat('Gierer-Meinhardt patterns t = ', sprintf(' %d ',
ceil(t(n)))));
            axis([0 L 0 max(max(max(sol(:,:,:),:)))+0.1])
        end
        ctr = ctr + 1;
    end
end
```

end


0.0500
0.0500
0.0100




0.0500
0.5500
0.0100




0.0500
1.0500
0.0100




0.0500
1.5500
0.0100




0.0500
2.0500
0.0100




0.0500
2.5500
0.0100




0.0500
3.0500
0.0100




0.0500
3.5500
0.0100




0.0500
4.0500
0.0100




0.0500
4.5500
0.0100




0.1500
0.0500
0.0100




0.1500
0.5500
0.0100




0.1500
1.0500
0.0100




0.1500
1.5500
0.0100




0.1500
2.0500
0.0100




0.1500
2.5500
0.0100




0.1500
3.0500
0.0100




0.1500
3.5500
0.0100




0.1500
4.0500
0.0100




0.1500
4.5500
0.0100




0.2500
0.0500
0.0100




0.2500
0.5500
0.0100




0.2500
1.0500
0.0100




0.2500
1.5500
0.0100




0.2500
2.0500
0.0100




0.2500
2.5500
0.0100




0.2500
3.0500
0.0100




0.2500
3.5500
0.0100




0.2500
4.0500
0.0100




0.2500
4.5500
0.0100




0.3500
0.0500
0.0100




0.3500
0.5500
0.0100




0.3500
1.0500
0.0100




0.3500
1.5500
0.0100




0.3500
2.0500
0.0100




0.3500
2.5500
0.0100




0.3500
3.0500
0.0100




0.3500
3.5500
0.0100




0.3500
4.0500
0.0100




0.3500
4.5500
0.0100




0.4500
0.0500
0.0100




0.4500
0.5500
0.0100




0.4500
1.0500
0.0100




0.4500
1.5500
0.0100




0.4500
2.0500
0.0100




0.4500
2.5500
0.0100




0.4500
3.0500
0.0100





0.4500
3.5500
0.0100





0.4500
4.0500
0.0100




0.4500
4.5500
0.0100





Question 5.



Question 6.

(ii)

Q6) From q_4 ,

$$k^2 = \frac{a}{Dh} - \frac{\omega}{2} - \frac{1}{2D}$$

When $\det J = 0$,

$$Dk^4 + \left(\omega D - \frac{2a}{h} + 1\right)k^2 + \left(1 - \frac{2a}{h}\right)\omega = 0$$

$$\underline{D\left(\frac{a}{Dh} - \frac{\omega}{2} - \frac{1}{2D}\right)^2 + \left(\omega D - \frac{2a}{h} + 1\right)\left(\frac{a}{Dh} - \frac{\omega}{2} - \frac{1}{2D}\right) + \left(1 - \frac{2a}{h}\right)\omega = 0.}$$

(i) Setting $k^2 = 0$ in $\det J$

$$\det J = \left(1 - \frac{2a}{h}\right)\omega$$

For -ve $\det J$,

$$\left(\frac{2a}{h} - 1\right)\omega > 0$$

For $\omega \neq 0$, at the boundary,

$$\boxed{2a = h}$$