

Error and Variance Analysis in the Fourier Domain

Gurprit Singh

Post doctoral researcher

Outline

Outline

- How error looks like in the Fourier domain

Outline


- How error looks like in the Fourier domain
- Bias in the Fourier domain


Outline


- How error looks like in the Fourier domain
- Bias in the Fourier domain
- Variance in the Fourier domain


Outline


- How error looks like in the Fourier domain
- Bias in the Fourier domain
- Variance in the Fourier domain
- Variance Convergence analysis of sampling patterns














Monte Carlo Integration

$$\tilde{\mu}_N =$$

Monte Carlo Integration

$$\tilde{\mu}_N = \int_D f(x) \mathbf{S}(x) dx =$$

Monte Carlo Integration

$$\tilde{\mu}_N = \int_D f(x) \mathbf{S}(x) dx =$$

$$\mathbf{S}(x) = \frac{1}{N} \sum_{k=1}^N \delta(x - x_k)$$

Monte Carlo Integration

$$\tilde{\mu}_N = \int_D f(x) \mathbf{S}(x) dx = \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$\mathbf{S}(x) = \frac{1}{N} \sum_{k=1}^N \delta(x - x_k)$$

Monte Carlo Integration

$$\tilde{\mu}_N = \int_D f(x) \mathbf{S}(x) dx = \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$\mathbf{S}(x) = \frac{1}{N} \sum_{k=1}^N \delta(x - x_k)$$

$$\hat{\mathbf{S}}(\omega) = \frac{1}{N} \sum_{k=1}^N e^{-i2\pi\omega x_k}$$

$$\text{Error} = \text{Bias}^2 + \text{Variance}$$

Error in Spatial Domain

$$I = \int_D f(x) dx$$

Error in Spatial Domain

$$I = \int_D f(x) dx$$

$$I - \tilde{\mu}_N = \int_D f(x) dx - \int_D f(x) \mathbf{S}(x) dx$$

Error in Spatial Domain

$$I - \tilde{\mu}_N = \int_D f(x) dx - \int_D f(x) \mathbf{S}(x) dx$$

Error in Spatial Domain

$$I - \tilde{\mu}_N = \int_D f(x) dx - \int_D f(x) \mathbf{S}(x) dx$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \int_D f(x) dx - \int_D f(x) \mathbf{S}(x) dx$$

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \int_D f(x) dx - \int_D f(x) \mathbf{S}(x) dx$$

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$E[I - \tilde{\mu}_N] = E[\hat{f}(0)] - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$E[I - \tilde{\mu}_N] = E[\hat{f}(0)] - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$E[I - \tilde{\mu}_N] = E[\hat{f}(0)] - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

Error in Fourier Domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$E[I - \tilde{\mu}_N] = E[\hat{f}(0)] - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - E\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain

$$E[I - \tilde{\mu}_N] = \hat{f}(0) - \int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain

$$\int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain

$$\int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain

$$\int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$

for frequencies other than zero

Bias in Fourier Domain



$$\int_{\Omega} E[\hat{f}^*(\omega)] E[\hat{\mathbf{S}}(\omega)] d\omega = \int_{\Omega} |E[\hat{f}^*(\omega)]| |E[\hat{\mathbf{S}}(\omega)]| e^{\Phi(E[\hat{f}^*(\omega)])} e^{\Phi(E[\hat{\mathbf{S}}(\omega)])} d\omega$$

To obtain an unbiased estimator:

$$E[\hat{\mathbf{S}}(\omega)] = 0$$



for frequencies other than zero

Phase Distribution





Phase Distribution

Multiple realizations





Phase Distribution

Multiple realizations





Phase Distribution



Multiple realizations





Phase Distribution



Phase Distribution




Phase Distribution





Phase Distribution

Multiple realizations





Phase Distribution

Multiple realizations





Phase Distribution

Multiple realizations



Phase Distribution

Multiple realizations



$$E[\hat{\mathbf{S}}(\omega)] = 0$$

Phase Distribution

- Uniform random shifting of samples change the phase of the samples' Fourier spectrum

Phase Distribution

- Uniform random shifting of samples change the phase of the samples' Fourier spectrum
- Uniform random shifting does not affect the amplitude

Phase Distribution

- Uniform random shifting of samples change the phase of the samples' Fourier spectrum
- Uniform random shifting does not affect the amplitude
- Zero expected Fourier spectrum of a sampling pattern implies translation invariant sample distribution

Phase Distribution

- Uniform random shifting of samples change the phase of the samples' Fourier spectrum
- Uniform random shifting does not affect the amplitude
- Zero expected Fourier spectrum of a sampling pattern implies translation invariant sample distribution
- The process is named Homogenization

$$\text{Error} = \text{Bias}^2 + \text{Variance}$$

$$\text{Error} = \text{Bias}^2 + \text{Variance}$$

$$\text{Error} = \text{Bias}^2 + \text{Variance}$$

Homogenization allows representation of error only in the variance form

Variance in the Fourier domain

Variance in the Fourier domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

Variance in the Fourier domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \text{Var}\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

Variance in the Fourier domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \text{Var}\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$I - \tilde{\mu}_N = \hat{f}(0) - \int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \text{Var}\left[\int_{\Omega} \hat{f}^*(\omega) \hat{\mathbf{S}}(\omega) d\omega\right]$$

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

This is a general form both for homogenised as well as non-homogenised sampling patterns

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 \text{Var}[\hat{\mathbf{S}}(\omega)] d\omega$$

For homogenised sampling pattern:

Variance in the Fourier domain

$$Var[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 Var[\hat{\mathbf{S}}(\omega)] d\omega$$

For homogenised sampling pattern:

$$Var[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain

$$Var[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 Var[\hat{\mathbf{S}}(\omega)] d\omega$$

For homogenised sampling pattern:

$$Var[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain


$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$


Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$



Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$



Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} P_f(\omega) E[P_{\mathbf{S}}(\omega)] d\omega$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} P_f(\omega) E[P_{\mathbf{S}}(\omega)] d\omega$$

In polar coordinates:

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} |\hat{f}^*(\omega)|^2 E[|\hat{\mathbf{S}}(\omega)|^2] d\omega$$

$$\text{Var}[\tilde{\mu}_N] = \int_{\Omega} P_f(\omega) E[P_{\mathbf{S}}(\omega)] d\omega$$

In polar coordinates:

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

For isotropic power spectra:

Variance in the Fourier domain

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \int_{\mathcal{S}^{d-1}} P_f(\rho \mathbf{n}) E[P_{\mathbf{S}}(\rho \mathbf{n})] d\mathbf{n} d\rho$$

For isotropic power spectra:

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_{\mathbf{S}}(\rho)] d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_{\mathbf{S}}(\rho)] d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Variance in the Fourier domain


$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$


Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Variance in the Fourier domain

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Variance Convergence Analysis

Power Spectra Product

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Power Spectra Product


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Power Spectra Product

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Power Spectra Product


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



For given number of Samples

Power Spectra Product


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



For given number of Samples

Power Spectra Product


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$




As we increase the number of samples

Radial Mean Power Spectra


Random




CCVT Balzer et al.[2009]



Jittered



Poisson Disk




Samples'

Radial Power Spectra Profiles

$$\text{Var}[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Samples' Radial Power Spectra Profiles

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Samples'

Radial Power Spectra Profiles

$$\tilde{P}_{\mathbf{S}}(\rho)_N = \begin{cases} \gamma \left(\frac{\rho}{\alpha \sqrt[d]{N}} \right)^b & \rho < \sqrt[d]{N} \\ \gamma & \text{otherwise} \end{cases}$$

Samples' Radial Power Spectra Profiles

$$\tilde{P}_{\mathbf{S}}(\rho)_N = \begin{cases} \gamma \left(\frac{\rho}{\alpha \sqrt[d]{N}} \right)^b & \rho < \sqrt[d]{N} \\ \gamma & \text{otherwise} \end{cases}$$

where,

$\tilde{P}_{\mathbf{S}}(\rho)_N = N \times \tilde{P}_{\mathbf{S}}(\rho)$ Normalized radial power spectra


$\gamma, \alpha \in \mathcal{R}^+ / 0$ Positive non-zero constant

d Dimensions



N Number of Samples

Bounds for Radial Mean Power Spectra


Random



CCVT Balzer et al.[2009]



Poisson Disk



Analytical Classification of Power Spectra


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_s(\rho)] d\rho$$

Analytical Classification of Power Spectra

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Analytical Classification of Power Spectra


$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Constant profiles
Quadratic profiles
Other profiles

Analytical Classification of Power Spectra

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Constant profiles
Quadratic profiles
Other profiles

Analytical Classification of Power Spectra

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

Best possible Integrand Power spectrum

⋮

Worst possible Integrand Power spectrum

Constant profiles


Quadratic profiles

Other profiles




Best and Worst Cases

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_s(\rho)] d\rho$$




Best and Worst Cases

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$




Best and Worst Cases

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$




Best and Worst Cases

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Best and Worst Cases

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$



Best and Worst Cases

Worst Case:

$$\tilde{P}_f(\rho) = \begin{cases} c_f & \rho < \rho_0 \\ \rho^{-d-1} & \text{otherwise} \end{cases}$$

Brandolini et al. [2001]

Best Case:

$$\tilde{P}_f(\rho) = \begin{cases} c_f & \rho < \rho_0 \\ 0 & \text{otherwise} \end{cases}$$

Convergence Tool

$$Var[\tilde{\mu}_N] = \mathcal{M}(\mathcal{S}^{d-1}) \int_0^\infty \tilde{P}_f(\rho) E[\tilde{P}_S(\rho)] d\rho$$

$$\tilde{P}_f(\rho) = \begin{cases} c_f & \rho < \rho_0 \\ \rho^{-d-1} & \text{otherwise} \end{cases}$$

$$\tilde{P}_f(\rho) = \begin{cases} c_f & \rho < \rho_0 \\ 0 & \text{otherwise} \end{cases}$$

$$\tilde{P}_S(\rho)_N = \begin{cases} \gamma \left(\frac{\rho}{\alpha \sqrt[d]{N}} \right)^b & \rho < \sqrt[d]{N} \\ \gamma & \text{otherwise} \end{cases}$$

Convergence rates

