


# OPTIMIZATION UNDER DECISION DEPENDENT UNCERTAINTY

Kartikey Sharma  
Omid Nohadani


Northwestern University  
Industrial Engineering and Management Sciences

July 4, 2018

# What is the problem?




# What is the problem?



$$d_e = \bar{d}_e(1 + 0.5\xi_e) \quad \xi \in \mathcal{U} = \{\xi \mid \xi_e \in [0, 1] \forall e\}$$

# What is the problem?




$$d_e = \bar{d}_e(1 + 0.5\xi_e) \quad \boldsymbol{\xi} \in \mathcal{U} = \{\boldsymbol{\xi} \mid \xi_e \in [0, 1] \forall e\}$$

$$\xi_e \sim \text{Unif}[0, 1]$$

$$\min_{\mathbf{y}} \mathbb{E}_{\boldsymbol{\xi} \in \mathcal{U}} [\mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y}]$$

$$\text{s.t. } \mathbf{y} \in Y$$

# What is the problem?



$$d_e = \bar{d}_e(1 + 0.5\xi_e) \quad \boldsymbol{\xi} \in \mathcal{U} = \{\boldsymbol{\xi} \mid \xi_e \in [0, 1] \forall e\}$$

$$\xi_e \sim \text{Unif}[0, 1]$$

$$\xi_e \in [0, 1]$$


$$\min_{\mathbf{y}} \mathbb{E}_{\boldsymbol{\xi} \in \mathcal{U}} [\mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y}]$$

$$\text{s.t. } \mathbf{y} \in Y$$

$$\min_{\mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y}$$

$$\text{s.t. } \mathbf{y} \in Y$$

# What is the problem?



$$d_e = \bar{d}_e(1 + 0.5\xi_e) \quad \boldsymbol{\xi} \in \mathcal{U} = \{\boldsymbol{\xi} \mid \xi_e \in [0, 1] \forall e\}$$

$$\xi_e \sim \text{Unif}[0, 1]$$

$$\xi_e \in [0, 1]$$


$$\xi_e \in [0, 1 - \gamma]$$

$$\begin{aligned} \min_{\mathbf{y}} \mathbb{E}_{\boldsymbol{\xi} \in \mathcal{U}} [\mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y}] \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

$$\begin{aligned} \min_{\mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y} \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

$$\begin{aligned} \min_{\mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y} \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

# What is the problem?



$$d_e = \bar{d}_e(1 + 0.5\xi_e) \quad \boldsymbol{\xi} \in \mathcal{U} = \{\boldsymbol{\xi} \mid \xi_e \in [0, 1] \forall e\}$$

$$\xi_e \sim \text{Unif}[0, 1]$$

$$\xi_e \in [0, 1]$$

$$\xi_e \in [0, 1 - \gamma \mathbf{x}_e]$$

$$\begin{aligned} \min_{\mathbf{y}} \mathbb{E}_{\boldsymbol{\xi} \in \mathcal{U}} [\mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y}] \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

$$\begin{aligned} \min_{\mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y} \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

$$\begin{aligned} \min_{\mathbf{y}, \mathbf{x}} \max_{\boldsymbol{\xi} \in \mathcal{U}(\mathbf{x})} \mathbf{d}(\boldsymbol{\xi})^\top \mathbf{y} + \mathbf{c}^\top \mathbf{x} \\ \text{s.t. } \mathbf{y} \in Y \end{aligned}$$

**Stochastic Opt:** Jonsbråten et al., 1998, Goel and Grossmann, 2004, 2006, Novoa et al., 2016, Gutin et al. 2015



**Stochastic Opt:** Jonsbråten et al., 1998, Goel and Grossmann, 2004, 2006, Novoa et al., 2016, Gutin et al. 2015

**Robust Opt:** Poss, 2013, 2014, Bertsimas and Vayanos, 2015, Vujanic et al., 2016, Zhang et al., 2016

**Stochastic Opt:** Jonsbråten et al., 1998, Goel and Grossmann, 2004, 2006, Novoa et al., 2016, Gutin et al. 2015

**Robust Opt:** Poss, 2013, 2014, Bertsimas and Vayanos, 2015, Vujanic et al., 2016, Zhang et al., 2016

$$\min_{\mathbf{x}, \mathbf{y}} \mathbf{c}^\top \mathbf{x} + \mathbf{d}^\top \mathbf{y}$$

$$\text{s.t. } \mathbf{Ax} + \mathbf{\Xi y} \leq \mathbf{b} \quad \forall \mathbf{\Xi} \in \mathcal{U}(\mathbf{x})$$

- ▶ Model interpretation.
  - ▶ Proactive uncertainty control
  - ▶ Natural effects
- ▶ Possible dependencies :
  - ▶  $\mathcal{U}(\mathbf{x}) = \{\mathbf{\Xi} \mid \mathbf{G} \cdot \text{vec}(\mathbf{\Xi}) \leq \mathbf{g} + \mathbf{\Delta x}\}$
  - ▶  $\mathcal{U}(\mathbf{x}) = \{\mathbf{\Xi} \mid \text{vec}(\mathbf{\Xi}) = \text{vec}(\bar{\mathbf{\Xi}}) + \mathbf{L}\mathbf{u}, \|\mathbf{u}\|_2 \leq 1 - \gamma x\}$

$$\min_{\mathbf{x}, \mathbf{y}} \mathbf{c}^\top \mathbf{x} + \mathbf{d}^\top \mathbf{y}$$

$$\text{s.t. } \mathbf{a}_i^\top \mathbf{x} + \boldsymbol{\xi}_i^\top \mathbf{y} \leq b_i \quad \forall \boldsymbol{\xi}_i \in \mathcal{U}_i^P(\mathbf{x})$$

$$\mathcal{U}_i^P(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}_i \boldsymbol{\xi} \leq \mathbf{g}_i + \boldsymbol{\Delta}_i \mathbf{x}\}$$

$$\begin{array}{ll} \min_{\mathbf{x}, \mathbf{y}} & \mathbf{c}^\top \mathbf{x} + \mathbf{d}^\top \mathbf{y} \\ \text{s.t.} & \mathbf{a}_i^\top \mathbf{x} + \boldsymbol{\xi}_i^\top \mathbf{y} \leq b_i \quad \forall \boldsymbol{\xi}_i \in \mathcal{U}_i^P(\mathbf{x}) \end{array}$$

$$\mathcal{U}_i^P(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}_i \boldsymbol{\xi} \leq \mathbf{g}_i + \boldsymbol{\Delta}_i \mathbf{x}\}$$

- ▶ Reformulation leads to a bilinear program.
- ▶ Indicates difficulty of problem.

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{y}} \quad & \mathbf{c}^\top \mathbf{x} + \mathbf{d}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{a}_i^\top \mathbf{x} + \boldsymbol{\xi}_i^\top \mathbf{y} \leq b_i \quad \forall \boldsymbol{\xi}_i \in \mathcal{U}_i^P(\mathbf{x}) \end{aligned}$$

$$\mathcal{U}_i^P(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}_i \boldsymbol{\xi} \leq \mathbf{g}_i + \boldsymbol{\Delta}_i \mathbf{x}\}$$

## THEOREM

*The robust linear problem with uncertainty set  $\mathcal{U}^P$  is NP-complete.*

- ▶ If  $x$  is binary, Big-M leads to MILP reformulation.
- ▶ Poor numerical performance
- ▶ Linearization does not leverage set structure
- ▶ Imposing structure allows better reformulations.

- ▶ If  $\mathbf{x}$  is binary, Big-M leads to MILP reformulation.
- ▶ Poor numerical performance
- ▶ Linearization does not leverage set structure
- ▶ Imposing structure allows better reformulations.

Let  $\mathbf{x} \in \{0, 1\}^n$

$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$

Constraint to be reformulated:

$$\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\overline{\Pi}}(\mathbf{x}).$$

$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$



$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$

$$\max_{\boldsymbol{\xi}} \mathbf{y}^\top \boldsymbol{\xi}$$

$$\text{s.t. } \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}$$

$$\boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x})$$

$$\boldsymbol{\xi} \geq \mathbf{0}$$

$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$

$$\max_{\boldsymbol{\xi}} \mathbf{y}^\top \boldsymbol{\xi}$$

$$\text{s.t. } \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}$$

$$\boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x})$$

$$\boldsymbol{\xi} \geq \mathbf{0}$$

$$\max_{\mathbf{z}, \boldsymbol{\zeta}} (\mathbf{y} - \overline{\Pi}\mathbf{x})^\top \mathbf{z} + \mathbf{y}^\top \boldsymbol{\zeta}$$

$$\text{s.t. } \mathbf{G}(\mathbf{z} + \boldsymbol{\zeta}) \leq \mathbf{g}$$

$$\mathbf{z} \leq \mathbf{W}\mathbf{e}$$

$$\boldsymbol{\zeta} \leq \mathbf{v}$$

$$\mathbf{z}, \boldsymbol{\zeta} \geq \mathbf{0}$$

$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$

$$\max_{\boldsymbol{\xi}} \mathbf{y}^\top \boldsymbol{\xi}$$

$$\text{s.t. } \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}$$

$$\boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x})$$

$$\boldsymbol{\xi} \geq \mathbf{0}$$

$$\max_{\mathbf{z}, \boldsymbol{\zeta}} (\mathbf{y} - \overline{\Pi}\mathbf{x})^\top \mathbf{z} + \mathbf{y}^\top \boldsymbol{\zeta}$$

$$\text{s.t. } \mathbf{G}(\mathbf{z} + \boldsymbol{\zeta}) \leq \mathbf{g}$$

$$\mathbf{z} \leq \mathbf{W}\mathbf{e}$$

$$\boldsymbol{\zeta} \leq \mathbf{v}$$

$$\mathbf{z}, \boldsymbol{\zeta} \geq \mathbf{0}$$

- $\overline{\Pi}$ : of upper bounds on dual variables. Similar to Big-M.

$$\mathcal{U}^{\overline{\Pi}}(\mathbf{x}) = \{\boldsymbol{\xi} \mid \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}, \quad \boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x}), \quad \boldsymbol{\xi} \geq \mathbf{0}\}$$

$$\max_{\boldsymbol{\xi}} \mathbf{y}^\top \boldsymbol{\xi}$$

$$\text{s.t. } \mathbf{G}\boldsymbol{\xi} \leq \mathbf{g}$$

$$\boldsymbol{\xi} \leq \mathbf{v} + \mathbf{W}(\mathbf{e} - \mathbf{x})$$

$$\boldsymbol{\xi} \geq \mathbf{0}$$

$$\max_{\mathbf{z}, \boldsymbol{\zeta}} (\mathbf{y} - \overline{\Pi}\mathbf{x})^\top \mathbf{z} + \mathbf{y}^\top \boldsymbol{\zeta}$$

$$\text{s.t. } \mathbf{G}(\mathbf{z} + \boldsymbol{\zeta}) \leq \mathbf{g}$$

$$\mathbf{z} \leq \mathbf{W}\mathbf{e}$$

$$\boldsymbol{\zeta} \leq \mathbf{v}$$

$$\mathbf{z}, \boldsymbol{\zeta} \geq \mathbf{0}$$

- ▶  $\overline{\Pi}$ : of upper bounds on dual variables. Similar to Big-M.
- ▶ Problem convex in  $\mathbf{x}$  and  $\mathbf{y}$ .
- ▶ Network interdiction (Cormican et al. 1996).

## THEOREM

*The constraint  $\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\overline{\Pi}}(\mathbf{x})$  can be reformulated as*

$$\mathbf{t}^\top \mathbf{g} + \mathbf{r}^\top \mathbf{W} \mathbf{e} + \mathbf{s}^\top \mathbf{v} \leq b$$

$$\mathbf{s}^\top + \mathbf{t}^\top \mathbf{G} \geq \mathbf{y}^\top$$

$$\mathbf{r}^\top + \mathbf{t}^\top \mathbf{G} \geq \mathbf{y}^\top - \mathbf{x}^\top \overline{\Pi}$$

$$\mathbf{r}, \mathbf{s}, \mathbf{t} \geq \mathbf{0}.$$

## THEOREM

*The constraint  $\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\overline{\Pi}}(\mathbf{x})$  can be reformulated as*

$$\mathbf{t}^\top \mathbf{g} + \mathbf{r}^\top \mathbf{W} \mathbf{e} + \mathbf{s}^\top \mathbf{v} \leq b$$

$$\mathbf{s}^\top + \mathbf{t}^\top \mathbf{G} \geq \mathbf{y}^\top$$

$$\mathbf{r}^\top + \mathbf{t}^\top \mathbf{G} \geq \mathbf{y}^\top - \mathbf{x}^\top \overline{\Pi}$$

$$\mathbf{r}, \mathbf{s}, \mathbf{t} \geq \mathbf{0}.$$

- ▶ Fewer constraints than Big-M reformulation.
- ▶ Convex problem : use of cut-generating methods.
- ▶ Better solution times.

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}^{SP}(\mathbf{x})} \quad & \sum_{(i,j) \in \mathcal{A}} c x_{ij} + \sum_{(i,j) \in \mathcal{A}} d_{ij}(\boldsymbol{\xi}) y_{ij} \\ \text{s.t.} \quad & \mathbf{x} \in \{0,1\}^{|\mathcal{A}|}, \mathbf{y} \in Y, \end{aligned}$$

$$\mathcal{U}^{SP}(\mathbf{x}) = \left\{ \boldsymbol{\xi} \mid \sum_{(i,j) \in \mathcal{A}} \xi_{ij} \leq \Gamma, \xi_{ij} \leq 1 - \gamma x_{ij}, \xi_{ij} \geq 0 \quad \forall (i,j) \in \mathcal{A} \right\}$$

# Appliction : Shortest Path

$c$  : cost of reduction

$$\min_{\mathbf{x}, \mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}^{SP}(\mathbf{x})} \sum_{(i,j) \in \mathcal{A}} c x_{ij} + \sum_{(i,j) \in \mathcal{A}} d_{ij}(\boldsymbol{\xi}) y_{ij}$$

s.t.  $\mathbf{x} \in \{0, 1\}^{|\mathcal{A}|}$ ,  $\mathbf{y} \in Y$ ,

$$\mathcal{U}^{SP}(\mathbf{x}) = \left\{ \boldsymbol{\xi} \mid \sum_{(i,j) \in \mathcal{A}} \xi_{ij} \leq \Gamma, \xi_{ij} \leq 1 - \gamma x_{ij}, \xi_{ij} \geq 0 \quad \forall (i,j) \in \mathcal{A} \right\}$$



# Application : Shortest Path

$c$  : cost of reduction


$$\min_{\mathbf{x}, \mathbf{y}} \max_{\boldsymbol{\xi} \in \mathcal{U}^{SP}(\mathbf{x})} \sum_{(i,j) \in \mathcal{A}} c x_{ij} + \sum_{(i,j) \in \mathcal{A}} d_{ij}(\boldsymbol{\xi}) y_{ij}$$

s.t.  $\mathbf{x} \in \{0, 1\}^{|\mathcal{A}|}$ ,  $\mathbf{y} \in Y$ ,

$$\mathcal{U}^{SP}(\mathbf{x}) = \left\{ \boldsymbol{\xi} \mid \sum_{(i,j) \in \mathcal{A}} \xi_{ij} \leq \Gamma, \xi_{ij} \leq 1 - \gamma x_{ij}, \xi_{ij} \geq 0 \quad \forall (i,j) \in \mathcal{A} \right\}$$


$\gamma$  : uncertainty reduction

# Network



$$\Gamma = 1, \gamma = 0.8$$


# Network



$$\Gamma = 1, \gamma = 0.8$$

**SP**      **Nominal** = 95      **Worst Case** = 127

# Network




$$\Gamma = 1, \gamma = 0.8$$

**SP**      **Nominal** = 95      **Worst Case** = 127

**RSP**      **Nominal** = 97.4      **Worst Case** = 110.15

# Network




$$\Gamma = 1, \gamma = 0.8$$

**SP**      **Nominal** = 95      **Worst Case** = 127

**RSP**      **Nominal** = 97.4      **Worst Case** = 110.15

**DDRSP**      **Nominal** =  $95.6 + c$       **Worst Case** =  $108.5 + c$

# Network



$$\Gamma = 1, \gamma = 0.8$$


**SP**      **Nominal** = 95      **Worst Case** = 127

**RSP**      **Nominal** = 97.4      **Worst Case** = 110.15

**DDRSP**      **Nominal** =  $95.6 + c$       **Worst Case** =  $108.5 + c$


# Numerical Results : Speed

100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ ,  $\Gamma = 2$



# Numerical Results : Speed


100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ ,  $\Gamma = 2$





# Numerical Results : Speed


100 random graphs,  $c = 1.0, \gamma = 0.2, \Gamma = 2$



$\Rightarrow \overline{\Pi}$  formulation better than Big-M


# Numerical Results : Performance

100 random graphs, 50 samples,  $|\mathcal{V}| = 30, \gamma = 0.2, \Gamma = 3$




# Numerical Results : Performance

100 random graphs, 50 samples,  $|\mathcal{V}| = 30, \gamma = 0.2, \Gamma = 3$





# Numerical Results : Performance

100 random graphs, 50 samples,  $|\mathcal{V}| = 30, \gamma = 0.2, \Gamma = 3$




# Numerical Results : Performance

100 random graphs, 50 samples,  $|\mathcal{V}| = 30, \gamma = 0.2, \Gamma = 3$





$\Rightarrow$  Performance of DDU improves with lower reduction costs.

100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ , nodes = 30





# Path Behavior

100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ , nodes = 30



# Path Behavior



100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ , nodes = 30





# Path Behavior


100 random graphs,  $c = 1.0$ ,  $\gamma = 0.2$ , nodes = 30



The number of arcs in the path increases and then decreases with increase in the total amount of uncertainty.

# Conclusion

- ▶ Decision-dependent uncertainty allows to reduce conservatism by proactive control or better models
- ▶ Problems with decision-dependent uncertainty are NP-complete.
- ▶ Leveraging set structure allows to improve performance.



Kartikey Sharma

kartikeysharma2014@u.northwestern.edu

Omid Nohadani and Kartikey Sharma. "Optimization under decision-dependent uncertainty." SIAM Journal on Optimization 28.2 (2018): 1773-1795.

## LEMMA

*If  $\mathbf{G}$  and  $\mathbf{y}$  are nonnegative, then  $\pi_i(\mathbf{x}, \mathbf{y}) \leq y_i \forall (\mathbf{x}, \mathbf{y})$ .*

- ▶ Consider an instance of the 3-Satisfiability problem (3-SAT) for a set  $N = \{1, 2, \dots, n\}$  of literals and  $m$  clauses, which tries to find a solution  $\mathbf{x} \in \{0, 1\}^n$  that satisfies

$$x_{i_1} + x_{i_2} + (1 - x_{i_3}) \geq 1 \quad \forall i = 1, \dots, m.$$

- ▶ Next, consider the following special decision dependent problem with  $\mathbf{x} \in \mathbb{R}^n$ ,  $\mathbf{y} \in \mathbb{R}^m$ ,  $z \in \mathbb{R}$

$$\min_{\mathbf{x}, \mathbf{y}, z \geq 0} \left\{ -z \mid z - \mathbf{a}^\top \mathbf{y} \leq 0, \quad \forall \mathbf{a} \in \mathcal{U}(\mathbf{x}), \quad \mathbf{x}, \mathbf{y} \leq \mathbf{1}, \quad -\mathbf{y} \leq -\mathbf{1} \right\},$$

$$\mathcal{U}(\mathbf{x}) = \{(a_1, \dots, a_m) \mid a_i \geq x_{i_1}, \quad a_i \geq x_{i_2}, \quad a_i \geq 1 - x_{i_3}, \quad a_i \leq 1\}$$

Note that the 3-SAT problem is embedded in this set.

## THEOREM

*The constraint  $\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\bar{\Pi}}(\mathbf{x})$  has the reformulation*

$$\mathbf{t}^\top \mathbf{d} + \mathbf{s}^\top \mathbf{v} + \mathbf{s}^\top \mathbf{W} \mathbf{e} - \sum_i r_i \leq b$$

$$\mathbf{s}^\top + \mathbf{t}^\top \mathbf{D} \geq \mathbf{y}^\top$$

$$w_i s_i - M(1 - x_i) \leq r_i \leq M x_i$$

$$r_i \leq w_i s_i$$

$$\mathbf{r}, \mathbf{s}, \mathbf{t} \geq \mathbf{0}$$

## THEOREM

*The constraint  $\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\overline{\Pi}}(\mathbf{x})$  has the reformulation*

$$\mathbf{t}^\top \mathbf{d} + \mathbf{s}^\top \mathbf{v} + \mathbf{s}^\top \mathbf{W} \mathbf{e} - \sum_i r_i \leq b$$

$$\mathbf{s}^\top + \mathbf{t}^\top \mathbf{D} \geq \mathbf{y}^\top$$

$$w_i s_i - M(1 - x_i) \leq r_i \leq M x_i$$

$$r_i \leq w_i s_i$$

$$\mathbf{r}, \mathbf{s}, \mathbf{t} \geq \mathbf{0}$$

- Large number of constraints and poor numerical performance

## THEOREM

*The constraint  $\mathbf{y}^\top \boldsymbol{\xi} \leq b \quad \forall \boldsymbol{\xi} \in \mathcal{U}^{\bar{\Pi}}(\mathbf{x})$  has the reformulation*

$$\mathbf{t}^\top \mathbf{d} + \mathbf{s}^\top \mathbf{v} + \mathbf{s}^\top \mathbf{W} \mathbf{e} - \sum_i r_i \leq b$$


$$\mathbf{s}^\top + \mathbf{t}^\top \mathbf{D} \geq \mathbf{y}^\top$$

$$w_i s_i - M(1 - x_i) \leq r_i \leq M x_i$$

$$r_i \leq w_i s_i$$

$$\mathbf{r}, \mathbf{s}, \mathbf{t} \geq \mathbf{0}$$

- ▶ Large number of constraints and poor numerical performance
- ▶ Does not leverage the structure of the uncertainty set





- Dimitris Bertsimas and Phebe Vayanos. Data-driven learning in dynamic pricing using adaptive optimization. 2015. URL [http://www.optimization-online.org/DB\\_HTML/2014/10/4595.html](http://www.optimization-online.org/DB_HTML/2014/10/4595.html).
- Vikas Goel and Ignacio E Grossmann. A stochastic programming approach to planning of offshore gas field developments under uncertainty in reserves. *Computers & Chemical Engineering*, 28(8):1409–1429, 2004.
- Vikas Goel and Ignacio E Grossmann. A class of stochastic programs with decision dependent uncertainty. *Mathematical Programming*, 108(2-3):355–394, 2006.
- Tore W Jonsbråten, Roger JB Wets, and David L Woodruff. A class of stochastic programs with decision dependent random elements. *Annals of Operations Research*, 82:83–106, 1998.
- Michael Poss. Robust combinatorial optimization with variable budgeted uncertainty. *4OR*, 11(1):75–92, 2013.
- Robin Vujanic, Paul Goulart, and Manfred Morari. Robust optimization of schedules affected by uncertain events. *Journal of Optimization Theory and Applications*, pages 1–22, 2016.