


Remote sensing system perspective on the Okavango

Thomas Gumbricht


WERRD Mid term meeting, Heja Lodge, Windhoek


25-28 August 2003





The integrated system perspective:


The Okavango Delta – Africa's last eden























































Rainfall and flooding - Okavango Delta








Annual sediment inflow
600 000 tonnes



Raises the Okavango Delta
1 meter in 50 000 years




**Annual sediment inflow
600 000 tonnes**




**Raises the Okavango Delta
1 meter in 50 000 years**

Population - Okavango Basin





**Total population:
400 000**





S


E














The Delta





Psst!

Ecoregion Classification methodology



**Water = 2.5 m below reference
level**



Permanent Swamp = 2.0 m below reference level



**Primary floodplain = 1.5 m below reference
level**



Secondary floodplain = 1.0 m below reference level



Grassland = reference level



Grassland

Salt pan = 0.5 m below reference level



Occasionally flooded grassland = 0.5 m below reference level



Grassland (with occasional flooding)

Salt pan = 0.5 m below reference level



Riverine forest = 1.2 m above reference level



Dry woodland = reference level




Dry Woodland (dominated by Mopane)

Dry woodland = reference level




Dry Woodland (dominated by Acacia)




Landcover ecoregions




Landcover ecoregions






Surveyed profiles




Microtopography




Relative microtopography of the Okavango Delta





Okavango Delta water balance




Classification of historical flood area


Unsupervised classification of ~ 400 satellite images (NOAA AVHRR, ERS-2 ATSR), and supervised classification of Landsat MSS / TM (subset of ~ 3000 images)



AVHRR




Unsupervised
classification




Manual
reclassification

Evaluation of AVHRR against Landsat TM & ATSR




AVHRR vs. Landsat TM




AVHRR vs. ATSR


Flooding, years (1985-2000)



Flooding, month (1985-2000)





Okavango Delta water balance 1992/93



3 ways to model the water flow in the Okavango Delta

1. Statistically by using historical data
2. Mathematical description of outflow and inflow of small cubes (like a checker-board)
3. Mathematical description of outflow and inflow of “natural” compartments


Statistical modelling of the Okavango Delta annual flooding





Data needed:


- Inflow at Panhandle
- Rainfall over the Delta
- Area of flooding (to be predicted)

Area of flooding =
Inflow at the Panhandle +
local precipitation +
previous years flood









4500 km²



5000 km²



5500 km²



6000 km²



6500 km²



7000 km²



7500 km²



8000 km²



8500 km²



9000 km²



9500 km²



10000 km²



10500 km²



11000 km²



11500 km²



12000 km²




12500 km²




13000 km²




1995




2000



1995








Fully distributed modelling of the Okavango Delta hydrology




Data needed:

- Inflow at Panhandle
- Rainfall over the Delta (in each checker board “cube”)
- Area of flooding (to be predicted)
- Regional and detailed topography (for each “cube”)
- Evapotranspiration in each “cube”
- Water storage in each “cube”
- Water flow resistance properties in each “cube”


Precipitation and evapotranspiration – driving variables of the Delta water balance




Remote sensing for estimating spatial distribution of precipitation and evapotranspiration



Field data for estimation of net recharge



Parameterisation of evapotranspiration, recharge and net exchange



Field data for accurate point measurements of surface energy balance (evapotranspiration)



Net radiometre




Microclimate station




At least 1 station for each land cover class

Vertical water balance of the Okavango Delta


Rain




PET



2 layer modflow model of the Okavango Delta



2 layer modflow model of the Okavango Delta Preliminary tests







Semi distributed modelling of the Okavango Delta hydrology




Data needed:


- Inflow at Panhandle
- Rainfall over the Delta (in each compartment)
- Area of flooding (to be predicted)
- Detailed topography (in each compartment)
- Evapotranspiration (for each compartment)
- Water storage in each compartment
- Water flow resistance between each compartment







Matter balance and islands – redirecting water flow on different time scales



Primary islands built from accumulation of clastic sediments

Island types

Inverted channel island



Primary islands built from accumulation of clastic sediments

Island types

Scroll bar island




Primary islands built from accumulation of clastic sediments

Island types

Anthill island



Evapotranspiration, salinity balance and island secondary growth



Secondary islands grown from precipitation of chemical sediments

Island types

Riparian forest island



Secondary islands grown from precipitation of chemical sediments

Island types


Salt islands




Detail of the Chitabe area



Detailed relief of the Chitabe area






Island max




Island core




Salt Balance: Coastline from Remote Sensing




Order of magnitude
correct



	A	B	C	D	E	F	G	H
Roundness	0.49	0.91	0.51	0.48	0.36	0.47	0.58	0.92
Regional salt position	distal*	na	na	proximal	distal	equal	proximal	na
Channel salt position	front	na	na	back	back	back	back*	na



Island orientation – interacting with water flow over the Delta surface



Summary

The Rift Valley and the Superswell created the Okavango and its Basin

The Basin feeds the Alluvial fan with sediments that keeps the surface flat

The Basin feeds clean water to the Delta that sits on the Alluvial fan

The living Wetland is sustained by the shifts in the Delta and Alluvial Fan

The Islands are born from the shifts in the Delta and Alluvial Fan

The Islands are the kidneys of the Okavango – removing salt from the Wetland

Conclusion

The **Superswell** will eventually divert the Okavango to the Zambezi

Clean water and sediments from the **Basin** is a must for the Okavango to live

Loss of water inflow would decrease the size of the **Delta** and **Wetland**

Loss of sediment inflow would disrupt the changes sustaining the living **Wetland**

Loss of sediment inflow would disrupt the **Island** birth and growth process

The growth of the **Alluvial Fan** preserves the flatness sustaining the **Delta**

The death and birth of channels is a creative destruction sustaining the living **Wetland**

Island birth and growth is needed for removing salt from the **Wetland**

Acknowledgements

This work was done in close collaboration with the University of the Witwatersrand in Johannesburg, South Africa, and was part of the SAFARI 2000 Research Initiative.

The studies were financially supported through

- Scholarship from Royal Swedish Academy of Sciences and The Swedish Foundation for International Cooperation in Research and Higher Education (STINT)
- Swedish International Development Agency (SIDA), research expenses (via KTH and Linköping University)

