Problem C
Three Kinds of Dice
Time limit: 1 second


Image from rawpixel, CC0

See how they roll! According to a famous story, Warren Buffett once challenged Bill Gates to a simple game of dice. He had three dice; the first player could examine them and choose one of the three. The second player would then choose one of the remaining dice, and both players would roll their dice against each other, aiming for the highest numbers. Warren offered to let Bill go first, but this made Bill suspicious so he opted to go second. It turned out to be a wise choice: these were *intransitive dice*. The first die had an advantage when rolling against the second, the second had an advantage when rolling against the third, but the first did not have an advantage when rolling against the third!

To formalize this: define a "die" as any shape with at least one face such that each face shows a positive integer. When a die is rolled, one of its faces is selected uniformly at random. When two dice roll against each other, the die whose selected face shows a higher number earns 1 point; if both numbers are equal, each die earns $\frac{1}{2}$ points. For dice D and D', define $score(D, D')$ as the expected number of points D earns from a single roll against D'. If $score(D, D') > \frac{1}{2}$, we say that D has an advantage over D'; if $score(D, D') = \frac{1}{2}$, the two dice are tied. For example, if D is the first die in the sample input and D' is the second, $score(D, D') = \frac{4}{9}$ and $score(D', D) = \frac{5}{9}$, so D' has an advantage over D.

Given two dice D_1 and D_2 such that D_1 has an advantage over D_2, you want a third die D_3 that forms an intransitive trio with the other two. Among all D_3 that have an advantage over or tie with D_1, compute the lowest possible $score(D_3, D_2)$. If this is less than $\frac{1}{2}$, you can make an intransitive trio! Similarly, among all D_3 such that D_2 has an advantage over or ties with D_3, compute the highest possible $score(D_3, D_1)$.

Input

The input contains two lines, each describing one die. One of the dice (the first or the second) has an advantage over the other. The die with the advantage is D_1 and the other is D_2.

The first integer on a line gives n ($1 \le n \le 10^5$), the number of faces on the die. Then follow n integers f_i ($1 \le f_i \le 10^9$ for each $1 \le i \le n$), giving the integer on each face.

Output

Output one line containing the lowest $score(D_3, D_2)$ and the highest $score(D_3, D_1)$ under the above conditions. The two scores do not need to use the same die D_3. Your answer should have an absolute error of at most 10^{-6}.

Sample Input 1	Sample Output 1
6 1 1 6 6 8 8 3 2 4 9	0.291666667 0.750000000

Sample Input 2	Sample Output 2
4 9 3 7 5 3 4 2 3	0.500000000 0.500000000