UNIT-IV
INVERSE LAPLACE TRANSFORM

Topic Learning Objectives:

Upon Completion of this unit, students will be able to:

- Obtain inverse Laplace transform of rational functions.
- Solve ordinary linear differential equations

Definition

Let $L\{f\ (t)\} = F(s)$, then f(t) is defined as the inverse Laplace transform of $F(s)$ and is denoted by $L^{-1}\{F(s)\}$. Thus $L^{-1}\{F(s)\} = f\ (t)$.

Linearity Property

Let $L^{-1}\{F(s)\} = f\ (t)$ and $L^{-1}\{G(s)\} = g(t)$ and a and b be any two constants,
Then $L^{-1}\ [a\ F(s) + b\ G(s)] = a\ L^{-1}\{F(s)\} + b\ L^{-1}\{\ G(s)\}$

Table of Inverse Laplace Transforms

F(s)	$f(t) = L^{-1}\{F(s)\}$
$\frac{1}{s}, s > 0$	1
$\frac{1}{s-a}, s > a$	e^{at}
$\frac{s}{s^2+a^2}, s > 0$	$\cos at$
$\frac{1}{s^2+a^2}, s > 0$	$\frac{\sin at}{a}$
$\frac{1}{s^2-a^2}, s > \lvert a \rvert$	$\frac{\sinh at}{a}$
$\frac{s}{s^2-a^2}, s > \lvert a \rvert$	$\cosh at$
$\frac{1}{s^{n+1}}, s > 0$ $n = 0, 1, 2, 3, \ldots$	$\frac{t^n}{n!}$
$\frac{1}{s^{n+1}}, s > 0$ $n \neq -1, -2, -3, \ldots..$	$\frac{t^n}{\Gamma(n+1)}$

Examples

1. Find the inverse Laplace transforms of the following:

$$(i)\ \frac{1}{2s-5} \quad (ii)\ \frac{s+b}{s^2+a^2} \quad (iii)\ \frac{2s-5}{4s^2+25}+\frac{4s-9}{9-s^2}$$

Solution:

$$(i)\ L^{-1}\left\{\frac{1}{2s-5}\right\}=\frac{1}{2}L^{-1}\left\{\frac{1}{s-5/2}\right\}=\frac{1}{2}e^{\frac{5t}{2}}$$

$$(ii)\ L^{-1}\left\{\frac{s+b}{s^2+a^2}\right\}=L^{-1}\left\{\frac{s}{s^2+a^2}\right\}+b\,L^{-1}\left\{\frac{1}{s^2+a^2}\right\}=\cos at+\frac{b}{a}\sin\ at$$

$$(iii)\ L^{-1}\left[\frac{2s-5}{4s^2+25}+\frac{4s-8}{9-s^2}\right]=\frac{2}{4}L^{-1}\left\{\frac{s-5/2}{s^2+\frac{25}{4}}\right\}-4L^{-1}\left\{\frac{s-9/2}{s^2-9}\right\}$$

$$=\frac{1}{2}\left[\cos\frac{5t}{2}-\sin\frac{5t}{2}\right]-4\left[\cos h3t-\frac{3}{2}\sin\ h3t\right]$$

Shifting property [Evaluation of $L^{-1}[F(s-a)]$]

If $L^{-1}[F(s)]=f(t)$, then $L^{-1}[F(s-a)]=e^{at}L^{-1}[F(s)]=e^{at}f(t)$

Examples:

1. Evaluate : $L^{-1}\left\{\frac{3s+1}{(s+1)^4}\right\}$.

Solution:

$$L^{-1}\left\{\frac{3(s+1-1)+1}{(s+1)^4}\right\}=3\,L^{-1}\left\{\frac{1}{(s+1)^3}\right\}-2\,L^{-1}\left\{\frac{1}{(s+1)^4}\right\}$$

$$=3e^{-t}\,L^{-1}\left\{\frac{1}{s^3}\right\}-2e^{-t}\,L^{-1}\left\{\frac{1}{s^4}\right\}$$

Using the formula

$L^{-1}\{\frac{1}{s^{n+1}}\}=\frac{t^n}{n!}$ and taking $n=2$ and 3, we get

$$=\frac{3e^{-t}t^2}{2}-\frac{e^{-t}t^3}{3}.$$

2. Evaluate : $L^{-1}\left\{\dfrac{s+2}{s^2\text{-}2s+5}\right\}$.

Solution:

$$\text{Given} = L^{-1}\left\{\frac{s+2}{(s-1)^2+4}\right\} = L^{-1}\left\{\frac{(s-1)+3}{(s-1)^2+4}\right\} = L^{-1}\left\{\frac{s-1}{(s-1)^2+4}\right\} + 3L^{-1}\left\{\frac{1}{(s-1)^2+4}\right\}$$

$$= e^t\, L^{-1}\left\{\frac{s}{s^2+4}\right\} + 3\,e^t\, L^{-1}\left\{\frac{1}{s^2+4}\right\}$$

$$= e^t \cos 2t + \frac{3}{2} e^t \sin 2t$$

3. Evaluate : $L^{-1}\left\{\dfrac{2s+1}{s^2+3s+1}\right\}$.

Solution:

$$\text{Given} = 2L^{-1}\left\{\frac{\left(s+\frac{3}{2}\right)-1}{\left(s+\frac{3}{2}\right)^2-5/4}\right\} = 2\left[L^{-1}\left\{\frac{\left(s+\frac{3}{2}\right)}{\left(s+\frac{3}{2}\right)^2-5/4}\right\} - L^{-1}\left\{\frac{1}{\left(s+\frac{3}{2}\right)^2-5/4}\right\}\right]$$

$$= 2\left[e^{\frac{-3t}{2}} L^{-1}\left\{\frac{s}{s^2-5/4}\right\} - e^{\frac{-3t}{2}} L^{-1}\left\{\frac{1}{s^2-5/4}\right\}\right]$$

$$= 2e^{\frac{-3t}{2}}\left[cosh\frac{\sqrt{5}}{2}t - \frac{2}{\sqrt{5}} sinh\frac{\sqrt{5}}{2}t\right].$$

4. Evaluate : $L^{-1}\left\{\dfrac{2s^2+5s-4}{s^3+s^2-2s}\right\}$.

Solution:

We have $\dfrac{2s^2+5s-4}{s^3+s^2-2s} = \dfrac{2s^2+5s-4}{s(s^2+s-2)} = \dfrac{2s^2+5s-4}{s(s+2)(s-1)} = \dfrac{A}{s} + \dfrac{B}{s+2} + \dfrac{C}{s-1}$ -------- (1)

Then $2s^2+5s-4 = A(s+2)(s-1) + B\,s\,(s-1) + Cs\,(s+2)$

For $s = 0$, $A = 2$, for $s = 1$, $C = 1$ and for $s = -2$, $B = -1$.

Using these values in (1),

$$L^{-1}\left\{\frac{2s^2+5s-4}{s^3+s^2-2s}\right\} = L^{-1}\left\{\frac{2}{s}\right\} - L^{-1}\left\{\frac{1}{s+2}\right\} + L^{-1}\left\{\frac{1}{s-1}\right\} = 2 - e^{-2t} + e^t$$

5. Use the method of partial fractions to find the time signals corresponding to the Laplace

transform function

$F(s) = \frac{4s+5}{(s+1)^2(s+2)}$

Solution:

Consider

$$\frac{4s+5}{(s+1)^2(s+2)} = \frac{A}{(s+1)^2} + \frac{B}{s+1} + \frac{C}{s+2}$$

Then $4s + 5 = A(s + 2) + B(s + 1)(s + 2) + C(s + 1)^2$

For $s = -1$, $A = 1$, for $s = -2$, $C = -3$

Comparing the coefficients of s^2 to get $B + C = 0$ so that $B = 3$.

Using these values in (1) to get

$$\frac{4s+5}{(s+1)^2+(s+2)} = \frac{1}{(s+1)^2} + \frac{3}{(s+1)} - \frac{3}{s+2}$$

Hence

$$L^{-1}\left\{\frac{4s+5}{(s+1)^2+(s+2)}\right\} = e^{-t}L^{-1}\left\{\frac{1}{s^2}\right\} + 3e^{-t}L^{-1}\left\{\frac{1}{s}\right\} - 3e^{-2t}L^{-1}\left\{\frac{1}{s}\right\}$$

$$= te^{-t} + 3e^{-t} - 3e^{-2t}$$

5. Evaluate : $L^{-1}\left\{\frac{s^3}{s^4-a^4}\right\}$.

Solution:

$$Let\ \frac{s^3}{s^4-a^4} = \frac{A}{s-a} + \frac{B}{s+a} + \frac{Cs+D}{s^2+a^2} \qquad \text{--------} \qquad (1)$$

Hence, $s^3 = A(s + a)(s^2 + a^2) + B(s - a)(s^2 + a^2) + (Cs + D)(s^2 - a^2)$

For $s = a, A = ¼$, *for* $s = -a$, $B = ¼$, comparing the constant terms to get

$D = a$, $(A-B) = 0$,

Comparing the coefficients of s^3 to get $1 = A + B + C$ and so $C = ½$.

Using these values in (1),

$$\frac{s^3}{s^4-a^4} = \frac{1}{4}\left[\frac{1}{s-a} + \frac{1}{s+a}\right] + \frac{1}{2}\frac{s}{s^2+a^2}$$

Taking inverse transforms,

$$L^{-1}\left\{\frac{s^3}{s^4-a^4}\right\} = \frac{1}{4}\left[e^{at} + e^{-at}\right] + \frac{1}{2}\cos at = \frac{1}{2}[\cosh at + \cos at]$$

6. Evaluate : $L^{-1}\left\{\frac{s}{s^4+s^2+1}\right\}$.

Solution:

Consider, $\dfrac{s}{s^4+s^2+1}=\dfrac{s}{(s^2+s+1)+(s^2-s+1)}=\dfrac{1}{2}\left[\dfrac{2s}{(s^2+s+1)(s^2-s+1)}\right]$

$$=\frac{1}{2}\left[\frac{(s^2+s+1)-(s^2-s+1)}{(s^2+s+1)(s^2-s+1)}\right]=\frac{1}{2}\left[\frac{1}{(s^2-s+1)}-\frac{1}{(s^2+s+1)}\right]$$

$$=\frac{1}{2}\left[\frac{1}{\left(s-\frac{1}{2}\right)^2+\frac{3}{4}}-\frac{1}{\left(s+\frac{1}{2}\right)^2+\frac{3}{4}}\right]$$

Therefore

$$L^{-1}\left\{\frac{s}{s^4+s^2+1}\right\}=\frac{1}{2}\left[e^{\frac{t}{2}}L^{-1}\left(\frac{1}{s^2+\frac{3}{4}}\right)-e^{-\frac{t}{2}}L^{-1}\left(\frac{1}{s^2+\frac{3}{4}}\right)\right]$$

$$=\frac{1}{2}\left[e^{\frac{t}{2}}\frac{\sin\frac{\sqrt{3}}{2}t}{\frac{\sqrt{3}}{2}}-e^{-\frac{t}{2}}\frac{\sin\frac{\sqrt{3}}{2}t}{\frac{\sqrt{3}}{2}}\right]$$

$$=\frac{2}{\sqrt{3}}\sin\left(\frac{\sqrt{3}}{2}t\right)\sin h\left(\frac{t}{2}\right)$$

Evaluation of $L^{-1}[\,e^{-as}F(s)]$

If $L[f(t)]=F(s)$, then $L[f(t-a)H(t-a)]=e^{-as}F(s)$

$$L^{-1}[\,e^{-as}F(s)]=f(t-a)H(t-a)$$

Examples

1. $Evaluate: L^{-1}\left\{\frac{e^{-5s}}{(s-2)^4}\right\}$.

Solution:

$a=5,\ F(s)=\dfrac{1}{(s-2)^4}$

$$f(t)=L^{-1}\{F(s)\}=L^{-1}\left\{\frac{1}{(s-2)^4}\right\}=e^{2t}L^{-1}\left\{\frac{1}{s^4}\right\}\quad=\frac{e^{2t}t^3}{6}$$

$$L^{-1}\left\{\frac{e^{-5s}}{(s-2)^4}\right\}=f(t-a)\,H(t-a)$$

$$=\frac{e^{2(t-5)}(t-5)^3}{6}H(t-5).$$

2. $Evaluate: L^{-1}\left[\dfrac{e^{-\pi s}}{s^2+1}+\dfrac{se^{-2\pi s}}{s^2+4}\right]$.

Solution:

Given $=\{f_1(t-\pi)H(t-\pi)+f_2(t-2\pi)H(t-2\pi)\}$ (1)

$$f_1(t)=L^{-1}\left\{\frac{1}{s^2+1}\right\}=\sin t$$

$$f_2(t)=L^{-1}\left\{\frac{s}{s^2+4}\right\}=\cos 2t$$

(1) leads to

$$\text{Given } = \sin(t-\pi)\,H(t-\pi)+\cos 2(t-2\pi)\,H(t-2\pi)$$
$$=-\cos t\,H(t-\pi)+\cos(2t)\,H(t-2\pi).$$

Inverse transform of logarithmic and trigonometric functions

If $L[tf(t)] = F(s)$ then $L[tf(t)] = L^{-1}\left[-\frac{d}{ds}F(s)\right] = tf(t)$

In general, $L^{-1}\left[(-1)^n\frac{d^n}{ds^n}F(s)\right] = t^n f(t)$

Examples

1. *Evaluate* $L^{-1}\left\{\log\left(\frac{s+a}{s+b}\right)\right\}$.

Let $F(s)=\log\left(\frac{s+a}{s+b}\right)=\log(s+a)-\log(s+b)$

Then $-\frac{d}{ds}F(s)=-\left[\frac{1}{s+a}-\frac{1}{s+b}\right]$

$$L^{-1}\left[-\frac{d}{ds}F(s)\right]=-\left[e^{-at}-e^{-bt}\right]$$

Or $t\,f(t)=e^{-bt}-e^{-at}$. Thus $f(t)=\frac{e^{-bt}-e^{-at}}{b}$.

2. Evaluate $L^{-1}\left\{\tan^{-1}\frac{a}{s}\right\}$.

Let $F(s)=\tan^{-1}\left(\frac{a}{s}\right)$

Then $-\frac{d}{ds}F(s)=\left[\frac{a}{s^2+a^2}\right]$ or $L^{-1}\left[-\frac{d}{ds}F(s)\right]=\sin at$ so that

or $t\,f(t)=\sin at$, $f(t)=\frac{\sin at}{a}$.

Inverse transform of $\frac{F(s)}{s}$

Since $L\int_0^t f(t)dt=\frac{F(s)}{s}$, we have $L^{-1}\left[\frac{F(s)}{s}\right]=\int_0^t f(t)dt$

Examples

1. Evaluate $L^{-1}\left[\frac{1}{s(s^2+a^2)}\right]$.

Solution:

Let $F(s)=\frac{1}{s^2+a^2}$ so that $f(t)=L^{-1}F(s)=\frac{\sin at}{a}$

Then $L^{-1}=\frac{1}{s(s^2+a^2)}=L^{-1}\left[\frac{F(s)}{s}\right]=\int_0^t \frac{\sin at}{a}dt=\frac{(1-\cos at)}{a^2}$

2. $L^{-1}\left[\frac{1}{s^2(s+a)^2}\right]$.

$$L^{-1}\left\{\frac{1}{(s+a)^2}\right\}=te^{-at}$$

$$L^{-1}\left\{\frac{1}{s(s+a)^2}\right\}=\int_0^t e^{-at}t\,dt$$

$$=\frac{1}{a^2}\left[1-e^{-at}(1+at)\right]$$

Now $L^{-1}\left\{\frac{1}{s^2(s+a)^2}\right\}=\frac{1}{a^2}\int_0^t\left[1-e^{-at}(1+at)\right]dt$

$$=\frac{1}{a^3}\left[at\left(1+e^{-at}\right)+2\left(e^{-at}-1\right)\right].$$

Inverse Laplace transform of F(s) using Convolution theorem

If $L^{-1}[F(s)]=f(t)$ and $L^{-1}[G(s)]=g(t)$, then
$L^{-1}[F(s)G(s)]=\int_0^t f(u)g(t-u)du=f(t)*g(t)$

This expression is called the convolution theorem for inverse Laplace transform.

Examples:

Employ convolution theorem to evaluate the following:

1. $L^{-1}\left\{\dfrac{1}{(s+a)(s+b)}\right\}$.

Solution:

Let $F(s) = \dfrac{1}{s+a}, G(s) = \dfrac{1}{s+b}$

Taking inverse transform, we get $f(t) = e^{-at}, g(t) = e^{-bt}$

By convolution theorem,

$$L^{-1}\left\{\frac{1}{(s+a)(s+b)}\right\} = \int_0^t e^{-a(t-u)}e^{-bu}du \quad = e^{-at}\int_0^t e^{(a-b)u}du$$

$$= e^{-at}\left[\frac{e^{(a-b)t}-1}{a-b}\right] \qquad = \frac{e^{-bt}-e^{-at}}{a-b}.$$

2. $L^{-1}\left\{\dfrac{s}{(s^2+a^2)^2}\right\}$.

Solution:

Let $F(s) = \dfrac{1}{s^2+a^2}, G(s) = \dfrac{s}{s^2+a^2}$, then $f(t) = \dfrac{\sin at}{a}, g(t) = \cos at$

By convolution theorem,

$$L^{-1}\left\{\frac{s}{(s^2+a^2)^2}\right\} = \int_0^t \frac{1}{a}\sin a(t-u)\cos au\, du$$

$$= \frac{1}{a}\int_0^t \frac{\sin at + \sin(at-2au)}{2}du,$$

$$= \frac{1}{2a}\left[u\sin at - \frac{\cos(at-2au)}{-2a}\right]_0^t = \frac{t\sin at}{2a}.$$

(3) $L^{-1}\left\{\dfrac{s}{(s-1)(s^2+1)}\right\}$.

Solution:

Let $F(s) = \dfrac{1}{s-1}, G(s) = \dfrac{s}{s^2+1}$

Then $f(t) = e^t, g(t) = sint$

$$L^{-1}\left\{\frac{1}{(s-1)(s^2+1)}\right\} = \int_0^t e^{t-u}\sin u\, du = e^t\left[\frac{e^{-u}}{2}(-\sin u - \cos u)\right]$$

$$= \frac{e^t}{2}\left[e^{-t}(-\sin t - \cos t) - (-1)\right] = \frac{1}{2}\left[e^t - \sin t - \cos t\right]$$

Exercise:

By employing convolution theorem evaluate the following:

1. $L^{-1}\left\{\frac{1}{(s+1)(s^2+1)}\right\}$.

2. $L^{-1}\left\{\frac{s}{(s+2)(s^2+9)}\right\}$.

3. $L^{-1}\left\{\frac{1}{(s^2+a^2)^2}\right\}$.

4. $L^{-1}\left\{\frac{s}{(s^2+a^2)(s^2+b^2)}\right\}, a \neq b.$

5. $L^{-1}\left\{\frac{1}{s^2(s+2)^2}\right\}$.

6. $L^{-1}\left\{\frac{4s+5}{(s-1)^2(s+2)}\right\}$.

Answers:

(ii) $\frac{1}{13}(2\cos 3t + 3\sin 3t - 2e^{-2t})$ (iii) $\frac{1}{2a^3}(\sin at - at\cos at)$ (iv) $\frac{1}{a^2-b^2}(\cos bt - \cos at)$ (v) $\frac{1}{4}(e^{-2t}(t+1)+t-1)$

Laplace Transform Method for Differential Equations

As noted earlier Laplace transform technique is employed to solve initial-value problems. The solution of such a problem is obtained by using the Laplace Transform of the derivatives of function and then the inverse Laplace Transform.

The following are the expressions for the derivatives derived earlier.

$L[f'(t)] = s\,L\,f(t) - f(0)$

$L[f''(t) = s^2\,L\,f(t) - s\,f(0) - f'(0)$

$L[f'''(t) = s^3\,L\,f(t) - s^2\,f(0) - s\,f'(0) - f''(0)$

Examples

1. $y'' + 2y' - 3y = \sin t,\ \ y(0) = y'(0) = 0$

Solution:

Taking the Laplace transform of the given equation,

$$[s^2Ly(t) - sy(0) - y'(0)] + 2[s\ Ly(t) - y(0)] - 3\ L\ y(t) = \frac{1}{s^2+1}$$

Using the given conditions, we get

$$[s\,L\{y(t)\} - y(0)] + L\{y(t)\} = \frac{1}{(s+1)^2}$$

Using the given condition, this becomes

$$(s+1)L\{y(t)\} - 2 = \frac{1}{(s+1)^2} \text{ so that } L\{y(t)\} = \frac{2s^2+4s+3}{(s+1)^3}$$

Taking the inverse Laplace transform, we get

$$Y(s) = L^{-1}\left\{\frac{2s^2+4s+3}{(s+1)^3}\right\}$$

$$= L^{-1}\left[\frac{2(s+1-1)^2+4(s+1-1)+3}{(s+1)^3}\right]$$

$$= L^{-1}\left[\frac{2}{s+1}+\frac{1}{(s+1)^3}\right] = \frac{1}{2}e^{-t}(t^2+4)$$

$$L[y(t)]\left[s^2+2s-3\right] = \frac{1}{s^2+1} \text{ or } L[y(t)] = \frac{1}{(s-1)(s+3)(s^2+1)} \text{ or}$$

$$y(t) = L^{-1}\left[\frac{1}{(s-1)(s+3)(s^2+1)}\right] = L^{-1}\left[\frac{A}{s-1}+\frac{B}{s+3}+\frac{Cs+D}{s^2+1}\right]$$

$$= L^{-1}\left[\frac{1}{8}\,\frac{1}{s-1}-\frac{1}{40}\,\frac{1}{s+3}+\frac{-\frac{s}{10}-\frac{1}{5}}{s^2+1}\right] \text{ (method of partial fractions)}$$

$$= \frac{1}{8}e^{t}-\frac{1}{40}e^{-3t}-\frac{1}{10}(\cos t+2\sin t).$$

2. Employ Laplace Transform method to solve the integral equation.

$$f(t) = 1+\int_0^t f(u)\sin(t-u)du$$

Solution:

Taking Laplace transform of the given equation, we get

$$L\,f(t) = \frac{1}{s}+L\int_0^t f(u)\sin(t-u)du$$

By using convolutio n theorem here .

$$L\,f(t) = \frac{1}{s}+Lf(t)\cdot L\sin t = \frac{1}{s}+\frac{L\,f(t)}{s^2+1} \text{ Thus}$$

$$L\,f(t) = \frac{s^2+1}{s^3} \quad \text{or} \quad f(t) = L^{-1}\left(\frac{s^2+1}{s^3}\right) = 1+\frac{t^2}{2}.$$

3. A particle is moving along a path satisfying , the equation $\frac{d^2x}{dt^2}+6\frac{dx}{dt}+25x=0$ where x denotes the displaceme nt of the particle at time t. If the initial position of the particle is at $x=20$ and the initial speed is 10, find the displaceme nt of the particle at any time t using Laplace transform s.

Solution:

Given equation may be rewritten as

$$x''(t)+6x'(t)+25x(t) = 0$$

Here the initial conditions are $x(0) = 20$, $x'(0) = 10$.

Taking the Laplace transform of the equation, we get

$$L\{x(t)\}[s^2+6s+25]-20s-130=0 \quad or \quad L\{x(t)\}=\frac{20s+130}{s^2+6s+25}$$

so that

$$x(t)=L^{-1}\left[\frac{20s+130}{(s+3)^2+16}\right] \quad =L^{-1}\left[\frac{20(s+3)+70}{(s+3)^2+16}\right]$$

$$=20\,L^{-1}\frac{s+3}{(s+3)^2+16}+70\;\;L^{-1}\frac{1}{(s+3)^2+16}$$

$$=20\,e^{-3t}\cos 4t+35\,\frac{e^{-3t}\sin 4t}{2}.$$

4. A voltage Ee^{-at} is applied at $t=0$ to a circuit of inductance L and resistance R. Show that the current at any time t is $\frac{E}{R-aL}\left[e^{-at}-e^{-\frac{Rt}{L}}\right]$.

Solution:

The circuit is an LR circuit. The differential equation with respect to the circuit is

$$L\frac{di}{dt}+Ri=E(t)$$

Here, L denotes the inductance, i denotes current at any time t and E(t) denotes the E.M.F.
It is given that $E(t) = E\,e^{-at}$. With this, we have
Thus, we have

$$L\frac{di}{dt}+Ri=Ee^{-at} \;\; or \;\; Li'(t)+R\,i(t)=Ee^{-at}$$

Taking Laplace transform on both sides to get

$$L[L\{i'(t)\}]+R[L\,\{i(t)\}]=E\;L\,\{e^{-at}\}$$

$$L[s\,L\,\{i(t)\}-i(o)]+R[L\,\{i(t)\}]=E\,\frac{1}{s+a}$$

Since $i(0) = 0$, $\quad L\,\{i(t)\}[sL+R]=\frac{E}{s+a} \quad or$

$$L\,\{i(t)\}=\frac{E}{(s+a)(sL+R)}$$

Taking inverse transform $L^{-1}\{i(t)\}=L^{-1}\left\{\frac{E}{(s+a)(sL+R)}\right\}$

$$=\frac{E}{R-aL}\left[L^{-1}\left\{\frac{1}{s+a}\right\}-L\,L^{-1}\left\{\frac{1}{sL+R}\right\}\right]$$

Thus $i(t)=\frac{E}{R-aL}\left[e^{-at}-e^{-\frac{Rt}{L}}\right]$.

5. Mass spring damper system can be modeled using Newton's and Hooke's law. The differential equation representing this system is $\frac{d^2x}{dt^2}+6\frac{dx}{dt}+25x=4\sin\omega t$ with initial conditions $x(0)=x'(0)=0$.

Solution:

Given equation may be rewritten as

$$x''(t)+6x'(t)+25x(t)=4\sin\omega t$$

Here the initial conditions are $x(0)=20, x'(0)=10$.

Taking the Laplace transform of the equation, we get

$$L\{x(t)\}[s^2+6s+25]=\frac{4\omega}{s^2+\omega^2} \quad \text{or} L\{x(t)\}=\frac{4\omega}{(s^2+\omega^2)(s^2+6s+25)}$$

On resolving in to partial fractions (with $\omega=2$) leads to

$$x(t)=L^{-1}\left[\frac{8}{(s^2+4)(s^2+6s+25)}\right] \quad =L^{-1}\left[\frac{As+B}{s^2+4}+\frac{Cs+D}{s^2+6s+25}\right]$$

$$=$$

$$=\frac{4}{195}\,[7\sin 2t-4\cos 2t]+\frac{2}{195}\,[e^{-3t}(8\cos 4t-\sin 4t)].$$

Exercise:

Employ Laplace transform method to solve the following initial value problems.

(i) $y''+5y'+6y=e^{-2t},\ y(0)=y'(0)=1$

(ii) $y'''+2y''-y'-2y=0,\ y(0)=1,\ y'(0)=2=y''(0)$

(iii) $y''+2y'+5y=e^{-t}\sin t,\ y(0)=0,\ y'(0)=1.$

Answers:

(i) $3e^{-2t}-2e^{-3t}+te^{-2t}$ (ii) $\frac{1}{3}e^{-2t}+\frac{5}{3}e^{-t}-e^{t}$ (iii) $\frac{1}{3}e^{-t}(\sin t+\sin 2t)$

Video links:

https://www.youtube.com/watch?v=6MXMDrs6ZmA
https://www.youtube.com/watch?v=wnnnv4wt-Lw
https://www.youtube.com/watch?v=N-zd-T17uiE
https://www.youtube.com/watch?v=l7nzLD3t4Uc