

BAYESIAN NON-PARAMETRICS PRIORS WITH DENSITY ESTIMATION

Andi Wang Kaspar Märtens Ho Chung Leon Law

12th February 2016

Why non-parametrics, i.e. infinite-dimensional parameters?

Why non-parametrics, i.e. infinite-dimensional parameters?
Less assumptions involved.

Why non-parametrics, i.e. infinite-dimensional parameters?

Less assumptions involved.

Bayesian non-parametrics: prior on these objects.

$$G \sim \text{Prior}$$

is a random measure

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

$$G \sim \text{Prior}$$

is a random measure

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

Want a conjugate prior for G .

$$G \sim \text{Prior}$$

is a random measure

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

Want a conjugate prior for G .

One solution: Dirichlet process, parameters $\alpha_0 > 0$, G_0 , a probability measure.

Recall the stick-breaking construction:

Recall the stick-breaking construction:

$$\pi'_k \stackrel{iid}{\sim} \text{Beta}(1, \alpha_0) \text{ and } \phi_k \stackrel{iid}{\sim} G_0$$

Recall the stick-breaking construction:

$$\pi'_k \stackrel{iid}{\sim} \text{Beta}(1, \alpha_0) \text{ and } \phi_k \stackrel{iid}{\sim} G_0$$

$$\pi_k = \pi'_k \prod_{l=1}^{k-1} (1 - \pi'_l)$$


Recall the stick-breaking construction:

$$\pi'_k \stackrel{iid}{\sim} \text{Beta}(1, \alpha_0) \text{ and } \phi_k \stackrel{iid}{\sim} G_0$$

$$\pi_k = \pi'_k \prod_{l=1}^{k-1} (1 - \pi'_l)$$

$$G = \sum_{k=1}^{\infty} \pi_k \delta_{\phi_k}$$

DIRICHLET PROCESS



$$G \sim \text{DP}(\alpha_0, G_0)$$

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

$$G \sim \text{DP}(\alpha_0, G_0)$$

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

$$G | X_1, \dots, X_n \sim \text{DP}(\hat{\alpha}, \hat{G})$$

DIRICHLET PROCESS CONJUGACY

$$G \sim \text{DP}(\alpha_0, G_0)$$

$$X_1, \dots, X_n | G \stackrel{iid}{\sim} G$$

$$G | X_1, \dots, X_n \sim \text{DP}(\hat{\alpha}, \hat{G})$$

$$\hat{\alpha} = \alpha + n$$

$$\hat{G} = \hat{\alpha}^{-1} \left(\alpha G_0 + \sum_{i=1}^n \delta_{X_i} \right)$$

Idea: use the Dirichlet process as a prior on the parameters of a mixture model.

Idea: use the Dirichlet process as a prior on the parameters of a mixture model.

$$G \sim \text{DP}(\alpha_0, G_0)$$

$$\phi_i \mid G \sim G$$

$$x_i \mid \phi_i \sim F(\phi_i)$$

where F is a parametric distribution with density $p(x|\phi)$.

Recalling

$$G = \sum_{k=1}^{\infty} \pi_k \delta_{\phi_k}$$

the Dirichlet process mixture specifies

$$p(x) = \sum_{k=1}^{\infty} \pi_k p(x|\phi_k)$$



DIRICHLET PROCESS MIXTURE

Recalling


$$G = \sum_{k=1}^{\infty} \pi_k \delta_{\phi_k}$$

the Dirichlet process mixture specifies

$$p(x) = \sum_{k=1}^{\infty} \pi_k p(x|\phi_k)$$



DIRICHLET PROCESS MIXTURE: FITTED TO DATA



INFERENCE FOR THE NUMBER OF COMPONENTS

When data is generated from a finite mixture,

When data is generated from a finite mixture,

Dirichlet process mixture is a misspecified model

INFERENCE FOR THE NUMBER OF COMPONENTS


When data is generated from a finite mixture,

Dirichlet process mixture is a misspecified model
and it is inconsistent for the number of components

INFERENCE FOR THE NUMBER OF COMPONENTS

When data is generated from a finite mixture,

Dirichlet process mixture is a misspecified model
and it is inconsistent for the number of components



Generalisation of the Dirichlet process.

Generalisation of the Dirichlet process.


Able to place positive mass on continuous measures.

Generalisation of the Dirichlet process.


Able to place positive mass on continuous measures.

Popular in survival analysis, where censored data is common.

PÓLYA TREE DIAGRAM



PÓLYA TREE DIAGRAM



P on Ω have a Pólya tree distribution, with parameters $(\Pi = \{\Pi_m; m = 0, 1, \dots\}, \mathcal{A})$ is written $P \sim \text{PT}(\Pi, \mathcal{A})$

THE PARAMETERS

We must specify: a set of nested partitions Π and constants \mathcal{A} for the beta random variables.

We must specify: a set of nested partitions Π and constants \mathcal{A} for the beta random variables.

Wish to center the tree around a specific G_0 , with cdf $F_0 : \mathbb{R} \rightarrow [0, 1]$.

THE PARAMETERS

We must specify: a set of nested partitions Π and constants \mathcal{A} for the beta random variables.

Wish to center the tree around a specific G_0 , with cdf $F_0 : \mathbb{R} \rightarrow [0, 1]$.

Take the beta parameters \mathcal{A} to be constant on each level, with value $c\rho(m)$ on level m .

THE PARAMETERS

We must specify: a set of nested partitions Π and constants \mathcal{A} for the beta random variables.

Wish to center the tree around a specific G_0 , with cdf $F_0 : \mathbb{R} \rightarrow [0, 1]$.

Take the beta parameters \mathcal{A} to be constant on each level, with value $c\rho(m)$ on level m .

If $\rho(m)$ increases sufficiently fast, the distribution is continuous.

THE PARAMETERS

We must specify: a set of nested partitions Π and constants \mathcal{A} for the beta random variables.

Wish to center the tree around a specific G_0 , with cdf $F_0 : \mathbb{R} \rightarrow [0, 1]$.

Take the beta parameters \mathcal{A} to be constant on each level, with value $c \rho(m)$ on level m .

If $\rho(m)$ increases sufficiently fast, the distribution is continuous.

When $\rho(m) = 2^{-m}$ we obtain the Dirichlet process.

One choice of Π is *dyadic quantiles*, of the form

$$\left(F_0^{-1}\left(\frac{j-1}{2^m}\right), F_0^{-1}\left(\frac{j}{2^m}\right) \right]$$

One choice of Π is *dyadic quantiles*, of the form

$$\left(F_0^{-1}\left(\frac{j-1}{2^m}\right), F_0^{-1}\left(\frac{j}{2^m}\right) \right]$$

We showed this is valid if and only if F_0 is strictly increasing on \mathbb{R} .

One choice of Π is *dyadic quantiles*, of the form

$$\left(F_0^{-1}\left(\frac{j-1}{2^m}\right), F_0^{-1}\left(\frac{j}{2^m}\right) \right]$$

We showed this is valid if and only if F_0 is strictly increasing on \mathbb{R} .

With this choice, we have that for each B , $\mathbb{E}[P(B)] = G_0(B)$, where $P \sim \text{PT}(\Pi, \mathcal{A})$.

Let us assume the following:

$$x|P \sim P \qquad P \sim \text{PT}(\Pi, \mathcal{A}) \qquad (1)$$


Let us assume the following:

$$x|P \sim P \qquad P \sim \text{PT}(\Pi, \mathcal{A}) \qquad (1)$$

Then the posterior is given by a Pólya tree, $P|x \sim \text{PT}(\Pi, \mathcal{A}^*)$ with:

$$\alpha_\epsilon^* = \begin{cases} \alpha_\epsilon + 1, & \text{if } x \in B_\epsilon \\ \alpha_\epsilon, & \text{otherwise} \end{cases} \qquad (2)$$

FINITE PÓLYA TREE DIAGRAM



DENSITY ESTIMATION WITH PÓLYA TREE




Figure: Pólya trees with two base measures, $G_0 = \mathcal{N}(0, 3^2)$ (left) and $G_0 = \mathcal{N}(0, 10^2)$ (right) with $\alpha_\epsilon = \alpha m^2$, where $\alpha \sim \Gamma(1, 0.01)$.

We have explored Bayesian non-parametric approaches for density estimation:

1. Dirichlet process mixtures
2. Pólya trees