

# Capítulo 7

## Confidence intervals

### 7.1. Introduction

**Definition 7.1.1** (Confidence interval). *Let  $X_1, \dots, X_n, \dots$  be a sequence of independent identically distributed random variables with the same distribution as  $X \sim F \in \mathcal{F}_M$ . Let  $\theta = \Phi(F)$  be the parameter of interest. Consider two statistics*

$$A_n = A_n(X_1, \dots, X_n), B_n = B_n(X_1, \dots, X_n)$$

*We say that the interval  $(A_n, B_n)$  is a  $(1 - \alpha)$  confidence interval for  $\theta$  if*

$$\Pr(A_n \leq \theta \leq B_n) = 1 - \alpha$$

*for all  $F \in \mathcal{F}_M$ . We write*

$$\text{CI}_{1-\alpha}(\theta) \equiv (A_n, B_n).$$

**Example 7.1.1.**

---

Confidence interval for  $\mu$  when  $X \sim N(\mu, \sigma^2)$  and  $\sigma^2$  is known.

- Complete derivation of the confidence interval

$$\text{CI}_{1-\alpha}(\mu) \equiv \left( \bar{X}_n \mp z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right).$$

- Meaning of  $\text{CI}_{1-\alpha}(\mu) = (A_n, B_n)$ , a random interval.
- Meaning of  $\text{CI}_{1-\alpha}(\mu) = (a_n, b_n)$ , a realization of the random interval  $(A_n, B_n)$ .

---

□

## 7.2. Confidence intervals: A “general” method

Assume that there exist an estimator  $\hat{\theta}$  of the parameter  $\theta$  such that

$$\hat{\theta} \sim N(\theta, \text{s.e.}(\hat{\theta}))$$

with known standard error  $\text{s.e.}(\hat{\theta})$ . Then

$$\left( \hat{\theta} \mp z_{\alpha/2} \text{s.e.}(\hat{\theta}) \right)$$

is a  $(1 - \alpha)$  confidence interval for  $\theta$ .

### Example 7.2.1.

---

Confidence interval for the difference of means in two independent samples, normal case, known variances:

Assume that  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$ , and that  $\sigma_X^2$  and  $\sigma_Y^2$  are known. Let  $\theta = \mu_X - \mu_Y$  be the parameter of interest.

Let  $X_1, \dots, X_n$  and  $Y_1, \dots, Y_m$  be two independent simple random samples of  $X$  and  $Y$ , respectively. Then

$$\bar{X}_n \sim N\left(\mu_X, \frac{\sigma_X^2}{n}\right), \bar{Y}_m \sim N\left(\mu_Y, \frac{\sigma_Y^2}{m}\right)$$

and  $\bar{X}_n$  and  $\bar{Y}_m$  are independent. Therefore

$$\bar{X}_n - \bar{Y}_m \sim N\left(\mu_X - \mu_Y, \frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}\right).$$

Applying the general method,

$$\text{CI}_{1-\alpha}(\mu_X - \mu_Y) \equiv \left( \bar{X}_n - \bar{Y}_m \mp z_{\alpha/2} \sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}} \right).$$

---

□

### 7.3. Formulas for confidence intervals

- Confidence level  $1 - \alpha$ , level of significance  $\alpha$ .
- $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ , sample mean
- $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ , sample variance.
- $S_p^2 = \frac{(n_X-1)S_X^2 + (n_Y-1)S_Y^2}{n+m-2}$ , pooled sample variance of two samples.
- $z_a$  is the  $(1 - a)$ -quantile of  $N(0, 1)$
- $t_a$  is the  $(1 - a)$ -quantile of  $t$ -Student

Assumptions	Parameter	Statistic	Distribution	$(1 - \alpha)$ confidence interval
$X \sim N(\mu, \sigma^2)$ $\sigma^2$ known	Mean $\mu$	$\bar{X}$	$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$	$\bar{X} \pm z_{\alpha/2}\sigma/\sqrt{n}$
$X \sim N(\mu, \sigma^2)$ $\sigma^2$ unknown	Mean $\mu$	$\bar{X}$	$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t\text{-Student}, n - 1 \text{ df}$	$\bar{X} \pm t_{\alpha/2}S/\sqrt{n}$
$X \sim N(\mu_X, \sigma_X^2),$ $Y \sim N(\mu_Y, \sigma_Y^2),$ $\sigma_X^2, \sigma_Y^2$ known, two independent samples	Difference of means $\mu_X - \mu_Y$	$\bar{X} - \bar{Y}$	$\frac{\bar{X} - \bar{Y}}{\sqrt{\frac{\sigma_X^2}{n_X} + \frac{\sigma_Y^2}{n_Y}}} \sim N(0, 1)$	$\bar{X} - \bar{Y} \pm z_{\alpha/2}\sqrt{\frac{\sigma_X^2}{n_X} + \frac{\sigma_Y^2}{n_Y}}$
$X \sim N(\mu_X, \sigma_X^2),$ $Y \sim N(\mu_Y, \sigma_Y^2),$ $\sigma_X^2 = \sigma_Y^2$ unknown, two independent samples	Difference of means $\mu_X - \mu_Y$	$\bar{X} - \bar{Y}$	$\frac{\bar{X} - \bar{Y}}{S_p\sqrt{\frac{1}{n_X} + \frac{1}{n_Y}}} \sim t\text{-Student},$ $n_X + n_Y - 2 \text{ df}$	$\bar{X} - \bar{Y} \pm t_{\alpha/2}S_p\sqrt{\frac{1}{n_X} + \frac{1}{n_Y}}$
$X$ r.v. $\mathbb{E}(X) = \mu$ $\text{Var}(X) = \sigma^2$ unknown, large sample size $n$	Mean $\mu$	$\bar{X}$	$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim N(0, 1)$	$\bar{X} \pm z_{\alpha/2}S/\sqrt{n}$
$X, Y$ r.v. $\mathbb{E}(X) = \mu_X, \mathbb{E}(Y) = \mu_Y$ $V(X) = \sigma_X^2, V(Y) = \sigma_Y^2$ unknown, two independent samples of large sizes $n_X, n_Y$	Difference of means $\mu_X - \mu_Y$	$\bar{X} - \bar{Y}$	$\frac{\bar{X} - \bar{Y}}{\sqrt{\frac{\sigma_X^2}{n_X} + \frac{\sigma_Y^2}{n_Y}}} \sim N(0, 1)$	$\bar{X} - \bar{Y} \pm z_{\alpha/2}\sqrt{\frac{\sigma_X^2}{n_X} + \frac{\sigma_Y^2}{n_Y}}$
$X \sim B(n, p)$	Proportion $p$	$\hat{p} = \frac{X}{n}$	$\frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}} \sim N(0, 1)$	$\hat{p} \pm z_{\alpha/2}\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$
$X \sim B(n_X, p_X),$ $Y \sim B(n_Y, p_Y)$ large values of $n_X, n_Y$	Difference of proportions $p_X - p_Y$	$\hat{p}_X - \hat{p}_Y = \frac{X}{n_X} - \frac{Y}{n_Y}$	$\frac{\hat{p}_X - \hat{p}_Y}{\sqrt{\frac{\hat{p}_X(1-\hat{p}_X)}{n_X} + \frac{\hat{p}_Y(1-\hat{p}_Y)}{n_Y}}} \sim N(0, 1)$	$\hat{p}_X - \hat{p}_Y \pm z_{\alpha/2}\sqrt{\frac{\hat{p}_X(1-\hat{p}_X)}{n_X} + \frac{\hat{p}_Y(1-\hat{p}_Y)}{n_Y}}$
$X \sim N(\mu, \sigma^2)$	Variance $\sigma^2$	$S^2$	$\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$	$\left[ \frac{(n-1)S^2}{\chi_{\alpha/2}^2}, \frac{(n-1)S^2}{\chi_{1-\alpha/2}^2} \right]$
$X \sim N(\mu_X, \sigma_X^2),$ $Y \sim N(\mu_Y, \sigma_Y^2)$	Quotient of variances $\sigma_X^2/\sigma_Y^2$	$\frac{S_X^2}{S_Y^2}$	$\frac{S_X^2/\sigma_X^2}{S_Y^2/\sigma_Y^2} \sim F(n_X - 1, n_Y - 1)$	$\left[ \frac{S_X^2/S_Y^2}{F_{\alpha/2}(n_X-1, n_Y-1)}, \frac{S_X^2/S_Y^2}{F_{1-\alpha/2}(n_X-1, n_Y-1)} \right]$

Cuadro 7.1: Summary of statistics and confidence intervals.