REVIVING CIRCULANT PRECONDITIONERS FOR ADAPTIVE MESH REFINEMENT *

K. WALL †

In collaboration with: James Adler, Xiaozhe Hu, Misha Kilmer

Abstract. We present a preconditioner for solving fractional partial differential equations (PDEs) on an adaptive mesh. Adaptive refinement of the problem domain results in a stiffness matrix with Toeplitz blocks along the main diagonal, while the fractional PDE yields a dense stiffness matrix, where off-diagonal blocks are stored as low-rank approximations. Our preconditioner utilizes ideas from the circulant preconditioner of Chan and Strang [SIAM Journal on Scientific Computing, 1989], which takes advantage of the Toeplitz blocks on the diagonal and also accounts for the low-rank nature of the off-diagonal blocks. We demonstrate its effectiveness at accelerating convergence for our systems and emphasize its efficient application. This work presents theoretical results about the spectral clustering of the preconditioned system. In order to prove these results, special consideration is taken on how the low-rank blocks perturb the eigenvalues of the Toeplitz block-diagonal system. Numerical tests for various fractional orders are used to inspect any assumptions and validate our results.

Key words. Preconditioner, Adaptive Refinement, Toeplitz, Circulant, DST, DCT

1. Introduction.

problem statement, describe A block by block, visually? decay, dependence on alpha

There has long been interest in solving Toeplitz linear systems efficiently. A matrix A is called Toeplitz if $a_{ij} = a_{i-j}$, in other words, A has constant diagonals. Arbitrary $n \times n$ matrices have up to n^2 unique entries and are solved directly by traditional techniques in $\mathcal{O}(n^3)$ time. Since a Toeplitz matrix has just at most $2n-1$ unique entries, we may expect to be able to solve it in $\mathcal{O}(n^2)$ time. This is indeed the case via techniques such as Levinson's algorithm [7]. Even this improvement, however, is infeasible for large systems. Instead we turn to iterative Krylov and multigrid methods. For these methods we can still take advantage of Toeplitz structure by using circulant preconditioners.

A circulant matrix is a Toeplitz matrix, that additionally has the "wrap-around" property where the last entry each row is the first entry of the subsequent row.

$$L = \begin{pmatrix} \ell_0 & \ell_{-1} & \cdots & \ell_{-(n-1)} \\ \ell_1 & \ell_0 & \cdots & \ell_{-(n-2)} \\ \vdots & \vdots & \ddots & \vdots \\ \ell_{n-1} & \ell_{n-2} & \cdots & \ell_0 \end{pmatrix} \quad C = \begin{pmatrix} c_0 & c_1 & \cdots & c_{n-1} \\ c_{n-1} & c_0 & \cdots & c_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ c_1 & c_2 & \cdots & c_0 \end{pmatrix} \tag{1.1}$$

Equation 1: L is a Toeplitz matrix and C is circulant.

Toeplitz matrices commonly arise in PDE discretization, signal processing, and

*This work was funded by NSF.

†Tufts University, (kate.wall@tufts.edu, https://katejeanw.github.io/).

control theory. Often the Toeplitz matrices are also symmetric positive definite (SPD). Given an SPD Toeplitz system $Lx = b$, the idea introduced by Strang and Chan is to use certain circulant preconditioners C so that $C^{-1}Tx = C^{-1}b$ is solved in fewer iterations [4].

We leverage this idea to build a preconditioner for stiffness matrices generated from the adaptive finite element method (AFEM) for fractional PDEs. In this setting the problem is discretized on a nonuniform mesh, and the resulting stiffness matrix is dense. In the usual finite element method (FEM) setting an uniform mesh results in a Toeplitz stiffness matrix. In the adaptive setting, after an initial solve on a uniform grid, the error on each element is estimated and the elements with largest error are refined via bisection. It is often the case that neighboring elements are refined the same number of times. So although the mesh is not globally uniform, there are areas of local uniformity. To build an effective preconditioner, we will take advantage of these locally uniform areas and their corresponding Toeplitz blocks in the stiffness matrix.

TODO: citations for facts about FEM and Toeplitz matrices

Although dense, the stiffness matrix can be effectively stored as a hierarchical matrix ($\mathcal{H}$-matrix). Due to weaker interaction between elements that are further apart in the domain, off-diagonal blocks are well-suited for low rank approximation. (See [8] for more stiffness matrix details.) The low rank representation makes for fast computations, but complicates both the implementation of the preconditioner and the spectral clustering of the preconditioned system.

In this paper we investigate how to precondition such systems using circulant matrices. Our investigation is focused on $\mathcal{H}$-matrices as in [8], but the same methods could be used on any matrix with Toeplitz blocks on the diagonal. We prove the preconditioned system has eigenvalues clustered around 1 and demonstrate numerical results with superlinear convergence.

We emphasize that our unique contributions are:

- building circulant preconditioners for adaptive meshes
- proving the preconditioned system has eigenvalues clustered around 1
- something else? numerical results? dealing with low-rank blocks?

2. Background. To understand the need for our preconditioner, we must give a bit more detail about AFEM. We restrict our attention to problems on a one-dimensional domain, $[a, b]$ with the discretization $a = x_0, x_1, \ldots, x_n = b$. Often FEM is done on a uniform mesh, that is each element $[x_i, x_{i+1}]$ is size $x_{i+1} - x_i$ for all $0 \leq i \leq n - 1$.

If the mesh is not fine enough to give the desired accuracy, one approach is to increase the number of elements, n. While this approach preserves uniformity, it usually requires recomputing the stiffness matrix entirely. Alternatively, if the level of refinement gives sufficiently small error for some parts of the domain, we can leave those unchanged and only refine in areas of larger error. This approach allows us to take advantage of computations that have already been performed, but the mesh is no longer uniform.

In practice we find that adjacent elements are often refined to the same level, so a group of elements forms a locally uniform mesh. Since uniform meshes give rise to SPD Toeplitz systems, we can see that if we formed the stiffness matrix for just a locally uniform subdomain we would have an SPD Toeplitz matrix. So wherever there are adjacent elements of the same size we can find a corresponding Toeplitz block on the main diagonal of our stiffness matrix, A. The size of this block depends

on how many adjacent elements are the same size. We have also observed that the boundary of the domain almost always requires the greatest level of refinement. In general we have larger Toeplitz blocks from locally uniform subdomains in the middle of the matrix, and smaller blocks—or indeed 1×1 blocks—near the boundary.

To build an effective preconditioner and investigate its properites, we have to take full advantage of these SPD Toeplitz blocks. Toeplitz matrices have many unique properties that give rise to efficient algorithms (see for example [1]). For our purposes we focus on their connection to functions in the Wiener class. This will allow us to take our problem from matrix operator theory to function theory. Suppose we have a singly infinite, symmetric Toeplitz matrix

$$L = \begin{pmatrix} \ell_0 & \ell_1 & \ell_2 & \\ \ell_1 & \ell_0 & \ell_1 & \ddots \\ \ell_2 & \ell_1 & \ell_0 & \ddots \\ & \ddots & \ddots & \ddots \end{pmatrix}.$$

Assume $\sum_{k=-\infty}^{\infty} |\ell_k| < \infty$. Then the function $\ell(z) = \sum_{k=-\infty}^{\infty} \ell_k z^k$ is real, positive, and in the Wiener class for $|z| = 1$. It will be convenient to define the corresponding truncated function for a finite subsection of the infinite matrix:

Definition 2.1. *The $m \times m$ finite subsection of the singly infinite matrix T is denoted T_m and defined as*

$$L_m = \begin{pmatrix} \ell_0 & \ell_1 & \cdots & \ell_{m-1} \\ \ell_1 & \ell_0 & \cdots & \ell_{m-2} \\ \vdots & \vdots & \ddots & \vdots \\ \ell_{m-1} & \ell_{m-2} & \cdots & \ell_0 \end{pmatrix}.$$

Similarly this matrix induces a function from the truncated series, $\ell_m(z) = \sum_{k=-(m-1)}^{m-1} \ell_k z^k$.

Previous work on Toeplitz systems offers a few options for circulant preconditioners. Although their construction differs, the spectrum of the preconditioned systems are asymptotically the same [3]. We use the construction given by Bini and Benedetto [2]. Given a symmetric Toeplitz matrix L we build a Hankel correction, H where

$$L = \begin{pmatrix} \ell_0 & \ell_1 & \ell_2 & \cdots & \ell_{n-2} & \ell_{n-1} \\ \ell_1 & \ell_0 & \ell_1 & \cdots & \ell_{n-3} & \ell_{n-2} \\ \ell_2 & \ell_1 & \ell_0 & \cdots & \ell_{n-4} & \ell_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \ell_{n-2} & \ell_{n-3} & \ell_{n-4} & \cdots & \ell_0 & \ell_1 \\ \ell_{n-1} & \ell_{n-2} & \ell_{n-3} & \cdots & \ell_1 & \ell_0 \end{pmatrix} H = \begin{pmatrix} \ell_2 & \ell_3 & \ell_4 & \cdots & \ell_{n-1} & 0 & 0 \\ \ell_3 & \ell_4 & \ell_5 & \cdots & 0 & 0 & 0 \\ \ell_4 & \ell_5 & \ell_6 & \cdots & 0 & 0 & \ell_{n-1} \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \ell_5 & \ell_4 & \ell_3 \\ 0 & 0 & \ell_{n-1} & \cdots & \ell_4 & \ell_3 & \ell_2 \end{pmatrix}.$$

Then Bini and Benedetto's preconditioner is defined as $\boldsymbol{\tau} := L - H$. H is a Hankel matrix, that is it is constant on the antidiagonals. Both L and H are symmetric, so they can be completely represented by the at most n unique values: the diagonals of $L : \ell_0, \ell_1, \ldots, \ell_{n-1}$, and the antidiagonals of $H : \ell_2, \ell_3, \ldots, \ell_{n-1}, 0, 0$.

We summarize the important properties of $\boldsymbol{\tau}$ (see [2] for details):

- $\boldsymbol{\tau}$ is diagonalized by the type-I discrete sine transform (DST) matrix, S. We write $\boldsymbol{\tau} = S\Lambda S^{-1} = S\Lambda S$.
- $\boldsymbol{\tau}$ can be applied in $n \log n$ time .

- The k-th eigenvalue of $\boldsymbol{\tau}$ is proportional to the k-th entry of $S\ell_1$ where ℓ_1 is the first column of L. Specifically, define $c_k := \sqrt{\frac{n+1}{2}}\frac{1}{\sin(\frac{\pi k}{n+1})}$, then

$$\lambda_k(\boldsymbol{\tau}) = c_k[S\ell_1]_k \tag{2.1}$$

.

- Each eigenvalue of $\boldsymbol{\tau}$ can be written as the truncated function t_m evaluated somewhere on the unit circle, IE $\lambda_k(\boldsymbol{\tau}) = \ell_m(z_k)$ where $|z_k| = 1$.

These preconditioner can also be thought of as coming from the kernel of a displacement operator. This framework is useful for generating yet other circulant preconditioners, see [5].

3. Our Preconditioner. In this section we set forth the properties we require from a preconditioner, how we use $\boldsymbol{\tau}$ in building our preconditioner, and how to build and apply our preconditioner efficiently.

A good preconditioner for an iterative method must in general decrease the total number of iterations, without increasing the cost of a single iteration. We borrow Kailath's [5] criteria for preconditioners, though similar criteria has been established in literature (for example [2]).

1. Complexity of constructing applying $\boldsymbol{\tau}$ should be $\mathcal{O}(m \log m)$.
2. A linear system with $\boldsymbol{\tau}$ should be solved in $\mathcal{O}(m \log m)$ operations.
3. The spectrum of $\boldsymbol{\tau}^{-1}A$ should be clustered around 1

How tightly the eigenvalues cluster around 1 will determine the speed of convergence. In proving results about the clustering we will refer to the infinite matrix framework established previously. First we summarize results established previously about the spectral clustering of $\boldsymbol{\tau}$ applied to a single Toeplitz block L. This result will then be used as a lemma in proving the spectral clustering for our stiffness matrix from the adaptive mesh. We will show that for m large enough

We make this last point more precise:

Definition 3.1 (Eigenvalue Clustering). *For any $\varepsilon > 0$ we say the eigenvalues of a matrix $C^{-1}L_m$ are clustered around a real number ρ if there exists N_1 and N_2 such that for all $m > N_1$ there are at most N_2 eigenvalues of $C^{-1}L_m$ that do not lie within $[\rho - \varepsilon, \rho + \varepsilon]$.*

Assume our stiffness matrix has k Toeplitz blocks $L_1, L_2, \ldots, L_k$ of respective sizes $m_1, m_2, \ldots, m_k$. Assume additionally these are ordered as they appear along the main diagonal. Each block can be thought of as a singly infinite matrix with a corresponding generating function, $\ell_1(z), \ell_2(z), \ldots \ell_k(z)$. Assume that for all $1 \leq i \leq k, \ell_i(z) \sum_{j=-\infty}^{\infty} |t_j| < \infty$ and $\ell_i(z) > 0$ for z on the unit circle.

We can now construct a preconditioner, $\mathcal{T}$ for the adaptive system. To explicitly construct $\mathcal{T}$ we can calculate the Hankel correction H_i for every Toeplitz block, L_i. Then we define $\boldsymbol{\tau}_i = L_i - H_i$, resulting in k matrices $\boldsymbol{\tau}_1, \boldsymbol{\tau}_2, \ldots, \boldsymbol{\tau}_k$. Finally we assemble $\mathcal{T}$ as the block diagonal matrix with $\boldsymbol{\tau}_i$ as the i-th block.

$$\mathcal{T} = \begin{pmatrix} \boldsymbol{\tau}_1 & & & \\ & \boldsymbol{\tau}_2 & & \\ & & \ddots & \\ & & & \boldsymbol{\tau}_m \end{pmatrix}.$$

This explicit construction is not the most efficient, instead we can apply $\mathcal{T}^{-1}$ implicitly in $m \log m$ time where $m = \sum_{j=1}^{k} m_j$.

Let S_{m_i} denote the type-I DST matrix of size m_i. Using equation 2.1 to calculate the eigenvalues of $\boldsymbol{\tau}_i$, we can write $\boldsymbol{\tau}_i = S_{m_i}\Lambda_i S_{m_i}$.

$$\mathcal{T}^{-1} = \begin{pmatrix} S_{m_1} & & & \\ & S_{m_2} & & \\ & & \ddots & \\ & & & S_{m_k} \end{pmatrix} \begin{pmatrix} \Lambda_1^{-1} & & & \\ & \Lambda_2^{-1} & & \\ & & \ddots & \\ & & & \Lambda_m^{-1} \end{pmatrix} \begin{pmatrix} S_{m_1} & & & \\ & S_{m_2} & & \\ & & \ddots & \\ & & & S_{m_k} \end{pmatrix}.$$

Multiplication of the $m \times m$ matrices that consist of the DST blocks can be done via the fast Fourier transform in $m \log m$ time. The eigenvalue matrix is diagonal and can be inverted in m operations. The scaling done by this diagonal matrix could in the worst case cost $\mathcal{O}(m^2)$, but as we will see in the next section, we need not scale our entire stiffness matrix.

How to apply diagonal? FFT w toeplitz multiplication?, maybe just cite it away

This establishes criteria 1 and 2 for $\mathcal{T}$.

Bullet point properties of $\mathcal{T}$

4. Theoretical Results. In this section we will prove that the spectral clustering in criteria 3 holds for $\mathcal{T}$. Proofs for clustering of circulant preconditioned problems int his area tend to follow a similar structure. First it's shown that the preconditioned system having a spectrum clustered around 1 is equivalent to the clustering of a related system around 0. This related system is then split into the sum of a "low-rank" term and a "small-norm" term. The small-norm term has spectrum clustered around 0, forcing the clustering of the entire system around 0 with the number of outliers bounded by the rank of the low-rank term.

We present our version of the spectral clustering proof for the case of $\boldsymbol{\tau}$ acting on a single Toeplitz block. This result is then used to help us prove the clustering of our preconditioner on the adaptive grid, $\mathcal{T}^{-1}A$.

hyphenation of low rank and small norm

For convenence we state Weyl's inequality and one other lemma that will be useful to us later. Both these results are corollaries to the min-max theorem.

LEMMA 4.1 (Weyl's Inequality).
Let M and E be Hermitian $m \times m$ matrices. Then for $A := M + E$ we have

$$|\lambda_k(A) - \lambda_k(M)| \leq ||E||_2 \ , 1 \leq k \leq m.$$

That is the eigenvalues of A are at most $||E||_2$ away from the eigenvalues of M.

LEMMA 4.2. *Let $T \in \mathbb{R}^{n\times n}$ be an invertible matrix. Let $A \in \mathbb{R}^{n\times n}$, then*

$$\lambda_k(T^{-1}A) \leq \frac{\lambda_k(A)}{\lambda_{\min}(T)}$$

.

Proof. Using the min-max theorem,

$$\begin{aligned}\lambda_k(T^{-1}A) &= \min_{\dim V=k}\max_{x\in V}\left(\frac{(Ax,x)}{(Tx,x)}\right)\\ &\le \min_{\dim V=k}\left[\max_{x\in V}\left(\frac{(Ax,x)}{(x,x)}\right)\max_{x\in V}\left(\frac{(x,x)}{(Tx,x)}\right)\right]\\ &\le \left[\min_{\dim V=k}\max_{x\in V}\left(\frac{(Ax,x)}{(x,x)}\right)\right]\max_{x\in\mathbb{R}^n}\left(\frac{(x,x)}{(Tx,x)}\right)\\ &= \lambda_k(A)\max_{x\in\mathbb{R}^n}\left(\frac{(x,x)}{(Tx,x)}\right) \le \lambda_k(A)\frac{1}{\lambda_{\min}(T)}.\end{aligned}$$

∎

4.1. Spectral Clustering of a Toeplitz Block.

LEMMA 4.3 (Eigenvalue Clustering on a Toeplitz Block). *Let L be a singly infinite, symmetric, Toeplitz matrix with diagonals $\ell_0, \ell_1, \ldots$. Then L_m is its $m \times m$ truncation with diagonals $\ell_0, \ell_1, \ldots, \ell_{m-1}$. Assume the generating function $\ell(z)$ is in the Wiener class and positive on the unit circle. Let $\boldsymbol{\tau}$ be the corresponding preconditioner, then for m large enough the spectrum of $\boldsymbol{\tau}^{-1}L$ is clustered around 1.*

Proof. Since $\ell(z) > 0$ for $|z| = 1$, compactness of the unit circle implies that for some $\varepsilon > 0$, $\ell(z) > 2\varepsilon$ when $|z| = 1$. By the Wiener class assumption we can choose N such that $\sum_{j=N}^{\infty} \ell_j z^j \le \sum_{j=N}^{\infty} |\ell_j| < \varepsilon$. So for all $m \ge N$,

$$2\varepsilon < \ell(z) = \ell_m(z) + \sum_{j=m}^{\infty} \ell_j z^j < \ell_m(z) + \varepsilon.$$

Thus $\ell_m(z) > \varepsilon$ on the unit circle.

Recall from the construction of $\boldsymbol{\tau}$,

$$\boldsymbol{\tau} = L - H \tag{4.1}$$

$$L = \boldsymbol{\tau} + H \tag{4.2}$$

$$\boldsymbol{\tau}^{-1}L = I + \boldsymbol{\tau}^{-1}H \tag{4.3}$$

where H is the Hankel matrix with antidiagonals $\ell_2, \ell_3, \ldots, \ell_{m-1}, 0, 0$. Therefore it suffices to show that the eigenvalues of $\boldsymbol{\tau}^{-1}H$ are clustered around 0.

We now split H into a low-rank matrix H_{LR} that contains the antidiagonals $\ell_0, \ldots, \ell_N$ and a small-norm matrix H_{SN} such that $H = H_{LR} + H_{SN}$. Let $s := \operatorname{rank}(H_{LR}) << m$. The small-norm descriptor is justified since $H_{SN} = H - H_{LR}$ is a hermitian $m \times m$ matrix with at most two copies of $\ell_N, \ldots, \ell_{m-1}$ in each row/column. Thus $||H_{SN}||_2 = \sqrt{||H_{SN}||_1||H_{SN}||_\infty} = ||H_{SN}||_1 < 2\varepsilon$. Hence by Weyl's Inequality at least $m - s$ of the eigenvalues of H_{SN} are clustered within 2ε of zero.

Now by lemma 4.2,

$$\begin{aligned}\lambda_k(\boldsymbol{\tau}^{-1}H) &\le \lambda_k(H)\frac{1}{\lambda_{\min}(\boldsymbol{\tau})}\\ &= \lambda_k(H)\frac{1}{a_m(z_{\min})} \le \lambda_k(H)\frac{1}{\varepsilon}.\end{aligned}$$

Thus the clustering of the spectrum of H around 0 implies the same for $\boldsymbol{\tau}^{-1}H$. ∎

4.2. Spectral Clustering of the Full System.

THEOREM 4.4 (Full Matrix Clustering). *Let A be the stiffness matrix as described in section/equation TODO and $\mathcal{T}$ as described in section 3. Then the spectrum of $\mathcal{T}^{-1}A$ is clustered around 1.*

Proof. Recall that our stiffness matrix is dense with Toeplitz blocks on the diagonal. The ij entry of A reflects the strength of the interaction between elements i and j. Thus we observe off-diagonal decay, as elements that are physically separated in the domain have weaker interaction. The rate of decay depends on the fractional order of the PDE.

In order to prove spectral clustering of $\mathcal{T}^{-1}A$ we will handle the Toeplitz blocks as in the previous section. Blocks that are distant from the diagonal represent the weakest entries and have small norm. Blocks that are adjacent to the diagonal are not small norm; we will resort to splitting them into a low rank and a small norm term. To represent these classifications we will use A_D for the Toeplitz blocks, A_E for blocks adjacent to the diagonal, and A_F for the rest so that $A = A_D + A_E + A_F$. Let H denote the block diagonal Hankel Correction matrix, observing then that $A_D = \mathcal{T} + H$ we see that $A = \mathcal{T} + H + A_E + A_F$. Our approach is to split $H + A_E + A_F$ into a small-norm term A_{SN} and a low-rank term A_{LR}. Thus

$$\mathcal{T}^{-1}A = \mathcal{T}^{-1}(\mathcal{T} + A_{SN} + A_{LR}) = I + \mathcal{T}^{-1}A_{SN} + \mathcal{T}^{-1}A_{LR}. \tag{4.4}$$

Hence it suffices to show that the spectrum of $\mathcal{T}^{-1}A_{SN}$ is clustered around 0 and that $\mathcal{T}^{-1}A_{LR}$ contributes a bounded number of eigenvalue outliers.

Recall that blocks in A_E can be well-approximated as a low-rank matrix. To that end, we begin by examining a single block in A_E, call it E. Let E be a $m_E \times m_E$ matrix with approximation rank r_E. Then we can split E via its singular value decomposition (SVD):

$$E = \left(\sum_{i=1}^{r_E} \sigma_i^{(E)} \mathbf{u}_i^{(E)} \mathbf{v}_i^{(E)} * \right) + \left(\sum_{i=r_E+1}^{m_E} \sigma_i^{(E)} \mathbf{u}_i^{(E)} \mathbf{v}_i^{(E)} * \right).$$

In the above equation $\mathbf{u}$ and $\mathbf{v}$ are length m_E vectors. With a slight abuse of notation, we will embed these in length m vectors to express the block matrix A_E as the SVD of its blocks. Let off-diag represent the set of off-diagonal blocks in A_E. Define $r := \max_{E \in \text{off-diag}} r_E$ and $M := \max_{E \in \text{off-diag}} m_E$. We can now express A_E as the sum of the SVD of its blocks, and then regroup terms into a low-rank sum and a small-norm sum.

$$\begin{aligned} A_E &= \sum_{E \in \text{off-diag}} \left[\left(\sum_{i=1}^{r} \sigma_i^{(E)} \mathbf{u}_i^{(E)} \mathbf{v}_i^{(E)} * \right) + \left(\sum_{i=r+1}^{m_E} \sigma_i^{(E)} \mathbf{u}_i^{(E)} \mathbf{v}_i^{(E)} * \right) \right] \\ &= \left(\sum_{i=1}^{r} \sigma_i \mathbf{u}_i \mathbf{v}_i * \right) + \left(\sum_{i=r+1}^{M} \sigma_i \mathbf{u}_i \mathbf{v}_i * \right). \end{aligned}$$

As before, we have $\varepsilon > 0$ and choose N much smaller than m so that for every Toeplitz block the truncated function for the block $a_{m_i}(z) > \varepsilon$ on the unit circle and $\sum_{i=N}^{\infty} |a_i| < \varepsilon$.

Possible comment: since we don't really choose block size in practice the actual block size dictates the size of ε. Over all the blocks we can take the max ε for a uniform bound, but many will be clustered tighter than that. Supports argument that bigger Toeplitz blocks = better clustering

We additionally split H by separating the anti-diagonals with coefficients $a_0, \dots, a_N$ and the anti-diagonals comprising of $a_{N+1}, \dots a_m$. So we have two splittings,

$$A_E = \left(\sum_{i=1}^{r} \sigma_i \mathbf{u}_i \mathbf{v}_i * \right) + \left(\sum_{i=r+1}^{M} \sigma_i \mathbf{u}_i \mathbf{v}_i * \right)$$
$$H = H|_{a_2,\dots,a_N} + H|_{a_{N+1},\dots,a_m}.$$

The first term in each sum can be thought of as the low-rank equivalent from before and similarly the second term is the small-norm summand. Finally we can make the splitting $A = \mathcal{T} + A_{SN} + A_{LR}$ where

$$A_{SN} = H|_{a_{N+1},\dots,a_m} + \sum_{i=r+1}^{M} \sigma_i \mathbf{u}_i \mathbf{v}_i * + A_F$$
$$A_{LR} = H|_{a_2,\dots,a_N} + \sum_{i=1}^{r} \sigma_i \mathbf{u}_i \mathbf{v}_i * .$$

Hankel notation needs to be better, diagonals come from different blocks, what is m? etc

Now $s := \operatorname{rank}(\mathcal{T}^{-1} A_{LR}) \leq \operatorname{rank}(A_{LR}) \leq 2(N-1)+r$ bounds the number of outlier eigenvalues. It remains to show $||\mathcal{T}^{-1} A_{SN}||_2 \leq \varepsilon$. Define $\tilde{A_E} = \sum_{i=r_{A_E}+1}^{n_{A_E}} \sigma_i \mathbf{u}_i \mathbf{v}_i *$ and $H_{SN} = H|_{a_{N+1},\dots,a_m}$, so that $A_{SN} = H_{SN} + \tilde{A_E} + A_F$.

$$||\mathcal{T}^{-1} A_{SN}||_2 \leq ||\mathcal{T}^{-1} H_{SN}||_2 + ||\mathcal{T}^{-1} \tilde{A_E}||_2 + ||\mathcal{T}^{-1} A_F||_2$$

Since the block multiplication $\mathcal{T}^{-1} H_{SN}$ is each $\boldsymbol{\tau}$ acting on its corresponding Hankel correction we can bound it by 4.3. We can bound $||\mathcal{T}^{-1} A_F||_2$ with Weyl's inequality:

$$||\mathcal{T}^{-1} A_F||_2 \leq ||\mathcal{T}^{-1}||_2 ||A_F||_2 = \sigma_{\max}(\mathcal{T}^{-1}) \sigma_{\max}(A_F) = \frac{\sigma_{\max}(A_F)}{\lambda_{\min}(\mathcal{T})}.$$

Since A_F are small-norm this value is small. Finally we bound $||\mathcal{T}^{-1} \tilde{A_E}||_2$. Using lemma 4.2

$$\begin{aligned} \lambda_k(\mathcal{T}^{-1} \tilde{A_E}) &\leq \lambda_k(\tilde{A_E}) \frac{1}{\lambda_{\min}(\mathcal{T})} \\ &= \lambda_k(\tilde{A_E}) \min_{n \in n_k} \min_{1 \leq i \leq n} \frac{\sin(\frac{\pi i}{n+1})}{\sum_{j=1}^{n} t_j \sin(\frac{\pi i j}{n+1})}. \end{aligned}$$

Where to stop with $\lambda_{\min} \mathcal{T}$? can take it to generating function or bound on gen func or expression for eigs (like above) or leave as is

To finish the proof we need to relate the k-th eigenvalue of the preconditioned matrices to their 2-norm,

$$\begin{aligned} ||\mathcal{T}^{-1} A_{SN}||_2 &\leq ||\mathcal{T}^{-1} H_{SN}||_2 + ||\mathcal{T}^{-1} A||_2 + ||\mathcal{T}^{-1} A_F||_2 \\ &= \lambda_{\max}(\mathcal{T}^{-1} H_{SN}) + \lambda_{\max}(\mathcal{T}^{-1} \tilde{A_E}) + \lambda_{\max}(\mathcal{T}^{-1} A_F) \\ &\leq \frac{1}{\lambda_{\min}(\mathcal{T})} \left(\lambda_{\max}(H_{SN}) + \lambda_{\max}(\tilde{A_E}) + \lambda_{\max}(A_F) \right) \end{aligned}$$

□

SPSD justifying eig to sv to norm

Possible comments to add:
- technically lots of 1×1 blocks at boundaries, these get jacobi? inverse so are clustered around 1
- all problems come from boundaries
- Extend proof to different kinds of circulant preconditioner

5. Numerical Results.

5.1. Validating Assumptions. In order to prove the spectral clustering, we made two assumptions beyond the standard literature in the field. First, that the blocks adjacent to the diagonal are well-approximated by low-rank matrices. Second, that the other off-diagonal blocks are small-norm. We tested those assumptions at the ? level of refinement, for fractional order ?.

one matrix as an example? one for all alpha?

- numerical test confirming off-diag low rank
- Explanation and tests showing off-off-diag are small norm
- enough info to reproduce

5.2. Improved Convergence. In figure ?? we can see some example results of the preconditioner on the spectrum of A. These tests are for problem TODO where $\alpha = 1.7$. The three different matrix sizes are $n = 31, 71, 2795$ and reflect $0, 3$ and 11 levels of refinement respectively.

The left-most column represents a matrix that consists of a single Toeplitz block, so the preconditioner reduces to the case of $\boldsymbol{\tau}$. In this case the clustering of the spectrum around 1 is very strong. In the $n = 71$ case the biggest Toeplitz blocks are size 11, and there are two such blocks. The eigenvalue clustering around 1 is still observable, but the size of the cluster has grown. Finally for $n = 2795$, the preconditioner reduces the spectral radius by several orders of magnitude. The spectrum is concentrated around 1, but the distribution of eigenvalues between 0 and 2 is more uniform. For this matrix the biggest Toeplitz block is size 939 and there are two other blocks just bigger than 200. On the other hand, there are 245 1×1 blocks. For all cases, the strength of the clustering is explained by the amount of smaller Toeplitz blocks in A. We can see how the clustered spectrum translates to accelereted convergence. In figure ?? the same matrices are used for preconditioned conjugate gradient (PCG), comparing our preconditioner and a Jacobi preconditioner with a tolerance of 10^{-8}. For the two smaller matrices the preconditioner clearly accelerates the point of superlinear convergence [6]. For the final matrix the Jacobi-preconditioned system fails to converge within the maximum of 600 iterations, but with the $\mathcal{T}$ preconditioner converges in about 300 iterations.

6. Conclusion. Future work: how to build adaptive mesh to increase block size, other circulant preconditioners, tensor preconditioners, higher dimension domain, mixed precision

REFERENCES

[1] Sheldon Axler, *Paul Halmos and Toeplitz Operators*, Springer New York, New York, NY, 1991, pp. 257–263.
[2] D. Bini and F. Benedetto, *A new preconditioner for the parallel solution of positive definite toeplitz systems*, (1990), p. 220–223.
[3] Raymond H. Chan, *The spectrum of a family of circulant preconditioned toeplitz systems*, SIAM Journal on Numerical Analysis, 26 (1989), pp. 503–506.



Fig. 5.1: Eigenvalue clustering before and after preconditioning. Each column represent a different matrix size. The first row shows the spectrum before preconditioning, the second shows the spectrum at the same scale after preconditioning. The third row shows the preconditioned spectrum on a smaller interval.



Fig. 5.2: Comparing the residual vector norm across PCG iterations. The blue line uses a Jacobi preconditioner and the orange line uses our $\mathcal{T}$ preconditioner.

[4] Raymond H. Chan and Gilbert Strang, *Toeplitz equations by conjugate gradients with circulant preconditioner*, SIAM Journal on Scientific and Statistical Computing, 10 (1989), pp. 104–119.

[5] Thomas Kailath and Vadim Olshevsky, *Displacement structure approach to discrete-trigonometric-transform based preconditioners of g.strang type and of t.chan type*, SIAM Journal on Matrix Analysis and Applications, 26 (2006), pp. 706–734.

[6] V. Simoncini and D. B. Szyld, *On the superlinear convergence of minres*, (2013).

[7] Norbert Wiener, *The Wiener RMS (Root Mean Square) Error Criterion in Filter Design and Prediction*, 1964, pp. 129–148.

[8] Xuan Zhao, Xiaozhe Hu, Wei Cai, and George Em Karniadakis, *Adaptive finite element method for fractional differential equations using hierarchical matrices*, Computer Methods in Applied Mechanics and Engineering, 325 (2017), pp. 56–76.