

On Biadjoint Triangles

K

May 11, 2026

CONTENTS

1	Monads	1
1.1	Monads and their Algebras	1
1.2	Beck's Monadicity Theorem	4

1 MONADS

1.1 Monads and their Algebras

It would be shameful not to begin our exploration without a good background in monads, which are described in detail in this section. Monads, loosely speaking, answer the question of what it means for a category to possess some *algebraic* structure. This quality is made more explicit in the definition of *monadicity*, which cements the idea that a category is *monadic* over another category if it has some algebraic quality in relation to the latter category. This intuition becomes all the more clear once examples of monads and monadicity are studied. Much of this comes directly from Mac Lane, and will begin with a quick remark on composeability of functors and natural transformations.

For a category $\mathbb{X}$ and endofunctor $T : \mathbb{X} \rightarrow \mathbb{X}$, there are composite endofunctors $T^2 = T \circ T$ and $T^3 = T^2 \circ T$. If, additionally, there is a natural transformation $\mu : T^2 \rightarrow T$, then, for each object x in $\mathbb{X}$, the natural transformations $T\mu, \mu T : T^3 \rightarrow T^2$ have, respectively, components $(T\mu)_x = T\mu_x$, and $(\mu T)_x = \mu_{Tx}$.

Definition 1.1. [1] A **monad** $T = (T, \eta, \mu)$ on a category $\mathbb{X}$ is a triple in which $T : \mathbb{X} \rightarrow \mathbb{X}$ is an endofunctor on $\mathbb{X}$ and η, μ are two natural transformations

$$\eta : 1_{\mathbb{X}} \longrightarrow T, \quad \mu : T^2 \longrightarrow T \quad (1.1)$$

such that the following diagram commutes:

$$\begin{array}{ccccc} T^3 & \xrightarrow{\mu T} & T^2 & \xleftarrow{\eta T} & T \\ T\mu \downarrow & & \mu \downarrow & \swarrow & \downarrow T\eta \\ T^2 & \xrightarrow{\mu} & T & \xleftarrow{\mu} & T^2 \end{array} \quad (1.2)$$

Replace in Theorem 1.1 the endofunctor $T : \mathbb{X} \rightarrow \mathbb{X}$ with a set M , T^2 and T^3 with the cartesian product $M \times M$ and $M \times M \times M$ respectively, and lastly the natural transformations $\eta : 1_{\mathbb{X}} \rightarrow T$ with $\eta : 1 \rightarrow M$ and $\mu : T^2 \rightarrow T$ with the binary operation $\mu : M \times M \rightarrow M$. Then the diagram in Equation (1.2) expresses the usual axioms of a monoid. This is what motivates the naming of the natural transformations η as the **identity** and μ as the **multiplication** of a monad (T, η, μ) . This is also where we recover the famous “a monad is just a monoid in ...”

Definition 1.2. Let $T = (T, \eta, \mu)$ be a monad on $\mathbb{X}$. A **T -algebra** is a pair (x, ξ) in which x is an object in $\mathbb{X}$ and $\xi : Tx \rightarrow x$ is a morphism in $\mathbb{X}$ making the diagram

$$\begin{array}{ccccc} T^2x & \xrightarrow{\mu_x} & Tx & \xleftarrow{\eta_x} & x \\ T\xi \downarrow & & \xi \downarrow & \swarrow & \\ Tx & \xrightarrow{\xi} & x & & \end{array} \quad (1.3)$$

commute. A **morphism of T -algebras** $h : (x, \xi) \rightarrow (x', \xi')$ is a morphism $h : x \rightarrow x'$ in $\mathbb{X}$ making the diagram below commute:

$$\begin{array}{ccc} Tx & \xrightarrow{Th} & Tx' \\ \xi \downarrow & & \downarrow \xi' \\ x & \xrightarrow{h} & x' \end{array} \quad (1.4)$$

Theorem 1.3. Every adjunction $(F, U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$ determines a monad $(UF, \eta, U\epsilon F)$ on $\mathbb{X}$.

Proof. Firstly, $T = UF$ is an endofunctor on $\mathbb{X}$, η is a natural transformation $\eta : 1_{\mathbb{X}} \rightarrow UF = T$ and $\mu = U\epsilon F : UFUF = T^2 \rightarrow T = UF$ is a natural transformation. For an object x in $\mathbb{X}$, the commutativity of the diagram below must be checked:

$$\begin{array}{ccccc}
 UFUFUFx & \xrightarrow{U\epsilon_FUFx} & UFUFx & \xleftarrow{\eta_{UFx}} & UFx \\
 \downarrow UFU\epsilon_{Fx} & & \downarrow U\epsilon_{Fx} & \swarrow & \downarrow UF\eta_x \\
 UFUFx & \xrightarrow{U\epsilon_{Fx}} & UFx & \xleftarrow{U\epsilon_{Fx}} & UFUFx
 \end{array} \tag{1.5}$$

The left square commutes because of the naturality of $\epsilon : FU \rightarrow 1_{\mathbb{A}}$ applied to $\epsilon_{Fx} : FUFx \rightarrow Fx$, which is preserved under U . The right square commutes because of the triangle identities for the adjunction $(F, U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$. The identities are recalled:

$$\begin{array}{ccc}
 F & \xrightarrow{F\eta} & FUF \\
 & \searrow & \downarrow \epsilon_F \\
 & & F
 \end{array}
 \quad
 \begin{array}{ccc}
 UFU & \xleftarrow{\eta U} & U \\
 U\epsilon \downarrow & \swarrow & \\
 U & &
 \end{array} \tag{1.6}$$

This is shown by first considering the bottom triangle in Equation (1.5), which is just the x component of the left triangle of Equation (1.6) under U , and so it commutes. The top triangle in Equation (1.5) is the Fx component of the right triangle in Equation (1.5). This shows that $(UF, \eta, U\epsilon F)$ is a monad on $\mathbb{X}$. This monad is called the **monad defined by the adjunction** $(F, U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$. $\square$

Then, naturally, the question arises of whether every monad induces an adjunction. The answer is yes, and is discussed in the next theorem.

Theorem 1.4. Every monad $T = (T, \eta, \mu)$ on $\mathbb{X}$ determines an adjunction $(F^T, U^T, \eta^T, \epsilon^T) : \mathbb{X} \rightarrow \mathbb{X}^T$

To each monad T we associate structures call T -algebras defined as follows.

In this section, we are interested in a relationship between monads and adjunctions: that being every monad produces an adjunction and *vice versa*.

Theorem 1.5. Every monad $T = (T, \eta, \mu)$ on a category $\mathbb{X}$ determines an adjunction $(F^T, U^T, \eta^T, \epsilon^T) : \mathbb{X} \rightarrow \mathbb{X}^T$ from $\mathbb{X}$ to the category of its T -algebras, $\mathbb{X}^T$. The monad defined by this adjunction is the same monad $T = (T, \eta, \epsilon)$.

Proof. We define

1. the right adjoint $U^T : \mathbb{X}^T \rightarrow \mathbb{X}$ to be the forgetful functor, defined by $U^T(X, \xi) = X$;
2. the left adjoint $F^T : \mathbb{X} \rightarrow \mathbb{X}^T$ to be the functor defined by $F^T(X) = (T(X), \mu_X)$;
3. the unit $\eta^T : 1_{\mathbb{X}} \rightarrow U^T F^T$ to be the same η of the monad;
4. the counit $\epsilon^T : F^T U^T \rightarrow 1_{\mathbb{X}^T}$ to be defined by $\epsilon^T_{(T(X), \mu_X)} = \mu_X$.

$\square$

So we have that a monad determines an adjunction, which, in return determines the same monad again. The reverse is not always true: an adjunction determines a monad which does not always determine the same adjunction. When an adjunction induces a monad that induces the same adjunction back, we call that adjunction *monadic*. We get a sense of how monadic an adjunction is (or measure its *monadicity* with the following construction.

Theorem 1.6. *Let $(F, U, \eta, \epsilon) : \mathbb{X} \longrightarrow \mathbb{A}$ be an adjunction and $T = (UF, \eta, U\epsilon F)$ be the monad defined by the adjunction. Then there is a unique functor $K : \mathbb{A} \longrightarrow \mathbb{X}^T$ with $U^T K = U$ and $KF = F^T$.*

Proof. We define $K : \mathbb{A} \longrightarrow \mathbb{X}^T$ by $K(A) = (U(A), U(\epsilon_A))$, and on morphisms $K(f) = U(f)$. $\square$

There is one last detail that will make our diagram complete: the functor K as defined above has a left adjoint, the functor $L : \mathbb{X}^T \longrightarrow \mathbb{A}$.

Theorem 1.7. *There is an adjunction $(L, K, \bar{\eta}, \bar{\epsilon}) : \mathbb{X}^T \longrightarrow \mathbb{A}$.*

Proof. We define

1. the left adjoint $L : \mathbb{X}^T \longrightarrow \mathbb{A}$, where $L(X, \xi)$ is defined as the coequaliser of

$$FUF(X) \begin{array}{c} \xrightarrow{\epsilon_{F(X)}} \\ \xrightarrow{F(\xi)} \end{array} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$


2. the right adjoint $K : \mathbb{A} \longrightarrow \mathbb{X}^T$, defined by $K(A) = (U(A), U(\epsilon_A))$ on objects and $K(f) = U(f)$ on morphisms;
3. the unit $\bar{\eta} : 1_{\mathbb{X}^T} \longrightarrow KL$ defined by $\bar{\eta}_{(X, \xi)} = U(\pi_{(X, \xi)})\eta_X$;
4. the counit $\bar{\epsilon} : LK \longrightarrow 1_{\mathbb{A}}$ is defined for each $\bar{\epsilon}_A$ as the unique morphism making the diagram

$$\begin{array}{ccc} FUFU(A) & \begin{array}{c} \xrightarrow{\epsilon_{FU(A)}} \\ \xrightarrow{FU(\epsilon_A)} \end{array} & FU(\bar{A}) \xrightarrow{\pi_{(U(A), U(\epsilon_A))}} L(U(A), U(\epsilon_A)) \\ & & \searrow \epsilon_A \quad \downarrow \bar{\epsilon}_A \\ & & A \end{array}$$

commute.

$\square$

Now our picture is complete, and looks like below, with the F - and U -triangles commuting.



Definition 1.8. Let $(F, U, \eta, \epsilon) : \mathbb{X} \longrightarrow \mathbb{A}$ be an adjunction and $T = (T, \eta, \mu)$ be its associated monad. Then

1. the functor $K : \mathbb{A} \longrightarrow \mathbb{X}^T$ as defined in the previous result is called the *comparison functor*;
2. the functor $U : \mathbb{A} \longrightarrow \mathbb{X}$ is said to be *monadic* when the comparison functor is an equivalence.

Monadic functors are very useful, because we can recover from them the adjunction that induces them. Moreover, monadicity means that, in some sense, a category is algebraic because it is equivalent to a category of algebras.

1.2 Beck's Monadicity Theorem

Theorem 1.9. *Beck's Monadicity Theorem* Let $(F, U, \eta, \epsilon) : \mathbb{X} \longrightarrow \mathbb{A}$ be an adjunction, and $T = (T, \eta, \mu)$ be its associated monad. The following conditions are equivalent:

1. the functor $U : \mathbb{A} \longrightarrow \mathbb{X}$ is monadic;
2. the functor U preserves coequaliser diagrams of the form:

$$FUF(X) \xrightarrow[F(\xi)]{\epsilon_{F(X)}} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$

for every T -algebra (X, ξ) , and for every object A in $\mathbb{A}$, the diagram

$$FUFU(A) \xrightarrow[FU(\epsilon_A)]{\epsilon_{FU(A)}} FU(A) \xrightarrow{\epsilon_A} A$$

is a coequaliser;

3. U preserves coequaliser diagrams of the form:

$$FUF(X) \xrightarrow[F(\xi)]{\epsilon_{F(X)}} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$

for every T -algebra (X, ξ) , and U reflects isos;

4. every diagram $A \xrightarrow[g]{f} B$ in $\mathbb{A}$ - where $U(A) \xrightarrow[U(g)]{U(f)} U(B)$ has an absolute coequaliser - has a coequaliser preserved by U , and U reflects isos.

Proof. Since it has been extensively covered that $(1) \iff (2) \iff (3)$, we will only prove $(1) \implies (4)$. Let $\mathcal{D}$ be a diagram in $\mathbb{A}$ such that $U(\mathcal{D}) = X \xrightarrow[v]{u} X'$ with in $\mathbb{X}$. Suppose $U(\mathcal{D})$ has an absolute coequaliser q .

$$\begin{array}{ccc} T(X) & \xrightarrow[T(v)]{T(u)} & T(X') \\ \xi \downarrow & & \downarrow \xi' \\ X & \xrightarrow[v]{u} & X' \xrightarrow{q} Q \end{array}$$

Since q is an absolute coequaliser of u and v in $\mathbb{X}$, then $T(q)$ is a coequaliser of $T(u)$ and $T(v)$ in $\mathbb{A}$. Since the u - and v -diagrams commute, then $q\xi'$ also coequalises $T(u)$ and $T(v)$. Then there is a unique morphism $k : T(Q) \rightarrow T(Q')$ such that $kT(q) = q\xi'$. Since this square commutes, then q is also a morphism in $\mathbb{X}^T$. Now we want to show that q is the coequaliser of u and v in $\mathbb{X}^T$. Let $q : (X', \xi') \rightarrow (Q', k')$ be a morphism of T -algebras such that $q'u = q'v$ in $\mathbb{X}^T$. Then $q'u = q'v$ in $\mathbb{X}$, and $T(q')T(u) = T(q')T(v)$ in $\mathbb{X}$, which, respectively, induce l and $T(l)$ in the diagram below, by universal property of the coequalisers.

$$\begin{array}{ccccc}
T(X) & \xrightleftharpoons[T(v)]{T(u)} & T(X') & \xrightarrow{T(q)} & T(Q) \\
\downarrow \xi & & \downarrow \xi' & & \downarrow k \\
X & \xrightleftharpoons[v]{u} & X' & \xrightarrow{q} & Q \\
& & & \searrow q' & \downarrow k' \\
& & & & Q'
\end{array}$$

$\xrightarrow{T(l)}$ (from $T(Q)$ to $T(Q')$)
 $\xrightarrow{l}$ (from Q to Q')

Now we just need to show that l is also a morphism of T -algebras, which is to show that $k'T(l) = lk$. Consider a segment of the diagram:

$$\begin{array}{ccccc}
T(X') & \xrightarrow{T(q)} & T(Q) & \xrightarrow{T(l)} & T(Q') \\
\xi' \downarrow & & \downarrow k & & \downarrow k' \\
X' & \xrightarrow{q} & Q & \xrightarrow{l} & Q'
\end{array}$$

The outer square commutes because $k'T(l)T(q) = k'T(q') = q'\xi' = lq\xi'$. The left square also commutes by assumption, and $T(q)$ is an epi since it is a coequaliser. Then by a pasting lemma, the right square commutes, too. Then l is indeed a T -algebra morphism, which shows that q is the coequaliser of u, v in $\mathbb{X}^T$. From (1), $\mathbb{X}^T \sim \mathbb{A}$, which means there is also a coequaliser of the diagram $\mathcal{D}$ in $\mathbb{A}$, which is trivially preserved by U . $\square$

Now we want a simpler way to check monadicity, which will become useful in the next example.

Theorem 1.10. *Crude Monadicity Theorem Let $U : \mathbb{A} \rightarrow \mathbb{X}$ be a functor. U is monadic if*

1. U has a left adjoint;
2. $\mathbb{A}$ has, and U preserves, reflexive coequalisers;
3. U reflects isomorphisms.

Proof. Let us only prove the second point. $\square$

REFERENCES

- [1] S. Eilenberg and S. Mac Lane. *Categories for the Working Mathematician*. Springer, 1971.