

On Biadjoint Triangles

KB

June 23, 2026

CONTENTS

1	Monads	1
1.1	Monads and their Algebras	1
1.2	Beck's Monadicity Theorem	5
1.3	The Adjoint Triangle Theorem	7

1 MONADS

1.1 Monads and their Algebras

It would be shameful not to begin our exploration without a good background in monads, which are described in detail in this section. Monads, loosely speaking, answer the question of what it means for a category to possess some *algebraic* structure. This quality is made more explicit in the definition of *monadicity*, which cements the idea that a category is *monadic* over another category if it has some algebraic quality in relation to the latter category. This intuition becomes all the more clear once examples of monads and monadicity are studied. Much of this comes directly from Mac Lane, and will begin with a quick remark on composeability of functors and natural transformations.

For a category $\mathbb{X}$ and endofunctor $T : \mathbb{X} \rightarrow \mathbb{X}$, there are composite endofunctors $T^2 = T \circ T$ and $T^3 = T^2 \circ T$. If, additionally, there is a natural transformation $\mu : T^2 \rightarrow T$, then, for each object X in $\mathbb{X}$, the natural transformations $T\mu, \mu T : T^3 \rightarrow T^2$ have, respectively, components $(T\mu)_X = T\mu_X$, and $(\mu T)_X = \mu_{T(X)}$.

Definition 1.1. [2] A **monad** $T = (T, \eta, \mu)$ on a category $\mathbb{X}$ is a triple in which $T : \mathbb{X} \rightarrow \mathbb{X}$ is an endofunctor on $\mathbb{X}$ and η, μ are two natural transformations

$$\text{id}_{\mathbb{X}} \xrightarrow{\eta} T \xleftarrow{\mu} T^2 \quad (1.1)$$

such that the following diagram commutes:

$$\begin{array}{ccccc} T^3 & \xrightarrow{\mu T} & T^2 & \xleftarrow{\eta T} & T \\ T\mu \downarrow & & \mu \downarrow & \swarrow & \downarrow T\eta \\ T^2 & \xrightarrow{\mu} & T & \xleftarrow{\mu} & T^2 \end{array} \quad (1.2)$$

Replace in Definition 1.1 the endofunctor $T : \mathbb{X} \rightarrow \mathbb{X}$ with a set M , T^2 and T^3 with the cartesian product $M \times M$ and $M \times M \times M$ respectively, the natural transformations $\eta : \text{id}_{\mathbb{X}} \rightarrow T$ with $\eta : \{\cdot\} \rightarrow M$ ($\{\cdot\}$ is any singleton set), and $\mu : T^2 \rightarrow T$ with the binary operation $\mu : M \times M \rightarrow M$. Then Diagram 1.2 expresses the usual axioms of a monoid. This is what motivates the naming of the natural transformations η as the **identity** and μ as the **multiplication** of a monad (T, η, μ) . This is also where we recover the famous “a monad is just a monoid in ...”

Definition 1.2. [2] Let $T = (T, \eta, \mu)$ be a monad on $\mathbb{X}$. A **T -algebra** is a pair (X, ξ) in which X is an object in $\mathbb{X}$ and $\xi : T(X) \rightarrow X$ is a morphism in $\mathbb{X}$ making the diagram

$$\begin{array}{ccccc} T^2(X) & \xrightarrow{\mu_X} & T(X) & \xleftarrow{\eta_X} & X \\ T\xi \downarrow & & \xi \downarrow & \swarrow & \\ T(X) & \xrightarrow{\xi} & X & & \end{array} \quad (1.3)$$

commute. A **morphism of T -algebras** $h : (X, \xi) \rightarrow (X', \xi')$ is a morphism $h : X \rightarrow X'$ in $\mathbb{X}$ making the diagram below commute:

$$\begin{array}{ccc} T(X) & \xrightarrow{T(h)} & T(X') \\ \xi \downarrow & & \downarrow \xi' \\ X & \xrightarrow{h} & X' \end{array} \quad (1.4)$$

Definition 1.3. For a monad $T = (T, \eta, \mu)$ on $\mathbb{X}$, the **Eilenberg-Moore category of T -algebras**, denoted by **$T\text{-Alg}$** , is the category of T -algebras with T -algebra morphisms.

Theorem 1.4. Every adjunction $(F \dashv U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$ determines a monad $(UF, \eta, U\epsilon F)$ on $\mathbb{X}$.

Proof. Firstly, $T = UF$ is an endofunctor on $\mathbb{X}$, η is a natural transformation $\eta : \text{id}_{\mathbb{X}} \rightarrow UF = T$ and $\mu = U\epsilon F : UFUF = T^2 \rightarrow T = UF$ is a natural transformation. For an object X in $\mathbb{X}$, the commutativity of the diagram below must be checked:

$$\begin{array}{ccccc}
 UFUFUF(X) & \xrightarrow{U\epsilon_{FUF(X)}} & UFUF(X) & \xleftarrow{\eta_{UF(X)}} & UF(X) \\
 \downarrow UFU\epsilon_{F(X)} & & \downarrow U\epsilon_{F(X)} & \swarrow & \downarrow UF\eta_X \\
 UFUF(X) & \xrightarrow{U\epsilon_{F(X)}} & UF(X) & \xleftarrow{U\epsilon_{F(X)}} & UFUF(X)
 \end{array} \tag{1.5}$$

The left square commutes because of the naturality of $\epsilon : FU \rightarrow \text{id}_{\mathbb{A}}$ applied to $\epsilon_{F(X)} : FUF(X) \rightarrow F(X)$, which is preserved under U . The right square commutes because of the triangle identities for the adjunction $(F \dashv U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$. The identities are recalled:

$$\begin{array}{ccc}
 F & \xrightarrow{F\eta} & FUF \\
 & \searrow & \downarrow \epsilon_F \\
 & & F
 \end{array}
 \quad
 \begin{array}{ccc}
 UFU & \xleftarrow{\eta U} & U \\
 \downarrow U\epsilon & & \swarrow \\
 U & &
 \end{array} \tag{1.6}$$

This is shown by first considering the bottom triangle in Diagram 1.5, which is just the X -component of the left triangle of Diagram 1.6 under U , and so it commutes. The top triangle in Diagram 1.5 is the $F(X)$ -component of the right triangle in Diagram 1.5. This shows that $(UF, \eta, U\epsilon F)$ is a monad on $\mathbb{X}$. This monad is called the **monad defined by the adjunction** $(F \dashv U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$. $\square$

Then, naturally, the question arises of whether every monad induces an adjunction. The answer is *yes*, and is discussed in the next theorem.

Theorem 1.5. Every monad $T = (T, \eta, \mu)$ on $\mathbb{X}$ determines an adjunction $(F^T \dashv U^T, \eta^T, \epsilon^T) : \mathbb{X} \rightarrow T\text{-Alg}$. The monad defined by this adjunction is the same monad $T = (T, \eta, \epsilon)$.

Proof. Define

1. the right adjoint $U^T : T\text{-Alg} \rightarrow \mathbb{X}$ to be the forgetful functor, $U^T(X, \xi) := X$;
2. the left adjoint $F^T : \mathbb{X} \rightarrow T\text{-Alg}$ to be the free functor, $F^T(X) := (T(X), \mu_X)$;
3. the unit $\eta^T : \text{id}_{\mathbb{X}} \rightarrow U^T F^T$ to be the same η of the monad;
4. the counit $\epsilon^T : F^T U^T \rightarrow \text{id}_{T\text{-Alg}}$ to be $\epsilon^T_{(T(X), \mu_X)} := \mu_X$.

Now it is checked that this is indeed the desired adjunction. Firstly, note that $(T(X), \mu_X)$ is a T -algebra. That is, the diagram from Equation (1.3) commutes:

$$\begin{array}{ccc}
 T^3(X) & \xrightarrow{\mu_{T(X)}} & T^2(X) \xleftarrow{\eta_{T(X)}} T(X) \\
 \downarrow T\mu_X & & \downarrow \mu_X \\
 T^2(X) & \xrightarrow{\mu_X} & T(X)
 \end{array}$$

But this is just the X -component of Diagram 1.2, which commutes because T is a monad.

Now to check naturality of η^T and ϵ^T . The naturality of η^T is just the naturality of η . This is shown by taking X an object in $\mathbb{X}$, then the X -component of η^T is the morphism in $\mathbb{X}$, $\eta_X^T : X \rightarrow U^T F^T(X) = U^T(T(X), \mu_X) = T(X)$, which is the same as η_X . Now for ϵ^T : take (X, ξ) an object in $T\text{-Alg}$. Then $F^T U^T(X, \xi) = F^T(X) = (T(X), \mu_X)$. Then the (X, ξ) -component of ϵ^T is $\epsilon_{(X, \xi)}^T : (T(X), \mu_X) \rightarrow (X, \xi)$. Taking a morphism of $h : (X, \xi) \rightarrow (X', \xi')$ in $T\text{-Alg}$ gives a naturality square:

$$\begin{array}{ccc} (X, \xi) & (T(X), \mu_X) & \xrightarrow{\mu_X} (X, \xi) \\ h \downarrow & F^T U^T(h) \downarrow & \downarrow h \\ (X', \xi') & (T(X)', \mu_{X'}) & \xrightarrow{\mu_{X'}} ((X')', \xi') \end{array} \quad (1.7)$$

For some morphism $f : X \rightarrow X'$ in $\mathbb{X}$, $F^T(f) : (T(X), \mu_X) \rightarrow (T(X)', \mu_{X'})$ must be a morphism of T -algebras, which means $F^T(f) : T(X) \rightarrow T(X')$ must be defined as $F^T(f) := T(f)$. Then $F^T U^T(h) = T(h)$. So $F^T U^T(h)$ in Diagram 1.7 becomes $T(h)$, and the square thus commutes by naturality of μ , and because h is a morphism of T -algebras. To check that there is an adjunction is to check the triangles in Diagram 1.6 commute. For (X, ξ) an object in $T\text{-Alg}$ and X an object in $\mathbb{X}$, the left triangle is the (X, ξ) -component of the left triangle of Diagram 1.7, and the right is the X -component of the right triangle of Diagram 1.7:

$$\begin{array}{ccc} T(X) & \xrightarrow{T\eta_X} T^2(X) & T(X) \xleftarrow{\eta_X} X \\ & \searrow \downarrow \mu_X & \downarrow \xi \swarrow \\ & T(X) & X \end{array}$$

The left and right triangles commute respectively from the commutativity of Diagram 1.2 and Diagram 1.3.

Lastly, notice that for an object X in $\mathbb{X}$, $U^T F^T(X) = T(X)$, $\eta^T = \eta$ and $U^T \epsilon^T F^T(X) = \mu_X$, and so the monad defined by this adjunction is $T = (T, \eta, \mu)$, as desired. $\square$

So we have that a monad determines an adjunction, which, in return determines the same monad again. The reverse is not always true: an adjunction determines a monad which does not always determine the same adjunction. When an adjunction induces a monad that induces the same adjunction back, we call that adjunction *monadic*. We get a sense of how monadic an adjunction is (or measure its *monadicity* with the following construction.

Theorem 1.6. *Let $(F \dashv U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$ be an adjunction and $T = (UF, \eta, U\epsilon F)$ be the monad defined by the adjunction. Then there is a unique functor $K : \mathbb{A} \rightarrow T\text{-Alg}$ with $U^T K = U$ and $KF = F^T$.*

Proof. Define $K : \mathbb{A} \rightarrow T\text{-Alg}$ by $K(A) := (U(A), U\epsilon_A)$, and on morphisms $K(f) := U(f)$. $\square$

There is another arrow that will make the triangle complete: the functor K as defined above has a left adjoint, the functor $L : T\text{-Alg} \rightarrow \mathbb{A}$.

Theorem 1.7. *There is an adjunction $(L \dashv K, \bar{\eta}, \bar{\epsilon}) : T\text{-Alg} \rightarrow \mathbb{A}$.*

Proof. We define

1. the left adjoint $L : T\text{-Alg} \rightarrow \mathbb{A}$, where $L(X, \xi)$ is defined as the coequaliser of

$$FUF(X) \xrightleftharpoons[F(\xi)]{\epsilon_{F(X)}} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$

2. the right adjoint $K : \mathbb{A} \rightarrow T\text{-Alg}$, defined by $K(A) := (U(A), U(\epsilon_A))$ on objects and $K(f) := U(f)$ on morphisms;
3. the unit $\bar{\eta} : \text{id}_{T\text{-Alg}} \rightarrow KL$ defined by $\bar{\eta}_{(X, \xi)} = U(\pi_{(X, \xi)})\eta_X$;
4. the counit $\bar{\epsilon} : LK \rightarrow \text{id}_{\mathbb{A}}$ is defined for each $\bar{\epsilon}_A$ as the unique morphism making the diagram

$$\begin{array}{ccc} FUFU(A) & \xrightleftharpoons[FU\epsilon_A]{\epsilon_{FU(A)}} & FU(A) \xrightarrow{\pi_{(U(A), U\epsilon_A)}} (LU(A), U\epsilon_A) \\ & & \searrow \epsilon_A \quad \downarrow \bar{\epsilon}_A \\ & & A \end{array}$$

commute.

□

Now our picture is complete, and looks like below, with the F - and U -triangles commuting.

$$\begin{array}{ccc} & & \mathbb{X} \\ & \nearrow U & \uparrow \\ \mathbb{A} & & \\ & \searrow F & \downarrow F^T \\ & & T\text{-Alg} \\ & \nwarrow K & \uparrow U^T \\ & \nearrow L & \end{array} \quad (1.8)$$

Definition 1.8. Let $(F \dashv U, \eta, \epsilon) : \mathbb{X} \rightarrow \mathbb{A}$ be an adjunction and $T = (T, \eta, \mu)$ be its associated monad. Then

1. the functor $K : \mathbb{A} \rightarrow T\text{-Alg}$ as defined in the previous result is called the **comparison functor**;
2. the functor $U : \mathbb{A} \rightarrow \mathbb{X}$ is said to be **monadic** when the comparison functor is an equivalence.

Monadic functors are very useful, because they recall the adjunction from which they came. Moreover, monadicity means that, in some sense, a category is algebraic because it is equivalent to a category of algebras.

As with any free-forgetful adjunction to a category of algebras, it is useful to know what the free algebras of the category are. In the case of the adjunction

Definition 1.9.

Theorem 1.10. *there is an adhybc*

1.2 Beck's Monadicity Theorem

Theorem 1.11 (Beck's Monadicity Theorem). [2] Let $(F, U, \eta, \epsilon) : \mathbb{X} \longrightarrow \mathbb{A}$ be an adjunction, and $T = (T, \eta, \mu)$ be its associated monad. The following conditions are equivalent:

1. the functor $U : \mathbb{A} \longrightarrow \mathbb{X}$ is monadic;
2. the functor U preserves coequaliser diagrams of the form:

$$FUF(X) \xrightarrow[F(\xi)]{\epsilon_{F(X)}} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$

for every T -algebra (X, ξ) , and for every object A in $\mathbb{A}$, the diagram

$$FUFU(A) \xrightarrow[FU(\epsilon_A)]{\epsilon_{FU(A)}} FU(A) \xrightarrow{\epsilon_A} A$$

is a coequaliser;

3. U preserves coequaliser diagrams of the form:

$$FUF(X) \xrightarrow[F(\xi)]{\epsilon_{F(X)}} F(X) \xrightarrow{\pi_{(X, \xi)}} L(X, \xi)$$

for every T -algebra (X, ξ) , and U reflects isos;

4. every diagram $A \xrightarrow[g]{f} B$ in $\mathbb{A}$ - where $U(A) \xrightarrow[U(g)]{U(f)} U(B)$ has an absolute coequaliser - has a coequaliser preserved by U , and U reflects isos.

Lemma 1.12. In the situation of Equation (1.8), the functor U^T reflects isomorphisms.

Proof. Working in the context of Diagram 1.8, let (X, ξ) and (X', ξ') be objects in $T\text{-Alg}$ such that $(X, \xi) \cong (X', \xi')$. There is an isomorphism $f : X \rightarrow X'$ in $\mathbb{X}$, $U^T(X, \xi) = X \cong X' = U^T(X', \xi')$, and so U^T reflects isos. $\square$

Proof of Theorem 1.11. Consider the situation of Theorem 1.11:

(1) $\Leftrightarrow$ (2): Using the labels of Diagram 1.8, note that since $U^T K = U$ and U^T reflects isos by Theorem 1.12, then U reflects isos iff so does K . Consider the diagram:

$$\begin{array}{ccc} UFUL(X) & \xrightarrow[UF(\xi)=T(\xi)]{U\epsilon_{F(X)}=\mu_X} & UF(X) \xrightarrow{\xi} X \\ & & \searrow \downarrow \eta'_X = U\pi_{(X, \xi)}\eta_X \\ & & UL(X, \xi) \end{array}$$

Let $\mathcal{D}$ be a diagram in $\mathbb{A}$ such that $U(\mathcal{D}) = X \rightrightarrows^{u,v} X'$ with in $\mathbb{X}$. Suppose $U(\mathcal{D})$ has an absolute coequaliser q .

$$\begin{array}{ccc} T(X) & \rightrightarrows^{T(u)}_{T(v)} & T(X') \\ \xi \downarrow & & \downarrow \xi' \\ X & \rightrightarrows^{u}_{v} & X' \end{array} \xrightarrow{q} Q$$

Since q is an absolute coequaliser of u and v in $\mathbb{X}$, then $T(q)$ is a coequaliser of $T(u)$ and $T(v)$ in $\mathbb{A}$. Since the u - and v -diagrams commute, then $q\xi'$ also coequalises $T(u)$ and $T(v)$. Then there is a unique morphism $k : T(Q) \rightarrow Q$ such that $kT(q) = q\xi'$. Since this square commutes, then q is also a morphism in $T\text{-Alg}$. Now we want to show that q is the coequaliser of u and v in $T\text{-Alg}$. Let $q : (X', \xi') \rightarrow (Q', k')$ be a morphism of T -algebras such that $q'u = q'v$ in $T\text{-Alg}$. Then $q'u = q'v$ in $\mathbb{X}$, and $T(q')T(u) = T(q')T(v)$ in $\mathbb{X}$, which, respectively, induce l and $T(l)$ in the diagram below, by universal property of the coequalisers.

$$\begin{array}{ccccc} T(X) & \rightrightarrows^{T(u)}_{T(v)} & T(X') & \xrightarrow{T(q)} & T(Q) \\ \downarrow \xi & & \downarrow \xi' & & \downarrow k \\ X & \rightrightarrows^{u}_{v} & X' & \xrightarrow{q} & Q \end{array} \begin{array}{ccc} & & T(Q') \\ & \searrow^{T(l)} & \downarrow k' \\ & & Q' \end{array}$$

Now we just need to show that l is also a morphism of T -algebras, which is to show that $k'T(l) = lk$. Consider a segment of the diagram:

$$\begin{array}{ccccc} T(X') & \xrightarrow{T(q)} & T(Q) & \xrightarrow{T(l)} & T(Q') \\ \xi' \downarrow & & \downarrow k & & \downarrow k' \\ X' & \xrightarrow{q} & Q & \xrightarrow{l} & Q' \end{array}$$

The outer square commutes because $k'T(l)T(q) = k'T(q') = q'\xi' = lq\xi'$. The left square also commutes by assumption, and $T(q)$ is an epi since it is a coequaliser. Then by a pasting lemma, the right square commutes, too. Then l is indeed a T -algebra morphism, which shows that q is the coequaliser of u, v in $T\text{-Alg}$. From (1), $T\text{-Alg} \sim \mathbb{A}$, which means there is also a coequaliser of the diagram $\mathcal{D}$ in $\mathbb{A}$, which is trivially preserved by U . $\square$

Now we want a simpler way to check monadicity, which will become useful in the next example.

Theorem 1.13. *Crude Monadicity Theorem Let $U : \mathbb{A} \rightarrow \mathbb{X}$ be a functor. U is monadic if*

1. U has a left adjoint;
2. $\mathbb{A}$ has, and U preserves, reflexive coequalisers;
3. U reflects isomorphisms.

Proof. Let us only prove the second point. $\square$

1.3 The Adjoint Triangle Theorem


Definition 1.14. [1] Given a functor $\mathbb{A} \xrightarrow{J} \mathbb{B}$, an **adjoint triangle for J** consists of a category $\mathbb{C}$, and adjunctions

$$(E \dashv R, \rho, \mu) : \mathbb{A} \rightarrow \mathbb{C}, \quad (L \dashv U, \eta, \varepsilon) : \mathbb{B} \rightarrow \mathbb{C} \quad (1.9)$$

$$\rho : \text{id}_{\mathbb{A}} \rightarrow RE, \quad \eta : \text{id}_{\mathbb{B}} \rightarrow UL \quad (1.10)$$

$$\mu : ER \rightarrow \text{id}_{\mathbb{C}}, \quad \varepsilon : LU \rightarrow \text{id}_{\mathbb{C}} \quad (1.11)$$

with $LJ = E$, displayed in the diagram:



. The equality $LJ = E$ is sometimes written as the natural transformation $LJ \xrightarrow{1} E$ and *vice versa*.


Lemma 1.15. [1] In the notation above, given an adjoint triangle for a functor J , there is a natural transformation $JR \xrightarrow{\Sigma} U$ which makes the following diagram commute:

$$\begin{array}{ccc} JRLJ & \xrightarrow{\Sigma LJ} & ULJ \\ \parallel & & \uparrow \eta J \\ JRE & \xleftarrow{J\rho} & J \end{array}$$

Proof. Define Σ as the composite:

$$\begin{array}{ccc} ULJR & \xlongequal{\quad} & UER \\ \uparrow \eta JR & & \downarrow U\mu \\ JR & \xrightarrow{\quad \Sigma \quad} & U \end{array}$$

Then we fill in the data of the diagram:



The commutativity of the diagram is given by the following observations:

- squares (b) and (e) commute by naturality of η ;
- (c) and (d) commute by naturality of 1 under the image of U ;
- (g) by naturality of μ under the image of U ;
- (f) by Diagram 1.6 under the image of U ;
- (a) by the definition of Σ : for some object A in $\mathbb{A}$, take $LJ(A)$ as an object in $\mathbb{C}$, then apply Σ .

The commutativity of the smaller diagrams show that the full diagram commutes. $\square$

Theorem 1.16 (Adjoint Triangle Theorem). *[1] Consider an adjoint triangle for a functor $J : \mathbb{A} \rightarrow \mathbb{B}$ using the notation of Theorem 1.14. Assume that η is the equaliser of the diagram*

$$\text{id}_{\mathbb{B}} \xrightarrow{\eta} UL \xrightleftharpoons[\eta UL]{UL \eta} ULUL$$

. If the equaliser of the diagram

$$RL \xrightleftharpoons[RL\Sigma L\rho RL]{RL\eta} RLUL$$

exists in the category $\mathbb{A}^{\mathbb{B}}$, then J has a right adjoint G . In this case, the equaliser gives the value of G .

Proof. The proof will follow a strategy similar to [3], therefore the theorem will first be translated into the form it takes in [3]. Firstly note that for some object Y in $\mathbb{B}$, the Y -component of $RL\Sigma L\rho RL$ is calculated as:

$$(RL\Sigma L\rho RL)_Y = RL(\Sigma_{L(Y)})\rho_{RL(Y)} = RL(U\mu \cdot \eta JR)_{L(Y)}\rho_{RL(Y)} = RL(U(\mu_{L(Y)})\eta_{JRL(Y)})\rho_{RL(Y)}$$

which is how the theorem is stated in [3]. For each pair of objects B in $\mathbb{B}$ and C in $\mathbb{C}$, let

$$\mathbb{C}(L(B), C) \xrightleftharpoons[\psi_{(B,C)}]{\chi_{(B,C)}} \mathbb{B}(B, U(C))$$

denote the natural isomorphism associated to the adjunction $L \dashv U$. The maps are given by

$$\chi_{(B,C)} = \mathbb{B}(\eta_B, U(C)) \circ U_{L(B), C} \text{ and } \psi_{(B,C)} = \mathbb{C}(L(B), \epsilon_C) \circ L_{B, U(C)}.$$

Next, we will prove there is a natural isomorphism $\mathbb{A}(-, G-) \cong \mathbb{B}(J-, -)$. Assuming the situation of the theorem. Diagram 1.16 is an equaliser iff for every pair of objects X, Y in $\mathbb{B}$, the diagram

$$\mathbb{B}(X, Y) \xrightarrow{\mathbb{B}(X, \eta_Y)} \mathbb{B}(X, UL(Y)) \xrightleftharpoons[\mathbb{B}(X, \eta_{UL(Y)})]{\mathbb{B}(X, UL(\eta_Y))} \mathbb{B}(X, ULUL(Y))$$

is an equaliser (by the fully-faithful *Yoneda embedding*). Now let $X = J(A)$ for some object A in $\mathbb{A}$ and consider the diagram:

$$\begin{array}{ccccc}
\mathbb{B}(J(A), Y) & \xrightarrow{\mathbb{B}(J(A), \eta_Y)} & \mathbb{B}(J(A), UL(Y)) & \xrightarrow[\mathbb{B}(J(A), \eta_{UL(Y)})]{\mathbb{B}(J(A), UL(\eta_Y))} & \mathbb{B}(J(A), ULUL(Y)) \\
\downarrow & & \uparrow \chi_{(J(A), L(Y))} \downarrow \psi_{(J(A), L(Y))} & & \downarrow \psi_{(J(A), ULUL(Y))} \\
\mathbb{B}(J(A), Y) & \xrightarrow{L_{J(A), Y}} & \mathbb{C}(LJ(A), L(Y)) & \xrightarrow[\gamma_Y]{\mathbb{C}(LJ(A), L(\eta_Y))} & \mathbb{C}(LJ(A), LUL(Y))
\end{array}$$

Where $\gamma_Y = \psi_{(J(A), ULUL(Y))} \circ \mathbb{B}(J(A), \eta_{UL(Y)}) \circ \chi_{(J(A), L(Y))}$. We verify the diagram commutes:

- The left rectangle commutes: some $f \in \mathbb{B}(J(A), Y)$ is sent to $\epsilon_{L(Y)} \cdot L(\eta_Y \cdot f) \in \mathbb{C}(LJ(A), L(Y))$ via $\psi_{(J(A), L(Y))} \circ \mathbb{B}(J(A), \eta_Y)$. But $\epsilon_{L(Y)} \cdot L(\eta_Y \cdot f) = \epsilon_{L(Y)} \cdot L(\eta_Y) \cdot L(f) = L(f)$ by Theorem 1.6. So the left rectangle commutes.
- The upper right rectangle commutes via naturality of $\psi_{(J(A), -)}$.
- The lower right rectangle commutes by construction.

Then we complete the commutative diagram:

$$\begin{array}{ccccc}
\mathbb{B}(J(A), Y) & \xrightarrow{\mathbb{B}(J(A), \eta_Y)} & \mathbb{B}(J(A), UL(Y)) & \xrightarrow[\mathbb{B}(J(A), \eta_{UL(Y)})]{\mathbb{B}(J(A), UL(\eta_Y))} & \mathbb{B}(J(A), ULUL(Y)) \\
\downarrow & & \uparrow \chi_{(J(A), L(Y))} \downarrow \psi_{(J(A), L(Y))} & & \downarrow \psi_{(J(A), ULUL(Y))} \\
\mathbb{B}(J(A), Y) & \xrightarrow{L_{J(A), Y}} & \mathbb{C}(LJ(A), L(Y)) & \xrightarrow[\gamma_Y]{\mathbb{C}(LJ(A), L(\eta_Y))} & \mathbb{C}(LJ(A), LUL(Y)) \\
\downarrow & & \downarrow & & \downarrow \\
\mathbb{B}(J(A), Y) & \xrightarrow{L_{J(A), Y}} & \mathbb{C}(E(A), L(Y)) & \xrightarrow[\gamma_Y]{\mathbb{C}(E(A), L(\eta_Y))} & \mathbb{C}(E(A), LUL(Y))
\end{array}$$

A morphism $g \in \mathbb{C}(E(A), L(Y))$ is sent to $\epsilon_{LUL(Y)} \cdot L(\eta_{UL(Y)}) \cdot LU(g) \cdot L(\eta_{J(A)}) = L(U(g) \cdot \eta_{J(A)})$ under γ_Y . This means that $\gamma_Y = L_{J(A), UL(Y)} \circ \chi_{(J(A), L(Y))}$. So the bottom line of the diagram is now as it appears in [3]. □

Corollary 1.17.

Because the Yoneda embedding preserves limits, the *Adjoint Triangle Theorem* may equivalently be formulated in terms of hom-sets.

REFERENCES

- [1] E. Dubuc. “Adjoint Triangles”. In: *Reports of the Midwest Category Seminar II*. Lecture Notes in Mathematics, vol 61., pp. 69-91. Springer Verlag, 1968.
- [2] S. Mac Lane. *Categories for the Working Mathematician*. Springer, 1971.
- [3] FL. Nunes. “On Biadjoint Triangles”. In: *Theory and Applications of Categories*, Vol. 31, No. 9, 2016, pp. 217–256. (2016).