





Distribution of positive IC50 titers over time by age
Age: 1
Age: 2
Age: 3
Age: 4
Age: 5
Age: 6
Age: 7
log2 IC50
0
5
10
2007
2008
2009
2010
2011
2012
Norovirus year
Strain
DH−2006
FH−2002
NO−2009
SY−2012



Distribution of positive IC50 titers by age and sample year
Year of sample: 2007
Year of sample: 2008
Year of sample: 2009
Year of sample: 2010
Year of sample: 2011
Year of sample: 2012
log2 IC50
0
5
10
1
2
3
4
5
6
7
Age (years)
Strain
DH−2006
FH−2002
NO−2009
SY−2012



Distribution of positive IC50 titers by age and birth year
Birth year: 2001
Birth year: 2002
Birth year: 2003
Birth year: 2004
Birth year: 2005
Birth year: 2006
Birth year: 2007
Birth year: 2008
Birth year: 2009
Birth year: 2010
Birth year: 2011
log2 IC50
Age (years)
Strain
DH−2006
FH−2002
NO−2009
SY−2012



Mean IC50 titers over time by age
Age: 1
Age: 2
Age: 3
Age: 4
Age: 5
Age: 6
Age: 7
log2 IC50
0
5
10
2007
2008
2009
2010
2011
2012
Norovirus year
Strain
DH−2006
FH−2002
NO−2009
SY−2012
Subset of measurements
All
Positives



Mean IC50 titers by age and sample year
Year of sample: 2007
Year of sample: 2008
Year of sample: 2009
Year of sample: 2010
Year of sample: 2011
Year of sample: 2012
log2 IC50
0
5
10
1
2
3
4
5
6
7
Age (years)
Strain
DH−2006
FH−2002
NO−2009
SY−2012
Subset of measurements
All
Positives


Mean IC50 titers by age and birth year






Seroprevalence with age by time
Year of sample: 2007
Year of sample: 2008
Year of sample: 2009
Year of sample: 2010
Year of sample: 2011
Year of sample: 2012
Proportion positive
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
1.0
1
2
3
4
5
6
7
Age (years)
Strain
DH−2006
FH−2002
NO−2009
SY−2012



Seroprevalence over time by age
Age: 1
Age: 2
Age: 3
Age: 4
Age: 5
Age: 6
Age: 7
Proportion positive
Norovirus year
Strain
DH−2006
FH−2002
NO−2009
SY−2012



Seroprevalence with age by birth year
Birth year: 2001
Birth year: 2002
Birth year: 2003
Birth year: 2004
Birth year: 2005
Birth year: 2006
Birth year: 2007
Birth year: 2008
Birth year: 2009
Birth year: 2010
Birth year: 2011
Proportion positive
Age (years)
Strain
DH−2006
FH−2002
NO−2009
SY−2012



3
2
1
0
cumu_infections
0
2
4
6
age



Year of birth: 2001
Year of birth: 2002
Year of birth: 2003
Year of birth: 2004
Year of birth: 2005
Year of birth: 2006
Year of birth: 2007
Year of birth: 2008
Year of birth: 2009
Year of birth: 2010
Year of birth: 2011
Cumulative number of infections
Age (years)
0
1
2
3
0
2
4
6



3
2
1
0
med
0
2
4
6
age



4
3
2
1
0
cumu_infections
0
2
4
6
age



A
1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0
Cumulative incidence
(proportion seroconverted)
0
1
2
3
4
5
6
7
Age (years)
Posterior draw
Posterior mean



A
Cumulative incidence
(total infections)
2.0
1.8
1.6
1.4
1.2
1.0
0.8
0.6
0.4
0.2
0.0
0
1
2
3
4
5
6
7
Age (years)
Posterior draw
Posterior mean



B
N=656
N=656
N=596
N=491
N=380
N=265
N=145
N=34
0.4
0.3
0.2
0.1
0.0
Incidence per year
(infections per capita)
0
1
2
3
4
5
6
7
Age (years)


A
1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0
Cumulative incidence
(proportion seroconverted)
0
1
2
3
4
5
6
7
Age (years)


B
N=114
N=206
N=253
N=256
N=239
N=191
N=124
N=34
0.4
0.3
0.2
0.1
0.0
Incidence per year
(infections per capita)
0
1
2
3
4
5
6
7
Age (years)


A
log IC50 against DH−2006
Age (years) at first virus isolation


C
Grimsby 1996
Farmington Hills 2002
Hunter 2004
Den Haag 2006
New Orleans 2009
Sydney 2012
Overall AR (95% CrI)
Overall AR (posterior median)
Attack rate (infections per capita)
Norovirus year


D
N=656
N=656
N=596
N=491
N=380
N=265
N=145
N=34
Incidence per year (infections per capita)
Age (years)