


pEGFP-1 Vector Information

GenBank Accession #: U55761

PT3026-5

Catalog #6086-1



Restriction Map and Multiple Cloning Site (MCS) of pEGFP-1. All restriction sites shown are unique. The *Not*I site follows the EGFP stop codon. The *Xba*I site (*) is methylated in the DNA provided by BD Biosciences. If you wish to digest the vector with this enzyme, you will need to transform the vector into a *dam*⁻ host and make fresh DNA.

Description

pEGFP-1 encodes a red-shifted variant of wild-type GFP (1–3) which has been optimized for brighter fluorescence and higher expression in mammalian cells. (Excitation maximum = 488 nm; emission maximum = 507 nm.) pEGFP-1 encodes the GFPmut1 variant (4) which contains the double-amino-acid substitution of Phe-64 to Leu and Ser-65 to Thr. The coding sequence of the EGFP gene contains more than 190 silent base changes which correspond to human codon-usage preferences (5). Sequences flanking EGFP have been converted to a Kozak consensus translation initiation site (6) to further increase the translation efficiency in eukaryotic cells. pEGFP-1 is a promoterless EGFP vector which can be used to monitor transcription from different promoters and promoter/enhancer combinations inserted into the MCS located upstream of the EGFP coding sequence. SV40 polyadenylation signals downstream of the EGFP gene direct proper processing of the 3' end of the EGFP mRNA. The vector backbone also contains an SV40 origin for replication in mammalian cells expressing the SV40 T antigen. A neomycin-resistance cassette (Neo^r) allows stably transfected eukaryotic cells to be selected using G418. The Neo^r cassette consists of the SV40 early promoter, the neomycin/kanamycin resistance gene of Tn5, and polyadenylation signals from the Herpes simplex virus thymidine kinase (HSV TK) gene. A bacterial promoter upstream of this cassette confers kanamycin resistance in *E. coli*. The pEGFP-1 backbone also provides a pUC origin of replication for propagation in *E. coli* and an f1 origin for single-stranded DNA production.



Clontech

United States/Canada
800.662.2566

Asia Pacific
+1.650.919.7300

Europe
+33.(0)1.3904.6880

Japan
+81.(0)77.543.6116

Clontech Laboratories, Inc.
A Takara Bio Company
1290 Terra Bella Ave.
Mountain View, CA 94043
Technical Support (US)
E-mail: tech@clontech.com
www.clontech.com

(PR29973; published 03 October 2002)

Use

EGFP can be used as an *in vivo* reporter of gene expression (2). Promoters should be cloned into the pEGFP-1 MCS upstream from the EGFP coding sequences. Without the addition of a functional promoter, this vector will not express EGFP. The recombinant EGFP vector can be transfected into mammalian cells using any standard transfection method. If required, stable transformants can be selected using G418 (7).

Location of features

- MCS: 12–89
- Enhanced green fluorescent protein (EGFP) gene
 - Kozak consensus translation initiation site: 90–100
 - Start codon (ATG): 97–99; Stop codon: 814–816
 - Insertion of Val at position 2: 100–102
 - GFPmut1 chromophore mutations (Phe-64 to Leu; Ser-65 to Thr): 289–294
 - His-231 to Leu mutation (A→T): 791
- SV40 early mRNA polyadenylation signal
 - Polyadenylation signals: 970–975 & 999–1004
 - mRNA 3' ends: 1008 & 1020
- f1 single-strand DNA origin: 1067–1522
(Packages noncoding strand of EGFP.)
- Ampicillin resistance (β -lactamase) promoter
 - 35 region: 1584–1589; –10 region: 1607–1612
 - Transcription start point: 1619
- SV40 origin of replication: 1863–1998
- SV40 early promoter
 - Enhancer (72-bp tandem repeats): 1694–1767 & 1768–1839
 - 21-bp repeats: 1843–1863, 1864–1884 & 1886–1906
 - Early promoter element: 1919–1925
 - Major transcription start points: 1915, 1953, 1959 & 1964
- Kanamycin/neomycin resistance gene
 - Neomycin phosphotransferase coding sequences:
 - Start codon (ATG): 2047–2049; stop codon: 2839–2841
 - G→A mutation to remove *Pst* I site: 2229
 - C→A (Arg→Ser) mutation to remove *Bss*H II site: 2575
- Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
 - Polyadenylation signals: 3077–3082 & 3090–3095
- pUC plasmid replication origin: 3426–4069

Primer Locations

- EGFP-N Sequencing Primer (#6479-1): 163–142
- EGFP-C Sequencing Primer (#6478-1): 750–771

Propagation in *E. coli*

- Suitable host strains: DH5 α , HB101 and other general purpose strains. Single-stranded DNA production requires a host containing an F plasmid such as JM109 or XL-1 Blue.
- Selectable marker: plasmid confers resistance to kanamycin (30 μ g/ml) to *E. coli* hosts.
- *E. coli* replication origin: pUC
- Copy number: \approx 500
- Plasmid incompatibility group: pMB1/ColE1

References

1. Prasher, D. C., *et al.* (1992) *Gene* **111**:229–233.
2. Chalfie, M., *et al.* (1994) *Science* **263**:802–805.
3. Inouye, S. & Tsuji, F. I. (1994) *FEBS Letters* **341**:277–280.
4. Cormack, B., *et al.* (1996) *Gene* (in press).
5. Haas, J., *et al.* (1996) *Curr. Biol.* **6**:315–324.
6. Kozak, M. (1987) *Nucleic Acids Res.* **15**:8125–8148.
7. Gorman, C. (1985). In *DNA cloning: A practical approach*, vol. II. Ed. D.M. Glover. (IRL Press, Oxford, U.K.), pp. 143–190.

Note: The attached sequence file has been compiled from information in the sequence databases, published literature, and other sources, together with partial sequences obtained by BD Biosciences Clontech. This vector has not been completely sequenced.

Notice to Purchaser

Use of BD Biosciences Clontech's Living Colors™ products containing DNA sequences coding for mutant *Aequorea victoria* green fluorescent protein (GFP) variants or proteins thereof requires a license from Amersham Biosciences under U.S. Patent Nos. 5,625,048; 5,777,079; 6,054,321 and other pending U.S. and foreign patent applications. In addition, certain BD Biosciences Clontech products are made under U.S. Patent No. 5,804,387 licensed from Stanford University.

Not-For-Profit research institutes or entities are granted an automatic license with the purchase of this product for use in non-commercial internal research purposes, the terms of which are disclosed in detail in the license that accompanies the shipment of this product. Such license specifically excludes the right to sell or otherwise transfer this product or its components to third parties.

For-Profit research institutes or entities must obtain a license from Amersham Biosciences. E-mail: gfp@amershambiosciences.com

Please contact BD Biosciences Clontech directly for any other assistance, including purchasing and technical support. All companies and institutions purchasing Living Colors™ products will be included in a quarterly report to Aurora Biosciences, as required by the BD Biosciences Clontech/Aurora Biosciences license agreement.

This product is intended to be used for research purposes only. It is not to be used for drug or diagnostic purposes nor is it intended for human use. BD Biosciences Clontech products may not be resold, modified for resale, or used to manufacture commercial products without written approval of BD Biosciences Clontech.

© 2002, Becton, Dickinson and Company