

# Sensory processing during mate choice impacts the dynamics of sexual selection



Kathryn Bullough

kb574@exeter.ac.uk

Dr Bram Kuijper


Dr Laura Kelley

@KathrynBullough

# 1 ) Background


magnitude of sexual signals often influences mating preferences, where females can compare male stimuli in a non-linear way. An example of this is Weber's Law, where discrimination between signals is based upon their proportional difference rather than

Discriminating clusters is easier the larger the proportional difference between stimuli



their absolute difference. This means that when comparing males, a fixed absolute difference is easier to perceive at smaller magnitudes than at larger magnitudes.

## 2 How could Weber's Law impact sexual selection?




## 3 Agent-based models


Simulations of Fisherian sexual selection were run for populations of 5000 individuals over 150000 generations.

Individuals were diploid organisms, with loci for male trait values (t) and female preference values (p). Mate choice was run using either open-ended or Weber preferences.


#### Fisherian open-ended preferences




## Fisherian Weber preferences





## 4 Unidimensional traits





### 5 Multidimensional traits





# 6 Conclusions

- Using Weber-based preferences in models of sexual selection limits the exaggeration of both male traits and female preferences
- When multiple traits and preferences are allowed to evolve, Weber-based preferences cause them to diverge to two different equilibria