

BISHOP SCORE (PELVIC SCORE)

Pre-Induction Cervical Ripening and Labor Success Predictor

1. PATIENT IDENTIFICATION

Patient Name: _____

Unit / Ward / Bed: _____

Age / DOB: ____/____/____

Surgery / Diagnosis: _____

MRN / ID: _____

Date / Time: ____/____/____ : ____

CERVICAL EXAMINATION PARAMETERS

Clinical Parameter	Options & Scoring Criteria	Score
1. Cervical Dilation Internal os diameter in cm	☐ Closed (0 cm) (0) ☐ 1 - 2 cm (1) ☐ 3 - 4 cm (2) ☐ ≥ 5 cm (3)	[]
2. Cervical Effacement Length of cervix shortened (%)	☐ 0 - 30% (thick) (0) ☐ 40 - 50% (1) ☐ 60 - 70% (2) ☐ ≥ 80% (paper thin) (3)	[]
3. Fetal Station Position relative to ischial spines (cm)	☐ -3 (floating/high) (0) ☐ -2 (1) ☐ -1 / 0 (at spines) (2) ☐ +1 / +2 (engaged/low) (3)	[]
4. Cervical Consistency Palpatory tissue firmness	☐ Firm (like tip of nose) (0) ☐ Medium (like chin) (1) ☐ Soft (like lips) (2)	[]
5. Cervical Position Direction of os relative to pelvic axis	☐ Posterior (0) ☐ Mid-position (1) ☐ Anterior (2)	[]

TOTAL SCORE: [] POINTS (Maximum: 13 points)

Assessor (RN / License #): _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Unfavorable Cervix (≤ 5 points)

Ripening agent indicated (dinoprostone / misoprostol / Foley balloon).

Intermediate Cervix (6 - 7 points)

Equivocal ripeness. Reassess or low-dose mechanical ripening vs oxytocin.

Favorable Cervix (≥ 8 points)

High vaginal delivery probability similar to spontaneous labor. Proceed.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment