

CAPURRO METHOD FOR GESTATIONAL AGE

Gestational Maturity Assessment in Newborns (Somatic & Neurological)

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____/____/____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____/____/____ : ____

SOMATIC AND NEUROLOGICAL CRITERIA

Clinical Parameter	Options & Scoring Criteria	Score
1. Ear Form / Pinna Curvature Incurving of pinna edge	☐ Flat, shapeless, no curve (0) ☐ Incurving of part of superior pinna (8) ☐ Well-defined incurving of entire superior pinna (16) ☐ Fully curved upper pinna, firm cartilage (24)	[]
2. Mammary Gland Size Palpation of breast tissue nodule	☐ Non-palpable nodule (0) ☐ Palpable nodule < 5 mm diameter (5) ☐ Palpable nodule 5 - 10 mm diameter (10) ☐ Palpable nodule > 10 mm diameter (15)	[]
3. Nipple Formation Areola elevation and border stippling	☐ Barely visible, no areola (0) ☐ Nipple visible, areola flat and smooth (5) ☐ Areola raised, non-stippled border (10) ☐ Areola raised, stippled border with definite bud (15)	[]
4. Skin Texture Inspection and palpation of epidermis	☐ Very thin, gelatinous, smooth (0) ☐ Smooth, medium thickness, superficial peeling (10) ☐ Superficial cracks, pale, definite peeling hands/feet (15) ☐ Thick parchment-like, deep cracked skin (20)	[]
5. Plantar Creases Sole creases on foot extension	☐ No creases visible on sole (0) ☐ Faint red marks over anterior sole only (5) ☐ Definite creases over anterior half of sole (10) ☐ Deep indentations over more than anterior half (15) ☐ Deep creases over entire sole surface (20)	[]
6. Scarf Sign (if neuro+somatic) Elbow position across chest	☐ Elbow across to opposite axillary line (0) ☐ Elbow between opposite axilla and midline (6) ☐ Elbow reaches midline (12) ☐ Elbow does not reach midline (18)	[]

TOTAL SCORE: [] POINTS (Maximum: Formula: [204 + Score] ÷ 7)

Assessor (RN / License #): _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Post-Term Infant (≥ 42 weeks)

Placental insufficiency risk, meconium surveillance, monitor neonatal blood glucose.

Full-Term Infant (37 - 41 weeks)

Mature infant. Routine rooming-in, skin-to-skin contact, exclusive breastfeeding.

Moderate Preterm (32 - 36 weeks)

Thermoregulation support, feeding assessment, phototherapy readiness.

Extreme Preterm (< 32 weeks)

NICU admission, respiratory support (surfactant), radiant warmer, minimal handling.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment