

## ELPO SCALE

Risk Assessment of Perioperative Surgical Positioning Injury

## 1. PATIENT IDENTIFICATION

Patient Name: \_\_\_\_\_

Age / DOB: \_\_\_\_/\_\_\_\_/\_\_\_\_

MRN / ID: \_\_\_\_\_

Unit / Ward / Bed: \_\_\_\_\_

Surgery / Diagnosis: \_\_\_\_\_

Date / Time: \_\_\_\_/\_\_\_\_/\_\_\_\_ : \_\_\_\_

## SEVEN PERIOPERATIVE RISK FACTORS

| Clinical Parameter                                       | Options & Scoring Criteria                                                                                                                                                                                          | Score |
|----------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <b>1. Surgical Position</b><br>Main surgical posture     | <input type="checkbox"/> Supine (1)<br><input type="checkbox"/> Prone (2)<br><input type="checkbox"/> Lateral (3)<br><input type="checkbox"/> Lithotomy / Trendelenburg (4)                                         | [   ] |
| <b>2. Surgical Duration</b><br>Total operating room time | <input type="checkbox"/> ≤ 1 hour (1)<br><input type="checkbox"/> 1 - 2 hours (2)<br><input type="checkbox"/> 2 - 4 hours (3)<br><input type="checkbox"/> 4 - 6 hours (4)<br><input type="checkbox"/> > 6 hours (5) | [   ] |
| <b>3. Anesthesia Type</b><br>Anesthetic technique used   | <input type="checkbox"/> Local / MAC (1)<br><input type="checkbox"/> Regional (Spinal/Epidural) (2)<br><input type="checkbox"/> General anesthesia (3)<br><input type="checkbox"/> Combined general + regional (4)  | [   ] |
| <b>4. Support Surface</b><br>Mattress and padding        | <input type="checkbox"/> Viscoelastic / Gel pad (1)<br><input type="checkbox"/> Foam mattress (2)<br><input type="checkbox"/> Standard OR table pad only (3)                                                        | [   ] |
| <b>5. Limb Positioning</b><br>Angle of extremities       | <input type="checkbox"/> Anatomical (≤90°) (1)<br><input type="checkbox"/> Arms extended >90° (2)<br><input type="checkbox"/> Limbs flexed / elevated (3)                                                           | [   ] |
| <b>6. Comorbidities</b><br>Underlying medical conditions | <input type="checkbox"/> None (1)<br><input type="checkbox"/> Vascular disease / Diabetes (2)<br><input type="checkbox"/> Malnutrition / Obesity (3)<br><input type="checkbox"/> Multiple comorbidities (4)         | [   ] |
| <b>7. Patient Age</b><br>Age risk category               | <input type="checkbox"/> 18 - 39 years (1)<br><input type="checkbox"/> 40 - 59 years (2)<br><input type="checkbox"/> 60 - 79 years (3)<br><input type="checkbox"/> ≥ 80 years / Pediatric (4)                       | [   ] |

TOTAL SCORE: [       ] POINTS (Maximum: 35 points (7 to 35))

Assessor (RN / License #): \_\_\_\_\_

## CLINICAL INTERPRETATION &amp; RECOMMENDED NURSING ACTIONS

## Low Risk of Injury (≤ 19 points)

Standard perioperative care, padded armboards, neutral alignment.

## High Risk of Injury (≥ 20 points)

Pressure-redistributing gel pads, nerve protectors, frequent posture checks.

Registered Nurse (RN) Signature &amp; Professional Stamp

Date &amp; Time of Assessment