

FLACC BEHAVIORAL PAIN SCALE

Pain Assessment for Infants, Young Children, and Non-Verbal Patients

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____/____/____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____/____/____ : ____

FIVE BEHAVIORAL CATEGORIES

Clinical Parameter	Options & Scoring Criteria	Score
F - Face Facial expression and grimace	☐ No particular expression or smile (0) ☐ Occasional grimace or frown, withdrawn, uninterested (1) ☐ Frequent to constant quivering chin, clenched jaw (2)	[]
L - Legs Muscular tension in lower limbs	☐ Normal position or relaxed (0) ☐ Uneasy, restless, tense (1) ☐ Kicking, or legs drawn up (2)	[]
A - Activity Body movement and positioning	☐ Lying quietly, normal position, moves easily (0) ☐ Squirming, shifting back and forth, tense (1) ☐ Arched, rigid or jerking (2)	[]
C - Cry Vocal complaints and weeping	☐ No cry (awake or asleep) (0) ☐ Moans or whimpers; occasional complaint (1) ☐ Crying steadily, screams or sobs, frequent complaints (2)	[]
C - Consolability Response to comforting measures	☐ Content, relaxed (0) ☐ Reassured by occasional touching, hugging, talking to (1) ☐ Difficult to console or comfort (2)	[]

TOTAL SCORE: [] POINTS (Maximum: 10 points (0 to 10))**Assessor (RN / License #):** _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Relaxed and Comfortable (0 points)

No evidence of pain. Maintain routine pediatric comfort measures.

Mild Discomfort (1 - 3 points)

Non-pharmacological soothing (distraction, swaddling, parent presence).

Moderate Pain (4 - 6 points)

Mild analgesia (paracetamol / ibuprofen) plus comforting interventions.

Severe Discomfort / Pain (7 - 10 points)

Prompt analgesic intervention (IV opioid / physician notification). Reassess in 15-30m.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment