

HENDRICH II FALL RISK MODEL

Screening for Fall Risk in Acute and Inpatient Healthcare Settings

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____/____/____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____/____/____ : ____

EIGHT CLINICAL RISK FACTORS

Clinical Parameter	Options & Scoring Criteria	Score
1. Confusion / Disorientation Impaired cognitive awareness and judgment	☐ No (0) ☐ Yes (4)	[]
2. Symptomatic Depression Presence of active depressive signs	☐ No (0) ☐ Yes (2)	[]
3. Altered Elimination Urgency, frequency, or incontinence	☐ No (0) ☐ Yes (1)	[]
4. Dizziness / Vertigo Reported or documented lightheadedness	☐ No (0) ☐ Yes (1)	[]
5. Male Gender Epidemiological risk weighting	☐ Female (0) ☐ Male (1)	[]
6. Antiepileptic Medications Administered anticonvulsants	☐ No (0) ☐ Yes (2)	[]
7. Benzodiazepine Medications Administered sedatives/hypnotics	☐ No (0) ☐ Yes (1)	[]
8. Get-Up-and-Go Test Rising from seated position	☐ Able to rise in single movement without hands (0) ☐ Pushes up in one attempt (1) ☐ Multiple attempts but succeeds (3) ☐ Unable to rise without assistance (4)	[]

TOTAL SCORE: [] POINTS (Maximum: 16 points (Risk if $\geq$ **Assessor (RN / License #):** _____ 5))

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Low Fall Risk (0 - 4 points)

Standard universal fall precautions, call light within reach, safe footwear.

High Fall Risk ($\geq$ 5 points)

Fall prevention protocol: yellow armband, bed alarm, scheduled toileting, assist ambulation.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment