

ONTARIO FEMALE RISK ASSESSMENT SCALE (OFRAS)

Obstetric Fall Risk Assessment Tool in Antepartum and Postpartum Care

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____/____/____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____/____/____ : ____

SIX MATERNAL RISK DOMAINS

Clinical Parameter	Options & Scoring Criteria	Score
1. History of Falls Falls during pregnancy or previous year	☐ No falls during pregnancy or past 12 months (0) ☐ History of fall during current pregnancy (2)	[]
2. Mobility & Gait Impairment Balance and ambulatory stability	☐ Normal, steady independent gait (0) ☐ Uses assistive device or steadying on furniture (1) ☐ Impaired balance, weak, unsteady gait (2)	[]
3. Medications & Anesthesia Administered sedating drugs / blocks	☐ No sedating meds / No anesthesia (0) ☐ Epidural / Spinal anesthesia within past 12 hours (2) ☐ Magnesium sulfate, opioids, or sedatives within past 24 hours (2)	[]
4. Elimination Needs Urinary frequency and urgency	☐ Independent, normal continent urination (0) ☐ Frequency, urgency, or urinary retention needing assistance (1)	[]
5. Mental & Sensory Status Alertness and dizziness	☐ Alert, oriented, cooperative (0) ☐ Dizziness, orthostatic hypotension, or blurred vision (1) ☐ Sedated, confused, or non-compliant with safety instructions (2)	[]
6. Fluid Balance & Blood Loss Hemodynamic impact of delivery	☐ Normal postpartum / antepartum status (0) ☐ Postpartum hemorrhage / Symptomatic anemia (Hb < 8 g/dL) (1)	[]

TOTAL SCORE: [] POINTS (Maximum: 10 points (Risk if ≥ **Assessor (RN / License #):** 3))
_____**CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS****Low Obstetric Fall Risk (0 - 2 points)**

Standard postpartum safety precautions, call bell within reach, bed in lowest position.

High Obstetric Fall Risk (≥ 3 points)

Fall alert sign, assist on first 3 ambulations, mandatory assist for bathroom transfers.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment