

PERROCA PATIENT CLASSIFICATION SYSTEM

Assessment of Nursing Care Complexity and Workforce Sizing

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____ / ____ / ____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____ / ____ / ____ : ____

NINE NURSING CARE COMPLEXITY DOMAINS

Clinical Parameter	Options & Scoring Criteria	Score
1. Planning of Nursing Care Coordination and nursing process	☐ Routine ward protocol (1) (1) ☐ Moderate coordination required (2) (2) ☐ Individualized complex nursing plan (3) (3) ☐ Intensive multi-team coordination (4) (4)	[]
2. Investigation & Monitoring Vital signs and clinical surveillance	☐ Routine monitoring q8h (1) (1) ☐ q6h monitoring (2) (2) ☐ q4h monitoring / continuous ECG (3) (3) ☐ Continuous invasive monitoring / hemodynamics (4) (4)	[]
3. Body Hygiene Hygiene and self-care dependence	☐ Independent self-care (1) (1) ☐ Assisted shower / bath (2) (2) ☐ Bed bath performed by 1 nurse (3) (3) ☐ Bed bath requiring 2+ nurses (4) (4)	[]
4. Nutrition & Hydration Nutritional intake method	☐ Oral feeding independently (1) (1) ☐ Assisted oral feeding (2) (2) ☐ Enteral nutrition (tube/stoma) (3) (3) ☐ Total parenteral nutrition (TPN) (4) (4)	[]
5. Elimination Sphincter control and care	☐ Independent toileting (1) (1) ☐ Commode / urinal assistance (2) (2) ☐ Incontinent / diaper changes (3) (3) ☐ Catheter / ostomy irrigation (4) (4)	[]
6. Locomotion / Activity Mobility in bed and ward	☐ Walks independently (1) (1) ☐ Walks with aid / staff assist (2) (2) ☐ Chair / wheelchair transfers (3) (3) ☐ Bedbound / dependent transfers (4) (4)	[]
7. Skin & Mucous Integrity Wound care and dressings	☐ Intact skin / clean (1) (1) ☐ Stage 1-2 injury / dressing 1x daily (2) (2) ☐ Extensive dressing 2x daily (3) (3) ☐ Complex wound dressing ≥3x daily (4) (4)	[]
8. Therapeutics Medication complexity	☐ Oral medications only (1) (1) ☐ Intermittent IV / IM meds (2) (2) ☐ Continuous IV infusions (3) (3) ☐ Vasoactive drugs / chemotherapy (4) (4)	[]
9. Emotional Support Psychosocial interaction	☐ Calm, receptive (1) (1) ☐ Anxious, needs reassurance (2) (2) ☐ Agitated, uncooperative (3) (3) ☐ Severe crisis / conflictive (4) (4)	[]

TOTAL SCORE: [] POINTS (Maximum: 36 points (9 to 36)) **Assessor (RN / License #):** _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Minimal Care (9 - 13 points)

Autonomous patient in stable condition. Basic ward nursing staffing.

Intermediate Care (14 - 20 points)

Moderate dependency for hygiene and medication administration.

High Dependency Care (21 - 26 points)

High nursing workload, partial clinical instability.

Semi-Intensive Care (27 - 31 points)

Step-down care, high risk of acute decompensation.

Intensive Care (32 - 36 points)

Full critical care, multi-organ support, continuous 1:1 or 1:2 nursing.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment