

PEDIATRIC EARLY WARNING SCORE (PEWS)

Inpatient Pediatric Deterioration Screening and Rapid Response Trigger

1. PATIENT IDENTIFICATION

Patient Name: _____

Age / DOB: ____/____/____

MRN / ID: _____

Unit / Ward / Bed: _____

Surgery / Diagnosis: _____

Date / Time: ____/____/____ : ____

THREE CORE PHYSIOLOGICAL DOMAINS

Clinical Parameter	Options & Scoring Criteria	Score
1. Behavior Mental state and irritability	☐ Playing / Appropriate response (0) ☐ Sleeping / Irritable but consolable (1) ☐ Inconsolable / Lethargic / Weak (2) ☐ Reduced response to pain / Comatose (3)	[]
2. Cardiovascular Skin color and capillary refill	☐ Pink / Cap refill 1-2s (0) ☐ Pale / Cap refill 3s (1) ☐ Grey / Cap refill 4s / Tachycardia >20bpm above normal (2) ☐ Grey, mottled / Cap refill ≥5s / Tachycardia >30bpm or Bradycardia (3)	[]
3. Respiratory Ventilatory effort and rate	☐ Normal rate, no retractions (0) ☐ Tachypnea >10bpm above normal, accessory muscles, or 30%+ FiO2 (1) ☐ Tachypnea >20bpm above normal, retractions, or 40%+ FiO2 (2) ☐ Tachypnea >30bpm above normal, grunting, sternal recession, or 50%+ FiO2 (3)	[]
4. Special Modifiers Clinical treatments	☐ Quarter-hourly nebulizers: Yes [] (2 pts) (2) ☐ Persistent postoperative vomiting: Yes [] (2 pts) (2)	[]

TOTAL SCORE: [] POINTS (Maximum: 9 points (+ modifiers))

Assessor (RN / License #): _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Low Risk (Green) (0 - 2 points)

Routine 4-hourly observations. Continue planned pediatric ward care.

Medium Risk (Amber) (3 - 4 points)

Repeat vitals within 1 hour, bedside nurse in charge review, notify resident physician.

High Risk (Red) (≥ 5 points (or single 3))

Immediate Medical Emergency Team (MET) / Pediatric Critical Care response.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment