

RAMSAY SEDATION SCALE

Assessment of Sedation Depth and Responsiveness in Critical Care

1. PATIENT IDENTIFICATION

Patient Name: _____

Unit / Ward / Bed: _____

Age / DOB: ____/____/____

Surgery / Diagnosis: _____

MRN / ID: _____

Date / Time: ____/____/____ : ____

SIX CLINICAL SEDATION LEVELS

Clinical Parameter	Options & Scoring Criteria	Score
Level 1 (Anxious / Agitated) Inadequate sedation	☐ Patient anxious, agitated, or restless, or both (1)	[]
Level 2 (Cooperative / Calm) Target sedation	☐ Patient cooperative, oriented, and tranquil (2)	[]
Level 3 (Responds to Verbal Only) Light sedation	☐ Patient responds to verbal commands only (3)	[]
Level 4 (Brisk Response to Stimulus) Moderate sedation	☐ Brisk response to light glabellar tap or loud auditory stimulus (4)	[]
Level 5 (Sluggish Response) Deep sedation	☐ Sluggish response to light glabellar tap or loud auditory stimulus (5)	[]
Level 6 (Unresponsive) Extreme / Over-sedation	☐ No response to glabellar tap or loud auditory stimulus (6)	[]

TOTAL SCORE: [] POINTS (Maximum: 6 points (Levels 1 to 6))

Assessor (RN / License #): _____

CLINICAL INTERPRETATION & RECOMMENDED NURSING ACTIONS

Under-Sedated (Level 1)

Risk of unplanned extubation, agitation. Optimize analgesia/sedation.

Optimal Sedation (Levels 2 - 3)

Target for awake, non-intubated or ventilating patients. Calm, cooperative.

Moderate Sedation (Level 4)

Acceptable during acute synchronization with mechanical ventilation.

Deep Sedation (Levels 5 - 6)

Risk of prolonged ventilation, ICU weakness. Plan daily sedation interruption.

Registered Nurse (RN) Signature & Professional Stamp

Date & Time of Assessment