



# Is the Speed of Ocean Surface Currents Weibull Distributed?

Kaushal Gianchandani<sup>1</sup>, S. Campisi-Pinto<sup>2</sup> & Yosef Ashkenazy<sup>3</sup>

<sup>1</sup>The Institute of Earth Sciences, Hebrew University of Jerusalem; <sup>2,3</sup>Solar Energy and Environmental Physics, BIDR, Ben-Gurion University



## INTRODUCTION

- Ocean surface currents are crucial for:
  - regulating Earth's climate by **transporting heat**.
  - movement of marine life** across the world and regulating temperature of marine ecosystems.
  - determining the **course of navigation** while traversing the ocean.
- Underlying causes: wind stress, buoyancy fluctuations, tides, wave dynamics etc.
- Nontrivial** to understand piece-wise contribution of each.
- Very little known about the shape and behaviour of their statistical distribution in time. But WHY would we want to know anything about it?


## MOTIVATION

- Ocean surface currents are also a potential source of electric energy, possibly **better than winds**.
- Accurate information about these distributions can be utilised as a bench mark for improved ocean and climate modelling


## AGENDA

- Test the long standing hypothesis that the pdfs under consideration are **Weibull** distributed.
- If the hypothesis is rejected, test whether the alternate, the **Generalised Gamma** distribution is better suited for the pdfs.


## WEIBULL DISTRIBUTION



## GENREALISED GAMMA DISTRIBUTION




## METHOD



The analysis starts by assuming


- the sample at hand ( $x$ ) are Weibull (WBL) /Gen. Gamma (GG) distributed; therefore,
- calculate the MLEs  $\lambda, k, (\epsilon)$  of  $x$ .
- using the estimated parameters, generate a set of random WBL/GG surrogate series  $\{S_{WBL/GG}^i\}$
- calculate up to the 50th moment for each surrogate distribution; v) estimate the confidence interval for each moment of the surrogates.
- calculate up to the 50th moment of the original sample  $x$ ; finally;
- benchmark the values of the moments of  $x$  against the boundary levels of the corresponding confidence interval if the moment is within its confidence interval, then conclude that  $x$  is WBL/GG for that moment else it is not.

## RESULTS



Currents  $\mu_3$  within Weibull C.I. ~ 78%

Currents  $\mu_3$  within Gen. Gamma C.I. ~ 94%



## CONCLUSION

- Successful statistical analysis of the available time-series.
- Surface Current speeds are not ALWAYS Weibull distributed** over the global marine surface.
- Generalised Gamma distribution is a better approximation** for a larger number of the global marine surface time series in comparison to the Weibull.
- The **Generalised Gamma distribution gives a better estimate of  $\mu_3$**  (the third moment) for time-series of the Currents.
- This is relevant because for many applications, for example, the electric power available from these currents ~  $A\rho\langle U^3 \rangle/2$ .

## DISCUSSION

- A similar analysis was carried out by Dr. Salvatore Campisi-Pinto for time-series of winds over the global surface and the results were consistent.
- Other methods were also used to further validate the hypothesis for both the winds and the currents and the results were robust.

### Contact:

Kaushal Gianchandani  
 Institute of Earth Sciences, Hebrew University of Jerusalem  
 kaushal.g@mail.huji.ac.il  
 kaushalgianchandani.github.io  
 +91-8280118710