

# Quantum Entanglement and Thermodynamics in Spin Systems

## Project Report: Phase 1

**Kaushal Joshi**

Department of Physics  
Visvesvaraya National Institute of Technology, Nagpur

11, Dec. 2024



# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work

## Boltzmann Entropy:

In Statistical Mechanics, Boltzmann's Equation is given by:

$$S = k_B \ln(\omega)$$

# Entropy

## Boltzmann Entropy:

In Statistical Mechanics, Boltzmann's Equation is given by:

$$S = k_B \ln(\omega)$$

## Shannon Entropy:

The entropy of a random variable  $X$ , that represents a system is given by:

$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

# Entropy

## Boltzmann Entropy:


In Statistical Mechanics, Boltzmann's Equation is given by:

$$S = k_B \ln(\omega)$$

## Shannon Entropy:

The entropy of a random variable  $X$ , that represents a system is given by:

$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$




# Entropy

## Boltzmann Entropy:

In Statistical Mechanics, Boltzmann's Equation is given by:

$$S = k_B \ln(\omega)$$



## Shannon Entropy:

The entropy of a random variable  $X$ , that represents a system is given by:

$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

## von Neumann Entropy:

In a quantum Mechanical system represented by a Density matrix  $\rho$ , von Neumann entropy is:

$$S = -\text{tr}(\rho \ln \rho)$$

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work

## Closed System:

Let us consider a pure state  $|\psi\rangle$ , Now if we let it evolve with time,

$$U(t)|\psi\rangle \rightarrow |\phi\rangle$$

Then,  $|\phi\rangle$  will be a pure state.



# Decoherence

## Closed System:

Let us consider a pure state  $|\psi\rangle$ , Now if we let it evolve with time,

$$U(t)|\psi\rangle \rightarrow |\phi\rangle$$

Then,  $|\phi\rangle$  will be a pure state.

## Open System:

Now let us consider the same state  $|\psi\rangle$  interacting with the environment. So the state of the system will now become  $|\psi'\rangle|e\rangle$  Now if we let it evolve with time,

$$U(t)|\psi'\rangle|e\rangle \rightarrow |F\rangle$$

Then,  $|F\rangle$  may not be a pure state.

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work

# Dephasing Channel

## Representation:

- To remove the coherence caused by the interaction of the system with the environment, we require a Dephasing channel, and the process is called Decoherence.
- For Computational Basis (i.e.  $|0\rangle$  &  $|1\rangle$ ), the Dephasing operation is given by:

$$D \rightarrow (1 - p)\rho_{AB} + p\sigma_z\rho_{AB}\sigma_z,$$

where  $\rho_{AB}$  is a density matrix

# Dephasing Channel


## Representation:

- To remove the coherence caused by the interaction of the system with the environment, we require a Dephasing channel, and the process is called Decoherence.
- For Computational Basis (i.e.  $|0\rangle$  &  $|1\rangle$ ), the Dephasing operation is given by:

$$D \rightarrow (1 - p)\rho_{AB} + p\sigma_z\rho_{AB}\sigma_z,$$

where  $\rho_{AB}$  is a density matrix

## Operation:



# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT**
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work


# Controlled NOT Gate

## Representation:

It is a 2-qubit gate represented as

$$|A, B\rangle \rightarrow |A, A \oplus B\rangle$$

It flips the  $2^{nd}$  qubit, if the  $1^{st}$  qubit is  $|1\rangle$ .




# Controlled NOT Gate

## Representation:

It is a 2-qubit gate represented as

$$|A, B\rangle \rightarrow |A, A \oplus B\rangle$$

It flips the  $2^{nd}$  qubit, if the  $1^{st}$  qubit is  $|1\rangle$ .



## Operation:

$$CNOT = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

**Truth Table:**

Input	Output
$ 00\rangle$	$ 00\rangle$
$ 01\rangle$	$ 01\rangle$
$ 10\rangle$	$ 11\rangle$
$ 11\rangle$	$ 10\rangle$

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics**
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work




# First Law of Thermodynamics

## First Law:

It is just another form of conservation of energy.

$$\Delta U = Q - W$$




# First Law of Thermodynamics

## First Law:

It is just another form of conservation of energy.

$$\Delta U = Q - W$$



## Isothermal Process:

For an Isothermal Process,  $\Delta U = 0$ .

$$\text{So, } Q = W \text{ and } Q_{rev} = T\Delta S$$

$$W = T\Delta S = K_B T \ln(\omega)$$

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations**
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work

# Thermodynamical Approach to Quantifying Quantum Correlations

## TAQQC Paper:

VOLUME 89, NUMBER 18

PHYSICAL REVIEW LETTERS

28 OCTOBER 2002

### Thermodynamical Approach to Quantifying Quantum Correlations

Jonathan Oppenheim,<sup>1,2</sup> Michał Horodecki,<sup>2</sup> Paweł Horodecki,<sup>3</sup> and Ryszard Horodecki<sup>2</sup>

<sup>1</sup>*Theoretical Physics Institute, University of Alberta, 412 Avadh Bhatia Physics Laboratory, Edmonton, Alberta, Canada T6G 2J1*

<sup>2</sup>*Institute of Theoretical Physics and Astrophysics, University of Gdańsk, 80–952 Gdańsk, Poland*

<sup>3</sup>*Faculty of Applied Physics and Mathematics, Technical University of Gdańsk, 80–952 Gdańsk, Poland*  
(Received 11 February 2002; published 11 October 2002)

We consider the amount of work which can be extracted from a heat bath using a bipartite state  $\rho$  shared by two parties. In general it is less than the amount of work extractable when one party is in possession of the entire state. We derive bounds for this “work deficit” and calculate it explicitly for a number of different cases. In particular, for pure states the work deficit is exactly equal to the distillable entanglement of the state. A form of complementarity exists between physical work which can be extracted and distillable entanglement. The work deficit is a good measure of the quantum correlations in a state and provides a new paradigm for understanding quantum nonlocality.

DOI: 10.1103/PhysRevLett.89.180402

PACS numbers: 03.65.Ud, 03.67.–a, 05.70.–a, 89.70.+c

# Thermodynamical Approach to Quantifying Quantum Correlations

## TAQQC Paper:

VOLUME 89, NUMBER 18

PHYSICAL REVIEW LETTERS

28 OCTOBER 2002

### Thermodynamical Approach to Quantifying Quantum Correlations

Jonathan Oppenheim,<sup>1,2</sup> Michał Horodecki,<sup>2</sup> Paweł Horodecki,<sup>3</sup> and Ryszard Horodecki<sup>2</sup>

<sup>1</sup>*Theoretical Physics Institute, University of Alberta, 412 Avadh Bhatia Physics Laboratory, Edmonton, Alberta, Canada T6G 2J1*

<sup>2</sup>*Institute of Theoretical Physics and Astrophysics, University of Gdańsk, 80–952 Gdańsk, Poland*

<sup>3</sup>*Faculty of Applied Physics and Mathematics, Technical University of Gdańsk, 80–952 Gdańsk, Poland*  
(Received 11 February 2002; published 11 October 2002)

We consider the amount of work which can be extracted from a heat bath using a bipartite state  $\rho$  shared by two parties. In general it is less than the amount of work extractable when one party is in possession of the entire state. We derive bounds for this “work deficit” and calculate it explicitly for a number of different cases. In particular, for pure states the work deficit is exactly equal to the distillable entanglement of the state. A form of complementarity exists between physical work which can be extracted and distillable entanglement. The work deficit is a good measure of the quantum correlations in a state and provides a new paradigm for understanding quantum nonlocality.

DOI: 10.1103/PhysRevLett.89.180402


PACS numbers: 03.65.Ud, 03.67.–a, 05.70.–a, 89.70.+c

**Work deficit is defined as the difference between the work extracted from taking the entire state with one party and the work extracted from taking the state shared with two parties.**

# Outline


- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model**
- 8 Quantum Model
- 9 Summary
- 10 Future Work

# Classical Model



# Classical Model

## Work Deficit:




$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$



# Classical Model

## Work Deficit:




$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$

- If we don't know the position of particles in  $n$  bits, then the entropy will be  $n$ .

# Classical Model

## Work Deficit:




$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$

- If we don't know the position of particles in  $n$  bits, then the entropy will be  $n$ .
- Now, if we know the position of any particle, then the entropy will be  $H(X)$ .

# Classical Model

## Work Deficit:




$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$

- If we don't know the position of particles in  $n$  bits, then the entropy will be  $n$ .
- Now, if we know the position of any particle, then the entropy will be  $H(X)$ .
- So the Work Extracted,  $W_c = (n - H(X))T = n - H(X)$ .

# Classical Model

## Work Deficit:




$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$

$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$

- If we don't know the position of particles in  $n$  bits, then the entropy will be  $n$ .
- Now, if we know the position of any particle, then the entropy will be  $H(X)$ .
- So the Work Extracted,  $W_c = (n - H(X))T = n - H(X)$ .
- Now if we replace Shannon Entropy with von Neumann Entropy,  $W_t = n - S(\rho)$ .

# Classical Model

## Work Deficit:



$$H(X) = - \sum_{x=1}^n p_x \log_2(p_x)$$


$$H(X) = - \sum_{x=1}^n \frac{1}{2^n} \log_2\left(\frac{1}{2^n}\right) = n$$

- If we don't know the position of particles in  $n$  bits, then the entropy will be  $n$ .
- Now, if we know the position of any particle, then the entropy will be  $H(X)$ .
- So the Work Extracted,  $W_c = (n - H(X))T = n - H(X)$ .
- Now if we replace Shannon Entropy with von Neumann Entropy,  $W_t = n - S(\rho)$ .
- Now, If  $W_l$  is the work extracted by Alice and Bob after performing LOCC operations.  
Then work deficit,  $\Delta = W_t - W_l$ .

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model**
- 9 Summary
- 10 Future Work

# Quantum Model (LOCC)



**Now Since Bob's qubit is known, so 1 bit of work can be extracted from Bob's qubit.**

# Fortran Program

```
!Generate Random Density Matrix
call RANDDMR(d,state)
!Compute Eigen Values
call DSYEV(JOBZ, UPLO, N, state, LDA, W,
           WORK, LWORK, INFO)
! Calculate the von Neumann entropy
sum = 0.0D0
DO i = 0, d - 1
    IF (W(i) > 0.0D0) THEN
        sum = sum - W(i) *
            LOG(W(i))/LOG(2.0d0)
    END IF
END DO
!Applying CNOT on rho
csys=0.0d0
call cnot(sys, csys)
```



# Fortran Program

```
!Initialising parameters for PTRDMR
site(0) = 1
call PTRDMR(s, s1, site, csys, rdm)
! Call DSYEV to compute eigenvalues and
  eigenvectors
call DSYEV(JOBZ, UPLO, M, rdmb, LDA1, RW,
  RWORK, LWORK1, INFO)
! Calculate the von Neumann entropy
sum = 0.0D0
DO i = 0, m - 1
    IF (RW(i) > 0.0D0) THEN
        sum = sum - RW(i) *
            LOG(RW(i))/LOG(2.0d0)
    END IF
END DO
c_ent = ent_rho - ent_red
write(100,*) c_ent
```

# Output

```
RHO :
  4.8770983126898163E-002   3.2789837842840362E-002   -8.2253410255965928E-002   -3.7497881267235578E-002
  3.2789837842840362E-002   0.20979556004068756         -0.14060519364323271       -0.10482907089261517
  -8.2253410255965928E-002  -0.14060519364323271        0.19387716076973360       0.17690209306155602
  -3.7497881267235578E-002  -0.10482907089261517        0.17690209306155602       0.54755629606268064

EV of RHO :
  1.3720352237903246E-003   8.0458945030873377E-002   0.23726620104507512       0.68090281870026104

Entropy of RHO :
  1.1755379513478428

CNOT_RHO :
  4.8770983126898163E-002   3.2789837842840362E-002   -3.7497881267235578E-002   -8.2253410255965928E-002
  3.2789837842840362E-002   0.20979556004068756         -0.10482907089261517       -0.14060519364323271
  -3.7497881267235578E-002  -0.10482907089261517        0.54755629606268064       0.17690209306155602
  -8.2253410255965928E-002  -0.14060519364323271        0.17690209306155602       0.19387716076973360

EV of CNOT_RHO :
  1.3720352237903068E-003   8.0458945030873349E-002   0.23726620104507512       0.68090281870026115

Entropy of CNOT_RHO :
  1.1755379513478423

RED_CNOT_RHO :
  0.59632727918957884       0.20969193090439639
  0.20969193090439639       0.40367272081042116

EV of RHO_B :
  0.26924114187645748       0.73075885812354247

Entropy of RHO_B :
  0.84037361376372455

Change in Entropy :
  0.33516433758411823
```

# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary**
- 10 Future Work

# Summary

## Work done so far:

- Literature review for basics of Quantum Information Theory.

# Summary

## Work done so far:

- Literature review for basics of Quantum Information Theory.
- Literature review for the basics of Linear Algebra.

# Summary

## Work done so far:

- Literature review for basics of Quantum Information Theory.
- Literature review for the basics of Linear Algebra.
- Fortran/ Python codes (i.e., 1-D Ising Model, Schmidt Decomposition for  $n$  qubits, Used Lapack to calculate the Eigen values and Eigen vectors for matrix).

# Summary

## Work done so far:

- Literature review for basics of Quantum Information Theory.
- Literature review for the basics of Linear Algebra.
- Fortran/ Python codes (i.e., 1-D Ising Model, Schmidt Decomposition for  $n$  qubits, Used Lapack to calculate the Eigen values and Eigen vectors for matrix).
- TAQQC Paper review.

## Work done so far:

- Literature review for basics of Quantum Information Theory.
- Literature review for the basics of Linear Algebra.
- Fortran/ Python codes (i.e., 1-D Ising Model, Schmidt Decomposition for  $n$  qubits, Used Lapack to calculate the Eigen values and Eigen vectors for matrix).
- TAQQC Paper review.
- Made code for calculating the change in entropy and then Work deficit for 2 qubit cases.



# Outline

- 1 Entropy
- 2 Decoherence
- 3 Dephasing Operation
- 4 Controlled Not Gate: CNOT
- 5 First Law of Thermodynamics
- 6 Thermodynamical Approach to Quantify Quantum Correlations
- 7 Classical Model
- 8 Quantum Model
- 9 Summary
- 10 Future Work**

# Future Work

## Work to be done:

- Quantifying entanglement for spin 1 - Heisenberg Model (AKLT)  
(i.e. **Affleck-Kennedy-Lieb-Tasaki Model**) using work deficit and other thermodynamical models.

# Future Work

## Work to be done:

- Quantifying entanglement for spin 1 - Heisenberg Model (AKLT)  
(i.e. **Affleck-Kennedy-Lieb-Tasaki Model**) using work deficit and other thermodynamical models.
- AKLT model is the extension of the Heisenberg Spin Model. Ground state is a one-dimensional state with a valence bond between every pair of electrons.

# Future Work

## Work to be done:

- Quantifying entanglement for spin 1 - Heisenberg Model (AKLT)  
(i.e. **Affleck-Kennedy-Lieb-Tasaki Model**) using work deficit and other thermodynamical models.
- AKLT model is the extension of the Heisenberg Spin Model. Ground state is a one-dimensional state with a valence bond between every pair of electrons.
- Because this leads to two spin half's for every site, the result must be the wavefunction of a spin 1 system.

# Future Work

## Work to be done:

- Quantifying entanglement for spin 1 - Heisenberg Model (AKLT) (i.e. **Affleck-Kennedy-Lieb-Tasaki Model**) using work deficit and other thermodynamical models.
- AKLT model is the extension of the Heisenberg Spin Model. Ground state is a one-dimensional state with a valence bond between every pair of electrons.
- Because this leads to two spin half's for every site, the result must be the wavefunction of a spin 1 system.
- Understanding the implications of work deficit in multiqubit scenarios.

# References

## References:

- Thermodynamical Approach to Quantifying Quantum Correlations: Jonathan Oppenheim, Michael Horodecki, Pawel Horodecki, and Ryszard Horodecki, Phys. Rev. Lett. 2002
- Numerical Recipes in Quantum Information Theory and Quantum Computing: M. S. Ramkarthik, and Payal D. Solanki, CRC Press 2022
- Quantum Computing: From Linear Algebra to Physical Realizations: Mikio Nakahara, and Tetsuo Ohmi, CRC Press 2008
- Maxwell's Demon 2: Entropy, Classical and Quantum Information, Computing, Leff and Rex, CRC Press 2002
- Statistical Thermodynamics: An Information Theory Approach, Aubin, John Wiley, 2024
- Quantum Information Theory, Riccardo Manenti and Mario Motta, Oxford University Press, 2023

*Thanks For Your Attention!*