

Statistical Computing with R: Masters in Data Sciences 503 (S20) Third Batch, SMS, TU, 2024

Shital Bhandary

Associate Professor

Statistics/Bio-statistics, Demography and Public Health Informatics

Patan Academy of Health Sciences, Lalitpur, Nepal

Faculty, Data Analysis and Decision Modeling, MBA, Pokhara University, Nepal

Faculty, FAIMER Fellowship in Health Professions Education, India/USA.

Review Preview

- One-way ANOVA
- Multiple proportion test
- Linear relationship
- Covariance
- Correlation
- Simple Linear regression model fit, interpretation and residual analysis
- Simple Linear Regression prediction and Machine Learning

Comparing means of an outcome variable across another variable with more than two categories:

- **One-way ANOVA**

- $H_0: \mu_1 = \mu_2 = \mu_3$
- H_1 : At least one pair of means are not equal
- **If H_1 is accepted, pairwise comparison (post-hoc) test must be done to find the significant pairs!**

- Compare mpg (miles per gallon) by cars with different gear (numbers of gears) using “mtcars” data
- Dependent variable = mpg
- Independent variable = gears

Assumptions of 1-way ANOVA:

- Same as two-samples t-test:
- Normally distributed:
 - Test of normality by each category
- Dependent variable must be “normally distributed” **for each categories**
- Homogenous variance:
 - **var.test is not useful (>2 groups)**
 - Levene’s Variance test is preferred
 - It is available in the “car” package
 - `library(car)`
 - `leveneTest(y~x, data=data)`
 - **x must be categorical i.e. factor!**
- Variance across categories must be same

1-way ANOVA assumptions checks:

Normality by categories:

- `with(mtcars, shapiro.test(mpg[gear == 3]))`

W = 0.95833, p-value = 0.6634

- `with(mtcars, shapiro.test(mpg[gear == 4]))`

W = 0.90908, p-value = 0.2076

- `with(mtcars, shapiro.test(mpg[gear == 5]))`

W = 0.90897, p-value = 0.4614

Equal variance among categories:

`library(car)`

`leveneTest(mpg ~ gear, data=mtcars)`

Result:

Levene's Test for Homogeneity of Variance (center = median)

	Df	F value	Pr(>F)
group 2		1.4886	0.242429

Levene's Test is a GOF test, so group variances are equal as p-value>0.05.

So, Classical 1-way ANOVA can be used now!

- `summary(aov(mpg ~ gear, data = mtcars))`
- Since F-test p-value < 0.05 , we accept H_1 . At least one of the mean pairs are not equal!
- This means, post-hoc test or pairwise comparison is required!
- **Fisher's LSD uses pairwise t-tests (not good)!**
- For classical 1-way ANOVA, Tukey HSD is the best post-hoc test!
- `TukeyHSD(aov(mpg ~ gear, data = mtcars))`

	Df	SumSq	MeanSq	Fvalue	Pr(>F)
gear	2	483.2	241.62	10.9	0.000295
Residuals	29	642.8	22.17		

Tukey multiple comparisons of means

95% family-wise confidence level

Fit: `aov(formula = mpg ~ gear, data = mtcars)`

\$gear	diff	lwr	upr	p adj
4-3	8.426667	3.9234704	12.929863	0.0002088
5-3	5.273333	-0.7309284	11.277595	0.0937176
5-4	-3.153333	-9.3423846	3.035718	0.4295874

Check this result with the simple linear model (regression):

- `summary(lm(mpg ~ gear, data = mtcars))`
 - P-value are reported without correcting them i.e. simple t-test were used, which can be checked with this command in R/R Studio:
 - `pairwise.t.test(mtcars$mpg, mtcars$gear, p.adj = "none")`
- | | |
|----------------------|----------------|
| • 3 | 4 |
| • 4 7.3e-05 (3 vs 4) | -- |
| • 5 0.038 (3 vs 5) | 0.218 (4 vs 5) |
- What is the interpretation?
 - **Why gear = 3 category is omitted in the result?**

Coefficients:

- Estimate Std. Error t value Pr(>|t|)
- | | | | | |
|-------------|--------------|-------|--------|------------------------|
| (Intercept) | 16.107 | 1.216 | 13.250 | 7.87e-14 *** |
| gear[T.4] | 8.427 | 1.823 | 4.621 | 7.26e-05 *** |
| gear[T.5] | 5.273 | 2.431 | 2.169 | 0.0384 * (why?) |
-
- Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
- R automatically creates 3 dummy variables for 3 categories of gear variable i.e. 3, 4 and 5 and uses only last two of them in the model and takes the first one as reference!
 - gear[T.3] = 1 if gear = 3, else 0
 - gear[T.4] = 1 if gear = 4, else 0
 - gear[T.5] = 1 if gear = 5, else 0

Multiple proportion test

- $H_0: P_1 = P_2 = P_3 \dots = P_n$
- H_1 : At least one of the proportion pairs are not equal
- Lets do it for gear variable of mtcars data
- `table(mtcars$gear)`
- `prop.test(x=c(15,12,5),
n=c(32,32,32)) #Correct=F?`
- `pairwise.prop.test(x=c(15,12,5),
n=c(32,32,32), correct=F)`

Question/queries so far?

Measures of linear relationship:


- Two continuous variables
- Assumption:
- Two continuous variables have linear or “tentative” linear relationship
- Assessed using “scatterplot”
- Measures of linear relationship:
- Covariance
 - Limitations
- Pearson’s Correlation Coefficient
 - Limitations
- Simple Linear regression

Covariance

- It measures the linear relationship between two quantitative variables.
 1. Positive values indicate a positive linear relationship; negative, a negative linear relationship.
 2. Close to zero means there is not much of a linear relationship.
 3. The magnitude of covariance is difficult to interpret.
 4. Covariance has problems with units (like feet compared to inches).

Example: which one is more linear?

- `plot(mtcars$wt, mtcars$mpg)`



There is a “tentative” linear relationship between mpg and weight variables! So, we can use measures of linear relationship for these variables!

Covariance between WT and MPG variables:

```
cov(mtcars$wt, mtcars$mpg)
```

- **-5.116685**

Do as follows now:

- Convert the weight (wt) variable measured in pound to kilogram and store it a new variable **wt2**
- Compute the covariance of weight in KG and MPG now!
- **-2.325766**

Sample covariance for a sample of size n with the observations:

$$s_{xy} = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

Population covariance:

$$\sigma_{xy} = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{N}$$

Pearson's Correlation Coefficient (r) to measure linear relationship:

- Measure the strength and direction of the linear relationship between two quantitative variables.
- A relative measure of strength of association (relationship) between 2 variables or a measure of strength per unit of standard deviation, $s_x * s_y$.
- **Solves “units” and “magnitude” problems of covariance.**

$$r_{xy} = \frac{s_{xy}}{s_x s_y}$$

$$s_{xy} = \text{sample covariance} = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

$$s_x = \text{sample standard deviation of } x = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

$$s_y = \text{sample standard deviation of } y = \sqrt{\frac{\sum (y_i - \bar{y})^2}{n - 1}}$$

Correlation of WT, WT2 and MPG variables:

```
cor(mtcars$wt, mtcars$mpg)
```

- **-0.8676594**

```
cor(mtcars$wt2, mtcars$mpg)
```

- **-0.8676594**

Interpretation (Pearson):

- Low degree: <0.25
- Medium degree: $0.25-0.75$
- High degree: >0.75

- How to check if this correlation is a valid linear correlation?

- We need to do the hypothesis testing:

- H_0 : Linear correlation is zero i.e. $\rho = 0$.
- H_1 : Linear correlation is NOT zero i.e. $\rho \neq 0$.

Test of “true” linear correlation of WEIGHT and MPG variables:

- `cor.test(mtcars$wt, mtcars$mpg)`
- `cor.test(mtcars$wt2, mtcars$mpg)`

Interpretation (two parts, always!):

- Since $p\text{-value} < 0.05$, we accept H_1
(**Decision**)
- This means the true linear correlation coefficient is NOT zero so computed sample estimate of this correlation coefficient as -0.87 is a valid estimate
(**Conclusion**)

Pearson's product-moment correlation

data: mtcars\$wt and mtcars\$mpg

$t = -9.559$, $df = 30$, **p-value = $1.294e-10$**

alternative hypothesis: true correlation is not equal to 0

95 percent confidence interval:

-0.9338264 -0.7440872

sample estimates:

cor

-0.8676594

Limitation of Linear correlation coefficient:

- It provides the magnitude and direction of the relationship between two linearly related quantitative variables
- It does not provide the estimate of change in dependent variable with respect to the change in the independent variable
- Thus, it is required to use a simple linear regression i.e.
$$y = a + bx$$
- Simple linear regression is an extension of the simple linear correlation
- **But it come with many assumptions!**

Simple Linear Regression:

A simple linear regression model of Y on X in stochastic form (population) in statistics is written as:

$$Y = \alpha + \beta X + u$$


where α and β are parameters called y-intercept and slope respectively, and u is called error or disturbance term, which is erratic or random in nature.

- For given n pairs of data values $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$ of (X, Y) , the estimated model is written as:

$$\hat{y} = a + bx$$

- \hat{y} is estimated value of Y based on a and b , which are least square estimates of α and β respectively.
- We need to calculate best solutions of n equations each containing two unknown parameters α and β using **OLS method**.

LINE OF BEST FIT for minimizing error - OLS
(OLS = Ordinary Least Square Method)



Simple Linear Regression Assumptions:

- Dependent variable: Normal
- **Dependent and Independent (continuous) variables: Linear**

Regression Model:

- Coefficient of determination > 0.50
- Regression ANOVA must be significant statistically
- Y-intercept (a) and slope (b) must be statistically significant

If these conditions are satisfied then it is called a BLUE estimate!

- Regression Model Residuals or Errors i.e. " $y - \hat{y}$ " must be:
 - **Linear** - Linearity of residuals
 - **Independent** - Independence of residuals (for time series)
 - **Normal** - Normality of residuals
 - **Equal variance** - Homoscedasticity of residuals
- Also known as LINE test
 - Each of these assumptions must be checked with graphs and statistical methods

Simple Linear Regression between MPG and WT variables:

- Dependent variable MPG follows normal distribution (checked!)
- We need to check after fitting the simple linear regression:
- Dependent variable MPG and independent variable WT has “tentative” linear relationship
- R-square > 0.50 (why?)
- Regression ANOVA p-value < 0.05 (why?)
- We can move forward!
- Regression coefficients i.e. a and b p-values < 0.05 .

Let's fit the model and get the summary:

```
lm1 <- lm(mtcars$mpg ~  
mtcars$wt)  
lm1
```

Call:

```
lm(formula = mtcars$mpg ~  
mtcars$wt)
```

The outputs shows the
“minimum” results for the model

Coefficients:

(Intercept)	mtcars\$wt
37.285	-5.344

R gives the “minimalist” output!

Let's ask R to provide summary of lm1:

- `summary(lm1)`
- The coefficient of determination (R-square) = 0.7528, which means the independent variable (wt) is able to explain around 75.28% of variance (variability) in the dependent variable (mpg)

The regression ANOVA, hypothesis:

- H0: Intercept only model ($y = a$) is better
- H1: Intercept only model is significantly reduced than the full model ($y = a + bx$)
- Regression ANOVA (given by F-Test) p-value < 0.05, we accept H1.
- It confirms that intercept only model is significantly reduced than the full model!

Residuals:

- Min 1Q Median 3Q Max
- -4.5432 -2.3647 -0.1252 1.4096 6.8727

Coefficients:

- Estimate Std. Error t value Pr(>|t|)
- (Intercept) 37.2851 1.8776 19.858 < 2e-16 ***
- mtcars\$wt -5.3445 0.5591 -9.559 1.29e-10 ***
- ---
- Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 3.046 on 30 degrees of freedom

Multiple R-squared: 0.7528, Adjusted R-squared: 0.7446

F-statistic: 91.38 on 1 and 30 DF, p-value: 1.294e-10

Question/queries?

- Next class
 - Residual Standard Error (RSE) or Standard Error of (Regression) Estimation
 - Use of RSE/SEE for testing regression constant i.e. a
 - Use of RSE/SEE for testing regression constant i.e. b
- Residual Analysis
 - L = Linearity of Residuals
 - I = Independence of Residuals
 - N = Normality of Residuals
 - E = Equal variance (homoscedasticity) of residuals

Thank you!

@shitalbhandary