

SC223 - Linear Algebra

Aditya Tatu

Lecture 30



October 18, 2023

Summary of Lecture 29

- **Similar matrices and Similarity transformation:** We say two matrices A and B are similar if there exists an invertible matrix, say S such that $B = SAS^{-1}$. The transformation $A \mapsto SAS^{-1}$ is said to be a similarity transformation of A by S .
- Let $T \in \mathcal{L}(U)$ be a linear operator. It is preferable to work with a basis of U , say β such that $[T]_{\beta}^{\beta}$ is diagonal/block-diagonal.

$$\begin{aligned} T &\in \mathcal{L}(U) & \beta &- \text{basis of } U \\ [T]_{\beta}^{\beta} &\in \mathbb{F}^{n \times n} & \alpha &- \text{basis of } U \\ [T]_{\alpha}^{\alpha} &\in \mathbb{F}^{n \times n} \\ [T]_{\alpha}^{\alpha} &= \underbrace{N_{\beta}^{\alpha}}_S [T]_{\beta}^{\beta} \underbrace{(N_{\beta}^{\alpha})^{-1}}_{S^{-1}} \\ [T]_{\gamma}^{\gamma} &= S_1 [T]_{\beta}^{\beta} (S_1)^{-1} \end{aligned}$$

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.

$$u \in U, \quad u = v_1 + v_2.$$

$$Tu = Tv_1 + Tv_2, \quad \begin{array}{l} v_1 \in V_1 \\ v_2 \in V_2. \end{array}$$

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.
- If $T(V_1) \subseteq V_1, T(V_2) \subseteq V_2$, then in a basis $B = B_{V_1} \cup B_{V_2}$, $[T]_B^B = ?$

Ex: Let $\dim(U) = 5$, $\dim(V_1) = 2$,
 $\dim(V_2) = 3$.

$$[T]_B^B \in \mathbb{F}^{5 \times 5} =$$

$$B = B_{V_1} \cup B_{V_2}$$

$$B_{V_1} = \{p_1, p_2\}$$

$$B_{V_2} = \{q_1, q_2, q_3\}.$$

$$\begin{bmatrix} \boxed{\begin{matrix} \star & \star \\ \star & \star \end{matrix}} & \begin{matrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{matrix} \\ \begin{matrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{matrix} & \boxed{\begin{matrix} \star & \star & \star \\ \star & \star & \star \\ \star & \star & \star \end{matrix}} \end{bmatrix}_{5 \times 5}$$

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.
- If $T(V_1) \subseteq V_1, T(V_2) \subseteq V_2$, then in a basis $B = B_{V_1} \cup B_{V_2}$, $[T]_B^B = ?$

$$[T]_B^B = \begin{bmatrix} [T_{V_1}]_{B_{V_1}}^{B_{V_1}} & 0 \\ 0 & [T_{V_2}]_{B_{V_2}}^{B_{V_2}} \end{bmatrix},$$

$[T]_B^B$ becomes block-diagonal.

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.
- If $T(V_1) \subseteq V_1, T(V_2) \subseteq V_2$, then in a basis $B = B_{V_1} \cup B_{V_2}$, $[T]_B^B = ?$

$$[T]_B^B = \begin{bmatrix} [T_{V_1}]_{B_{V_1}}^{B_{V_1}} & 0 \\ 0 & [T_{V_2}]_{B_{V_2}}^{B_{V_2}} \end{bmatrix},$$

$[T]_B^B$ becomes block-diagonal.

- **Definition:** Let $T \in \mathcal{L}(U)$. If V is a subspace of U such that $\forall u \in V, Tu \in V$, then V is said to be an *invariant subspace* of T , or *T -invariant subspace*.

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.
- If $T(V_1) \subseteq V_1, T(V_2) \subseteq V_2$, then in a basis $B = B_{V_1} \cup B_{V_2}$, $[T]_B^B = ?$

$$[T]_B^B = \begin{bmatrix} [T_{V_1}]_{B_{V_1}}^{B_{V_1}} & 0 \\ 0 & [T_{V_2}]_{B_{V_2}}^{B_{V_2}} \end{bmatrix},$$

$[T]_B^B$ becomes block-diagonal.

● **Definition:** Let $T \in \mathcal{L}(U)$. If V is a subspace of U such that $\forall u \in V, Tu \in V$, then V is said to be an *invariant subspace* of T , or *T-invariant subspace*.

- U and $\{\theta\}$ are trivial examples of invariant subspaces.

$$A \in \mathbb{R}^{n \times n}, A \in \mathcal{L}(\mathbb{R}^n)$$

$$C(A), \text{ let } y \in C(A), Ay \in C(A).$$

$$\underline{C(A^T)} : \text{ let } y \in C(A^T) \Rightarrow y = A^T x. Ay = A^T z.$$

Invariant Subspaces

- Let $T \in \mathcal{L}(U)$, and $U = V_1 \oplus V_2$.
- Consider $T|_{V_1}, T|_{V_2}$.
- If $T(V_1) \subseteq V_1, T(V_2) \subseteq V_2$, then in a basis $B = B_{V_1} \cup B_{V_2}$, $[T]_B^B = ?$

$$[T]_B^B = \begin{bmatrix} [T_{V_1}]_{B_{V_1}}^{B_{V_1}} & 0 \\ 0 & [T_{V_2}]_{B_{V_2}}^{B_{V_2}} \end{bmatrix},$$

$[T]_B^B$ becomes block-diagonal.

- **Definition:** Let $T \in \mathcal{L}(U)$. If V is a subspace of U such that $\forall u \in V, Tu \in V$, then V is said to be an *invariant subspace* of T , or *T -invariant subspace*.
- U and $\{\theta\}$ are trivial examples of invariant subspaces.
- Other examples include $N(T), R(T)$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}, u \neq \theta$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x =$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx =$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$,

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u =$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}, u \neq \theta$. Let $x \neq \theta, x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.

$$\left. \begin{array}{l} \\ \\ \\ \end{array} \right\} (T - \lambda I)(u) = \theta$$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is not invertible, and \det

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is not invertible, and $\det([T]_{\beta}^{\beta} - \lambda I_n) =$

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is not invertible, and $\det([T]_{\beta}^{\beta} - \lambda I_n) = 0$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is not invertible, and $\det([T]_\beta^\beta - \lambda I_n) = 0$.
- Define $c(\lambda) = \det([T]_\beta^\beta - \lambda I)$. Eigenvalues of T are roots of the polynomial $c(\lambda)$, and the eigenvector is a non-zero vector belonging to $\mathcal{N}(T - \lambda I)$.

1 dimensional Invariant Subspaces

- Let V be a 1-dimensional invariant subspace for $T \in \mathcal{L}(U)$.
- $V = \{ku \mid \forall k \in \mathbb{F}\}$, $u \neq \theta$. Let $x \neq \theta$, $x \in V$, i.e., $x = a_1 u$ for some non-zero $a_1 \in \mathbb{F}$.
- Since V is T -invariant, $Tx \in V$, therefore $Tx = a_2 u \Rightarrow Tu = \frac{a_2}{a_1} u$.
- We usually write this as $Tu = \lambda u$, and we call λ **as the eigenvalue associated with eigenvector u** .
- Note that if $Tu = \lambda u$, then $(T - \lambda I)u = \theta$.
- Thus $(T - \lambda I)$ is not invertible, and $\det([T]_{\beta}^{\beta} - \lambda I_n) = 0$.
- Define $c(\lambda) = \det([T]_{\beta}^{\beta} - \lambda I)$. Eigenvalues of T are roots of the polynomial $c(\lambda)$, and the eigenvector is a non-zero vector belonging to $\mathcal{N}(T - \lambda I)$.
- The polynomial c is called the **characteristic polynomial** of T .

End of class.

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e =$

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.
- Note that for any other basis β , $[T]_\beta^\beta [v_i]_\beta =$

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.
- Note that for any other basis β , $[T]_\beta^\beta [v_i]_\beta = \lambda_i [v_1]_\beta$.

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.
- Note that for any other basis β , $[T]_\beta^\beta [v_i]_\beta = \lambda_i [v_i]_\beta$.
- Thus,

$$[T]_\beta^\beta \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ [v_1]_\beta & [v_2]_\beta & \dots & [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} = \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ \lambda_1 [v_1]_\beta & \lambda_2 [v_2]_\beta & \dots & \lambda_n [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

=

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.
- Note that for any other basis β , $[T]_\beta^\beta [v_i]_\beta = \lambda_i [v_i]_\beta$.
- Thus,

$$\begin{aligned}
 [T]_\beta^\beta \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ [v_1]_\beta & [v_2]_\beta & \dots & [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} &= \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ \lambda_1 [v_1]_\beta & \lambda_2 [v_2]_\beta & \dots & \lambda_n [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \\
 &= \underbrace{\begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ [v_1]_\beta & [v_2]_\beta & \dots & [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}}_E \underbrace{\begin{bmatrix} \lambda_1 & 0 & \dots & \vdots \\ 0 & \lambda_2 & \dots & 0 \\ 0 & \dots & \vdots & \lambda_n \end{bmatrix}}_\Lambda
 \end{aligned}$$

- For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if a diagonal matrix representation is preferred, we need an *eigenbasis* of the linear operator: $e = \{v_1, \dots, v_n\}$.
- Since $Tv_i = \lambda_i v_i$, $[T]_e^e = \text{diag}(\lambda_1, \dots, \lambda_n)$.
- Note that for any other basis β , $[T]_\beta^\beta [v_i]_\beta = \lambda_i [v_i]_\beta$.
- Thus,

$$\begin{aligned}
 [T]_\beta^\beta \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ [v_1]_\beta & [v_2]_\beta & \cdots & [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} &= \begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ \lambda_1 [v_1]_\beta & \lambda_2 [v_2]_\beta & \cdots & \lambda_n [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \\
 &= \underbrace{\begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ [v_1]_\beta & [v_2]_\beta & \cdots & [v_n]_\beta \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}}_E \underbrace{\begin{bmatrix} \lambda_1 & 0 & \cdots & \vdots \\ 0 & \lambda_2 & \cdots & 0 \\ 0 & \cdots & \vdots & \lambda_n \end{bmatrix}}_\Lambda \\
 [T]_\beta^\beta E = E \Lambda &\Rightarrow \Lambda = [T]_E^E = E^{-1} [T]_\beta^\beta E
 \end{aligned}$$

- Eigenvalues of T are roots of $\det([T]_\beta^\beta - \lambda I_n)$, and corresponding eigenvectors are linearly independent vectors in $\mathcal{N}(T - \lambda I)$.

Examples

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

Results on Eigenvectors and Eigenvalues

- Proposition 19: Let $\lambda_1, \dots, \lambda_m$ be distinct eigenvalues of $T \in \mathcal{L}(U)$. Then the eigenvectors v_1, \dots, v_m associated with these eigenvalues are linearly independent.