

SC223 - Linear Algebra

Aditya Tatu

Lecture 24



September 29, 2023

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) =$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.

Proof:
Since $U \cap W$ is a FDVS, it has a basis.

Let $B_{U \cap W} = \{\underbrace{p_1, \dots, p_k}_{\hookrightarrow \textcircled{1}}\}$ be a basis of $U \cap W$.

$\{p_1, \dots, p_k\}$ in U is LI.

" in W is LI.

Let $B_U = \{p_1, \dots, p_k, u_1, \dots, u_n\}$ be a basis of U $\xrightarrow{\textcircled{2}}$

Let $B_W = \{p_1, \dots, p_k, w_1, \dots, w_m\}$ " of W $\hookrightarrow \textcircled{3}$

$$\beta_{U+W} = \beta_U \cup \beta_W$$

$$u_i = \sum_{i=2}^n a_i u_i + \sum_{i=1}^k b_i p_i + \sum_{i=1}^m c_i w_i \Rightarrow \begin{matrix} a_i = 0 \\ b_i = 0 \\ c_i = 0 \end{matrix} \quad \forall i$$

$$\left. \begin{array}{l} \Rightarrow \beta_U \cup \beta_W \text{ is LI.} \\ \text{span}(\beta_U \cup \beta_W) = U+W \end{array} \right\} \Rightarrow \beta_{U+W} = \beta_U \cup \beta_W$$

$$\dim(U+W) = |\beta_U \cup \beta_W| = n+k+m$$

$$\dim(U) = n+k$$

$$\dim(W) = m+k$$

$$\dim(U \cap W) = k$$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) =$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) =$

$$\text{span}(\emptyset) = \{\emptyset\}.$$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.


Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.
Similarly, $N(A^T) \oplus C(A) = \mathbb{R}^m$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.
Similarly, $N(A^T) \oplus C(A) = \mathbb{R}^m$.

$$A: \mathbb{R}^n \longrightarrow \mathbb{R}^m$$




Linear Transformations

Linear transformations

- Let U and V be vector spaces over the same field \mathbb{F} . A function $f : U \rightarrow V$ is said to be **Linear transformation** from U to V if

Additive : $\forall x, y \in U, f(x + y) = f(x) + f(y)$

Homogeneous : $\forall a \in \mathbb{F}, \forall x \in U, f(a \cdot x) = a \cdot f(x)$.



Linear transformations

- Let U and V be vector spaces over the same field \mathbb{F} . A function $f : U \rightarrow V$ is said to be **Linear transformation** from U to V if

$$\textit{Additive} : \forall x, y \in U, f(x + y) = f(x) + f(y)$$

$$\textit{Homogeneous} : \forall a \in \mathbb{F}, \forall x \in U, f(a \cdot x) = a \cdot f(x).$$

- If domain and co-domain vector spaces are same, we usually call Linear transformations, **Linear Operators**.

Linear transformations

- Let U and V be vector spaces over the same field \mathbb{F} . A function $f : U \rightarrow V$ is said to be **Linear transformation** from U to V if

$$\text{Additive : } \forall x, y \in U, f(x + y) = f(x) + f(y)$$

$$\text{Homogeneous : } \forall a \in \mathbb{F}, \forall x \in U, f(a \cdot x) = a \cdot f(x).$$

- If domain and co-domain vector spaces are same, we usually call Linear transformations, **Linear Operators**.
- Examples:

$$\textcircled{1} \quad U. \quad I : U \rightarrow U \quad \left. \begin{array}{l} \\ \forall x \in U, I(x) = x. \end{array} \right\} \text{Identity mapping}$$

$$\textcircled{2} \quad U = \mathcal{L}_2(\mathbb{R}) \quad \text{Let } h \in \mathcal{L}_2(\mathbb{R}) \quad h(x) = 0 \quad |x| \geq a$$
$$y(t) = \int_{-\infty}^{\infty} h(z) x(t-z) dz$$

$$(3) U = \mathbb{R}^n, V = \mathbb{R}^m.$$

$$A \in \mathbb{R}^{m \times n}, \quad A: x \rightarrow A \cdot x.$$

$$(4) U = V = \mathbb{R}^2.$$

$$\theta \in [0, 2\pi).$$

$$R_\theta: \mathbb{R}^2 \rightarrow \mathbb{R}^2.$$

$$R_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$(5) U = P_n(\mathbb{R}). \quad \forall p \in P_n(\mathbb{R}), \quad \frac{d}{dx}(p) \in P_n(\mathbb{R})$$

Is $\frac{d}{dx}$ a L.T?

④. $U = P_n(\mathbb{R})$, $V = P_{n+1}(\mathbb{R})$

$\int : U \rightarrow V$ a L.T.?

$$\int (p + q)(x) dx = \int p(x) dx + \int q(x) dx$$

Vector space Homomorphism & Isomorphism

- We say that two vector spaces over \mathbb{F} , U and V are **homomorphic** if there exists a linear transformation between them.

Vector space Homomorphism & Isomorphism

- We say that two vector spaces over \mathbb{F} , U and V are **homomorphic** if there exists a linear transformation between them.
- We say that two vector spaces over \mathbb{F} , U and V are **isomorphic** if there exists an **invertible** linear transformation between them.

Vector space Homomorphism & Isomorphism

- We say that two vector spaces over \mathbb{F} , U and V are **homomorphic** if there exists an linear transformation between them.
- We say that two vector spaces over \mathbb{F} , U and V are **isomorphic** if there exists an **invertible** linear transformation between them.
- If T is an invertible linear transformation between U and V , then we say that T is an **isomorphism** between U and V .

Vector space Homomorphism & Isomorphism

- We say that two vector spaces over \mathbb{F} , U and V are **homomorphic** if there exists a linear transformation between them.
- We say that two vector spaces over \mathbb{F} , U and V are **isomorphic** if there exists an **invertible** linear transformation between them.
- If T is an invertible linear transformation between U and V , then we say that T is an **isomorphism** between U and V .
- **Proposition 19:** Show that two vector spaces U and V over \mathbb{F} are isomorphic iff they have the same dimensions.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i), i = 1, \dots, n$ is enough to allow us to compute $T(x), \forall x \in U$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i)$, $i = 1, \dots, n$ is enough to allow us to compute $T(x)$, $\forall x \in U$.
- Now, $T(u_i) \in V$, thus $T(u_i) = \sum_{j=1}^m c_{ji} v_j$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i)$, $i = 1, \dots, n$ is enough to allow us to compute $T(x)$, $\forall x \in U$.
- Now, $T(u_i) \in V$, thus $T(u_i) = \sum_{j=1}^m c_{ji} v_j$.
- Then, $y = \sum_{i=1}^n a_i (\sum_{j=1}^m c_{ji} v_j) = \sum_{i=1}^n \sum_{j=1}^m a_i c_{ji} v_j$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i)$, $i = 1, \dots, n$ is enough to allow us to compute $T(x)$, $\forall x \in U$.
- Now, $T(u_i) \in V$, thus $T(u_i) = \sum_{j=1}^m c_{ji} v_j$.
- Then, $y = \sum_{i=1}^n a_i (\sum_{j=1}^m c_{ji} v_j) = \sum_{i=1}^n \sum_{j=1}^m a_i c_{ji} v_j$.
- Also, since $y \in V$, $y = \sum_{j=1}^m b_j v_j$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i)$, $i = 1, \dots, n$ is enough to allow us to compute $T(x)$, $\forall x \in U$.
- Now, $T(u_i) \in V$, thus $T(u_i) = \sum_{j=1}^m c_{ji} v_j$.
- Then, $y = \sum_{i=1}^n a_i (\sum_{j=1}^m c_{ji} v_j) = \sum_{i=1}^n \sum_{j=1}^m a_i c_{ji} v_j$.
- Also, since $y \in V$, $y = \sum_{j=1}^m b_j v_j$.
- Thus, $\sum_{j=1}^m b_j v_j = \sum_{j=1}^m \sum_{i=1}^n c_{ji} a_i v_j$.

Representation of Linear Transformations between FDVS

- Let $T : U \rightarrow V$ be a LT, and let $\beta_U := \{u_1, \dots, u_n\}$ and $\beta_V = \{v_1, \dots, v_m\}$ denote the basis of U and V respectively.
- For an $x \in U$, $x = \sum_{i=1}^n a_i u_i$,
 $y = Tx = T(\sum_{i=1}^n a_i u_i) = \sum_{i=1}^n a_i T(u_i)$.
- Thus, $T(u_i)$, $i = 1, \dots, n$ is enough to allow us to compute $T(x)$, $\forall x \in U$.
- Now, $T(u_i) \in V$, thus $T(u_i) = \sum_{j=1}^m c_{ji} v_j$.
- Then, $y = \sum_{i=1}^n a_i (\sum_{j=1}^m c_{ji} v_j) = \sum_{i=1}^n \sum_{j=1}^m a_i c_{ji} v_j$.
- Also, since $y \in V$, $y = \sum_{j=1}^m b_j v_j$.
- Thus, $\sum_{j=1}^m b_j v_j = \sum_{j=1}^m \sum_{i=1}^n c_{ji} a_i v_j$.

- For $k \in \{1, \dots, m\}$, $b_k = \sum_{i=1}^n c_{ki} a_i$,

- For $k \in \{1, \dots, m\}$, $b_k = \sum_{i=1}^n c_{ki} a_i$, or,

$$\underbrace{\begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}}_{[y]_{\beta_V}} = \underbrace{\begin{bmatrix} c_{11} & \dots & c_{1n} \\ \vdots & \vdots & \vdots \\ c_{m1} & \dots & c_{mn} \end{bmatrix}}_{[T]_{\beta_U}^{\beta_V}} \underbrace{\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}}_{[x]_{\beta_U}}$$

- The matrix $[T]_{\beta_U}^{\beta_V}$ is called the matrix representation of the linear transformation T with respect to the basis β_U and β_V .

Examples

- $\frac{d}{dx} : \mathcal{P}_3(\mathbb{R}) \rightarrow \mathcal{P}_3(\mathbb{R})$.

Examples

- $\frac{d}{dx} : \mathcal{P}_3(\mathbb{R}) \rightarrow \mathcal{P}_3(\mathbb{R})$.
- $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Examples

- $\frac{d}{dx} : \mathcal{P}_3(\mathbb{R}) \rightarrow \mathcal{P}_3(\mathbb{R})$.
- $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$.
- Let $p \in \mathcal{P}_3(\mathbb{R})$ be such that $p(x) = p_0 + p_1x + p_2x^2 + p_3x^3$. Define $T_p : \mathcal{P}_3(\mathbb{R}) \rightarrow \mathcal{P}_6(\mathbb{R})$ by $T_p(q) = p \cdot q, \forall q \in \mathcal{P}_3(\mathbb{R})$, where \cdot represents multiplication between polynomials.