

SC223 - Linear Algebra

Aditya Tatu

Lecture 21



September 21, 2023

Summary of Lecture 20

- Let V be a VS, and $U = \{v_1, \dots, v_n\}$.
- **Definition:** (Span of a set) We define the **Span of U** as

$$\text{span}(U) := \{a_1 v_1 + \dots + a_n v_n \mid \forall a_1, \dots, a_n \in \mathbb{F}\},$$

i.e., the set of all possible linear combinations of elements from U .

- If $|U| = \infty$, $\text{span}(U)$ is the set of **all** possible linear combinations of **all** possible finite subsets of U .
- **Proposition 10:** Let $U \subseteq V$. Then $\text{span}(U)$ is a subspace of V .
- Let V be a VS, and let $W \subset V$. If $\text{span}(W) = V$, we say that W is a **spanning set** of V , or W **spans** V .
- We say that a vector space V is **finite dimensional** if there exists a finite spanning set. Notation: FDVS.
- **Linearly independent set:** Let V be a vector space and let $W = \{v_1, \dots, v_n\} \subset V$. We say that the set W is a set of linear independent vectors, if

$$a_1 v_1 + \dots + a_n v_n = \theta \Rightarrow a_i = 0, i = 1, \dots, n$$

Summary of Lecture 20

- Let V be a VS, and $U = \{v_1, \dots, v_n\}$.
- **Definition:** (Span of a set) We define the **Span of U** as

$$\text{span}(U) := \{a_1 v_1 + \dots + a_n v_n \mid \forall a_1, \dots, a_n \in \mathbb{F}\},$$

i.e., the set of all possible linear combinations of elements from U .

- If $|U| = \infty$, $\text{span}(U)$ is the set of **all** possible linear combinations of **all** possible finite subsets of U .
- **Proposition 10:** Let $U \subseteq V$. Then $\text{span}(U)$ is a subspace of V .
- Let V be a VS, and let $W \subset V$. If $\text{span}(W) = V$, we say that W is a **spanning set** of V , or W **spans** V .
- We say that a vector space V is **finite dimensional** if there exists a finite spanning set. Notation: FDVS.
- **Linearly independent set:** Let V be a vector space and let $W = \{v_1, \dots, v_n\} \subset V$. We say that the set W is a set of linear independent vectors, if

$$a_1 v_1 + \dots + a_n v_n = \theta \Rightarrow a_i = 0, i = 1, \dots, n$$

- What if $|W| = \infty$?

$$\text{Let } W_1 = \{v_1, \dots, v_5\}.$$

$$v_1 = \sum_{i=2}^5 v_i$$

$$W_1 \rightarrow \text{LD}.$$

$$W_2 = W_1 \cup \{v_6, \dots, v_{10}\}.$$

$$W_2 \rightarrow \text{LD}$$

$$W_3 = W_1 \cup \{p_i, i \in \mathbb{Z}\}.$$

$$\hookrightarrow \text{LD}$$

$$W, \quad |W| \equiv \infty$$

W is said to be LD if there is a finite subset of W which is LD.

● Example: $V = \mathcal{P}(\mathbb{R})$, $U = \{x^i, i = 0, 1, \dots\}$.

$$W = \{x^{k_i}, k_i \in \mathbb{N}, i = 1, \dots, m\}.$$

$$p(x) = a_1 x^{k_1} + a_2 x^{k_2} + \dots + a_m x^{k_m} = 0 \rightarrow \textcircled{1}$$

$$p(x_1) = a_1 x_1^{k_1} + \dots + a_m x_1^{k_m} = 0$$

$$\Rightarrow \boxed{a_1 = a_2 = \dots = a_m = 0}$$

Assume $k_j = \max_i \{k_i\}$.

$\Rightarrow \deg(p) = k_j \Rightarrow \# \text{ of roots of } p = k_j$

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● **Proof:** Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

① $\{w_1, u_1, \dots, u_n\}$ is LD

$$\text{span}(\{w_1, u_1, \dots, u_n\}) = V$$

$$u_j \in \text{span}(\{w_1, u_1, \dots, u_{j-1}, u_{j+1}, \dots, u_n\})$$

$$w_1 = \sum_{i=1}^n a_i u_i$$

$$\text{span}(\{w_1, u_1, \dots, u_{j-1}, u_{j+1}, \dots, u_n\}) = V$$

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD,

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) =$

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD i.e.,

$\exists u_k \in U, k \neq j, u_k \in \text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\})$.

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD i.e.,

$\exists u_k \in U, k \neq j, u_k \in \text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\})$.

● $\text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}) = V$

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD i.e.,

$\exists u_k \in U, k \neq j, u_k \in \text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\})$.

● $\text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}) = V$

● The number of elements in the set $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}$ remains n .

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD i.e.,

$\exists u_k \in U, k \neq j, u_k \in \text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\})$.

● $\text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}) = V$

● The number of elements in the set $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}$ remains n .

● Is it possible that $m > n$?

● **Proposition 11:** In a FDVS, the number of elements in a linearly independent set of vectors is always less than equal to the number of elements in a spanning set.

● Proof: Let V be a VS and $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_m\}$ be its subsets such that $\text{span}(U) = V$ and W is LI.

● $\{w_1, u_1, \dots, u_n\}$ is LD, i.e.,

$\exists u_j \in U, u_j \in \text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\})$.

● $\text{span}(\{w_1, u_i, i = 1, \dots, n, i \neq j\}) = V$

● $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j\}$ is LD i.e.,

$\exists u_k \in U, k \neq j, u_k \in \text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\})$.

● $\text{span}(\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}) = V$

● The number of elements in the set $\{w_1, w_2, u_i, i = 1, \dots, n, i \neq j, k\}$ remains n .

● Is it possible that $m > n$?

● If so, after n iterations, we will reach a contradiction:

$\text{span}(\{w_1, w_2, \dots, w_n\}) = V$

Basis of a Vector space

- **Definition:** (Basis) Let V be a vector space. A subset $\beta \subset V$ is said to be a **Basis** of V if (1) $\text{span}(\beta) = V$, and (2) β is a set of linearly independent vectors.

Basis of a Vector space

- **Definition:** (Basis) Let V be a vector space. A subset $\beta \subset V$ is said to be a **Basis** of V if (1) $\text{span}(\beta) = V$, and (2) β is a set of linearly independent vectors.
- Examples:

● **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.
- **Proposition 15:** Every FDVS has a basis.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.
- **Proposition 15:** Every FDVS has a basis.
- **Proposition 16:** A subset $U = \{u_1, \dots, u_n\}$ of VS V is a basis of V if and only if every $v \in V$ can be uniquely written as $v = a_1 u_1 + a_2 u_2 + \dots + a_n u_n, a_i \in \mathbb{F}, i = 1, \dots, n$.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.
- **Proposition 15:** Every FDVS has a basis.
- **Proposition 16:** A subset $U = \{u_1, \dots, u_n\}$ of VS V is a basis of V if and only if every $v \in V$ can be uniquely written as $v = a_1 u_1 + a_2 u_2 + \dots + a_n u_n, a_i \in \mathbb{F}, i = 1, \dots, n$.
- **Proposition 17:** Any set of basis vectors of a FDVS contains the same number of elements.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.
- **Proposition 15:** Every FDVS has a basis.
- **Proposition 16:** A subset $U = \{u_1, \dots, u_n\}$ of VS V is a basis of V if and only if every $v \in V$ can be uniquely written as $v = a_1 u_1 + a_2 u_2 + \dots + a_n u_n, a_i \in \mathbb{F}, i = 1, \dots, n$.
- **Proposition 17:** Any set of basis vectors of a FDVS contains the same number of elements.
- **Dimension of a Vector Space:** Let V be a FDVS. For any set of basis vectors β of V , we define the dimension of V as $\dim(V) := |\beta|$.

- **Proposition 13:** For a FDVS, every spanning set can be reduced to a basis.
- **Proposition 14:** For a FDVS, every LI set can be extended to a basis.
- **Proposition 15:** Every FDVS has a basis.
- **Proposition 16:** A subset $U = \{u_1, \dots, u_n\}$ of VS V is a basis of V if and only if every $v \in V$ can be uniquely written as $v = a_1 u_1 + a_2 u_2 + \dots + a_n u_n, a_i \in \mathbb{F}, i = 1, \dots, n$.
- **Proposition 17:** Any set of basis vectors of a FDVS contains the same number of elements.
- **Dimension of a Vector Space:** Let V be a FDVS. For any set of basis vectors β of V , we define the dimension of V as $\dim(V) := |\beta|$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) =$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) =$

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.
Similarly, $N(A^T) \oplus C(A) = \mathbb{R}^m$.

Some beautiful results

- **Theorem 3:** (Rank-Nullity Theorem). Let $A \in \mathbb{R}^{m \times n}$, with $\text{rank}(A) = r$. Then $r + \dim(N(A)) = n$.
- **Proposition 18:** Let U, W be subspaces of FDVS V . Then, $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
- **Theorem 4:** Let $A \in \mathbb{R}^{m \times n}$. $N(A) + C(A^T) = N(A) \oplus C(A^T) = \mathbb{R}^n$.
Similarly, $N(A^T) \oplus C(A) = \mathbb{R}^m$.

$$A: \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

