

# SC223 - Linear Algebra

Aditya Tatu

Lecture 14



August 28, 2023

# Vector Spaces

$$(\mathbb{R}, +, \times)$$

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$  and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

►  $(V, +)$  is an **Abelian group**:

►  $\forall x, y \in V, x + y \in V.$

►  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x.$

►  $\forall x \in V, \exists y \in V, x + y = y + x = \theta.$  We will denote  $y$  by  $-x.$

►  $\forall x, y, z \in V, (x + y) + z = x + (y + z).$

►  $\forall x, y \in V, x + y = y + x.$

► **Closure with respect to Scalar multiplication:**  $\cdot : \mathbb{F} \times V \rightarrow V.$

► **Scalar Multiplication identity:**  $\exists 1 \in \mathbb{F}$  such that  $1 \cdot v = v, \forall v \in V.$

► **Distributivity:**  $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v,$  and  $\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u.$

► **Compatibility of field and scalar multiplication:**

$\forall a, b \in \mathbb{F}, \forall u \in V, (a \times b) \cdot u = a \cdot (b \cdot u).$

$\mathbb{F}$   $\uparrow$   $\mathbb{F}$

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:
  - ▶  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:
  - ▶  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.
  - ▶  $(\mathbb{F} - \{0\}, \times)$  is an **Abelian group**. The multiplicative identity will be denoted by 1.

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:
  - ▶  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.
  - ▶  $(\mathbb{F} - \{0\}, \times)$  is an **Abelian group**. The multiplicative identity will be denoted by 1.
  - ▶ **Distributivity:**  $\forall a, b, c \in \mathbb{F}, (a +_F b) \times c = a \times c +_F b \times c$ .

# Examples of Fields

►  $(\mathbb{Z}_2, +_2, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$



## Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$

$$(a+bi) \times (c+di) =$$

$$\frac{(ac-bd) + (bc+ad)i}{= 0}$$

$$\forall (a+bi) \in \mathbb{C} - \{0\} = 0$$

$$(a+bi) \times 1 = a+bi \Rightarrow 1 \text{ is the } \times\text{-identity}$$

$$(a+bi) \times (c+di) = \boxed{1}.$$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.

$$\frac{p_1(x)}{q_1(x)} + \frac{p_2(x)}{q_2(x)}, \quad q_1 \neq 0, q_2 \neq 0.$$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple  $(V, +, \cdot)$  with field  $(\mathbb{F}, +_F, \times)$  satisfies all vector space axioms, we say that  $(V, +, \cdot)$  **forms a vector space over  $\mathbb{F}$** .

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple  $(V, +, \cdot)$  with field  $(\mathbb{F}, +_F, \times)$  satisfies all vector space axioms, we say that  $(V, +, \cdot)$  **forms a vector space over  $\mathbb{F}$** .
- Any element of the vector space  $(V, +, \cdot)$  will be referred to as a **vector**, and any element  $a \in \mathbb{F}$  will be referred to as a **scalar**.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .

## Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .

$$V = \mathbb{R}^n, \quad \mathbb{F} = \mathbb{R}.$$

$$\forall u, w \in V, \quad \underline{u + w} := \begin{pmatrix} u_1 + w_1 \\ \vdots \\ u_n + w_n \end{pmatrix}$$

$$\forall a \in \mathbb{R}, \forall u \in V, \quad a \cdot u := \begin{pmatrix} a \times u_1 \\ a \times u_2 \\ \vdots \\ a \times u_n \end{pmatrix}$$

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .





## Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{F}^n, +, \cdot)$  over  $\mathbb{F}$ .

$$(\mathbb{Z}_2^n, +, \cdot) \text{ over } \mathbb{Z}_2$$

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{F}^n, +, \cdot)$  over  $\mathbb{F}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{F}^n, +, \cdot)$  over  $\mathbb{F}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathcal{P}(\mathbb{R})$  is the set of all polynomials of one variable with real coefficients.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{F}^n, +, \cdot)$  over  $\mathbb{F}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathcal{P}(\mathbb{R})$  is the set of all polynomials of one variable with real coefficients.
- $(\mathbb{L}_2(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{L}_2(\mathbb{R})$  denotes the set of all square-integrable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

$$\forall f \in \mathbb{L}_2(\mathbb{R}), \int_{-\infty}^{\infty} (f(t))^2 dt < \infty.$$

# Properties of Vector Spaces

- **Proposition 1:** Every Vector space has a unique additive identity.

# Properties of Vector Spaces

- **Proposition 1:** Every Vector space has a unique additive identity.
- **Proposition 2:** Every vector in a vector space has a unique additive inverse.

# Properties of Vector Spaces

- **Proposition 1:** Every Vector space has a unique additive identity.
- **Proposition 2:** Every vector in a vector space has a unique additive inverse.
- **Proposition 3:**  $\forall v \in V, 0 \cdot v = \theta$

# Properties of Vector Spaces

- **Proposition 1:** Every Vector space has a unique additive identity.
- **Proposition 2:** Every vector in a vector space has a unique additive inverse.
- **Proposition 3:**  $\forall v \in V, 0 \cdot v = \theta$
- **Proposition 4:**  $\forall a \in \mathbb{F}, a \cdot \theta = \theta$ .



# Properties of Vector Spaces

- **Proposition 1:** Every Vector space has a unique additive identity.
- **Proposition 2:** Every vector in a vector space has a unique additive inverse.
- **Proposition 3:**  $\forall v \in V, 0 \cdot v = \theta$
- **Proposition 4:**  $\forall a \in \mathbb{F}, a \cdot \theta = \theta$ .
- **Proposition 5:**  $\forall v \in V, (-1) \cdot v = -v$ .