

SC223 - Linear Algebra

Aditya Tatu

Lecture 34



November 1, 2023

Summary of Lecture 33

- Matrix Diagonalization: Existence of an eigenbasis, say E , is equivalent to $A = E\Lambda E^{-1}$, where Λ is the diagonal matrix.
- **Proposition 20:** Let $\lambda_1, \dots, \lambda_m$ be distinct eigenvalues of $T \in \mathcal{L}(U)$. Then the eigenvectors v_1, \dots, v_m associated with these eigenvalues are linearly independent.

Results on Eigenvectors and Eigenvalues

● **Proposition 21:** For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if $\sum_{i=1}^m AM(\lambda_i) = n$, and $GM(\lambda_i) = AM(\lambda_i)$, $i = 1, \dots, m$, then T is diagonalizable.

Let $\lambda \in \mathbb{F}$ s.t. $C_T(\lambda) = 0$ & $C_T(x) = (x - \lambda)^k p(x)$ with $p(x) \neq 0$.

$AM(\lambda) = k$; Assume $G.M(\lambda) = k+1 = \dim N(T - \lambda I)$

Let $\{v_1, \dots, v_{k+1}\}$ be LI vectors from $N(T - \lambda I)$.

$$Tv_i = \lambda v_i, i = 1, \dots, k+1.$$

Let $\beta = \{v_1, \dots, v_{k+1}, u_1, \dots, u_{n-k-1}\}$ be a basis of U .

$$[T]_{\beta}^{\beta} = \begin{bmatrix} \lambda & 0 & 0 & \star \\ 0 & \lambda & 0 & \star \\ 0 & 0 & 0 & \star \\ \hline (k+1) \times (k+1) & & & \star \\ 0 & 0 & 0 & \star \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \star \end{bmatrix} = \begin{bmatrix} \lambda I_{k+1} & A_{(k+1) \times (n-k-1)} \\ 0 & B_{(n-k-1) \times (n-k-1)} \end{bmatrix}$$

$$[T]_{\beta}^{\beta} = \begin{bmatrix} \lambda I_{k+1} & A \\ 0 & B \end{bmatrix}$$

$$C_{[T]_{\beta}^{\beta}}(x) = (x - \lambda)^{k+1} p(x)$$

$$[T]_{\beta}^{\beta} - x I_n = \begin{bmatrix} (x - \lambda) I_{k+1} & A \\ 0 & B - x I_{(n-k-1)} \end{bmatrix}$$

$$\det([T]_{\beta}^{\beta} - x I_n) = (x - \lambda)^{k+1} \det(B - x I_{n-k-1})$$

$$AM(\lambda) \geq k+1.$$

Results on Eigenvectors and Eigenvalues

- **Proposition 21:** For $T \in \mathcal{L}(U)$, $\dim(U) = n$, if $\sum_{i=1}^m AM(\lambda_i) = n$, and $GM(\lambda_i) = AM(\lambda_i)$, $i = 1, \dots, m$, then T is diagonalizable.
- **Proposition 22:** For $T \in \mathcal{L}(U)$, $\dim(U) = n$, for any eigenvalue λ of T , $GM(\lambda) \leq AM(\lambda)$

Applications

- Coupled Differential Equations:

Applications

- Coupled Differential Equations:
 - ▶ Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Applications

- Coupled Differential Equations:

- ▶ Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Assume that the rate of change of predator population is $a \in \mathbb{R}$ times the current predator population and $b \in \mathbb{R}$ times that of the prey population.

Applications

- Coupled Differential Equations:

► Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Assume that the rate of change of predator population is $a \in \mathbb{R}$ times the current predator population and $b \in \mathbb{R}$ times that of the prey population. Similarly, the rate of change of prey population is $c \in \mathbb{R}$ times the current predator population and $d \in \mathbb{R}$ times the prey population, compute the predator-prey population at any time t , given some initial population.

$$\frac{d}{dt} x(t) = ax(t) + by(t)$$

$$\frac{d}{dt} y(t) = cx(t) + dy(t).$$

Applications

- Coupled Differential Equations:

- ▶ Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Assume that the rate of change of predator population is $a \in \mathbb{R}$ times the current predator population and $b \in \mathbb{R}$ times that of the prey population. Similarly, the rate of change of prey population is $c \in \mathbb{R}$ times the current predator population and $d \in \mathbb{R}$ times the prey population, compute the predator-prey population at any time t , given some initial population.

$$\frac{d}{dt} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

Applications

- Coupled Differential Equations:

► Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Assume that the rate of change of predator population is $a \in \mathbb{R}$ times the current predator population and $b \in \mathbb{R}$ times that of the prey population. Similarly, the rate of change of prey population is $c \in \mathbb{R}$ times the current predator population and $d \in \mathbb{R}$ times the prey population, compute the predator-prey population at any time t , given some initial population.

$$\frac{d}{dt} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

$$\frac{d}{dt} X(t) = A X(t)$$

$$\frac{d}{dt} x(t) = a x(t)$$

$$x(t) = e^{at} x(0)$$

Applications

- Coupled Differential Equations:

► Let $x(t), y(t)$ denote the population of predators and prey in an environment at any time t .

Assume that the rate of change of predator population is $a \in \mathbb{R}$ times the current predator population and $b \in \mathbb{R}$ times that of the prey population. Similarly, the rate of change of prey population is $c \in \mathbb{R}$ times the current predator population and $d \in \mathbb{R}$ times the prey population, compute the predator-prey population at any time t , given some initial population.

$$\frac{d}{dt} e^{At} = A e^{At}$$

$$\frac{d}{dt} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

$$\frac{d}{dt} X(t) = A X(t)$$

$$X(t) = \exp(At) X(0)$$

$$e^{at} = 1 + at + \frac{a^2 t^2}{2!} + \dots$$

$$e^{At} = e^{\begin{bmatrix} at & bt \\ ct & dt \end{bmatrix}} \quad e^{At} := I_n + At + \frac{A^2 t^2}{2!} + \frac{A^3 t^3}{3!} + \dots$$

$$e^{At} = I + At + \frac{A^2 t^2}{2!} + \frac{A^3 t^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{A^n t^n}{n!}$$

$$\text{If } A = E \Lambda E^{-1}$$

$$e^{At} = E I E^{-1} + E A E^{-1} t + E A^2 E^{-1} \frac{t^2}{2!} + E A^3 E^{-1} \frac{t^3}{3!} + \dots = \sum_{n=0}^{\infty} E \Lambda^n E^{-1} \frac{t^n}{n!}$$

$$e^{At} = E \left(\sum_{n=0}^{\infty} \frac{\Lambda^n t^n}{n!} \right) E^{-1} = E e^{\Lambda t} E^{-1}$$

$$\Lambda = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \quad e^{\Lambda t} = \begin{bmatrix} e^{\lambda_1 t} & & 0 \\ & e^{\lambda_2 t} & \\ 0 & & \ddots & \\ & & & e^{\lambda_n t} \end{bmatrix}$$

$$\frac{d}{dt} X(t) = A X(t) \Rightarrow \frac{d}{dt} X(t) = E \Lambda E^{-1} X(t)$$

$$\frac{d}{dt} \underbrace{(E^{-1} X(t))}_{Y(t)} = \Lambda (E^{-1} X(t))$$

$$\frac{d}{dt} Y(t) = \Lambda Y(t)$$

Applications

- Coupled Differential Equations:




Figure: Source: Differential Equations and Linear Algebra, by GB Gustafson.

$$\begin{aligned}i_1' &= - \left(\frac{R_2}{L_1} \right) i_2 - \left(\frac{R_3}{L_1} \right) i_3, \\i_2' &= - \left(\frac{R_2}{L_2} + \frac{R_2}{L_1} \right) i_2 + \left(\frac{R_1}{L_2} - \frac{R_3}{L_1} \right) i_3, \\i_3' &= \left(\frac{R_1}{L_3} - \frac{R_2}{L_1} \right) i_2 - \left(\frac{R_1}{L_3} + \frac{R_3}{L_1} + \frac{R_3}{L_3} \right) i_3\end{aligned}$$

Applications

- Coupled Differential Equations:
- Epidemics Modeling:
 - ▶ Population can be divided into: Susceptible/Healthy (S), Infected (I), and Dead (D) classes.
 - ▶ Susceptible $\xrightarrow{-a}$ Infected \xrightarrow{b} Susceptible: $\frac{d}{dt}S = -aS(t) + rI(t)$
 - ▶ Infected $\xrightarrow{-d}$ Dead: $\frac{d}{dt}D = dI(t)$
 - ▶ Infected: $\frac{d}{dt}I = aS(t) - dI(t) - rI(t)$

End of Class-

Applications

- Page Rank/Importance:

Applications


- Page Rank/Importance:
 - ▶ Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.

Applications

- Page Rank/Importance:
 - ▶ Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.
 - ▶ Let $x_i, i = 1, \dots, 4$ denote the importance of these pages, or probability that a surfer will visit page P_i . (to be computed).

Applications

- Page Rank/Importance:
 - Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.
 - Let $x_i, i = 1, \dots, 4$ denote the importance of these pages, or probability that a surfer will visit page P_i . (to be computed).



Applications

- Page Rank/Importance:

- Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.
- Let $x_i, i = 1, \dots, 4$ denote the importance of these pages, or probability that a surfer will visit page P_i . (to be computed).




Figure: Model: If P_2 has links to P_3 and P_4 , a surfer go to these pages with equal probability. Source: pi.math.cornell.edu

- $\lim_{n \rightarrow \infty} A^n x_0 = y,$

Applications

- Page Rank/Importance:

- Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.
- Let $x_i, i = 1, \dots, 4$ denote the importance of these pages, or probability that a surfer will visit page P_i . (to be computed).




Figure: Model: If P_2 has links to P_3 and P_4 , a surfer go to these pages with equal probability. Source: pi.math.cornell.edu

- $\lim_{n \rightarrow \infty} A^n x_0 = y, Ay = y,$

Applications

- Page Rank/Importance:

- Let $P_i, i = 1, \dots, 4$ denote the 4 pages that are relevant to a search.
- Let $x_i, i = 1, \dots, 4$ denote the importance of these pages, or probability that a surfer will visit page P_i . (to be computed).




Figure: Model: If P_2 has links to P_3 and P_4 , a surfer go to these pages with equal probability. Source: pi.math.cornell.edu

- $\lim_{n \rightarrow \infty} A^n x_0 = y, Ay = y, y = [0.38, 0.12, 0.29, 0.19]^T$.

Projection

- Let V be a FDVS, and let U, W be subspaces of V such that $V = U \oplus W$.

Projection

- Let V be a FDVS, and let U, W be subspaces of V such that $V = U \oplus W$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V, v = u + w, u \in U, w \in W, P_U(v) = u$.
- Such a linear operator is called *Projection* on U along W , or on U with nullspace W .

Projection

- Let V be a FDVS, and let U, W be subspaces of V such that $V = U \oplus W$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V, v = u + w, u \in U, w \in W, P_U(v) = u$.
- Such a linear operator is called *Projection* on U along W , or on U with nullspace W .
- Properties:

Inner Products and Norms

- How do we measure length and angles between vectors in a Vector space?

Inner Products and Norms

- How do we measure length and angles between vectors in a Vector space?
- Typically, length of a vector in \mathbb{R}^2 is defined as $\|(x, y)\| = \sqrt{x^2 + y^2}$.

Inner Products and Norms

- How do we measure length and angles between vectors in a Vector space?
- Typically, length of a vector in \mathbb{R}^2 is defined as $\|(x, y)\| = \sqrt{x^2 + y^2}$.
- This is called the *Euclidean length* or *Euclidean norm* of the vector (x, y) .

Inner Products and Norms

- How do we measure length and angles between vectors in a Vector space?
- Typically, length of a vector in \mathbb{R}^2 is defined as $\|(x, y)\| = \sqrt{x^2 + y^2}$.
- This is called the *Euclidean length* or *Euclidean norm* of the vector (x, y) .
- Is this the only way to define length?

Inner Products and Norms

- How do we measure length and angles between vectors in a Vector space?
- Typically, length of a vector in \mathbb{R}^2 is defined as $\|(x, y)\| = \sqrt{x^2 + y^2}$.
- This is called the *Euclidean length* or *Euclidean norm* of the vector (x, y) .
- Is this the only way to define length?
- What are the necessary conditions for a function on vector space for it be called *length*?

Normed Vector space

- **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

Normed Vector space

● **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

1. Positive definiteness: $\|x\| \geq 0, \forall x \in V$ and $\|x\| = 0 \Leftrightarrow x = \theta$.

Normed Vector space

● **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

1. Positive definiteness: $\|x\| \geq 0, \forall x \in V$ and $\|x\| = 0 \Leftrightarrow x = \theta$.
2. Absolute homogeneity: $\forall x \in V, \forall a \in \mathbb{F}, \|a \cdot x\| = |a| \|x\|$.

Normed Vector space

● **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

1. Positive definiteness: $\|x\| \geq 0, \forall x \in V$ and $\|x\| = 0 \Leftrightarrow x = \theta$.
2. Absolute homogeneity: $\forall x \in V, \forall a \in \mathbb{F}, \|a \cdot x\| = |a| \|x\|$.
3. Triangular inequality: $\forall x, y \in V, \|x + y\| \leq \|x\| + \|y\|$.

Normed Vector space

● **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

1. Positive definiteness: $\|x\| \geq 0, \forall x \in V$ and $\|x\| = 0 \Leftrightarrow x = \theta$.
2. Absolute homogeneity: $\forall x \in V, \forall a \in \mathbb{F}, \|a \cdot x\| = |a| \|x\|$.
3. Triangular inequality: $\forall x, y \in V, \|x + y\| \leq \|x\| + \|y\|$.

● A vector space V with a valid norm $\|\cdot\|$ is called a **Normed vector space** and is denoted by $(V, \|\cdot\|)$.

Normed Vector space

● **Definition:** (Normed Vector Space) A normed vector space (NVS) is a vector space $(V, +, \cdot)$ over either \mathbb{R} or \mathbb{C} with a **norm**, a function $\|\cdot\| : V \rightarrow \mathbb{R}$ which satisfies the following properties:

1. Positive definiteness: $\|x\| \geq 0, \forall x \in V$ and $\|x\| = 0 \Leftrightarrow x = \theta$.
2. Absolute homogeneity: $\forall x \in V, \forall a \in \mathbb{F}, \|a \cdot x\| = |a| \|x\|$.
3. Triangular inequality: $\forall x, y \in V, \|x + y\| \leq \|x\| + \|y\|$.

● A vector space V with a valid norm $\|\cdot\|$ is called a **Normed vector space** and is denoted by $(V, \|\cdot\|)$.

● Also note that given a NVS $(V, \|\cdot\|)$, we can define distance between two vectors x and y as $d(x, y) := \|x - y\|$. Such a distance or metric is called the **induced metric**.

Examples of NVS

- 1-norm on \mathbb{R}^n : $\|x\| = \sum_{i=1}^n |x_i|$. This is denoted by $\|x\|_1$.

Examples of NVS

- 1-norm on \mathbb{R}^n : $\|x\| = \sum_{i=1}^n |x_i|$. This is denoted by $\|x\|_1$.
- max or the sup norm on \mathbb{R}^n : $\|x\| = \max_{i=1}^n \{|x_i|\}$. This is denoted by $\|x\|_{\text{sup}}$ or $\|x\|_{\text{max}}$.

Examples of NVS

- 1-norm on \mathbb{R}^n : $\|x\| = \sum_{i=1}^n |x_i|$. This is denoted by $\|x\|_1$.
- max or the sup norm on \mathbb{R}^n : $\|x\| = \max_{i=1}^n \{|x_i|\}$. This is denoted by $\|x\|_{\text{sup}}$ or $\|x\|_{\text{max}}$.
- L_2 norm on $\mathcal{P}_n([-1, 1])$: $\|x\|_{L_2} = \sqrt{\int_{-1}^1 (x(t))^2 dt}$.