

# SC223 - Linear Algebra

Aditya Tatu

Lecture 13

•



August 25, 2023

# Example of Structure in Math

- The examples with (a) Symmetries of a rectangle, (b) set of all two-bits with bitwise addition modulo-2, and others can be abstracted as  $S = \{e, a, b, a \cdot b\}$  and operation  $\cdot$  as

$$a^2 = e \Rightarrow$$

$\cdot$	$e$	$a$	$b$	$a \cdot b$
$e$	$e$	$a$	$b$	$a \cdot b$
$a$	$a$	$e$	$a \cdot b$	$b$
$b$	$b$	$a \cdot b$	$e$	$a$
$a \cdot b$	$a \cdot b$	$b$	$a$	$e$

1. Existence of identity
2. Closure
3. Inverse.

- The examples (a)  $\{1, 3, 5, 7\}$  with multiplication modulo-8, (b)  $\{1, i, -1, -i\}$  with complex number multiplication can be abstracted as  $S = \{e, a, b, a \cdot b\}$  with the operation  $\cdot$  and

$$a^2 = b.$$

$\cdot$	$e$	$a$	$b$	$a \cdot b$
$e$	$e$	$a$	$b$	$a \cdot b$
$a$	$a$	$b$	$a \cdot b$	$e$
$b$	$b$	$a \cdot b$	$e$	$a$
$a \cdot b$	$a \cdot b$	$e$	$a$	$b$

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set  $G$  with a *binary operation*, denoted by  $\cdot$ , that satisfy the following axioms:
  - **Closure:**  $\forall a, b \in G, a \cdot b \in G$ .

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set  $G$  with a *binary operation*, denoted by  $\cdot$ , that satisfy the following axioms:
  - **Closure:**  $\forall a, b \in G, a \cdot b \in G$ .
  - **Identity:** There exists an element  $e \in G$  such that  $\forall a \in G, a \cdot e = e \cdot a = a$ .

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set  $G$  with a *binary operation*, denoted by  $\cdot$ , that satisfy the following axioms:
  - **Closure:**  $\forall a, b \in G, a \cdot b \in G$ .
  - **Identity:** There exists an element  $e \in G$  such that  $\forall a \in G$ ,  
 $a \cdot e = e \cdot a = e$
  - **Inverse:** For each  $a \in G$ , there exists an element  $a^{-1} \in G$  such that  
 $a \cdot a^{-1} = a^{-1} \cdot a = e$ . The element  $a^{-1}$  is called the *inverse* of  $a$ .

$$a \cdot b = b \cdot a = e, \quad b = a^{-1}$$

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set  $G$  with a *binary operation*, denoted by  $\cdot$ , that satisfy the following axioms:
  - **Closure:**  $\forall a, b \in G, a \cdot b \in G$ .
  - **Identity:** There exists an element  $e \in G$  such that  $\forall a \in G, a \cdot e = e \cdot a = e$ .
  - **Inverse:** For each  $a \in G$ , there exists an element  $a^{-1} \in G$  such that  $a \cdot a^{-1} = a^{-1} \cdot a = e$ . The element  $a^{-1}$  is called the *inverse* of  $a$ .
  - **Associativity:**  $\forall a, b, c \in G, (a \cdot b) \cdot c = a \cdot (b \cdot c)$

# Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set  $G$  with a *binary operation*, denoted by  $\cdot$ , that satisfy the following axioms:
  - **Closure:**  $\forall a, b \in G, a \cdot b \in G$ .
  - **Identity:** There exists an element  $e \in G$  such that  $\forall a \in G, a \cdot e = e \cdot a = e$ .
  - **Inverse:** For each  $a \in G$ , there exists an element  $a^{-1} \in G$  such that  $a \cdot a^{-1} = a^{-1} \cdot a = e$ . The element  $a^{-1}$  is called the *inverse* of  $a$ .
  - **Associativity:**  $\forall a, b, c \in G, (a \cdot b) \cdot c = a \cdot (b \cdot c)$
- We denote the group by the tuple  $(G, \cdot)$ .
- If  $\cdot$  is commutative on  $G$ , then we call  $(G, \cdot)$  a commutative Group or Abelian Group



# Vector Spaces

- We have seen linear combinations of elements from

# Vector Spaces

- We have seen linear combinations of elements from

- ▶  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$



$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$



$\forall x, y \in \mathbb{R}^{m \times n},$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$

►

$$\forall x, y \in \mathbb{R}^\infty,$$

$$x \in \mathbb{R}^\infty, x: \mathbb{Z} \rightarrow \mathbb{R}.$$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$

►

$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

$$x: \mathbb{Z} \rightarrow \mathbb{R}$$



# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$



$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

►  $\forall f, g \in \{h : \mathbb{R} \rightarrow \mathbb{R}\},$

# Vector Spaces

- We have seen linear combinations of elements from

►  $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$



$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

►  $\forall f, g \in \{h : \mathbb{R} \rightarrow \mathbb{R}\}, \forall a, b \in \mathbb{R}, a \cdot f + b \cdot g, (a \cdot f + b \cdot g)(t) =$   
 $a \cdot f(t) + b \cdot g(t), \forall t \in \mathbb{R}.$

# Vector Spaces

- **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

 $\mathbb{R}^n$  $\mathbb{R}$

# Vector Spaces

- **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:
  - ▶  $(V, +)$  is an **Abelian group**:

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

- ▶  $\forall x, y \in V, x + y \in V$
- ▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$
- ▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta.$

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .



# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

▶  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

▶  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

▶  $\forall x, y \in V, x + y = y + x$ .

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

▶  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

▶  $\forall x, y \in V, x + y = y + x$ .

▶ **Closure with respect to Scalar multiplication:**  $\cdot : \mathbb{F} \times V \rightarrow V$ .

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

▶  $(V, +)$  is an **Abelian group**:

▶  $\forall x, y \in V, x + y \in V$

▶  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

▶  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

▶  $\forall x, y \in V, x + y = y + x$ .

▶ **Closure with respect to Scalar multiplication:**  $\cdot : \mathbb{F} \times V \rightarrow V$ .

▶ **Scalar Multiplication identity:**  $\exists 1 \in \mathbb{F}$  such that  $1 \cdot v = v, \forall v \in V$ .

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

►  $(V, +)$  is an **Abelian group**:

►  $\forall x, y \in V, x + y \in V$

►  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

►  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

►  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

►  $\forall x, y \in V, x + y = y + x$ .

► **Closure with respect to Scalar multiplication:**  $\cdot : \mathbb{F} \times V \rightarrow V$ .

► **Scalar Multiplication identity:**  $\exists 1 \in \mathbb{F}$  such that  $1 \cdot v = v, \forall v \in V$ .

► **Distributivity:**  $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$ , and

$\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$ .

# Vector Spaces

● **Definition:** A Vector space is a set  $V$  with a **field**  $(\mathbb{F}, +_F, \times)$ , and two binary operations, vector addition  $+$  and scalar multiplication  $\cdot$  that satisfy the following axioms:

►  $(V, +)$  is an **Abelian group**:

►  $\forall x, y \in V, x + y \in V$

►  $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

►  $\forall x \in V, \exists y \in V, x + y = y + x = \theta$ . We will denote  $y$  by  $-x$ .

►  $\forall x, y, z \in V, (x + y) + z = x + (y + z)$ .

►  $\forall x, y \in V, x + y = y + x$ .

► **Closure with respect to Scalar multiplication:**  $\cdot : \mathbb{F} \times V \rightarrow V$ .

► **Scalar Multiplication identity:**  $\exists 1 \in \mathbb{F}$  such that  $1 \cdot v = v, \forall v \in V$ .

► **Distributivity:**  $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$ , and

$\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$ .

► **Compatibility of field and scalar multiplication:**

$\forall a, b \in \mathbb{F}, \forall u \in V, (a \times b) \cdot u = a \cdot (b \cdot u)$ .

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:

# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:
  - ▶  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.



# Field

- **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:
  - ▶  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.
  - ▶  $(\mathbb{F} - \{0\}, \times)$  is an **Abelian group**. The multiplicative identity will be denoted by 1.

# Field

● **Definition:**(Field). A field is a set  $\mathbb{F}$  with two binary operations, addition  $+_F$  and multiplication  $\times$  that satisfy the following axioms:

►  $(\mathbb{F}, +_F)$  is an **Abelian group**. The additive identity will be denoted by 0.

►  $(\mathbb{F} - \{0\}, \times)$  is an **Abelian group**. The multiplicative identity will be denoted by 1.

► **Distributivity:**

$$\forall a, b, c \in \mathbb{F}, (a +_F b) \times c = a \times c +_F b \times c, a \times (b +_F c) = a \times b +_F a \times c$$

# Examples of Fields

►  $(\mathbb{Z}_2, +_2, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.

# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple  $(V, +, \cdot)$  with field  $(\mathbb{F}, +_F, \times)$  satisfies all vector space axioms, we say that  $(V, +, \cdot)$  forms a vector space over  $\mathbb{F}$ .



# Examples of Fields

- ▶  $(\mathbb{Z}_2, +_2, \times)$
- ▶  $(\mathbb{R}, +, \times)$
- ▶  $(\mathbb{C}, +, \times)$
- ▶  $(\mathbb{Q}, +, \times)$
- ▶  $(\mathbb{R}[x], +, \times)$ , where  $\mathbb{R}[x]$  is the set of all rational polynomials of the form  $\frac{p(x)}{q(x)}$ , with  $q \neq 0$ , and  $p$  and  $q$  are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple  $(V, +, \cdot)$  with field  $(\mathbb{F}, +_F, \times)$  satisfies all vector space axioms, we say that  $(V, +, \cdot)$  forms a vector space over  $\mathbb{F}$ .
- Any element of the vector space  $(V, +, \cdot)$  will be referred to as a **vector**, and any element  $a \in \mathbb{F}$  will be referred to as a **scalar**.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathcal{P}(\mathbb{R})$  is the set of all polynomials of one variable with real coefficients.

# Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{R}^n, +, \cdot)$  over  $\mathbb{R}$ .
- $(\mathbb{C}^n, +, \cdot)$  over  $\mathbb{C}$ .
- $(\mathbb{R}^\infty, +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{R}^\infty$  is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathcal{P}(\mathbb{R})$  is the set of all polynomials of one variable with real coefficients.
- $(\mathbb{L}_2(\mathbb{R}), +, \cdot)$  over  $\mathbb{R}$ , where  $\mathbb{L}_2(\mathbb{R})$  denotes the set of all square-integrable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ .