

SC223 - Linear Algebra

Aditya Tatu

Lecture 12



August 23, 2023

Structure


- What is *structure*?

Structure

- What is *structure*? Structure is the arrangement and relation between parts of an object, without getting into the particulars of an example or an instance.

Example of Structure in Math

- Consider the following rectangle.




- What transformations leave the rectangle (not the vertices) unchanged?

$$S = \{I, R_{180^\circ}, F_h, F_v\} \text{ with } \circ$$
$$(S, \circ), \quad \circ: S \times S \rightarrow ?$$

Example of Structure in Math


- Consider the following rectangle.



- What transformations leave the rectangle (not the vertices) unchanged?
- $S_r = \{I, F_h, F_v, R\}$.

Example of Structure in Math

- Consider the following rectangle.



- What transformations leave the rectangle (not the vertices) unchanged?
- $S_r = \{I, F_h, F_v, R\}$. With Composition \circ , we have (S_r, \circ) :

Cayley Table

\circ	I	F_h	F_v	R
I	I	F_h	F_v	R
F_h	F_h	I	R	F_v
F_v	F_v	R	I	F_h
R	R	F_v	F_h	I

\rightarrow Identity

- Consider the set $S = \{00, 01, 10, 11\}$

- Consider the set $S = \{00, 01, 10, 11\}$ with bitwise addition modulo-2, $+_2$.

- Consider the set $S = \{00, 01, 10, 11\}$ with bitwise addition modulo-2, $+_2$.

Identity ←

$+_2$	00	01	10	11
00	00	01	10	11
01	01	00	11	10
10	10	11	00	01
11	11	10	01	00

$$a \in S, b \in S, a \circ b = \text{Identity}.$$

1. S is closed
2. Identity
3. Inverse.

- Compare the two:

\circ	I	F_h	F_v	R
I	I	F_h	F_v	R
F_h	F_h	I	R	F_v
F_v	F_v	R	I	F_h
R	R	F_v	F_h	I

$+_2$	00	01	10	11
00	00	01	10	11
01	01	00	11	10
10	10	11	00	01
11	11	10	01	00

- Compare the two:

\circ	I	F_h	F_v	R
I	I	F_h	F_v	R
F_h	F_h	I	R	F_v
F_v	F_v	R	I	F_h
R	R	F_v	F_h	I

$+_2$	00	01	10	11
00	00	01	10	11
01	01	00	11	10
10	10	11	00	01
11	11	10	01	00

- Consider the set of integers $\{1, 3, 5, 7\}$ with multiplication modulo-8.

	1	3	5	7
1	1	3	5	7
3	3	1	7	5
5	5	7	1	3
7	7	5	3	1

- Compare the two:

\circ	I	F_h	F_v	R
I	I	F_h	F_v	R
F_h	F_h	I	R	F_v
F_v	F_v	R	I	F_h
R	R	F_v	F_h	I

$+_2$	00	01	10	11
00	00	01	10	11
01	01	00	11	10
10	10	11	00	01
11	11	10	01	00

- Consider the set of integers $\{1, 3, 5, 7\}$ with multiplication modulo-8.

- Consider the set of matrices

$\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right\}$, with matrix multiplication.

- Consider the set $\{e, a, b, a \cdot b\}$ with an operation \cdot .

\cdot	e	a	b	$a \cdot b$
e	e	a	b	$a \cdot b$
a	a	e	$a \cdot b$	b
b	b	$a \cdot b$	e	a
$a \cdot b$	$a \cdot b$	b	a	e

	e	a	b	ab
e	e	a	b	ab
a	a	e	ab	b
b	b			
ab	ab			

$$ab = b$$

$$ab = ab$$

$$a \cdot a = ?$$

$$a^2 = b$$

$$a^2 = a \times$$

$$a^2 = e \rightarrow \textcircled{1}$$

$$a^2 = b \rightarrow \textcircled{2}$$

$$a^2 = ab \times$$

- Consider the set $\{e, a, b, a \cdot b\}$ with an operation \cdot

\cdot	e	a	b	$a \cdot b$
e	e	a	b	$a \cdot b$
a	a	e	$a \cdot b$	b
b	b	$a \cdot b$	e	a
$a \cdot b$	$a \cdot b$	b	a	e

- Note that in these examples: $a^2 = b^2 = (ab)^2 = e$.

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4,
 $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4, $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$ with addition modulo-4.

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4, $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$ with addition modulo-4.

	[0]	[1]	[2]	[3]
[0]	[0]	[1]	[2]	[3]
[1]	[1]	[2]	[3]	[0]
[2]	[2]	[3]	[0]	[1]
[3]	[3]	[0]	[1]	[2]

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4, $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$ with addition modulo-4.

	[0]	[1]	[2]	[3]
[0]	[0]	[1]	[2]	[3]
[1]	[1]	[2]	[3]	[0]
[2]	[2]	[3]	[0]	[1]
[3]	[3]	[0]	[1]	[2]

- Consider the set of complex numbers $C = \{1, i, -1, i\}$ with the usual complex number multiplication \times .

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4, $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$ with addition modulo-4.

	[0]	[1]	[2]	[3]
[0]	[0]	[1]	[2]	[3]
[1]	[1]	[2]	[3]	[0]
[2]	[2]	[3]	[0]	[1]
[3]	[3]	[0]	[1]	[2]

- Consider the set of complex numbers $C = \{1, i, -1, i\}$ with the usual complex number multiplication \times .

\times	1	i	-1	$-i$
1	1	i	-1	$-i$
i	i	-1	$-i$	1
-1	-1	$-i$	1	i
$-i$	$-i$	1	i	-1

- Consider the set \mathbb{Z}_4 of integers equivalent modulo-4, $\mathbb{Z}_4 = \{[0], [1], [2], [3]\}$ with addition modulo-4.

	[0]	[1]	[2]	[3]
[0]	[0]	[1]	[2]	[3]
[1]	[1]	[2]	[3]	[0]
[2]	[2]	[3]	[0]	[1]
[3]	[3]	[0]	[1]	[2]

- Consider the set of complex numbers $C = \{1, i, -1, i\}$ with the usual complex number multiplication \times .

\times	1	i	-1	$-i$
1	1	i	-1	$-i$
i	i	-1	$-i$	1
-1	-1	$-i$	1	i
$-i$	$-i$	1	i	-1

- These examples can be written using the set $S = \{e, a, b, a \cdot b\}$ and operation \cdot as

\cdot	e	a	b	$a \cdot b$
e	e	a	b	$a \cdot b$
a	a	b	$a \cdot b$	e
b	b	$a \cdot b$	e	a
$a \cdot b$	$a \cdot b$	e	a	b

- These examples can be written using the set $S = \{e, a, b, a \cdot b\}$ and operation \cdot as

\cdot	e	a	b	$a \cdot b$
e	e	a	b	$a \cdot b$
a	a	b	$a \cdot b$	e
b	b	$a \cdot b$	e	a
$a \cdot b$	$a \cdot b$	e	a	b

- The first few examples can be abstracted as $S = \{e, a, b, a \cdot b\}$ with the operation \cdot and

\cdot	e	a	b	$a \cdot b$
e	e	a	b	$a \cdot b$
a	a	e	$a \cdot b$	b
b	b	$a \cdot b$	e	a
$a \cdot b$	$a \cdot b$	b	a	e

Group

- All the above examples are examples of an *algebraic* structure called **Group**.

Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set G with a *binary operation*, denoted by \cdot , that satisfy the following axioms:
 - **Closure:** $\forall a, b \in G, a \cdot b \in G$.

Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set G with a *binary operation*, denoted by \cdot , that satisfy the following axioms:
 - **Closure:** $\forall a, b \in G, a \cdot b \in G$.
 - **Identity:** There exists an element $e \in G$ such that $\forall a \in G, a \cdot e = e \cdot a = a$.

Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set G with a *binary operation*, denoted by \cdot , that satisfy the following axioms:
 - **Closure:** $\forall a, b \in G, a \cdot b \in G$.
 - **Identity:** There exists an element $e \in G$ such that $\forall a \in G$, $a \cdot e = e \cdot a = e$.
 - **Inverse:** For each $a \in G$, there exists an element $a^{-1} \in G$ such that $a \cdot a^{-1} = a^{-1} \cdot a = e$. The element a^{-1} is called the *inverse* of a .

Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set G with a *binary operation*, denoted by \cdot , that satisfy the following axioms:
 - **Closure:** $\forall a, b \in G, a \cdot b \in G$.
 - **Identity:** There exists an element $e \in G$ such that $\forall a \in G, a \cdot e = e \cdot a = e$.
 - **Inverse:** For each $a \in G$, there exists an element $a^{-1} \in G$ such that $a \cdot a^{-1} = a^{-1} \cdot a = e$. The element a^{-1} is called the *inverse* of a .
 - **Associativity:** $\forall a, b, c \in G, (a \cdot b) \cdot c = a \cdot (b \cdot c)$

Group

- All the above examples are examples of an *algebraic* structure called **Group**.
- A **Group** is a non-empty set G with a *binary operation*, denoted by \cdot , that satisfy the following axioms:
 - **Closure:** $\forall a, b \in G, a \cdot b \in G$.
 - **Identity:** There exists an element $e \in G$ such that $\forall a \in G, a \cdot e = e \cdot a = e$.
 - **Inverse:** For each $a \in G$, there exists an element $a^{-1} \in G$ such that $a \cdot a^{-1} = a^{-1} \cdot a = e$. The element a^{-1} is called the *inverse* of a .
 - **Associativity:** $\forall a, b, c \in G, (a \cdot b) \cdot c = a \cdot (b \cdot c)$
- We denote the group by the tuple (G, \cdot) .

Vector Spaces

- We have seen linear combinations of elements from

Vector Spaces

- We have seen linear combinations of elements from

- ▶ $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$

►

$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$



$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$



$\forall x, y \in \mathbb{R}^{m \times n},$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$

►

$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$

►

$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$

►

$$\forall x, y \in \mathbb{R}^\infty,$$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$



$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$



$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

► $\forall f, g \in \{h : \mathbb{R} \rightarrow \mathbb{R}\},$

Vector Spaces

- We have seen linear combinations of elements from

► $\forall x, y \in \mathbb{R}^2, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \end{bmatrix}$



$$\forall x, y \in \mathbb{R}^3, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ ax_2 + by_2 \\ ax_3 + by_3 \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^n, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := a \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + b \cdot \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} ax_1 + by_1 \\ \vdots \\ ax_n + by_n \end{bmatrix}$$



$$\forall x, y \in \mathbb{R}^{m \times n}, \forall a, b \in \mathbb{R}, [a \cdot x + b \cdot y]_{ij} := a[x]_{ij} + b[y]_{ij}, 1 \leq i \leq m, 1 \leq j \leq n$$



$$\forall x, y \in \mathbb{R}^\infty, \forall a, b \in \mathbb{R}, a \cdot x + b \cdot y := (\dots, ax_{-1} + by_{-1}, ax_0 + by_0, ax_1 + by_1, \dots)$$

► $\forall f, g \in \{h : \mathbb{R} \rightarrow \mathbb{R}\}, \forall a, b \in \mathbb{R}, a \cdot f + b \cdot g, (a \cdot f + b \cdot g)(t) =$
 $a \cdot f(t) + b \cdot g(t), \forall t \in \mathbb{R}.$

Vector Spaces

- **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

Vector Spaces

- **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:
 - ▶ $(V, +)$ is an **Abelian group**:

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

- ▶ $\forall x, y \in V, x + y \in V$
- ▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$
- ▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta.$

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

▶ $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

▶ $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

▶ $\forall x, y \in V, x + y = y + x$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

▶ $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

▶ $\forall x, y \in V, x + y = y + x$.

▶ **Closure with respect to Scalar multiplication:** $\cdot: \mathbb{F} \times V \rightarrow V$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

▶ $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

▶ $\forall x, y \in V, x + y = y + x$.

▶ **Closure with respect to Scalar multiplication:** $\cdot \mathbb{F} \times V \rightarrow V$.

▶ **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

▶ $(V, +)$ is an **Abelian group**:

▶ $\forall x, y \in V, x + y \in V$

▶ $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

▶ $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

▶ $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

▶ $\forall x, y \in V, x + y = y + x$.

▶ **Closure with respect to Scalar multiplication:** $\cdot \mathbb{F} \times V \rightarrow V$.

▶ **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.

▶ **Distributivity:** $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$, and

$\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

► $(V, +)$ is an **Abelian group**:

► $\forall x, y \in V, x + y \in V$

► $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

► $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by $-x$.

► $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

► $\forall x, y \in V, x + y = y + x$.

► **Closure with respect to Scalar multiplication:** $\cdot \mathbb{F} \times V \rightarrow V$.

► **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.

► **Distributivity:** $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$, and

$\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$.

► **Compatibility of field and scalar multiplication:**

$\forall a, b \in \mathbb{F}, \forall u \in V, (a \times b) \cdot u = a \cdot (b \cdot u)$.

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:
 - ▶ $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:
 - ▶ $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.
 - ▶ $(\mathbb{F} - \{0\}, \times)$ is an **Abelian group**. The multiplicative identity will be denoted by 1.

Field

● **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:

► $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.

► $(\mathbb{F} - \{0\}, \times)$ is an **Abelian group**. The multiplicative identity will be denoted by 1.

► **Distributivity:**

$$\forall a, b, c \in \mathbb{F}, (a +_F b) \times c = a \times c +_F b \times c, a \times (b +_F c) = a \times b +_F a \times c$$

Examples of Fields

► $(\mathbb{Z}_2, +_2, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple $(V, +, \cdot)$ with field $(\mathbb{F}, +_F, \times)$ satisfies all vector space axioms, we say that $(V, +, \cdot)$ forms a vector space over \mathbb{F} .

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple $(V, +, \cdot)$ with field $(\mathbb{F}, +_F, \times)$ satisfies all vector space axioms, we say that $(V, +, \cdot)$ forms a vector space over \mathbb{F} .
- Any element of the vector space $(V, +, \cdot)$ will be referred to as a **vector**, and any element $a \in \mathbb{F}$ will be referred to as a **scalar**.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathcal{P}(\mathbb{R})$ is the set of all polynomials of one variable with real coefficients.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathcal{P}(\mathbb{R})$ is the set of all polynomials of one variable with real coefficients.
- $(\mathbb{L}_2(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathbb{L}_2(\mathbb{R})$ denotes the set of all square-integrable functions $f : \mathbb{R} \rightarrow \mathbb{R}$.